

# 家族性高脂血症のアポ蛋白BおよびEの遺伝子DNAに関する研究

著者	馬淵 宏
著者別表示	Mabuchi Hiroshi
雑誌名	昭和61(1986)年度 科学研究費補助金 一般研究(B) 研究成果報告書概要
巻	1985 1986
ページ	2p.
発行年	1989-03-29
URL	<a href="http://doi.org/10.24517/00068051">http://doi.org/10.24517/00068051</a>



# 1986 Fiscal Year Final Research Report Summary

## Studies on apolipoprotein B and E genes in familial hyperlipidemias

Research Project

### Project/Area Number

59480198

### Research Category

Grant-in-Aid for General Scientific Research (B)

### Allocation Type

Single-year Grants

### Research Field

内科学一般

### Research Institution

University of Kanazawa

### Principal Investigator

**MABUCHI Hiroshi** Assistant Professor, School of Medicine, Univ. Of Kanazawa, 医学部, 助教授 (00019960)

### Co-Investigator(Kenkyū-buntansha)

KOIZUMI Junji Assistant, Hospital, School of Medicine, Univ. of Kanazawa, 医学部付属病院, 助手 (20161846)

### Project Period (FY)

1984 - 1986

### Keywords

Apolipoprotein E gene / Apolipoprotein E deficiency / Familial hypercholesterolemia / LDL receptor / LDL receptor gene / FH-Tonami / 遺伝子診断

### Research Abstract

To probe a human adult liver cDNA library for apo-E cDNA clones, an oligonucleotide corresponding to apo E amino acids 218 to 222 was synthesized. One of 8 apoE cDNA clone (pAPOE7) was used as an apo E DNA probe for study of genomic DNA from a patient with apo E deficiency. Five micrograms of genomic DNA of apo E deficiency patient digested with Hind III or EcoRI, produced hybridization bands of approximately 1.9 and 12 Kb, respectively. The sizes of the hybridization bands were the same for normal controls and the proband's DNA.

Full length cDNA (pLDLR-3) and partial cDNA (pLDLR-2HHI) for human LDL receptor (LDLR) were used as LDLR DNA probes for studying genomic DNA from patients with familial hypercholesterolemia (FH) and normal subjects. Restriction fragment length polymorphism (RFLP) study of LDLR by PvuII showed allele A (16.5kb and 3.5kb) and allele B (14.0kb, 3.5kb and 2.5kb). The frequencies of the RFLP were determined in 28 FH patients and 13 normal subjects. The frequencies of allele A and B in FH and normal subjects were 0.08 and 0.13, and 0.72 and 0.08, respectively.

Two patients (K.Y. and S.O.) with FH showed an abnormal fragment pattern of LDLR gene. After digestions with various restriction enzymes, we considered

that these two patients had a common mutant LDLR allele that had a 6kb deletion, and the deletion encompassed an area near the 5' end of exon 15 up to the Hind III site in intron 15. The FH phenotype and the abnormal fragment of LDLR completely cosegregated in all the 11 members of the family of K.Y. In the family of S.O, S.O and her two daughters (M.T. and C.M.) with FH showed an abnormal fragment of LDLR gene. The serum cholesterol levels of M.T.'s baby was 78 mg/dl. On Southern blotting analysis, the baby revealed an abnormal 10.5kb fragment. On the other hand, C.M.'s baby was normocholesterolemic (51 mg/dl) and showed no abnormal LDLR gene and was diagnosed as normal subject. We designated the patients with this new mutant gene as "FH-Tonami". ▲ Less

## Research Products (11 results)

All Other

All Publications (11 results)

- [Publications] 武田三昭,梶波康二,伊藤英章,坂井健二,道下一朗,嘉門信雄,吉村陽,若杉隆伸,馬淵宏,竹田亮祐: 動脈硬化. 13. 843-847 (1985) ▼
- [Publications] 梶波康二,武田三昭,伊藤英章,酒井泰征,道下一朗,嘉門信雄,坂井健志,若細隆伸,馬淵宏,竹田亮祐: 動脈硬化. 13. 1353-1356 (1986) ▼
- [Publications] 馬淵宏,伊藤英章,武田三昭,梶波康二,若杉隆伸,竹田亮祐: Therapeutic Research. 5. 306-311 (1986) ▼
- [Publications] 梶波康二,馬淵宏: Medical Way. 3. 83-90 (1986) ▼
- [Publications] 梶波康二,武田三昭,伊藤英章,道下一朗,嘉門信雄,酒井泰征,若杉隆伸,馬淵宏,竹田亮祐: 動脈硬化. 14. 485-489 (1986) ▼
- [Publications] Mabuchi,H, et al.: Atherosclerosis. 61. 1-6 (1986) ▼
- [Publications] Mitsuaki Takeda, et al.: "Cloning of human apolipoprotein E cDNA and studies on genomic DNA of human apolipoprotein E gene." J.Japan Atherosclerosis Society. 13. 843-847 (1985) ▼
- [Publications] Kouji Kajinami et al.: "Studies on bovine LDL receptor cDNA." J.Japan Atherosclerosis Society. 13. 1353-1356 (1986) ▼
- [Publications] Hiroshi Mabuchi, et al.: "Metabolism of serum lipoproteins in patients with apolipoprotein E deficiency or apolipoprotein E-3 deficiency." Therapeutic Research. 5. 306-311 (1986) ▼
- [Publications] Kouji Kajinami, et al.: "Diagnosis of familial hypercholesterolemia by LDL receptor gene analysis." Medical Way. 3. 83-90 (1986) ▼
- [Publications] Kouji Kajinami, et al.: "Studies on LDL receptor gene." J.Japan Atherosclerosis Society. 14. 485-489 (1986) ▼

URL: [https://kaken.nii.ac.jp/report/KAKENHI-PROJECT-59480198/594801981986kenkyu\\_seika\\_hokoku\\_](https://kaken.nii.ac.jp/report/KAKENHI-PROJECT-59480198/594801981986kenkyu_seika_hokoku_)

Published: 1989-03-29