



This is a self-archived – parallel published version of an original article. This version may differ from the original in pagination and typographic details. When using please cite the original.

AUTHOR Teemu J. Niiranen, Karri Suvila, Susan Cheng,

TITLE Early-Onset Hypertension: Under-Recognized, Under-Treated, and Under-Estimated in Risk*,

YEAR 2020.

DOI <https://doi.org/10.1016/j.jacc.2020.04.059>.

VERSION Author's accepted manuscript

COPYRIGHT Lisence: [CC BY NC ND](https://creativecommons.org/licenses/by-nc-nd/4.0/)

CITATION Teemu J. Niiranen, Karri Suvila, Susan Cheng, Early-Onset Hypertension: Under-Recognized, Under-Treated, and Under-Estimated in Risk*, -Journal of the American College of Cardiology, Volume 75, Issue 23, 2020, Pages 2931-2933, ISSN 0735-1097, <https://doi.org/10.1016/j.jacc.2020.04.059> (<http://www.sciencedirect.com/science/article/pii/S0735109720351603>)

**Early-Onset Hypertension:
Under-Recognized, Under-Treated, and Under-Estimated in Risk**

Authors:

Teemu J. Niiranen, M.D.^{1,2}, Karri Suvila, M.D.¹, Susan Cheng, M.D., M.P.H.³

Affiliations:

1. Department of Medicine, Turku University Hospital and University of Turku, Turku, Finland
2. Department of Public Health Solutions, Finnish Institute for Health and Welfare, Turku, Finland
3. Smidt Heart Institute, Cedars-Sinai Medical Center, Los Angeles, California

Corresponding author:

Teemu J. Niiranen, Department of Medicine, University of Turku, Kiinamyyllynkatu 4–8, 20014 University of Turku, Finland; Email: Teemu.Niiranen@thl.fi; Telephone/Fax: +358 2 313 0000, Twitter handle: BP_Teemu

Conflicts of interest:

None.

High blood pressure remains a leading modifiable risk factor for cardiovascular disease (CVD) in both the young and the old (1,2). Surprisingly few studies have examined the age at which hypertension *begins* and its relation to later-life risk for CVD outcomes (3–8). The evidence to date would suggest that an early age of hypertension onset is associated with not only increased risk of CVD but also transmission to offspring, whereas hypertension onset in later life is more benign in nature (3–8). Importantly, prior analyses have involved predominantly retrospective case-control designs that may be subject to selection or recall bias and are also limited in their ability to establish a temporal relationship between the exposure and outcome. In this issue of the *Journal of the American College of Cardiology*, Wang et al. effectively advance our understanding by investigating the prognosis of early compared to late hypertension onset in relation to CVD risk in the setting of a large cohort study design (9).

In their analyses of hypertension onset, Wang et al. used data from the prospective Kailuan cohort, which includes 101510 Chinese community-dwelling participants who were followed biennially between 2006 and 2017. For this study, the authors used a sample of 19887 cases with new onset hypertension and an equal number of age- and sex-matched controls without hypertension. The follow-up period for the cases started at the time of hypertension onset, and the follow-up period for the controls started during the same year when incident cases were identified. The primary CVD outcomes included myocardial infarction, stroke, and all-cause mortality. The researchers used Cox proportional hazards regression models to determine the risk of outcomes in subgroups according to hypertension onset age, compared with those without hypertension. The authors observed that individuals with hypertension onset age ≤ 45 years carried the highest risk for CVD (hazard ratio [HR], 2.26; 95% confidence interval [CI], 1.19–4.30) and all-cause mortality (HR, 2.59; 95% CI, 1.32–5.07).

The risk estimates of both CVD and all-cause mortality consistently decreased with each incrementally later age decade of hypertension onset. Consequently, participants with hypertension onset age ≥ 65 years had the lowest hazards of CVD (HR 1.33; 95% CI, 1.04–1.69) and all-cause mortality (HR 1.29; 95% CI, 1.11–1.51).

This report by Wang. et al has several strengths, perhaps the most important being the first time-to-event cohort study to assess the relation between age of hypertension onset and adverse outcomes. Additional reliability is offered by the substantially large cohort size with the number of outcome events including 1672 cases of incident CVD and a total of 2008 deaths. Importantly, the researchers were also able to objectively determine the exact timing between age of hypertension onset and the incidence of events in the participants. There were also some limitations that were noted by the authors. The mean follow-up duration for participants studied was relatively short, at 6.5 years. Furthermore, the initial cohort consisted of employees of a single company and only 65.3% agreed to participate in the study (10), contributing to some level of potential selection bias. Additionally, estimated hazards in the multivariable models were adjusted for fasting blood glucose but not the use of antihyperglycemic medication, which could lead to an underestimation of diabetes related risks. Similarly, blood pressure levels during and prior to the outcome assessments were not considered. Notwithstanding these limitations, the strengths of this study overall outweigh the weaknesses.

Notably, the results from the study by Wang. et al are consistent with the findings of prior case-control studies (3–6). In the Framingham and the Coronary Artery Risk Development in Young Adults studies, early-onset hypertension was related to hypertension-mediated organ damage (5,6) and CVD death (4) to a greater extent than late-onset hypertension. Extending

from prior work, Wang et al. also demonstrate that early-onset compared to late-onset hypertension is significantly associated with greater risk of all-cause mortality. Only a few studies have examined hypertension onset age as a risk factor for diseases other than CVD. Two previous studies with contrasting results have suggested that age of hypertension onset might be related to risk of developing dementia in the elderly (11,12). Thus, the association between hypertension onset age and non-CVD mortality warrants further investigation.

The implications of this work are potentially far reaching. The authors correctly speculate that assessment of hypertension onset age could improve the overall CVD risk stratification, and that individuals with early-onset hypertension might benefit from more intensive hypertension treatment, including lifestyle interventions and antihypertensive therapy. In addition, prior research suggests that parental age of hypertension could also be used for estimating risk of hypertension in offspring (4). However, implementing assessment of hypertension onset age into clinical practice may have its challenges. Namely, studies have mainly defined time of onset using serial blood pressure readings which may not be available in the regular clinical setting. To address this potential barrier to pragmatic implementation of new knowledge, we recently showed that the age of hypertension onset can be reliably determined using self-report (6). Thus, use of self-report could improve the feasibility of incorporating an assessment of hypertension onset age into clinical practice, particularly in situations where prior medical records data are not available.

In summary, Wang et al. provide additional compelling evidence on the adverse effects of early-onset hypertension in a large scale prospective cohort study. Although the rates of hypertension awareness, treatment, and control are lower among younger compared to older adults, current guidelines do not recommend using age of hypertension onset as part of a

CVD risk assessment; contemporary recommendations also do not yet emphasize the importance of adequate therapy in young patients (2,13,14). Future guidelines iterations could include revisions that specify treatment approaches for patients with early-onset hypertension, given their considerably greater lifetime CVD risk when compared to patients with late-onset hypertension (3–6,9).

References

1. Zhou B, Danaei G, Stevens GA et al. Long-term and recent trends in hypertension awareness, treatment, and control in 12 high-income countries: an analysis of 123 nationally representative surveys. **Lancet** 2019; 394:639–651.
2. Zhang Y, Moran AE. Trends in the prevalence, awareness, treatment, and control of hypertension among young adults in the United States, 1999 to 2014. **Hypertension** 2017; 70:736–742.
3. Buck C, Baker P, Bass M, Donner A. The prognosis of hypertension according to age at onset. **Hypertension** 1987; 9:204–208.
4. Niiranen TJ, McCabe EL, Larson MG et al. Heritability and risks associated with early onset hypertension: Multigenerational, prospective analysis in the Framingham Heart Study. **BMJ** 2017; 357:j1949.
5. Suvila K, McCabe EL, Lehtonen A et al. Early Onset Hypertension Is Associated With Hypertensive End-Organ Damage Already by MidLife. **Hypertension** 2019; 74:305–312.
6. Suvila K, McCabe EL, Lima JAC et al. Self-Reported Age of Hypertension Onset and Hypertension-Mediated Organ Damage in Middle-Aged Individuals. **Am J Hypertens** 2020 [E-pub ahead of print]; doi: 10.1093/ajh/hpaa055.
7. Wang N-YY, Young JH, Meoni LA, Ford DE, Erlinger TP, Klag MJ. Blood pressure change and risk of hypertension associated with parental hypertension: The Johns Hopkins precursors study. **Arch Intern Med** 2008; 168:643–648.
8. Niiranen TJ, McCabe EL, Larson MG et al. Risk for hypertension crosses generations in the community: a multi-generational cohort study. **Eur Heart J** 2017; 38:2300–2308.
9. Wang C, Yuan Y, Zheng M et al. Association of Hypertension Onset Age with

- Cardiovascular Diseases and All-cause Mortality: A Prospective Cohort Study. **J Am Coll Cardiol** 2020; XX:XXX-XXX.
10. Zhang Q, Zhou Y, Gao X et al. Ideal cardiovascular health metrics and the risks of ischemic and intracerebral hemorrhagic stroke. **Stroke** 2013; 44:2451–2456.
 11. Corrada MM, Hayden KM, Paganini-Hill A et al. Age of onset of hypertension and risk of dementia in the oldest-old: The 90+ Study. **Alzheimer's Dement** 2017; 13:103–110.
 12. Gilsanz P, Mayeda ER, Glymour MM et al. Female sex, early-onset hypertension, and risk of dementia. **Neurology** 2017; 89:1886–1893.
 13. Williams B, Mancia G, Spiering W et al. 2018 ESC/ESH Guidelines for the management of arterial hypertension. **Eur Heart J** 2018; 39:3021–3104.
 14. Whelton PK, Carey RM, Aronow WS et al. 2017 ACC/AHA/AAPA/ABC/ACPM/AGS/APhA/ASH/ASPC/NMA/PCNA Guideline for the Prevention, Detection, Evaluation, and Management of High Blood Pressure in Adults. **J Am Coll Cardiol** 2018; 71:e127–e248.