

[Click here to view linked References](#)

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## **Voriconazole more likely than posaconazole increases plasma exposure to sublingual buprenorphine causing a risk of a clinically important interaction**

Mari Fihlman<sup>1</sup>, Tuija Hemmilä<sup>1</sup>, Nora M. Hagelberg<sup>1</sup>, Kristiina Kuusniemi<sup>1</sup>, Janne T. Backman<sup>2</sup>, Jouko Laitila<sup>2</sup>, Kari Laine<sup>3,4</sup>, Pertti J Neuvonen<sup>2</sup>, Klaus T. Olkkola<sup>5</sup> and Teijo I. Saari<sup>1</sup>

<sup>1</sup>Department of Anaesthesiology, Intensive Care, Emergency Care and Pain Medicine, University of Turku and Turku University Hospital, 20520 Turku, Finland

<sup>2</sup>Department of Clinical Pharmacology, University of Helsinki and HUSLAB, Helsinki University Central Hospital, 00014 Helsinki, Finland

<sup>3</sup>Department of Pharmacology, Drug Development and Therapeutics, University of Turku, 20100 Turku, Finland

<sup>4</sup>Medbase Ltd, FI-20100 Turku, Finland

<sup>5</sup>Department of Anaesthesiology, Intensive Care and Pain Medicine, University of Helsinki and Helsinki University Hospital, 00014 Helsinki, Finland

#**Corresponding author:** Teijo Saari, Department of Anaesthesiology and Intensive Care, University of Turku; P.O. Box 52; Kiinamylynkatu 4-8; FI-20520 Turku; Finland. Phone: +358-2-313 0441. Fax: +358-2-313 3960. E-mail: [teijo.saari@utu.fi](mailto:teijo.saari@utu.fi)

### **Acknowledgements**

We thank Mrs. Elina Kahra (medical laboratory technologist, Clinical Pharmacology, TYKSLAB, Hospital District of Southwest Finland, Turku, Finland) for her skillful technical assistance. This study was supported financially by Turku University Hospital research fund (EVO 13821), Turku, Finland

1  
2  
3  
4 **ABSTRACT**  
5  
6

7 **Purpose:** This study aimed to determine possible effects of voriconazole and posaconazole on the  
8 pharmacokinetics and pharmacological effects of sublingual buprenorphine.  
9

10  
11 **Methods:** We used a randomized, placebo-controlled crossover study design with 12 healthy male  
12 volunteers. Subjects were given a dose of 0.4 mg (0.6 mg during placebo phase) sublingual  
13 buprenorphine after a 5-day oral pretreatment with either (i) placebo, (ii) voriconazole 400 mg twice  
14 daily on the first day and 200 mg twice daily thereafter or (iii) posaconazole 400 mg twice daily.  
15 Plasma and urine concentrations of buprenorphine and its primary active metabolite  
16 norbuprenorphine were monitored over 18 h and pharmacological effects were measured.  
17  
18

19 **Results:** Compared to placebo, voriconazole increased the mean area under the plasma concentration-  
20 time curve ( $AUC_{0-\infty}$ ) of buprenorphine 1.80-fold (90% confidence interval 1.45-2.24;  $P<0.001$ ), its  
21 peak concentration ( $C_{max}$ ) 1.37-fold ( $P<0.013$ ) and half-life ( $t_{1/2}$ ) 1.37-fold ( $P<0.001$ ). Posaconazole  
22 increased the  $AUC_{0-\infty}$  of buprenorphine 1.25-fold ( $P<0.001$ ). Most of the plasma norbuprenorphine  
23 concentrations were below the limit of quantification (0.05 ng/ml). Voriconazole, unlike  
24 posaconazole, increased the urinary excretion of norbuprenorphine 1.58-fold (90% confidence  
25 interval 1.18-2.12;  $P<0.001$ ) but there was no quantifiable parent buprenorphine in urine. Plasma  
26 buprenorphine concentrations correlated with the pharmacological effects but the effects did not differ  
27 significantly between the phases.  
28  
29

30 **Conclusions:** Voriconazole, and to a minor extent posaconazole, increase plasma exposure to  
31 sublingual buprenorphine, probably via inhibition of cytochrome P450 3A and/or P-glycoprotein.  
32 Care should be exercised in the combined use of buprenorphine with triazole antimycotics,  
33 particularly with voriconazole, because their interaction can be of clinical importance.  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

**Keywords:** buprenorphine, azole antifungals, voriconazole, posaconazole, pharmacokinetics, drug-  
drug interaction.

## INTRODUCTION

Buprenorphine is a semisynthetic partial  $\mu$ -opioid receptor agonist. The analgesic efficacy of buprenorphine is 20–40 times higher than that of morphine [1]. Buprenorphine acts also as an antagonist at the  $\kappa$ -opioid receptor and as an agonist at the  $\delta$ -opioid receptor and opioid receptor-like receptor. In low doses, it is used in the treatment of moderate acute and chronic pain whereas in high doses, it is used in the management of opioid withdrawal symptoms and opioid dependence [2, 3].

Buprenorphine can be administered in various formulations and there is a growing interest for its use as an alternative to methadone that could increase access to treatment and be more acceptable to patients [4, 5].

After an oral and sublingual administration of buprenorphine, variability in its absorption and disposition increases its susceptibility to drug interactions. Buprenorphine undergoes extensive metabolism, particularly during its first-pass and has an oral bioavailability of 15% only [6, 7]. Bioavailability following sublingual administration of buprenorphine is higher, but the estimates of absolute bioavailability vary from 15 to 30% [8, 9]. Some transporters such as P-glycoprotein can play a role in the pharmacokinetics of buprenorphine and/or its metabolites [10–12]. Peak plasma concentrations of buprenorphine are reached within 1-3 hours after sublingual administration [13–15]. Practically no unconjugated parent drug is excreted into urine [14]. The main metabolic pathway, N-dealkylation of buprenorphine, is catalyzed mainly by cytochrome P450 (CYP) 3A4, and also by CYP3A5 and CYP2C8, yielding an active metabolite norbuprenorphine [16], which is further glucuronidated to norbuprenorphine-3-glucuronide [6, 17–20]. In addition to the N-dealkylation, smaller fractions of buprenorphine are converted to other hydroxylated metabolites and to buprenorphine-3-glucuronide. Some *in vitro* data suggest that CYP2C9 and CYP2C19 are involved in

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

smaller oxidative pathways of buprenorphine [19]. Overall, about 80–90% of buprenorphine-derived compounds are excreted by the biliary system and can be subject to enterohepatic circulation [21].

Voriconazole and posaconazole are triazole antifungal agents clinically used in the treatment of disseminated fungal infections. Voriconazole inhibits the activities of CYP3A, CYP2C19 and CYP2C9 enzymes [22] and it has greatly increased plasma concentrations of orally administered CYP3A4 substrates, e.g., oxycodone [23] and midazolam [24]. Posaconazole ~~also~~ is a moderately strong inhibitor of CYP3A4 and CYP2C18, but is not a clinically relevant inhibitor of CYP1A2, CYP2C9, CYP2D6 or CYP2E1 [25, 26]. Concomitant use of posaconazole has increased the plasma concentrations of many drugs that are predominantly metabolized by CYP3A4 [27]. In addition, posaconazole is more potent than voriconazole as inhibitor of P-glycoprotein and breast cancer resistance protein [28].

Some interaction studies have been conducted using high-dose buprenorphine and antiretrovirals [29–32]. However, the effect of azole antifungals on the pharmacokinetics of buprenorphine is largely unknown. In particular, there seem to be no previous studies on the possible interaction of voriconazole and posaconazole with sublingual buprenorphine although their concomitant use is likely to occur commonly. We hypothesized that inhibition of CYP-mediated metabolism of buprenorphine by voriconazole or posaconazole leads to significant changes in the plasma concentrations of buprenorphine.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## MATERIALS AND METHODS

### *Study participants*

On the basis of our previous drug-drug interaction studies [33, 34], we calculated that 10 subjects were needed to detect a 30% difference in the area under the concentration-time curve ( $AUC_{0-\infty}$ ) of buprenorphine at a power of 80% and a level of significance of  $P < 0.05$ . To be prepared for dropouts, 12 healthy non-smoking male volunteers were recruited in the study. All subjects completed the study. Age and body mass index ranges were 19 to 23 years and 19.8 to 24.8 kg/m<sup>2</sup>, respectively. The inclusion criteria for the volunteers were age 18 to 40 years, body weight within  $\pm 15\%$  of the ideal weight, and blood pressure within normal limits. The criteria for exclusion included a previous history of intolerance to the study drugs; concomitant drug therapy; past or present significant disease or drug allergy, alcoholism, drug abuse, or psychological or emotional problems; blood donation within the 4 weeks prior to the study; lifestyle habits that would compromise the conditions of the study or interpretation of the results. Informed consent was obtained from all individual participants included in the study and the eligible volunteers were ascertained to be healthy by clinical examination, medical history, routine laboratory tests, and an electrocardiogram. It was ensured that urine drug screening was negative. The risk of participants to develop opioid dependency was considered low as evaluated by the Finnish translation of the Abuse Questions [35].

### *Study outline and drug administration*

The ethics committee of the Hospital District of Southwest Finland and by the Finnish National Agency for Medicines approved the study protocol, and it was registered in the EudraCT clinical trials register under code number 2010-020953-14. A placebo-controlled, single-blinded, randomized, three-phase crossover study design with 5-day pretreatment periods was used, separated by intervals

1  
2  
3  
4 of 4 weeks. The volunteers ingested orally one of the following treatments in a randomized order: (i)  
5  
6 voriconazole (Vfend 200 mg tablet; Pfizer, Sandwich, Great Britain) 400 mg at 8.00 and 20.00 on day  
7  
8 1, 200 mg at 8.00 and 20.00 on days 2-4, and 200 mg at 10.00 and 20.00 on day 5 (voriconazole  
9  
10 phase), (ii) posaconazole (Noxafil 40 mg/ml oral suspension; Merck Sharp & Dohme, Hoddesdon,  
11  
12 Great Britain) 400 mg at 8.00 and 20.00 for 4 days and on day 5, posaconazole was given at 10.00  
13  
14 and 20.00 (posaconazole phase), or (iii) placebo at 8:00 am and 8:00 pm for 4 days, and on day 5  
15  
16 placebo was given at 10.00 and 20.00 (control phase). We assessed the adherence to the  
17  
18 premedication schedule of mobile telephone text messages. On day 5, after  
19  
20  
21 fasting overnight, the subjects ingested the premedication with 100 ml of water followed by a single  
22  
23  
24 dose of 0.4 mg (0.6 mg during placebo phase) of sublingual buprenorphine (Temgesic 0.2 mg tablets;  
25  
26  
27 RB Pharmaceuticals Limited, Slough, Great Britain) with 20 ml of water 1 h later. Standardized meals  
28  
29  
30 were served 4 and 8 h after the buprenorphine challenge.  
31  
32

33  
34 On the test days, a forearm vein was cannulated, and blood samples (10 ml) for  
35  
36 pharmacokinetic measurements were collected into ethylenediaminetetraacetic acid-containing tubes  
37  
38 immediately before and at the following time points after the administration of buprenorphine: 30 min  
39  
40 and 1, 1.5, 2, 3, 4, 5, 6, 8, 10, 12, and 18 h. Plasma was separated within 30 min and stored at  $-70^{\circ}\text{C}$   
41  
42 until analysis. Another venous cannula was inserted to the opposite forearm for the possible  
43  
44 administration of naloxone (Naloxon B. Braun 0,4 mg/ml, B. Braun Melsungen AG, Melsungen,  
45  
46 Germany). Urine was collected up to 18 h after buprenorphine administration. Urine aliquots were  
47  
48 stored at  $-70^{\circ}\text{C}$  until analysis. Determination of drug concentrations has been described in detail in  
49  
50 Supplement file 1.  
51  
52

53  
54  
55  
56 *Pharmacokinetic measurements*  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4 We observed the peak plasma concentration ( $C_{\max}$ ) and time to  $C_{\max}$  ( $t_{\max}$ ) directly  
5  
6 from the data. The area under the plasma concentration–time curve (AUC) values from 0 to 18 h  
7  
8 ( $AUC_{0-18}$ ), as well as from 0 h to infinity ( $AUC_{0-\infty}$ ), were calculated for buprenorphine applying  
9  
10 noncompartmental methods using the WinNonlin pharmacokinetics program (version 4.1; Pharsight,  
11  
12 Mountain View, CA). After visual identification of the terminal log-linear part of each concentration–  
13  
14 time curve-the elimination rate constant ( $k_e$ ) was determined using linear  
15  
16 regression analysis. The  $t_{1/2}$  was calculated using the equation  $t_{1/2} = \ln 2/k_e$ . The AUC values were  
17  
18 calculated using a combination of the linear and log-linear trapezoidal rules with extrapolation to  
19  
20 infinity, when appropriate, by division of the last measured concentration by  $k_e$ . We also calculated  
21  
22 the cumulative amount of free, unconjugated norbuprenorphine excreted into urine from 0 to 18h ( $A_e$ ).  
23  
24  
25  
26  
27  
28

### 29 *Measurement of pharmacological effects*

30  
31  
32 Adverse effects were evaluated using a questionnaire before, and 3 and 6 hours after buprenorphine  
33  
34 administration. Other subjective effects of buprenorphine were evaluated using 100-mm visual analog  
35  
36 scales for the following items: drowsy/alert, very poor performance/very good performance, no  
37  
38 drug effect/very strong drug effect, relaxed/anxious, no nausea/very strong nausea, calm/restless. The  
39  
40 Maddox wing test [36] and Cogan’s pupillometer [37] was used to measure the central coordination  
41  
42 of extraocular muscles and pupil size, respectively. A digit symbol substitution test was used to  
43  
44 estimate central processing of sensory information by recording the number of correct symbols  
45  
46 substituted in 3 min [38]. The analgesic effect was evaluated using the cold pressor test as described  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65 earlier [33].



1  
2  
3  
4 Pharmacological effects were evaluated

5  
6 prior to and at 1, 2, 3, 4, 5, 6, 8, 10 and 12 h after buprenorphine administration. For each effect  
7  
8 variable, area under the response–time (AUEC) curve was determined using the trapezoidal rule.  
9

### 10 11 *Statistical analysis*

12  
13  
14  
15 The data were evaluated for normality of distribution with probit plots and the Shapiro–Wilk’s  
16  
17 W-test. Log-transformed data were analysed but nontransformed

18  
19 results are reported. The  $AUC_{0-\infty}$  of buprenorphine was the primary outcome variable in the study,  
20  
21 and all other pharmacokinetic and all pharmacodynamic parameters were secondary variables.  
22

23  
24 Geometric mean ratios with 90% CIs were calculated for the pharmacokinetic variables. Lack of  
25  
26 interaction was assumed if the 90% CI of the geometric mean ratios for pharmacokinetic variables  
27  
28 were within the acceptance limit of 0.8–1.25. Pharmacokinetic variables were compared also using  
29  
30 repeated-measures analysis of variance with a posteriori testing was performed using the Tukey test.  
31

32  
33 Values for  $t_{max}$  were analyzed using the Wilcoxon signed-ranks test. Differences were regarded as  
34  
35 statistically significant when  $P < 0.05$ . The Pearson product moment correlation coefficient was used  
36  
37 to investigate the possible relationship between the ratios of the  $AUC_{0-\infty}$  of buprenorphine during the  
38  
39 treatment phases (voriconazole or posaconazole) to the  $AUC_{0-\infty}$  of buprenorphine during the control  
40  
41 phase, as well as to the  $C_{trough}$  of voriconazole or posaconazole before the administration of  
42  
43 buprenorphine. The associations of plasma buprenorphine concentrations with psychomotor and  
44  
45 analgesic effects were also calculated using the Pearson’s product moment correlation coefficient.  
46  
47  
48

49  
50 The results are expressed as mean values  $\pm$  SD. All data were analyzed using SYSTAT for Windows  
51  
52 (version 10.2; Systat Software, Richmond, CA) and R software (version 3.2.0) ~~was also applied for~~  
53  
54 ~~statistical analysis~~ [39].  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## RESULTS

### *Pharmacokinetics*

The mean plasma concentrations of buprenorphine during the placebo, voriconazole and posaconazole phases are shown in Figure 1 and the individual amounts of norbuprenorphine excreted into urine in Figure 2. The concentrations of norbuprenorphine were around or below the LLQ in the most of the plasma samples during all three study phases. The effects of voriconazole and posaconazole on the pharmacokinetics of buprenorphine are summarized in Table 1 and Figure 3.

Effect of voriconazole: Compared to the placebo phase, voriconazole increased the mean  $AUC_{0-\infty}$  of buprenorphine by 1.80-fold (90% confidence interval (CI) 1.45-2.24;  $P<0.001$ ) and its  $C_{max}$  by 1.37-fold (90% CI 1.05-1.79;  $P<0.001$ ). Voriconazole increased the mean  $t_{1/2}$  of buprenorphine from 7.9 h to 11.0 h ( $P<0.001$ ). The mean  $C_{trough}$  of voriconazole on day 5 was 1522 ng/ml (range 668-4162 ng/ml).

Effect of posaconazole: Compared to the placebo phase, posaconazole increased the mean  $AUC_{0-\infty}$  of buprenorphine by 1.25-fold (90% CI 1.03-1.52;  $p=0.016$ ) and its  $C_{max}$  by 1.20-fold (0.97-1.48) but the latter effect was not statistically significant ( $p=0.206$ ). Posaconazole had no effect on the  $t_{1/2}$  of buprenorphine. The mean  $C_{trough}$  of posaconazole on day 5 was 967 ng/ml (range 367-1758 ng/ml).

Effects on the renal excretion of buprenorphine and norbuprenorphine: Voriconazole increased the cumulative amount of norbuprenorphine excreted in urine by 1.6-fold (90% CI 1.18-2.12;  $P<0.001$ ), when compared to placebo (Table 1). The concentrations of parent buprenorphine in urine were in no study phase reliably quantifiable being much lower than those of norbuprenorphine.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

Comparison of voriconazole and posaconazole: Voriconazole increased the mean plasma  $AUC_{0-18}$  and  $AUC_{0-\infty}$  of buprenorphine significantly more than posaconazole ( $P<0.001$ ). Similarly the cumulative amount of norbuprenorphine excreted to urine was significantly increased during voriconazole, compared to posaconazole ( $P<0.001$ ).

### 3.2 Pharmacological effects

Almost every subject experienced some mild or moderate adverse effects (Supplementary Table S1). The most frequent adverse effect was sedation, followed by ataxia, dizziness and nausea. These effects were transient and did not require any treatment.

There was a linear correlation between plasma buprenorphine concentration and pharmacological drug effect ( $P<0.001$ ). However, there were no statistically significant differences in the pharmacological effects of buprenorphine between the three phases (Supplementary Figure S1).

## DISCUSSION

This study was designed to investigate the effect of voriconazole and posaconazole on the pharmacokinetics and pharmacodynamics of sublingual buprenorphine in healthy volunteers. The strong CYP3A and CYP2C inhibitor voriconazole markedly increased the  $C_{max}$  and  $AUC_{0-\infty}$  of buprenorphine and prolonged its  $t_{1/2}$ . In some subjects, the exposure to buprenorphine was increased more than two-fold by voriconazole. On the other hand, the effects of posaconazole on buprenorphine exposure were minor.

We measured also the urinary excretion of norbuprenorphine. As an N-dealkylated metabolite, it is less lipophilic than the parent buprenorphine and can be excreted into urine to some extent also in the unconjugated form. Surprisingly, voriconazole caused a 58% increase in the amount of norbuprenorphine excreted into urine, although it is reasonable to assume that voriconazole as a strong inhibitor of CYP3A4 decreases the (CYP3A4-mediated) N-dealkylation of buprenorphine. On the other hand, the strong and more selective CYP3A4 inhibitor posaconazole actually tended to decrease the excretion of norbuprenorphine into urine, consistent with inhibition of its CYP3A4-mediated formation. Furthermore, norbuprenorphine is a substrate of P-glycoprotein [40]. Different effects of posaconazole (potent inhibitor) and voriconazole (weak inhibitor) on P-glycoprotein may partially explain their different effects on buprenorphine plasma concentrations and on urinary excretion of norbuprenorphine. Unfortunately, we could not determinate the renal clearances, due to low free, unconjugated concentrations of plasma norbuprenorphine and urine buprenorphine.

It is noteworthy that 400 mg posaconazole twice daily has increased the  $AUC$ s of sensitive CYP3A4 substrates several fold; for example, the  $AUC$  of oral midazolam was increased about 5-fold, consistent with roughly 80% inhibition of the total CYP3A4-mediated clearance of midazolam [41]. Thus, as posaconazole increased the  $AUC$  of buprenorphine by 25% only, equal to 20% reduction in

1  
2  
3  
4 oral clearance, it seems that after sublingual buprenorphine dosing, only a small fraction (less than  
5  
6 30-40%) of buprenorphine dose is metabolized by CYP3A4, and that other CYP-enzymes and UGTs  
7  
8 play a larger role. Accordingly, strong inhibition of CYP2C9 and CYP2C19 mediated alternative  
9  
10 pathways, in addition to inhibition of CYP3A4-mediated N-demethylation, by voriconazole may  
11  
12 partially explain why voriconazole increased the urinary excretion of norbuprenorphine, and  
13  
14 increased buprenorphine plasma concentrations more than did posaconazole. In addition, the  
15  
16 increased urinary excretion of the metabolite may involve unidentified membrane transporter  
17  
18 mechanisms.  
19  
20  
21  
22  
23

24 Previous studies on drug-drug-interactions between CYP-inhibitors and buprenorphine are  
25  
26 scarce. The effects of HIV protease inhibitors have been characterized most thoroughly on the  
27  
28 pharmacokinetics of high-dose sublingual buprenorphine. Ritonavir increased the *AUC* of  
29  
30 buprenorphine significantly (57%), while the other protease inhibitors studied had no effect [30–32,  
31  
32 42]. Similarly, the non-nucleoside reverse-transcriptase inhibitors efavirenz and delavirdine increased  
33  
34 the *AUC* of buprenorphine [29]. Atazanavir alone or together with ritonavir increased buprenorphine  
35  
36 *AUC*<sub>0-18</sub> 93% and 67%, respectively, and lead to significant increase in sedative effect [31]. These  
37  
38 interactions with buprenorphine are of the same order as those observed in our present study with  
39  
40 voriconazole and might necessitate a decrease in the buprenorphine dose during concomitant  
41  
42 treatment.  
43  
44  
45  
46  
47  
48

49 Compared to oral intake, sublingual administration has increased the bioavailability of  
50  
51 buprenorphine considerably, from 15% to up to 30-60% [9, 43]. Our present results suggest that  
52  
53 strong CYP-inhibitors may further increase the bioavailability of sublingual buprenorphine as both  
54  
55 the *C*<sub>max</sub> and *AUC* of buprenorphine were clearly increased after voriconazole pretreatment. Most  
56  
57 probably, voriconazole caused increased exposure by inhibiting both the intestinal and hepatic  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4 CYP3A increasing both the  $C_{max}$  and  $t_{1/2}$  of buprenorphine. In addition, other mechanisms may also be  
5  
6 involved. P-glycoprotein is an efflux transporter in the intestinal wall, blood–brain barrier and many  
7  
8 other tissues [44]. Several opioids, including morphine [45], fentanyl [46] and alfentanil [47], are  
9  
10 substrates of P-glycoprotein, and especially posaconazole P-glycoprotein inhibitor [28, 48, 49]. Previ-  
11  
12 ous studies conducted in transfected cells  
13  
14  
15 and mice indicate that P-glycoprotein mediated drug efflux influences brain access and  
16  
17  
18 antinociceptive effects of norbuprenorphine [12].  
19  
20

21  
22 The majority of a buprenorphine dose is excreted as different metabolites into the bile and  
23  
24 circulates in enterohepatic system [21]. Voriconazole as a strong inhibitor of many CYP enzymes  
25  
26 alters the pharmacokinetics of several drugs metabolized by CYP3A, including oral hypnotics [24, 50,  
27  
28 51] and opioids [33, 52–54]. Voriconazole has decreased the clearance of alfentanil by 83% [49] and  
29  
30 that of fentanyl by 23% [53], and increased the  $AUC_{0-24}$  of R-methadone by 47% [54]. In the present  
31  
32 study, voriconazole increased the exposure to sublingual buprenorphine by 1.8-fold, i.e., less than has  
33  
34 been its effect (10x-fold) on oral midazolam [24]. Sublingual buprenorphine partially bypasses the  
35  
36 intestinal first-pass metabolism and therefore seems to be less prone than, e.g., oral midazolam or  
37  
38 oxycodone to the effects of drugs affecting intestinal and hepatic CYP3A4 during the first-pass  
39  
40 metabolism [33, 55, 56].  
41  
42  
43  
44

45  
46 The half-life of buprenorphine increased significantly after voriconazole, but not after  
47  
48 posaconazole pretreatment, which is likely to reflect a decrease in buprenorphine systemic clearance  
49  
50 by voriconazole, although an increase in the volume of distribution cannot be excluded without  
51  
52 intravenous buprenorphine dosing. The route of administration may affect the apparent half-life of  
53  
54 buprenorphine since the estimates appear to be longer after sublingual administration compared to  
55  
56 intravenous administration [57, 58]. Furthermore, longer half-lives than observed in the present study  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

have been described after high doses of buprenorphine [15, 43, 57–59]. The terminal elimination phase can continue for a longer time than was the sampling time in the present study, where the limit of buprenorphine quantification after small doses did not allow a longer sampling. Thus, our studies should be interpreted with care in this regard.

In our earlier studies, we have characterized previously unrecognized, drastic increases in the *AUC* of several substrate drugs, which have a limited therapeutic index, such as midazolam, triazolam, quinidine, tizanidine and oxycodone, when potent inhibitors of their metabolism have been administered concomitantly [34, 56, 60–63]. Based on these experiences, and for ethical and safety reasons in general, only a single clinically relevant, small dose of buprenorphine was employed, because we studied healthy volunteers and we wanted to minimize the risk of adverse events. This may be the reason that the values for the pharmacodynamic variables differed only slightly between phases. Our study was designed mainly to evaluate the pharmacokinetics of buprenorphine, and its power was not sufficient for more precise pharmacodynamic analysis. Nonetheless, we detected a significant linear correlation between plasma buprenorphine concentrations and pharmacological effects, which proves that our methodology was sensitive enough to measure the effects of buprenorphine.

In conclusion, our results show that even a short treatment with clinically used doses of voriconazole increases the exposure to sublingual buprenorphine and this interaction may have a considerable clinical relevance in individual patients. When the interindividual variation in the pharmacokinetics of buprenorphine is taken into account, some individuals given voriconazole may have a more than 100% increase in the exposure to buprenorphine. Thus, at least the interaction between voriconazole and sublingual buprenorphine can be of clinical significance. It has been previously shown that the strong CYP3A-inhibitor ketoconazole had no effect on the

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

pharmacokinetics of transdermal buprenorphine [64]. Our results warrant careful patient monitoring when sublingual buprenorphine is used with triazole antifungals, especially voriconazole. Posaconazole is less likely to cause a clinically significant interaction with sublingual buprenorphine.



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

**Description of authors' roles**

Mari Fihlman took care of the clinical phase of the study and data collection, participated in data analysis and statistical analysis and wrote the manuscript. Klaus Olkkola and Kari Laine designed the study, wrote the protocol, supervised and coordinated the clinical implementation of the study, and participated in data analysis. Tuija Hemmilä and Kristiina Kuusniemi participated the clinical phase and data collection. Janne T. Backman, Jouko Laitila and Pertti J Neuvonen performed the analytical assays. Teijo Saari designed the study, analysed the data, performed statistical analysis, and wrote the manuscript. All authors materially participated in the research and/or manuscript preparation. All authors have contributed to and approved the final manuscript.

**Conflict of interest Declaration**

The authors declare no conflict of interest.

## REFERENCES

1. Cowan A (1995) Update on the general pharmacology of buprenorphine. In: Cowan A, Lewis J (eds) *Buprenorphine Combat. drug Abus. with a unique opioid*. Wiley-Liss, New York, pp 189–211
2. Mattick RP, Ali R, White JM, et al (2003) Buprenorphine versus methadone maintenance therapy: a randomized double-blind trial with 405 opioid-dependent patients. *Addiction* 98:441–52.
3. Petitjean S, Stohler R, Déglon JJ, et al (2001) Double-blind randomized trial of buprenorphine and methadone in opiate dependence. *Drug Alcohol Depend* 62:97–104.
4. Yarborough BJH, Stumbo SP, McCarty D, et al (2016) Methadone, buprenorphine and preferences for opioid agonist treatment: A qualitative analysis. *Drug Alcohol Depend* 160:112–8. doi: 10.1016/j.drugalcdep.2015.12.031
5. Gryczynski J, Jaffe JH, Schwartz RP, et al (2013) Patient perspectives on choosing buprenorphine over methadone in an urban, equal-access system. *Am J Addict* 22:285–91.
6. Cone EJ, Gorodetzky CW, Yousefnejad D, et al (1984) The metabolism and excretion of buprenorphine in humans. *Drug Metab Dispos* 12:577–81.
7. Elkader A, Sproule B (2005) Buprenorphine: clinical pharmacokinetics in the treatment of opioid dependence. *Clin Pharmacokinet* 44:661–80.
8. Mendelson J, Upton RA, Everhart ET, et al (1997) Bioavailability of sublingual buprenorphine. *J Clin Pharmacol* 37:31–7.
9. Nath RP, Upton RA, Everhart ET, et al (1999) Buprenorphine pharmacokinetics: relative bioavailability of sublingual tablet and liquid formulations. *J Clin Pharmacol* 39:619–623.
10. Alhaddad H, Cisternino S, Declèves X, et al (2012) Respiratory toxicity of buprenorphine results from the blockage of P-glycoprotein-mediated efflux of norbuprenorphine at the blood-brain barrier in mice. *Crit Care Med* 40:3215–23.
11. Hassan HE, Myers AL, Coop A, Eddington ND (2009) Differential Involvement of P-Glycoprotein (ABCB1) in Permeability, Tissue Distribution, and Antinociceptive Activity of Methadone, Buprenorphine, and Diprenorphine: In Vitro and In Vivo Evaluation. *J Pharm Sci*

- 1  
2  
3  
4 98:4928–4940.  
5  
6  
7 12. Brown SM, Campbell SD, Crafford A, et al (2012) P-glycoprotein is a major determinant of  
8 norbuprenorphine brain exposure and antinociception. *J Pharmacol Exp Ther* 343:53–61. doi:  
9 10.1124/jpet.112.193433  
10  
11  
12 13. McAleer SD, Mills RJ, Polack T, et al (2003) Pharmacokinetics of high-dose buprenorphine  
13 following single administration of sublingual tablet formulations in opioid naïve healthy male  
14 volunteers under a naltrexone block. *Drug Alcohol Depend* 72:75–83.  
15  
16  
17 14. Ciraulo D a, Hitzemann RJ, Somoza E, et al (2006) Pharmacokinetics and pharmacodynamics  
18 of multiple sublingual buprenorphine tablets in dose-escalation trials. *J Clin Pharmacol*  
19 46:179–92.  
20  
21  
22 15. Bullingham RE, McQuay HJ, Dwyer D, et al (1981) Sublingual buprenorphine used  
23 postoperatively: clinical observations and preliminary pharmacokinetic analysis. *Br J Clin*  
24 *Pharmacol* 12:117–22.  
25  
26  
27 16. Brown SM, Holtzman M, Kim T, Kharasch ED (2011) Buprenorphine metabolites,  
28 buprenorphine-3-glucuronide and norbuprenorphine-3-glucuronide, are biologically active.  
29 *Anesthesiology* 115:1251–60.  
30  
31  
32 17. Iribarne C, Picart D, Dréano Y, et al (1997) Involvement of cytochrome P450 3A4 in N-  
33 dealkylation of buprenorphine in human liver microsomes. *Life Sci* 60:1953–64.  
34  
35  
36 18. Kobayashi K, Yamamoto T, Chiba K, et al (1998) Human buprenorphine N-dealkylation is  
37 catalyzed by cytochrome P450 3A4. *Drug Metab Dispos* 26:818–21.  
38  
39  
40 19. Moody DE, Slawson MH, Strain EC, et al (2002) A liquid chromatographic-electrospray  
41 ionization-tandem mass spectrometric method for determination of buprenorphine, its  
42 metabolite, norbuprenorphine, and a coformulant, naloxone, that is suitable for in vivo and in  
43 vitro metabolism studies.  
44  
45  
46 20. Picard N, Cresteil T, Djebli N, Marquet P (2005) In vitro metabolism study of buprenorphine:  
47 evidence for new metabolic pathways. *Drug Metab Dispos* 33:689–95. doi:  
48 10.1124/dmd.105.003681  
49  
50  
51 21. Brewster D, Humphrey MJ, McLeavy MA (1981) Biliary excretion, metabolism and  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

- 1  
2  
3  
4 enterohepatic circulation of buprenorphine. *Xenobiotica* 11:189–96.  
5  
6  
7 22. Niwa T, Shiraga T, Takagi A (2005) Effect of antifungal drugs on cytochrome P450 (CYP)  
8 2C9, CYP2C19, and CYP3A4 activities in human liver microsomes. *Biol Pharm Bull* 28:1805–  
9 8.  
10  
11  
12 23. Hagelberg NM, Nieminen TH, Saari TI, et al (2009) Voriconazole drastically increases  
13 exposure to oral oxycodone. *Eur J Clin Pharmacol* 65:263–271.  
14  
15  
16 24. Saari TI, Laine K, Leino K, et al (2006) Effect of voriconazole on the pharmacokinetics and  
17 pharmacodynamics of intravenous and oral midazolam. *Clin Pharmacol Ther* 79:362–370.  
18  
19  
20 25. Wexler D, Courtney R, Richards W, et al (2004) Effect of posaconazole on cytochrome P450  
21 enzymes: a randomized, open-label, two-way crossover study. *Eur J Pharm Sci* 21:645–53.  
22  
23  
24 26. Moody DE, Liu F, Fang WB (2015) Azole antifungal inhibition of buprenorphine, methadone  
25 and oxycodone in vitro metabolism. *J Anal Toxicol* 39:374–86.  
26  
27  
28 27. Li Y, Theuretzbacher U, Clancy CJ, et al (2010) Pharmacokinetic/pharmacodynamic profile of  
29 posaconazole. *Clin Pharmacokinet* 49:379–96.  
30  
31  
32 28. Lempers VJC, van den Heuvel JJMW, Russel FGM, et al (2016) Inhibitory Potential of  
33 Antifungal Drugs on ATP-Binding Cassette Transporters P-Glycoprotein, MRP1 to MRP5,  
34 BCRP, and BSEP. *Antimicrob Agents Chemother* 60:3372–9.  
35  
36  
37 29. McCance-Katz EF, Moody DE, Morse GD, et al (2006) Interactions between buprenorphine  
38 and antiretrovirals. I. The nonnucleoside reverse-transcriptase inhibitors efavirenz and  
39 delavirdine. *Clin Infect Dis* 43 Suppl 4:S224–34.  
40  
41  
42 30. McCance-Katz EF, Moody DE, Smith PF, et al (2006) Interactions between buprenorphine and  
43 antiretrovirals. II. The protease inhibitors nelfinavir, lopinavir/ritonavir, and ritonavir. *Clin*  
44 *Infect Dis* 43 Suppl 4:S235–46.  
45  
46  
47 31. McCance-Katz EF, Moody DE, Morse GD, et al (2007) Interaction between buprenorphine and  
48 atazanavir or atazanavir/ritonavir. *Drug Alcohol Depend* 91:269–78.  
49  
50  
51 32. Hulskotte EGJ, Bruce RD, Feng H-P, et al (2015) Pharmacokinetic interaction between HCV  
52 protease inhibitor boceprevir and methadone or buprenorphine in subjects on stable  
53 maintenance therapy. *Eur J Clin Pharmacol* 71:303–11.  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

33. Nieminen TH, Hagelberg NM, Saari TI, et al (2009) Rifampin greatly reduces the plasma concentrations of intravenous and oral oxycodone. *Anesthesiology* 110:1371–8.

34. Saari TI, Grönlund J, Hagelberg NM, et al (2010) Effects of itraconazole on the pharmacokinetics and pharmacodynamics of intravenously and orally administered oxycodone. *Eur J Clin Pharmacol* 66:387–97.

35. Michna E, Ross EL, Hynes WL, et al (2004) Predicting aberrant drug behavior in patients treated for chronic pain: importance of abuse history. *J Pain Symptom Manage* 28:250–8.

36. Hannington-Kiff JG (1970) Measurement of recovery from outpatient general anaesthesia with a simple ocular test. *Br Med J* 3:132–5.

37. Cogan D (1941) A simplified entoptic pupillometer. *Am J Ophthalmol* 24:1431–1433.

38. Stone BM (1984) Pencil and paper tests--sensitivity to psychotropic drugs. *Br J Clin Pharmacol* 18 Suppl 1:15S–20S.

39. R Foundation for Statistical Computing (2015) R: A language and environment for statistical computing.

40. Tournier N, Chevillard L, Megarbane B, et al (2010) Interaction of drugs of abuse and maintenance treatments with human P-glycoprotein (ABCB1) and breast cancer resistance protein (ABCG2). *Int J Neuropsychopharmacol* 13:905–15.

41. Krishna G, Moton A, Ma L, et al (2009) Effects of oral posaconazole on the pharmacokinetic properties of oral and intravenous midazolam: a phase I, randomized, open-label, crossover study in healthy volunteers. *Clin Ther* 31:286–98.

42. Bruce RD, Altice FL, Moody DE, et al (2010) Pharmacokinetic interactions between buprenorphine/naloxone and once-daily lopinavir/ritonavir. *J Acquir Immune Defic Syndr* 54:511–4.

43. Mendelson J, Upton RA, Everhart ET, et al (1997) Bioavailability of sublingual buprenorphine. *J Clin Pharmacol* 37:31–37.

44. DuBuske LM (2005) The role of P-glycoprotein and organic anion-transporting polypeptides in drug interactions. *Drug Saf* 28:789–801.

45. Kharasch ED, Hoffer C, Whittington D, Sheffels P (2003) Role of P-glycoprotein in the

- 1  
2  
3  
4 intestinal absorption and clinical effects of morphine. *Clin Pharmacol Ther* 74:543–54.  
5  
6  
7 46. Kharasch ED, Hoffer C, Altuntas TG, Whittington D (2004) Quinidine as a probe for the role  
8 of p-glycoprotein in the intestinal absorption and clinical effects of fentanyl. *J Clin Pharmacol*  
9 44:224–33.  
10  
11  
12 47. Kalvass JC, Olson ER, Pollack GM (2007) Pharmacokinetics and pharmacodynamics of  
13 alfentanil in P-glycoprotein-competent and P-glycoprotein-deficient mice: P-glycoprotein  
14 efflux alters alfentanil brain disposition and antinociception. *Drug Metab Dispos* 35:455–9.  
15  
16  
17 48. Sansone-Parsons A, Krishna G, Simon J, et al (2007) Effects of age, gender, and race/ethnicity  
18 on the pharmacokinetics of posaconazole in healthy volunteers. *Antimicrob Agents Chemother*  
19 51:495–502.  
20  
21  
22  
23  
24 49. Ashbee HR, Gilleece MH (2012) Has the era of individualised medicine arrived for  
25 antifungals? A review of antifungal pharmacogenomics. *Bone Marrow Transplant* 47:881–894.  
26  
27  
28 50. Saari TI, Laine K, Leino K, et al (2006) Effect of voriconazole on the pharmacokinetics and  
29 pharmacodynamics of zolpidem in healthy subjects. *Br J Clin Pharmacol* 63:116–120.  
30  
31  
32 51. Saari TI, Laine K, Bertilsson L, et al (2007) Voriconazole and fluconazole increase the  
33 exposure to oral diazepam. *Eur J Clin Pharmacol* 63:941–949.  
34  
35  
36 52. Saari TI, Laine K, Leino K, et al (2006) Voriconazole, but not terbinafine, markedly reduces  
37 alfentanil clearance and prolongs its half-life. *Clin Pharmacol Ther* 80:502–508.  
38  
39  
40 53. Saari TI, Laine K, Neuvonen M, et al (2008) Effect of voriconazole and fluconazole on the  
41 pharmacokinetics of intravenous fentanyl. *Eur J Clin Pharmacol* 64:25–30.  
42  
43  
44 54. Liu P, Foster G, Labadie R, et al (2007) Pharmacokinetic interaction between voriconazole and  
45 methadone at steady state in patients on methadone therapy. *Antimicrob Agents Chemother*  
46 51:110–8.  
47  
48  
49 55. Olkkola KT, Backman JT, Neuvonen PJ (1994) Midazolam should be avoided in patients  
50 receiving the systemic antimycotics ketoconazole or itraconazole. *Clin Pharmacol Ther*  
51 55:481–5.  
52  
53  
54 56. Olkkola KT, Ahonen J, Neuvonen PJ (1996) The effects of the systemic antimycotics,  
55 itraconazole and fluconazole, on the pharmacokinetics and pharmacodynamics of intravenous  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4 and oral midazolam. *Anesth Analg* 82:511–6.  
5

- 6  
7 57. Bullingham RE, McQuay HJ, Moore A, Bennett MR (1980) Buprenorphine kinetics. *Clin*  
8 *Pharmacol Ther* 28:667–72.  
9  
10  
11 58. Kuhlman JJ, Lalani S, Magluilo J, et al (1996) Human pharmacokinetics of intravenous,  
12 sublingual, and buccal buprenorphine. *J Anal Toxicol* 20:369–78.  
13  
14  
15 59. Kuhlman JJ, Levine B, Johnson RE, et al (1998) Relationship of plasma buprenorphine and  
16 norbuprenorphine to withdrawal symptoms during dose induction, maintenance and withdrawal  
17 from sublingual buprenorphine. *Addiction* 93:549–59.  
18  
19  
20  
21 60. Kaukonen KM, Olkkola KT, Neuvonen PJ (1997) Itraconazole increases plasma concentrations  
22 of quinidine. *Clin Pharmacol Ther* 62:510–7.  
23  
24  
25 61. Varhe A, Olkkola KT, Neuvonen PJ (1994) Oral triazolam is potentially hazardous to patients  
26 receiving systemic antimycotics ketoconazole or itraconazole. *Clin Pharmacol Ther* 56:601–7.  
27  
28  
29 62. Backman JT, Kivistö KT, Olkkola KT, Neuvonen PJ (1998) The area under the plasma  
30 concentration-time curve for oral midazolam is 400-fold larger during treatment with  
31 itraconazole than with rifampicin. *Eur J Clin Pharmacol* 54:53–8.  
32  
33  
34  
35 63. Granfors MT, Backman JT, Neuvonen M, et al (2004) Fluvoxamine drastically increases  
36 concentrations and effects of tizanidine: a potentially hazardous interaction. *Clin Pharmacol*  
37 *Ther* 75:331–41.  
38  
39  
40  
41 64. Kapil RP, Cipriano A, Michels GH, et al (2012) Effect of ketoconazole on the pharmacokinetic  
42 profile of buprenorphine following administration of a once-weekly buprenorphine transdermal  
43 system. *Clin Drug Investig* 32:583–92.  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

**FIGURE LEGENDS**

**Figure 1.** Mean plasma (SD) concentrations of buprenorphine in 12 healthy volunteers after 0.4 mg (0.6 mg in placebo phase) sublingual buprenorphine on the fifth day of pretreatment with placebo (*open circles*), voriconazole 400 mg twice on the first day, thereafter 200 mg twice daily (*filled triangles*) or posaconazole 400 mg twice daily (*filled circles*) for 5 days. Right panel shows the same concentrations in semilogarithmic scale. Values are normalized for a sublingual dose of 1.0 mg.

**Figure 2.** The individual amounts of urinary norbuprenorphine excreted during 18 hours after 0.4 mg (0.6 mg in placebo phase) sublingual buprenorphine on the fifth day of pretreatment with placebo, voriconazole (400 mg twice on the first day, thereafter 200 mg twice daily) or posaconazole (400 mg twice daily) for 5 days in 12 healthy volunteers. Values are normalized for a sublingual dose of 1.0 mg.



**Figure 3.** Individual pharmacokinetic parameters after sublingual buprenorphine. Values for maximum concentration ( $C_{max}$ ), area under plasma concentration–time curve extrapolated to infinity ( $AUC_{0-\infty}$ ) and elimination half-life ( $t_{1/2}$ ) in 12 healthy volunteers after 0.4 mg (0.6 mg in placebo phase) sublingual buprenorphine on the fifth day of pretreatment with placebo, voriconazole (400 mg twice on the first day, thereafter 200 mg twice daily) or posaconazole (400 mg twice daily) for 5 days. Values are normalized for a sublingual dose of 1.0 mg.



**Table 1.** Pharmacokinetic parameters of buprenorphine after sublingual administration of 0.6 mg (control phase) or 0.4 mg (posaconazole and voriconazole phases) of buprenorphine on the fifth day of pre-treatment with oral posaconazole (400 mg twice daily for 5 days), voriconazole 400 mg twice daily on the first day and 200 mg for 4 days) or placebo to 12 healthy volunteers.

Parameter	Placebo	Posaconazole	Posaconazole / Placebo		Voriconazole	Voriconazole / Placebo		Voriconazole / Posaconazole	
			GMR (90% CI)	<i>p</i> -value		GMR (90% CI)	<i>p</i> -value	GMR (90% CI)	<i>p</i> -value
<i>Buprenorphine</i>									
$C_{\max}$ (ng/ml)	0.70 ± 0.20	0.84 ± 0.27	1.20 (0.95, 1.51)	0.159	0.96 ± 0.27	1.37 (1.05, 1.79)	0.013	1.15 (0.93, 1.42)	0.333
$t_{\max}$ (h)	2.0 (1.0-3.0)	2.0 (1.5-3.0)	-	0.667	2.0 (1.5-3.0)	-	0.40	-	0.40
AUC <sub>0-18</sub> (ng·h/ml)	4.1 ± 1.2	5.0 ± 1.3	1.22 (1.05, 1.41)	0.026	6.5 ± 1.7	1.58 (1.28, 1.94)	<0.001	1.30 (1.10, 1.53)	0.003
AUC <sub>0-∞</sub> (ng·h/ml)	4.9 ± 1.3	6.1 ± 1.6	1.25 (1.20, 1.43)	0.008	8.9 ± 2.5	1.80 (1.45, 2.24)	<0.001	1.44 (1.23, 1.68)	<0.001
Cl/F (l/min)	2.9 ± 0.9	2.3 ± 0.7	0.80 (0.70, 0.91)	0.007	1.7 ± 0.6	0.56 (0.48, 0.65)	<0.001	0.70 (0.60, 0.81)	<0.001
V <sub>z</sub> /F (l)	32 ± 12	27 ± 11	0.82 (0.63, 1.10)	0.430	25 ± 11	0.76 (0.52, 1.12)	0.22	0.93 (0.64, 1.36)	0.892
$t_{1/2}$ (h)	7.9 ± 2.8	8.1 ± 2.9	1.02 (0.77, 1.35)	0.991	11.0 ± 3.9	1.37 (0.88, 2.12)	<0.001	1.34 (0.90, 2.00)	0.230
<i>Norbuprenorphine</i>									
A <sub>e</sub> (μg)	5.7 ± 2.2	4.8 ± 2.2	0.81 (0.58, 1.14)	0.239	9.0 ± 3.1	1.58 (1.18, 2.12)	<0.001	1.95 (1.45, 2.61)	<0.001

GMR, geometric mean ratio; CI, confidence interval;  $C_{\max}$ , peak plasma concentration;  $t_{\max}$ , concentration peak time; AUC<sub>0-18</sub> and AUC<sub>0-∞</sub>, area under curve from 0 to 18 h and from 0 to infinity, respectively;  $t_{1/2}$ , elimination half-life; A<sub>e</sub>, amount excreted into urine within 18 h. Values are normalized for a sublingual dose of 1.0 mg. Data are shown as mean ± standard deviation (SD) and as the geometric mean ratios with the 90% confidence interval (CI) in parenthesis - except for  $t_{\max}$ , which is given as median and range.



- Buprenorphine + placebo
- Buprenorphine + posaconazole
- ▲ Buprenorphine + voriconazole

