

Self- reported fitness and objectively measured physical activity profile among older adults: a twin study

Katja Waller, PhD ¹, Henri Vähä-Ypyä, PhD ², Noora Lindgren, MSc ³, Jaakko Kaprio, MD, PhD ^{4,5}, Harri Sievänen, ScD ², Urho M. Kujala, MD, PhD ¹

¹ Faculty of Sport and Health Sciences, University of Jyväskylä, Finland

² The UKK Institute for Health Promotion Research, Tampere, Finland

³ Turku PET Centre, University of Turku, Turku

⁴ Department of Public Health, University of Helsinki, Helsinki

⁵ Institute for Molecular Medicine Finland FIMM, University of Helsinki, Helsinki

Corresponding Author: Katja Waller, PhD, Faculty of Sport and Health Sciences, P.O. Box 35

(LL), FIN-40014 University of Jyväskylä, Finland. Tel. +358 40 8053601, E-mail:

katja.waller@jyu.fi

Abstract

BACKGROUND: Maintaining good fitness and good level of physical activity are important factors for maintaining physical independence later in life. The aim was to investigate the relationship between self-reported fitness and objectively measured physical activity (PA) and sedentary behavior (SB) in the elderly.

METHODS: Same-sex twin pairs born 1940-1944 in Finland were invited to the study. Altogether 787 individuals (mean age 72.9 years), of whom 404 were female, used a hip-worn triaxial accelerometer for at least four days and answered a question on perceived fitness. First, individual differences were studied between four fitness categories. Secondly, pairwise differences were examined among twin pairs discordant for fitness.

RESULTS: Self-reported fitness explained moderately the variation in objectively measured PA parameters: R² for daily steps 26%, for daily mean MET 31%, for daily moderate-to-vigorous activity (MVPA) time 31%, and lower for SB time 14 % (all $p < 0.001$). Better self-reported fitness was associated with more steps taken on average (8558 daily steps (very good fitness) vs. 2797 steps (poor fitness), $p < 0.001$) and with a higher amount of MVPA (61 min vs. 12 min $p < 0.001$, respectively) in the adjusted multivariable model. Among 156 twin pairs discordant for self-reported fitness, co-twins with better fitness took more steps, did more MVPA and had less SB (all, $p < 0.05$) compared to their less fit co-twins, however difference was smaller among MZ than DZ pairs.

CONCLUSION: One simple question on self-reported fitness is associated with daily activity profile among community-dwelling older people. However, genetic factors modulate this association to some extent.

Key words: exercise, accelerometer, fitness, self-report, elderly

Introduction

Among the elderly, high physical activity (PA) and fitness are indicators of good physical health and functioning, both accounting for reduced probability of developing several chronic diseases and all-cause mortality (1-3). Both fitness and PA are important in maintaining Activities of Daily Living (ADL), which includes a spectrum activities ranging from self-care and basic mobility to physically demanding household chores (4,5). Physical fitness is also an important factor for maintaining physical independence later in life even irrespective of obesity status (6). Decreased aerobic fitness, assessed as self-reported feeling of fatigue and exhaustion, and low PA are both frailty-related criteria (7,8). Therefore, from the viewpoint of healthy aging, it is important that individuals maintain sufficient level of PA and optimal physical fitness.

As far as we are aware, no study has examined the association between self-reported fitness and objectively measured PA and sedentary behavior (SB) among older adults. However, mixed results have been obtained when objectively measured PA were associated with measured cardiorespiratory fitness (i.e., maximal oxygen uptake- VO_2). Among middle-aged people, correlation between accelerometer-measured vigorous activity and measured VO_2 was moderate ($r=0.50$), while no differences in sedentary time were found between fit and unfit persons (9). Among elderly people (mean age 75.5 years), a submaximal VO_2 field test exhibited moderately high correlations (0.61-0.69) with different accelerometer-measured PA parameters, while SB exhibited a moderately high negative correlation (-0.69) with VO_2 (7). Some studies have investigated the relationship between objectively measured PA and physical function or other measures of fitness among older adults, for example higher MVPA is associated with better strength and faster timed chair stand speed (10) and lower MVPA time with poorer functional fitness (11).

Objective fitness tests are time-consuming and difficult to conduct in primary health care, and therefore a simpler evaluation of physical fitness among older adults is desirable. Self-reported fitness might suffice to assess fitness in normal daily use. Associations between self-reported fitness and measured fitness among adults have been moderate to very strong correlations when a 22-item survey on self-reported fitness (multidimensional fitness) was used (12) and moderate associations when only one question was used (13,14). However, among older adults fewer studies have been conducted and multidimensional self-reported fitness question batteries have displayed only weak-to-moderate correlations with objectively-measured fitness (4,15). van Heuvelen et al. (1997) (13) suggested that older adults might not be able to evaluate their multidimensional fitness and this tended to reflect their general physical functioning, and therefore a single question about self-reported fitness question among older adults might capture the same information.

However, it is unknown whether perceived fitness, assessed as self-reported fitness, is associated with objectively measured PA or SB. Older adults who perceive themselves unfit might be less active and spend more time in sedentary activities than individuals considering themselves fit, regardless of their actual fitness. Therefore, the study assesses relationship between a single question about self-reported fitness and objectively measured PA and SB. Specifically, this study examines whether PA and SB differ between the self-reported fitness categories and whether the possible differences persist when genetic factors are taken into account within a twin-study design. It is important that the elderly remain fit and perform at least some physical activity so that they can enjoy independent living and avoid or at least postpone the development of frailty. Therefore, feasible and broadly accessible tools to identify those individuals at imminent risk of developing frailty are needed.

Methods

Study participants

The study is based on the older Finnish Twin cohort and is part of the MOBILETWIN study (16). In this study conducted during 2014-2016, a sub-cohort of twin pairs born between 1940 and 1944 were invited to participate in a telephone interview, use accelerometer and respond to an activity related questionnaire. Altogether 787 twin individuals (including 283 complete pairs; 119 monozygotic (MZ) pairs, 148 same-sex dizygotic (DZ) pairs and 16 same-sex pairs with uncertain zygosity) answered the required questions about self-reported fitness and wore an accelerometer for pre-determined time. The final analysis included 383 males and 404 females, whose mean age was 72.9 years (range 71.1-75.0).

Objectively measured physical activity and sedentary behavior

Physical activity was measured with a hip-worn, light triaxial accelerometer (Hookie AM20, Traxmeet Ltd, Espoo). The device and the instructions on how it should be worn were mailed to the participants who provided a written consent. The participants were instructed to use the accelerometer during waking hours for 7 consecutive days. Thereafter the participants mailed the accelerometer back to UKK Institute in a prepaid envelope for data analysis. The time criterion for adequate accelerometer data collection was at least four days with a minimum of 10 hours per day. The daily non-wear time during waking hours was defined as a sum of at least 30 minutes of consecutive zero acceleration.

The raw acceleration data were analysed by recently developed and validated MAD-APE algorithms (17-19). The acronym MAD stands for the mean amplitude deviation of the raw acceleration signal and the MAD value of the resultant acceleration has shown a strong correlation ($r > 0.9$) with directly measured incident VO_2 while walking or running on a track (19). The acronym APE, in turn, stands for the angle for posture estimation and it together with incident

MAD values provides about 90% accuracy in detecting the body posture not only in laboratory conditions but also in free-living conditions (17). These novel algorithms were used to differentiate between lying, sitting, standing and different PA intensities. For the analysis, MAD and APE values were determined for each 6 second epoch and a 1 minute moving average of the MAD values was calculated to estimate incident energy consumption expressed in metabolic equivalents (MET). The MET values were categorised as follows: 1.5-3 MET for light activities, 3-6 MET for moderate activities and over 6 MET for vigorous activities. Sedentary activities were defined as MET less than 1.5 while lying, sitting or standing. Exact explanations for all objectively measured variables can be seen in Supplementary Table 1.

In order to acquire a comprehensive understanding on the relationship between self-reported fitness and physical activity profile, a variety of PA and SB variables were determined from the processed acceleration data. Main PA and SB variables in this study were: mean daily times spent in lying, sitting or standing or in light, moderate or vigorous activities. Other variables were the number of daily steps, running steps, sit-to-stand transitions, activity bouts comprising at least 10 minutes consecutive activity, the mean daily MET for all recorded days (daily MET) and the most intense 10-minute period (Peak-10 min MET based on 10 minute moving exponential average of MAD data) during the monitoring week. In some analyses, lying and sitting were combined to represent sedentary behavior (SB) and moderate and vigorous physical activities were combined to designate activity at a moderate-to-vigorous activity level (MVPA).

Physical function questionnaire and covariates

Together with the accelerometer, the participants received a questionnaire about their physical function, self-reported health and fitness. In this study, self-reported fitness was assessed by their response to the question: “Is your current physical fitness in your opinion, 1) Very good, 2) Fairly good, 3) Satisfactory 4) Fairly poor, 5) Very poor. Due to the low number of responses in categories four and five, these two were combined for the analyses and called poor.

Age, sex, body mass index (BMI, kg/m²) and diseases restricting mobility were used as covariates in the study. BMI was calculated from reported height and weight and information on diseases was self-reported and asked as follows: Do you have a physician diagnosed disease that affect your ability to move or exercise? Participants answered either yes or no. The main diseases restricting mobility were musculoskeletal, cardiovascular, neurological and pulmonary diseases.

Statistical analyses

Data analyses were conducted using StataIC version 14. To determine R-squared (R²) for the associations linear regression was used. In the analyses, the twins were treated as individuals but since the observations from twin pairs may be correlated, robust estimators of variance (the cluster option in Stata) were used. All basic analyses calculating R² were adjusted for age and sex. In order to estimate R² only for self-reported fitness, it was entered into the model after the basic model with age and sex, and then the change in R² (ΔR^2) was determined. Subsequently, the Sidak multiple-comparison test was applied to identify differences between the four fitness categories. The same procedure was conducted when the regression analysis was additionally adjusted for BMI and self-reported disease. A multivariable model for R² including all covariates was also devised. Square root-transformation of lying, moderate activity, MVPA and 10 min bouts and logarithm-transformation of running steps, vigorous activity and Peak-10 min MET were done to make these variables more normally distributed in linear regression analyses. Means and standard deviations (SD) were calculated with bootstrapping option due to these not normally distributed variables.

Pairwise differences were studied among 156 twin pairs (90 DZ, 57 MZ and 9 uncertain zygosity) discordant for self-reported fitness, one co-twin having better self-reported fitness vs. their co-twin (for example 1 vs 2, 2 vs 3, 1 vs 3 etc.). Pairs with uncertain zygosity were included in the pairwise analysis of all discordant pairs, but they were not included when either MZ or DZ pairs were analysed separately. Wilcoxon Signed Rank test was used to study pairwise differences in PA and

SB variables. In another set of twin analyses, pairwise differences were also studied among 40 twin pairs (27 DZ, 11 MZ pairs, 2 uncertain zygosity) discordant for daily steps. Discordance was defined as one twin belonging to the lowest step count tertile (in the whole sample) and their co-twin to the highest step count tertile. The exact symmetry test was used to assess differences between self-reported fitness categories.

Results

In all of the 787 individuals for whom we had both the fitness and acceleration data, the accelerometers were worn for an average of 14.0 h/day (range 11.0-20.1 h/day, SD 1.2). An average 6 387 (range 380 – 18 362, SD 3 117) daily steps were taken and 31.8 (range 0 – 91.4, SD 15.1) sit-to-stand transitions were done per day. Participant characteristics broken down by self-rated fitness categories are shown in Table 1. Individuals who reported being the most fit had the lowest BMI, while the highest BMI values were present in those individuals with satisfactory fitness. Individuals with poor fitness reported having proportionally the highest number of diseases restricting their mobility. The recommended 10 000 daily steps for healthy adults to be classified as active (20) was achieved by 110 (14%) participants, the vast majority (97/110) belonged to one of the top two self-reported fitness categories (very good or fairly good).

Self-reported fitness explained moderately variation in different PA and SB variables (Table 2). Highest additional R^2 after age- and sex adjustment was 20.8% for daily MET and 20.4% for MVPA. Similar moderate explanation values were observed for daily steps and moderate activity alone, whereas lower additional R^2 values were seen for sedentary behavior variables. The multivariable model including self-reported fitness, age, sex, BMI and disease explained variation most (>30%) for Peak-10 min MET, daily MET, MVPA and vigorous activity while while less than 10% of the variance in SB variables was accounted for by the same multivariable model variables (Table 2).

The differences between self-reported fitness categories and objectively measured physical activity varied according to the activity variable. Figure 1 shows the overall daily activity patterns broken down by the self-reported fitness categories. Most of the SB variables revealed only small differences between the fitness categories (Table 3). In the fully adjusted model (age, sex, BMI and disease restricting mobility) no group level differences were seen in lying down and only one between-group difference in sitting time (Very good vs. Satisfactory) and in standing time (Fairly good vs. Poor).

Differences were evident between the self-reported fitness categories and variables describing PA, as participants with better self-reported fitness were clearly more active. For example, participants with very good fitness took an average of 8558 steps per day, while participants with fairly good (6724), satisfactory (5196) and poor (2797) fitness took substantially fewer steps with significant differences ($p < 0.001$ between all categories). Significant differences between all self-reported fitness categories were also seen in moderate activity, MVPA, daily MET, Peak-10 min MET and number of activity bouts longer than 10 minutes, all of these differences remained statistically significant even after adjustments (Table 3). Participants reporting very good fitness undertook some vigorous activity (3.3 min/day) while all other fitness groups did practically none (< 1 min/day, $p < 0.001$). Those with poor fitness had significantly less sit-to-stand transitions and undertook less light activity compared to the other fitness groups even after adjustments. Figure 2 illustrates the differences in PA and SB between self-reported fitness groups separately for men and women.

Pairwise analysis of 156 twin pairs discordant for self-reported fitness showed statistically significant differences between all variables (steps, Peak-10 min MET, SB, light activity and MVPA, $p < 0.05$) but standing ($p = 0.187$), table 4. Similar significant differences were seen among DZ pairs while no significant differences were observed among MZ pairs. Similar results were seen when limited number of pairs (9 MZ pairs, 25 DZ pairs) were analyzed with larger

discordance (difference ≥ 2 fitness categories). According to another set of discordant pairwise analyses including 40 pairs discordant for daily step, self-reported fitness was significantly better among co-twins who belonged to highest step count tertile ($p < 0.001$). No significant differences were seen among 11 discordant MZ pairs ($p = 0.63$).

Discussion

This study showed that self-reported fitness is associated with objectively measured physical activity in adults in their early 70s. Self-reported fitness explained at its best about 20% of the variation in different objectively measured PA variables. Statistically significant differences between all self-reported fitness categories were seen in PA variables, whereas some sedentary behaviors (lying and standing) did not differ between the different categories of self-reported fitness. The results suggest that an easily assessed self-reported fitness might be a relatively good indicator of habitual physical activity level in older adults whereas sedentary behavior does not vary so much by fitness level. Furthermore the perceived fitness might be an important factor in determining whether older adults decide to participate in physical activities.

Several objectively measured activity variables (Daily MET, Peak-10 min MET, steps, number of 10 min bouts and moderate activity) displayed the greatest differences according to self-reported fitness, as significant differences were seen between all fitness categories even after adjustments. Additionally, participants reporting very good fitness undertook some vigorous activity while other groups did hardly any at all. All of these variables describe activities with high intensity and therefore they can be expected to be closely related to fitness. As significant differences were observed between all fitness categories, even between those belonging to very good and fairly good categories, the single question about fitness seems to indicate current actual fitness and physical activity level among older adults. Egerton et al. (2016) (21) found that self-reported fatigue was also associated with objectively measured physical activity. However, in that study cardiorespiratory fitness attenuated the association and fitness was found to be the key predictor for physical activity among community-living 70-77-year old people.

Co-twins with better self-reported fitness were more active and had less sedentary behavior compared to their co-twins with lower fitness level; however, this was not seen among MZ pairs

even when the discordance was larger. Similar results were seen in the another set of twin analysis, as self-reported fitness was significantly better among co-twins who belonged to the highest step count tertile compared to their co-twin in the lowest tertile, again no difference was seen in this analysis of MZ pairs. This suggests that genetic factors explain some of the association between self-reported fitness and objectively measured physical activity, though the finding was in the same direction, but sample size was small in this second set of twin analysis. Also, it seems that twins from MZ pairs were less different from each other in these objectively measured PA and SB variables compared to twins in DZ pairs, as mean intrapair difference in most of the variables was significantly smaller among MZ pairs. This is shown by our additional discordance analyses (Supplementary Table 2). It is known that the heritability of physical activity (22) and cardiorespiratory fitness is at least moderate (23,24). In our recent paper (16), we presented some genetic modelling on these objectively measured physical activity variables, and showed that the broad sense heritability is 0.41 (95% CI 0.24-0.56) for sedentary behavior, 0.60 (0.49-0.70) for MVPA, 0.48 (0.35-0.60) for steps and 0.60 (0.45-0.73) for Peak-10min MET, indicating that the fitness related variables had more genetic influence than the sedentary and lower activity variables (16).

In the present study, the only differences observed in vigorous activity between the fitness categories were between very good fitness and the other categories. Only about a quarter of the participants (n=218, 28%) achieved a vigorous activity level, which might be partly explained by the cut point used (6 MET) for vigorous activity. In the current study, these MET cut-off points were based on studies conducted in younger individuals and they might not reflect appropriate MET values for vigorous activity older adults. Evenson et al. (2012) stated that the use of fixed cut points to determine MVPA seemed to be most problematic for older adults. Some older adults might not be able to undertake such vigorous activities (6 METS and above) because of their declined maximal cardiorespiratory fitness (25). Furthermore, it might be more difficult for older

adults to reach moderate and vigorous activities when activities are categorized as absolute intensities (26) as was seen in a recent study where older participants had lower total activity counts, as well as lower light, moderate, and vigorous activity minutes compared with younger participants when PA was measured as absolute intensity while they had more moderate and vigorous intensity minutes when PA was defined as relative intensity (27). Many older adults with low cardiorespiratory fitness might feel as if they are exercising vigorously, but they might not reach even a moderate activity level when the activity is assessed with accelerometers. On average, for adults aged 65 to 79, moderate-intensity activity perceived as “somewhat hard” corresponds to 3.2 to 4.7 METS, and vigorous-intensity activity perceived as “hard” corresponds to 4.8 to 6.7 METS (28). Therefore, the cut point for vigorous activity (6 MET) might have been too high for our population. So far there is currently no consensus on the optimal cut-off points for older adults.

The results showed that those participants who had poor fitness undertook significantly fewer light activities (MET level 1.5-3) and had fewer sit-to-stand transitions compared to those in the other fitness categories. No differences were observed between the other fitness categories. Light activities among older adults could pertain to household chores and self-care tasks, such as dressing (2.5 MET), washing dishes (1.8 MET), cooking and food preparation (2.5 MET) (29). These results indicate that older adults who consider that they have poor fitness are not able to maintain for a long time the intensity level required for ADL tasks, such as self-care activities or basic mobility-related household activities (4,5). Even though the poor category has relatively few participants (n=35), the observation of significantly lower activity level in the poor fitness group clearly shows that the easily administered self-reported fitness question can be used to identify people with poor fitness and low levels of physical activity. Therefore these individuals can be targeted with interventions and detailed examinations. However, self-reported fitness is not the only indicator of physical activity level among elderly as 80% of the variation is explained by

other factors; therefore to increase physical activity level of people with low fitness could be achieved by many different ways.

The decreased time spent in PA among participants with poor fitness has been substituted with increased sedentary behavior, both lying and sitting, although these were not all statistically significant when adjusted with BMI and disease. When one examines the mean times (Table 3), there were substantial differences in these activities between the least fit individuals and those with better fitness. For older adults with poor fitness, it might be important to address the issue of switching the activities from long-term sitting to interruptions of sitting, various light activities and then further on to moderate activities. This could improve both their perceived fitness and actual fitness gradually so that they start to believe that they are capable of undertaking physically more demanding daily activities. Achieving slightly more activities and feeling more fit might be important in preventing and reducing frailty and improving perceived health among older adults.

Strengths and limitations

To our knowledge, this is the first study that has examined the associations between self-reported fitness and objectively measured PA and SB in a twin study setting. Besides the twin-study setting, another major strength is the use of state-of-the-art and valid algorithms to determine a variety of PA and SB variables (17-19). Also, the total sample size was relatively large.

The main limitation is the cross-sectional design, preventing us from drawing conclusions about the direction of causality. Participants could have reduced their physical activity first due to some reason (e.g. disease or injury) and this reduced activity might have caused them to feel unfit or vice versa. Feeling of being unfit due to any reason is likely inactivate people. Another limitation of the study is the known limitations of accelerometers, such as inability to measure exercise intensity correctly during cycling, swimming and resistance training (30). However, most common type of exercise within this age group is walking, therefore the results would not be

affected very much by this limitation. Also, the lack of objective measurements of physical functioning and fitness are limitations.

Conclusion

A single-item question on self-reported fitness is associated with daily activity profile among community-dwelling older people. Elderly individuals who perceive their fitness poor would likely benefit from being identified and from appropriate measures to improve factors accounting for their low mobility and inactivity.

Acknowledgement

Author contributions: U.K and H.S designed the study, K.W drafted the manuscript and analysed the data, H.V-Y. and N.L collected data, J.K led the twin cohort collection. All authors contributed to the revision of the manuscript and approved the final version of the manuscript.

Conflict of interest

none

References

1. Blair SN, Cheng Y, Holder JS. Is physical activity or physical fitness more important in defining health benefits? *Med Sci Sports Exerc.* 2001;33(6 Suppl):S379-99; discussion S419-20.
2. Kodama S, Saito K, Tanaka S, et al. Cardiorespiratory fitness as a quantitative predictor of all-cause mortality and cardiovascular events in healthy men and women: a meta-analysis. *JAMA.* 2009;301(19):2024-2035. DOI: 10.1001/jama.2009.681
3. Booth FW, Roberts CK, Laye MJ. Lack of exercise is a major cause of chronic diseases. *Compr Physiol.* 2012;2(2):1143-1211. DOI: 10.1002/cphy.c110025.
4. Weening-Dijksterhuis E, de Greef MH, Krijnen W, van der Schans CP. Self-reported physical fitness in frail older persons: reliability and validity of the Self-Assessment of Physical Fitness (SAPF). *Percept Mot Skills.* 2012;115(3):797-810. DOI:10.2466/10.06.15
5. van Heuvelen MJ, Kempen GI, Brouwer WH, de Greef MH. Physical fitness related to disability in older persons. *Gerontology.* 2000;46(6):333-341. DOI:10.1159/000022187
6. Sardinha LB, Cyrino ES, Santos LD, Ekelund U, Santos DA. Fitness but not weight status is associated with projected physical independence in older adults. *Age (Dordr).* 2016;38(3):54-016-9911-4. DOI: 10.1007/s11357-016-9911-4
7. Bastone Ade C, Ferriolli E, Teixeira CP, Dias JM, Dias RC. Aerobic Fitness and Habitual Physical Activity in Frail and Nonfrail Community-Dwelling Elderly. *J Phys Act Health.* 2015;12(9):1304-1311. DOI: 10.1123/jpah.2014-0290
8. Walston J, Hadley EC, Ferrucci L, et al. Research agenda for frailty in older adults: toward a better understanding of physiology and etiology: summary from the American Geriatrics

Society/National Institute on Aging Research Conference on Frailty in Older Adults. *J Am Geriatr Soc.* 2006;54(6):991-1001. DOI: 10.1111/j.1532-5415.2006.00745.x

9. Dyrstad SM, Anderssen SA, Edvardsen E, Hansen BH. Cardiorespiratory fitness in groups with different physical activity levels. *Scand J Med Sci Sports.* 2016;26(3):291-298. DOI: 10.1111/sms.12425

10. Keevil VL, Cooper AJ, Wijndaele K, et al. Objective Sedentary Time, Moderate-to-Vigorous Physical Activity, and Physical Capability in a British Cohort. *Med Sci Sports Exerc.* 2016;48(3):421-429. DOI: 10.1249/MSS.0000000000000785

11. Santos DA, Silva AM, Baptista F, et al. Sedentary behavior and physical activity are independently related to functional fitness in older adults. *Exp Gerontol.* 2012;47(12):908-912. DOI: 10.1016/j.exger.2012.07.011

12. Keith NR, Clark DO, Stump TE, Miller DK, Callahan CM. Validity and reliability of the Self-Reported Physical Fitness (SRFit) survey. *J Phys Act Health.* 2014;11(4):853-859. DOI: 10.1123/jpah.2012-0264

13. Lerche L, Olsen A, Petersen KE, et al. Validity of physical activity and cardiorespiratory fitness in the Danish cohort "Diet, Cancer and Health-Next Generations". *Scand J Med Sci Sports.* 2017. Dec;27(12):1864-1872. DOI: 10.1111/sms.12873

14. Aadahl M, Kjær M, Kristensen JH, Møllerup B, Jørgensen T. Self-reported physical activity compared with maximal oxygen uptake in adults. *European Journal of Cardiovascular Prevention & Rehabilitation.* 2007;14(3):422-428. DOI: 10.1097/HJR.0b013e3280128d00

15. van Heuvelen MJG, Kempen GIJM, Ormel J, De Greef MHG. Self-reported physical-fitness of older persons - A substitute for performance-based measures of physical-fitness. *Journal of aging and physical activity*. 1997;5(4):298-310.
16. Waller K, Vähä-Ypyä H, Törmäkangas T, et al. Long-term leisure-time physical activity and other health habits as predictors of objectively monitored late-life physical activity - A 40-year twin study. *Sci Rep*. 2018;8(1):9400-018-27704-7. DOI: 10.1038/s41598-018-27704-7
17. Vähä-Ypyä H, Husu P, Suni J, Vasankari T, Sievänen H. Reliable recognition of lying, sitting, and standing with a hip-worn accelerometer. *Scand J Med Sci Sports*. 2018 Mar;28(3):1092-1102. DOI: 10.1111/sms.13017.
18. Vähä-Ypyä H, Vasankari T, Husu P, Suni J, Sievänen H. A universal, accurate intensity-based classification of different physical activities using raw data of accelerometer. *Clin Physiol Funct Imaging*. 2015;35(1):64-70. DOI: 10.1111/cpf.12127
19. Vähä-Ypyä H, Vasankari T, Husu P, et al. Validation of Cut-Points for Evaluating the Intensity of Physical Activity with Accelerometry-Based Mean Amplitude Deviation (MAD). *PLoS One*. 2015;10(8):e0134813. DOI: 10.1371/journal.pone.0134813
20. Tudor-Locke C, Bassett DR, Jr. How many steps/day are enough? Preliminary pedometer indices for public health. *Sports Med*. 2004;34(1):1-8.
21. Egerton T, Chastin SF, Stensvold D, Helbostad JL. Fatigue May Contribute to Reduced Physical Activity Among Older People: An Observational Study. *J Gerontol A Biol Sci Med Sci*. 2016;71(5):670-676. DOI: 10.1093/gerona/glv150
22. Stubbe JH, Boomsma DI, Vink JM, et al. Genetic influences on exercise participation in 37,051 twin pairs from seven countries. *PLoS ONE*. 2006;1(1).

<http://www.plosone.org/article/info:doi/10.1371/journal.pone.0000022>. DOI:

10.1371/journal.pone.0000022 .

23. Bouchard C, Dionne FT, Simoneau JA, Boulay MR. Genetics of aerobic and anaerobic performances. *Exerc Sport Sci Rev*. 1992;20:27-58.

24. Zadro JR, Shirley D, Andrade TB, Scurrah KJ, Bauman A, Ferreira PH. The Beneficial Effects of Physical Activity: Is It Down to Your Genes? A Systematic Review and Meta-Analysis of Twin and Family Studies. *Sports Med Open*. 2017;3(1):4-016-0073-9. DOI: 10.1186/s40798-016-0073-9

25. Evenson KR, Buchner DM, Morland KB. Objective measurement of physical activity and sedentary behavior among US adults aged 60 years or older. *Prev Chronic Dis*. 2012;9:E26.

26. Kujala UM, Pietila J, Myllymaki T, et al. Physical Activity: Absolute Intensity versus Relative-to-Fitness-Level Volumes. *Med Sci Sports Exerc*. 2017;49(3):474-481. DOI: 10.1249/MSS.0000000000001134

27. Schrack JA, Leroux A, Fleg JL, et al. Using Heart Rate and Accelerometry to Define Quantity and Intensity of Physical Activity in Older Adults. *J Gerontol A Biol Sci Med Sci*. 2018;73(5):668-675. DOI: 10.1093/gerona/gly029

28. Pollock M, Gaesser GA, Butcher J, et al. American College of Sports Medicine Position Stand. The recommended quantity and quality of exercise for developing and maintaining cardiorespiratory and muscular fitness, and flexibility in healthy adults. *Med Sci Sports Exerc*. 1998;30(6):975-991.

29. Ainsworth BE, Haskell WL, Herrmann SD, et al. The Compendium of Physical Activities Tracking Guide. <https://sites.google.com/site/compendiumofphysicalactivities/>7.6.2016.

30. Sievanen H, Kujala UM. Accelerometry-Simple, but challenging. *Scand J Med Sci Sports*. 2017;27(6):574-578. DOI: 10.1111/sms.12887

Table 1. Group characteristics broken down by the self-reported fitness categories.

	Very good (1)	Fairly good (2)	Satisfactory (3)	Poor (4)	
	n = 127	n = 389	n = 236	n = 35	p
Sex					0.033 ^b
Male, n (%)	75 (59.1%)	181 (46.5%)	106 (44.9%)	21 (60%)	
Female, n (%)	52 (40.9%)	208 (53.5%)	130 (55.1%)	14 (40%)	
Age, mean years (SD)	72.9 (0.9)	72.9 (1.0)	72.8 (0.9)	72.6 (1.0)	0.23 ^a
BMI, mean (SD)	24.3 (2.9)	25.7 (3.3)	27.6 (4.6)	27.3 (4.8)	<0.001 ^a
Disease restricting mobility					<0.001 ^b
No, n (%)	121 (96%)	325 (85%)	137 (59%)	12 (36%)	
Yes, n (%)	5 (4%)	58 (15%)	96 (41%)	21 (64%)	
Accelerometer wear time	14:08:50 (1:04:58)	14:01:19 (1:24:47)	14:00:31 (1:21:54)	13:50:56 (1:36:18)	0.27 ^a
Accelerometer wear days	6.85 (0.46)	6.72 (0.59)	6.70 (0.62)	6.6 (0.74)	0.027

^a Linear regression, sex and age adjusted

^b Rao-Scott symmetry test

Table 2. Explanatory value (R^2) of self-reported fitness alone and in the multivariable model explaining various objectively measured physical activity variables.

	Model 1, R² for self-reported fitness ^a	Model 2, Multivariable R² ^b
Activity variable		
Steps	18.5%	26.1%
Running steps	8.8%	11.6%
Sit-to-stand transitions	6.1%	19.0%
Daily MET	20.8%	30.5%
Peak-10 min MET	19.4%	31.9%
No. > 10 min activity bouts	12.8%	20.5%
Time spent in each activity level		
Lying	4.2%	6.4%
Sitting	3.2%	6.3%
Standing	4.3%	14.4%
Light activity	4.7%	8.2%
Moderate activity	19.4%	29.9%
Vigorous activity ^c	13.4%	33.1%
SB	8.8%	13.9%
MVPA	20.4%	31.4%

^a Model 1: basic regression analysis, cl(family), additional R^2 for self-reported fitness after age- and sex adjustment

^b Model 2: Multivariable model; activity variable + age, sex, BMI and disease, R^2 includes all the variables

^c Based on 218 subjects, who had vigorous activity

Table 3. Associations between self-reported fitness and accelerometer measured physical activity. ^a

	Very good (1)	Fairly good (2)	Satisfactory (3)	Poor (4)		
	n = 127	n = 389	n = 236	n = 35	sig. diff. between groups ^{b,} c	sig. diff. between groups ^{b,} d
Activity variable	mean (SD)	mean (SD)	mean (SD)	mean (SD)		
Steps	8 558 (2 913)	6 724 (2 937)	5 196 (2683)	2797 (1742)	a – f	a – f
Running steps	248 (564)	74 (179)	44 (87)	33 (25)	a – e	a – c, e
Sit-to-stand transitions	38.5 (11.7)	37.6 (12.7)	33.4 (15.4)	23.0 (16.2)	b – f	c, e, f
Overall MET per day	1.50 (0.15)	1.40 (0.13)	1.34 (0.13)	1.22 (0.08)	a – f	a – f
Peak-10 min MET	4.3 (1.18)	3.7 (0.81)	3.2 (0.75)	2.7 (0.44)	a – f	a – f
No. bouts ≥ 10 min activity	3.47 (1.6)	2.78 (1.7)	2.15 (1.6)	1.17 (1.1)	a – f	a – f
Time spent doing different activities	mean h:min:sec (SD)	mean h:min:sec (SD)	mean h:min:sec (SD)	mean h:min:sec (SD)		
Lying	1:00:01 (0:50:41)	1:07:57 (0:58:27)	1:24:32 (1:07:17)	2:08:10 (2:05:22)	b – e	-

Sitting	7:20:33 (1:19:41)	7:35:15 (1:30:12)	7:58:26 (1:28:13)	8:26:13 (2:04:17)	b – d	b
Standing	1:34:28 (0:42:13)	1:34:46 (0:44:01)	1:20:36 (0:43:28)	0:56:43 (0:45:35)	b - f	e
Light activity	3:12:53 (0:57:07)	3:01:33 (1:02:33)	2:47:52 (1:03:42)	2:08:13 (0:56:06)	b, c, e, f	c, e, f
Moderate activity	0:57:29 (0:29:43)	0:41:07 (0:24:21)	0:28:52 (0:21:57)	0:11:37 (0:08:59)	a – f	a – f
Vigorous activity	0:03:27 (0:07:21)	0:00:40,4 (0:03:12)	0:00:13,2 (0:01:29)	0:00:00,66 (0:00:02)	a – c, e, f	a – c
SB	8:20:16 (1:24:22)	8:41:38 (1:38:36)	9:21:45 (1:39:10)	10:33:29 (2:02:17)	b – f	b – e
MVPA	1:00:55 (0:31:41)	0:41:48 (0:24:56)	0:29:06 (0:22:20)	0:11:37 (0:08:59)	a – f	a – f

^a Mean (SD) calculated with bootstrapping based on 1000 samples

^b Linear regression followed with Sidak's multiple comparisons test, significant difference ($p < 0.05$) between groups represented by the following letters;

a = 1 vs. 2, b= 1 vs. 3, c= 1 vs. 4, d= 2 vs. 3, e = 2 vs. 4, f = 3 vs. 4

^c Adjusted for age and sex

^d Adjusted for age, sex, BMI and disease

Table 4. Objectively measured PA and SB in twin pairs discordant for self-reported fitness

	Mean time activity/day^a hours:minutes:seconds median (IQR) (95% CI)		Z and p value^b
	Co-twins with higher self-reported fitness	Co-twins with lower self-reported fitness	
Sedentary behavior time/day, h:min			
All pairs (n=156)	8:36 (2:12) (8:23 - 8:55)	9:12 (2:17) (8:48 - 9:29)	Z= -3.17 p=0.002
DZ pairs (n=90)	8:42 (2:25) (7:59 - 9:05)	9:21 (2:33) (8:47 - 9:48)	Z=-3.185 p=0.001
MZ pairs (n=57)	8:40 (1:53) 8:27 - 9:06)	9:10:22 (1:46:45) (8:43 - 9:29)	Z= -0.683 p=0.494
Standing time/day, h:min			
All pairs	1:29 (1:02) (1:14 - 1:36)	1:23 (1:02) (1:12 - 1:32)	Z= -1.320 p=0.187
DZ pairs	1:27 (1:08) (1:13 - 1:38)	1:12 (1:08) (1:01 - 1:27)	Z= -2.102 p=0.036
MZ pairs	1:32 (0:57) (1:07 - 1:42)	1:31 (0:50) (1:19 - 1:44)	Z= -0.624 p=0.533
Time of light physical activity/day, h:min			
All pairs	2:52 (1:09) (2:40 - 3:01)	2:40 (1:20) (2:27 - 2:58)	Z= -2.350 p=0.019

DZ pairs	3:01 (1:16) (2:43 - 3:16)	2:34 (1:22) (2:25 - 3:04)	Z= -3.261 p=0.001
MZ pairs	2:34 (0:55) (2:25 - 2:55)	2:56 (1:18) (2:24 - 3:09)	Z= -0.282 p=0.778
Time of moderate-to-vigorous physical activity/day, h:min:sec			
All pairs	0:38:14 (0:35:54) (0:33:22 - 0:42:57)	0:28:11 (0:32:12) (0:24:26 - 0:34:44)	Z= -4.093 p<0.001
DZ pairs	0:40:16 (0:40:57) (0:32:11 - 0:46:07)	0:23:24 (0:34:01) (0:17:30 - 0:29:51)	Z= -4.140 p<0.001
MZ pairs	0:38:37 (0:29:53) (0:28:50 - 0:45:31)	0:36:31 (0:28:30) (0:30:13 - 0:42:27)	Z= -0.973 p=0.330
Daily step count, number of steps			
All pairs	6513 (3943) (5928 - 7142)	4968 (3676) (4475 - 5759)	Z= -4.611 p<0.001
DZ pairs	6678 (4142) (6019 - 7998)	4652 (3049) (4128 - 5569)	Z= -4.589 p<0.001
MZ pairs	6506 (3909) (5690 - 7937)	6406 (4284) (4805 - 6804)	Z= -1.053 p=0.292
Peak-10min MET, MET			
All pairs	3.69 (1.31) (3.53 - 3.85)	3.21 (1.06) (3.14 - 3.37)	Z= -4.423 p<0.001
DZ pairs	3.65 (1.24) (3.44 - 3.83)	3.12 (0.94) (2.91 - 3.23)	Z= -4.243 p<0.001
MZ pairs	3.72 (1.29) (3.30 - 4.16)	3.43 (0.99) (3.26 - 3.83)	Z= -1.212 p=0.226

^aDescriptive analyses with bootstrapping (1000 samples)

^bZ-score and p by Wilcoxon matched-pairs signed-rank test




Figure 1. Average activity profile per day for each self-rated fitness categories.








Figure 2. Different objectively measured activities according to the self-reported fitness categories separately for males and females. Bars and the error bars represent the unadjusted means and 95% CI of the means.