

Brief Report

Adaptation of bacteria to glyphosate: a microevolutionary perspective of the enzyme 5-enolpyruvylshikimate-3-phosphate synthase

Miia J. Rainio,¹ Suvi Ruuskanen,¹ Marjo Helander,¹
Kari Saikkonen,² Irma Saloniemi¹ and Pere Puigbò^{1,3,4*}

 1,3,4*

¹Department of Biology, University of Turku, Turku, Finland.

²Biodiversity Unit, University of Turku, Turku, Finland.

³Nutrition and Health Unit, Eurecat Technology Centre of Catalonia, Reus, Catalonia, Spain.

⁴Department of Biochemistry and Biotechnology, Rovira i Virgili University, Tarragona, Catalonia, Spain.

Abstract

Glyphosate is the leading herbicide worldwide, but it also affects prokaryotes because it targets the central enzyme (5-enolpyruvylshikimate-3-phosphate, EPSP) of the shikimate pathway in the synthesis of the three essential aromatic amino acids in bacteria, fungi and plants. Our results reveal that bacteria may easily become resistant to glyphosate through changes in the 5-enolpyruvylshikimate-3-phosphate synthase active site. This indicates the importance of examining how glyphosate affects microbe-mediated ecosystem functions and human microbiomes.

Background

Evolutionarily conserved 5-enolpyruvylshikimate-3-phosphate synthase (EPSPS) is the key enzyme of the shikimate pathway in the synthesis of three essential aromatic amino acids (phenylalanine, tyrosine and tryptophan) across taxa, including bacteria, fungi and plants (Bentley, 1990; Richards *et al.*, 2006). The EPSPS enzyme is also involved in the production of the para-aminobenzoate (McConkey, 1999). Thus, glyphosate may have an effect on the synthesis of folate even in environments rich in phenylalanine, tyrosine and tryptophan. The EPSPS enzyme is also known as AroA (EC:2.5.1.19) and catalyses the reaction

from shikimate-3-phosphate to chorismate. Because the shikimate pathway does not occur in vertebrates, the use of glyphosate, targeting the EPSPS enzyme by competing for the binding site with the second substrate of the EPSPS (phosphoenol pyruvate, PEP) (Schönbrunn *et al.*, 2001), is regarded safe for use in food production.

Due to its affordable price, effectiveness and broad-spectrum ability to kill weeds, glyphosate has become the most commonly used herbicide worldwide (Woodburn, 2000; Helander *et al.*, 2012; Myers *et al.*, 2016). Although glyphosate antibiotic properties are known (Kurenbach *et al.*, 2015; Kurenbach *et al.*, 2018), its possible effects on microbiomes (Funke *et al.*, 2006; Motta *et al.*, 2018; Gómez-Gallego *et al.*, 2020; Ruuskanen *et al.*, 2020) have largely been neglected until recently (Motta *et al.*, 2018; Mesnage *et al.*, 2019). As microbes have driven eco-evolutionary processes since the origin of life, we note the importance of thoroughly understanding the possible undesirable effects of glyphosate on ecosystem structures, functions and services (Bote *et al.*, 2019). Here, we propose that the wide use of glyphosate may potentially affect microbial communities by (i) eradicating susceptible microbes and (ii) leading to evolution of resistance.

Results and discussion

We analysed 663 microbial genomes distributed across 32 alignable tight genomic clusters (ATGCs) (Novichkov *et al.*, 2009; Puigbò *et al.*, 2014), with the *EPSPSClass* server (<http://ppuigbo.me/programs/EPSPSClass>) (Leino *et al.*, 2020) to identify changes in sensitivity to glyphosate. The algorithm classifies EPSPS proteins based on the presence and absence of amino acid markers in the active site that determine whether the enzyme is putatively sensitive (class I) or resistant (class II, III and IV) to the glyphosate (see methods). Currently, the majority of species of prokaryotes belong to class I, thus are putatively affected by the herbicide (Leino *et al.*, 2020). We analysed 32 ATGCs that amply represented bacteria and one archaeal group. The EPSPS sequence was not found in ATGC033 (*Mycoplasma*), small-genome single-

Received 1 October, 2020; accepted 20 January, 2021. **For correspondence. E-mail pepuav@utu.fi.

© 2021 The Authors. *Environmental Microbiology Reports* published by Society for Applied Microbiology and John Wiley & Sons Ltd. This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

membrane bacteria (Meseguer *et al.*, 2003); ATGC046 (*Rickettsia*), obligate intracellular parasites (Zomorodipour and Andersson, 1999) and ATGC056 (*Lactobacillus*). Even though *Lactobacillus* strains in the ATGC dataset do not have the EPSPS sequence, other strains have the enzyme and present differential sensitivity to glyphosate. Previous studies have shown that some *Lactobacillus* species are susceptible to glyphosate (Shehata *et al.*, 2013).

A microevolutionary analysis of the EPSPS enzyme shows that phylogenetics best explains bacterial sensitivity to glyphosate (Fig. 1, Table 1 and Supplementary - Table 1). In general, Firmicutes are significantly more (putatively) resistant to glyphosate than Proteobacteria, whereas Actinobacteria is the most sensitive group to glyphosate. Furthermore, bacterial lifestyle is also associated with sensitivity, i.e. facultative host-associated (FHA) bacteria are more sensitive to glyphosate than free-living (FL) bacteria (Supplementary Tables 2 and 3; Supplementary Fig. 1). However, an analysis with a

greater number of samples would be necessary to test whether this association with lifestyle is independent of phylogeny. Intracellular parasites seem to be the most putatively sensitive group; however, due to the limited amount of data, we cannot determine statistical significance for this group's sensitivity. We hypothesize that sensitivity mirrors the strength of selective force of glyphosate on bacteria. FL bacteria are directly exposed to glyphosate spray at high concentrations, whereas host-associated microbes can be exposed to lower concentrations when the glyphosate is moving through the plant from leaves to roots, and intracellular parasites might be protected by the cell membrane. Genomes with %GC below 40% tend to be more resistant to glyphosate (Fig. 2A). Spearman correlation coefficients between sensitivity and genome parameters from the original ATGC dataset (Puigbò *et al.*, 2014) were not statistically significant (see methods and Supplementary Table 4). To statistically decouple the contribution of %GC, ATGC clusters




Fig. 1. Distribution of the predicted sensitivity to glyphosate across the species tree. Pie charts indicate the proportion of species that are putatively sensitive (green) or resistant (red) to glyphosate, and unclassified (black). See full description in Table 1.

Table 1. Summary table of the percentage of species sensitive/resistant to glyphosate.

ATGC	Genera	Taxonomy	Lifestyle	N	CDS	GC	P (%)	CI	CII	CIII	CIV	U	%S (%)
ATGC001	Enterobacteria	Proteobacteria	FHA	109	4652	0.528	8.24	109	1	0	0	0	99
ATGC002	Enterobacter– Klebsiella	Proteobacteria	FHA	11	5375	0.583	8.16	10	0	0	0	1	91
ATGC003	<i>Streptococcus</i>	Firmicutes	FL	22	2114	0.408	9.37	0	22	0	0	0	0
ATGC004	<i>Streptococcus</i>	Firmicutes	FL	22	1865	0.395	8.97	0	22	0	0	0	0
ATGC005	Streptococcus	Firmicutes	FL	16	1977	0.422	9.21	0	15	0	0	4	0
ATGC014	Bacillus	Firmicutes	FL	31	5704	0.361	9.42	0	30	0	0	2	0
ATGC015	<i>Bacillus</i>	Firmicutes	FL	24	4089	0.459	8.91	0	24	0	0	0	0
ATGC021	<i>Chlamydia</i>	Chlamydia	P	44	898	0.417	8.87	44	0	0	0	0	100
ATGC022	Chlamydia– Chlamydophila	Chlamydia	P	19	1126	0.397	8.93	3	0	0	0	17	15
ATGC025	<i>Mycobacterium</i>	Actinobacteria	FHA	32	3981	0.659	6.53	34	0	0	0	0	100
ATGC052	<i>Helicobacter</i>	Proteobacteria	FHA	51	1564	0.397	9.80	0	0	0	0	51	0
ATGC054	<i>Staphylococcus</i>	Firmicutes	FL	41	2624	0.337	9.15	0	41	0	0	0	0
ATGC067	Corynebacterium	Actinobacteria	FL	15	2089	0.530	7.06	12	0	0	0	3	80
ATGC068	Corynebacterium	Actinobacteria	FL	12	2262	0.543	7.05	11	0	0	0	1	92
ATGC072	<i>Pseudomonas</i>	Proteobacteria	FL	12	5386	0.633	7.52	0	12	0	0	0	0
ATGC082	<i>Clostridium</i>	Firmicutes	FL	10	3691	0.292	9.20	0	10	0	0	0	0
ATGC089	<i>Burkholderia</i>	Proteobacteria	FL	13	5743	0.688	7.30	0	0	0	0	25	0
ATGC090	Burkholderia	Proteobacteria	FL	12	6697	0.678	7.43	12	0	0	0	2	86
ATGC094	<i>Sulfolobus</i>	Archaea	FL	12	2735	0.356	10.20	12	0	0	0	0	100
ATGC105	<i>Bifidobacterium</i>	Actinobacteria	FHA	11	1972	0.609	7.52	11	0	0	0	0	100
ATGC106	<i>Bifidobacterium</i>	Actinobacteria	FHA	10	1567	0.615	7.47	10	0	0	0	0	100
ATGC109	Listeria	Firmicutes	FL	31	2891	0.385	8.93	0	29	0	0	2	0
ATGC121	<i>Shewanella</i>	Proteobacteria	FL	14	4360	0.475	8.22	14	0	0	0	0	100
ATGC128	<i>Yersinia</i>	Proteobacteria	FHA	19	4114	0.489	8.12	19	0	0	0	0	100
ATGC135	<i>Xanthomonas</i>	Proteobacteria	FHA	13	4447	0.654	7.19	0	13	0	0	0	0
ATGC137	<i>Brucella– Ochrobactrum</i>	Proteobacteria	FHA	19	3268	0.584	7.59	0	19	0	0	0	0
ATGC138	Neisseria	Proteobacteria	FHA	18	1961	0.537	8.27	17	0	0	0	1	94
ATGC139	<i>Francisella</i>	Proteobacteria	FHA	18	1634	0.333	9.93	18	0	0	0	0	100
ATGC144	Campylobacter	Proteobacteria	FHA	11	1646	0.309	10.31	0	10	0	0	2	0
ATGC153	<i>Acinetobacter</i>	Proteobacteria	FL	14	3759	0.403	8.82	0	13	0	0	0	0
ATGC163	Propionibacterium	Actinobacteria	FL	11	2280	0.608	6.51	9	0	0	0	2	82
ATGC186	<i>Legionella</i>	Proteobacteria	FHA	10	3166	0.391	9.16	0	10	0	0	0	0

ATGCs containing species with potential differential sensitivity to glyphosate are highlighted in bold. FHA: facultative host-associated; FL: free-living; P: intracellular parasite. CDS: number of protein coding genes; N: number of genomes; GC: median GC; P: percentage of aromatic amino acids. CI: class I; CII: class II; CIII: class III; CIV: class IV; %S: percentages of sensitive species.

(closely related phylogenetic species), number of genes, taxonomic groups (e.g., Actinobacteria, Proteobacteria, Firmicutes) and lifestyle (FL, FHA and intracellular parasites) to the bacterial sensitivity to glyphosate (sensitive, resistant, unknown), we carried out a multiple correspondence analysis (MCA) and random forest (RF) classification model. The MCA analysis revealed that Firmicutes and FL bacteria are mostly resistant, whereas Actinobacteria, Proteobacteria and FHA appeared to be sensitive to glyphosate (Fig. 2B and V). The RF suggests that ATGC clusters (i.e. phylogenetics) are the most important definer of sensitivity to glyphosate (Fig. 2D).

EPSPS is the central enzyme in the metabolic pathway for the synthesis of three essential amino acids in bacteria (Steinrücken and Amrhein, 1980); thus, its protein sequence is presumably evolutionarily conserved (Novick and Doolittle, 2020). Indeed, the EPSPS sequence is highly conserved within 17 ATGCs that preserve sensitivity status across species. However, 12 ATGCs contained

species with different sensitivities to glyphosate (Fig. 3, Table 1 and Supplementary Figs 2–13). Although the active site of the EPSPS is highly conserved in prokaryotes, in some species, the sensitivity to glyphosate may be altered with few changes in the active site (Fig. 3). In Enterobacteria (ATGC001), only *Enterobacter* R4 368 (out of 109 species) is putatively resistant to glyphosate after recently acquiring a second copy of the EPSPS via horizontal gene transfer (HGT) from a resistant strain of *Klebsiella pneumoniae* (Fig. 4). Other Proteobacteria clusters have species with differential sensitivity status after few amino acid changes in the active site (Fig. 3). This microevolutionary analysis shows that some bacteria may become putatively resistant to glyphosate through small changes, analogous to the antibiotic resistance mechanism (Kurenbach *et al.*, 2018; Bote *et al.*, 2019). Thus, the wide use of glyphosate may have a very large impact on the species diversity and composition of microbial communities not only because of a




Fig. 2. Importance of variables to determine sensitivity to glyphosate. A. Kernel density distribution of predicted sensitivity in the ATGCs by %GC. Genomes with low %GC (<40%) are more resistant to glyphosate (chi test p -value <0.001). B. Multiple correspondence analysis (MCA) plot from 656 genomes and the variables sensitivity, ATGC, taxonomic group, %GC content, number of genes and lifestyle. Additional plots are available in Supplementary Fig. 15. C. Plot of loadings in the MCA. Values are available in Supplementary Table 12. D. Variable importance plot from a random forest classification model (number of trees = 1000) based on mean decrease accuracy and mean decrease Gini. The higher the value of mean decrease accuracy and mean decrease Gini, the higher the importance of the variable in the model.

potential purifying selection effect against sensitive bacteria but also because (i) some bacterial groups may adapt rapidly to become resistant to glyphosate and (ii) glyphosate-based herbicides may enhance multidrug resistance in bacteria (Liu *et al.*, 2020). Previous studies have shown that glyphosate can affect the microbiome composition of insects, including bees (Motta *et al.*, 2018), potato beetles (Gómez-Gallego *et al.*, 2020) and birds (Ruuskanen *et al.*, 2020). Moreover, preliminary results suggest a putative effect in the human (Leino *et al.*, 2020) and rat (Mesnage *et al.*, 2019) gut microbiomes. In addition, glyphosate may affect ecosystem functions including nutrient cycle and carbon balance, plant growth-promoting microbes (e.g. mycorrhizae and nitrogen-fixing bacteria), microbially mediated plant-herbivore and pathogen interactions, and pollination biology depending on the related microbial community composition and its sensitivity to glyphosate (Helander *et al.*, 2018; Gómez-Gallego *et al.*, 2020).

An analysis of variability in the amino acid landscape (see methods) shows that the amino acids in the active site of the EPSPS that bind PEP are 23% more conserved than the rest of the amino acid residues

(Supplementary Fig. 14), in agreement with results from previous studies (Schönbrunn *et al.*, 2001). Moreover, amino acids flanking binding sites (± 5 residues) also show higher levels of conservation than average, possibly due to selection preserving the function of the EPSPS. Thus, additional residues within the active site may determine sensitivity to the herbicide. Bacteria potentially more resistant to glyphosate tend to have fewer amino acid substitutions than sensitive ones, e.g. the binding amino acids of actinobacteria and FL bacteria (more resistant to glyphosate) tend to be more conserved (Fig. 5 and Supplementary Tables 5 and 6). Although there are homogeneous and heterogeneous clusters in relation to glyphosate sensitivity, the active site is equally conserved (Supplementary Table 7). Further analyses of amino acid variabilities in EPSPS (Supplementary Table 8) should help elucidate the effects of glyphosate on microbial communities. Moreover, predictions based on amino acid markers may partially explain resistance to glyphosate. However, glyphosate resistance may also occur due to alternative mechanisms, such as efflux pumps or elevated expression of the EPSPS (Staub *et al.*, 2012).




Fig. 3. Examples of the differential predicted sensitivity to glyphosate in three Proteobacteria ATGCs. Phylogenetic species trees of three ATGCs. Filled boxes show bacteria putatively sensitive (green) and resistant (red) to glyphosate, and unclassified (black). Each box from left to right corresponds to Class I alpha, Class I beta, Class II, Class III, Class IV and unclassified. The amino acids in the active site are shown on the right side of the figure. A. Change from sensitive to resistant in *Enterobacter* R4 368 through putative horizontal gene transfer from *Klebsiella pneumoniae* (Fig. 4). B. Two independent changes from sensitive to unclassified in propionibacterium with single amino acid changes. C. Change from resistant to unclassified in *Campylobacter* with several amino acid changes.

Conclusions

In conclusion, this study shows that phylogenetic groups and bacterial lifestyles are key factors determining sensitivity to glyphosate. Our results suggest that the sensitivity of bacterial species to glyphosate varies and can change in the short evolutionary time of ATGC. Moreover, microbes more exposed to the herbicide (FL bacteria) are putatively more resistant. Nevertheless, because single mutations in the EPSPS active site and rapid HGT events may change the status of sensitivity, heavy use of glyphosate may impact microbial biodiversity independently of taxonomy and lifestyle. The central and unanswered questions are (i) what are the effects of glyphosate-modulated microbiomes on ecosystem functions and services and (ii) on human wellbeing.

Methods

Protein sequences

Protein data were obtained from the ATGCs database, which contains data on >4.5 million proteins encoded in

>1500 genomes of prokaryotes (approximately 60% of proteins and 62% of genomes from RefSeq as of June 2013) that met the same criteria as the original ATGCs (Novichkov *et al.*, 2009), as described in Puigbò *et al.* (2014). The ATGC database provides an ideal environment for a microevolutionary analysis, as it provides a widespread representation of genes from a representative set of prokaryotes. However, other databases (e.g. the Clusters of Orthologous Groups (Galperin *et al.*, 2019)) would be more suitable for a comprehensive macroevolutionary analysis of the EPSPS. EPSPS protein sequences were annotated through BLAST mapping onto the Clusters of Orthologous Genes (COG) database (Galperin *et al.*, 2015). EPSPS belongs to COG0128 (category E, amino acid transport and metabolism). We analysed 32 ATGCs that amply represented bacteria and one archaeal group. Each ATGC contains at least 10 species by the definition in the original study (Puigbò *et al.*, 2014), and up to 109 species are present in the enterobacterial cluster (ATGC001). The final ATGC dataset used in this study includes 332 FHA species (13 ATGCs), 312 FL species (17 ATGCs) and

A

CLUSTAL multiple sequence alignment by MUSCLE (3.8)

```

Enterobacter_R4_368_uid208672|YP
Klebsiella_pneumoniae|WP_0658106
Enterobacter_638_uid58727|YP_001
Enterobacter_R4_368_uid208672|YP

Enterobacter_R4_368_uid208672|YP
Klebsiella_pneumoniae|WP_0658106
Enterobacter_638_uid58727|YP_001
Enterobacter_R4_368_uid208672|YP

Enterobacter_R4_368_uid208672|YP
Klebsiella_pneumoniae|WP_0658106
Enterobacter_638_uid58727|YP_001
Enterobacter_R4_368_uid208672|YP

Enterobacter_R4_368_uid208672|YP
Klebsiella_pneumoniae|WP_0658106
Enterobacter_638_uid58727|YP_001
Enterobacter_R4_368_uid208672|YP

Enterobacter_R4_368_uid208672|YP
Klebsiella_pneumoniae|WP_0658106
Enterobacter_638_uid58727|YP_001
Enterobacter_R4_368_uid208672|YP

Enterobacter_R4_368_uid208672|YP
Klebsiella_pneumoniae|WP_0658106
Enterobacter_638_uid58727|YP_001
Enterobacter_R4_368_uid208672|YP

Enterobacter_R4_368_uid208672|YP
Klebsiella_pneumoniae|WP_0658106
Enterobacter_638_uid58727|YP_001
Enterobacter_R4_368_uid208672|YP

Enterobacter_R4_368_uid208672|YP
Klebsiella_pneumoniae|WP_0658106
Enterobacter_638_uid58727|YP_001
Enterobacter_R4_368_uid208672|YP

Enterobacter_R4_368_uid208672|YP
Klebsiella_pneumoniae|WP_0658106
Enterobacter_638_uid58727|YP_001
Enterobacter_R4_368_uid208672|YP
    
```

B

Species	Class I	Class II
Klebsiella pneumoniae (WP_065810684)		X
Enterobacter_R4_368 (YP_008098272.1)		X
Enterobacter_638 (YP_001176159.1)	X	
Enterobacter_R4_368_uid208672 (YP_008110400.1)	X	

C

Species	CLASS	IDENTITY
Klebsiella pneumoniae (WP_065810684)	Class I (alpha)	0.800
	Class I (beta)	0.889
	Class II	1.000
	Class III	0.346
	Class IV	0.167
Enterobacter_R4_368 (YP_008098272.1)	CLASS	IDENTITY
	Class I (alpha)	0.800
	Class I (beta)	0.889
	Class II	1.000
	Class III	0.335
Enterobacter_638 (YP_001176159.1)	CLASS	IDENTITY
	Class I (alpha)	1.000
	Class I (beta)	1.000
	Class II	0.605
	Class III	0.470
Enterobacter_R4_368 (YP_008110400.1)	CLASS	IDENTITY
	Class I (alpha)	1.000
	Class I (beta)	1.000
	Class II	0.605
	Class III	0.454
Class IV	0.233	

Fig. 4. Putative acquisition of glyphosate resistance in *Enterobacter* R4 68 (ATGC001) via horizontal gene transfer from *Klebsiella pneumoniae*. A. Multiple sequence alignment of proteins. The sequence WP_065810684 from *K. pneumoniae* was obtained as a result of the best BLAST hit (identity 100%). B. Classification of the EPSPS enzyme into sensitive (class I) and resistant (class II) to glyphosate. C. Identity values of the four proteins to the reference EPSPS sequences in EPSPSClass server (<https://ppuigbo.me/programs/EPSPSClass/>).




Fig. 5. Amino acid substitutions in the active site and sensitivity to glyphosate by taxonomy and lifestyle. AS: amino acid substitution in the active site; No_AS: amino acid substitutions in other residues.

19 intracellular parasites (two ATGCs). The EPSPS sequence was not found in ATGC033 (*Mycoplasma*), ATGC046 (*Rickettsia*) and ATGC056 (*Lactobacillus*). Even though *Lactobacillus* strains in the ATGC dataset do not have the EPSPS sequence, the enzyme is present in other strains (Supplementary Table 9).

Classification of the EPSPS

Glyphosate inhibits the shikimate pathway by inhibiting the enzyme EPSPS (Schönbrunn *et al.*, 2001; Abraham, 2020). The EPSPS class has been classified with the EPSPSClass server (<http://ppuigbo.me/programs/>

EPSPSClass), which assesses the type of potential sensitivity of the EPSPS enzyme to the herbicide glyphosate (Leino *et al.*, 2020). The amino acid markers to determine potential glyphosate sensitivity to glyphosate (Supplementary Table 10) are also available at Leino *et al.* (2020)) and in the web server. EPSPS enzymes can be classified as class I (alpha or beta) (Light *et al.*, 2016; Firdous *et al.*, 2018), class II (Barry *et al.*, 1997; Priestman *et al.*, 2005), III (Carozzi *et al.*, 2006) or IV (Lira *et al.*, 2013) based on the presence of known amino acid markers (class I, class II and class IV) and motifs (class III) (Barry *et al.*, 1997; Priestman *et al.*, 2005; Carozzi *et al.*, 2006; Lira *et al.*, 2013; Light *et al.*, 2016; Firdous *et al.*, 2018). In addition, a large portion of microbial species are yet unclassified, and their potential sensitivities to glyphosate are unknown (Leino *et al.*, 2020). Different uses of amino acid markers distinguish the EPSPS enzymes into four classes using the reference EPSPS enzymes from *Vibrio cholerae serotype O1* (vcEPSPS, class I), *Coxiella burnetii* (cbEPSPS, class II), *Brevundimonas vesicularis* (bvEPSPS, class III) and *Streptomyces davawensis* (sdEPSPS, class IV).

Analysis of the amino acid substitution landscape

We have performed a protein sequence alignment of the EPSPS sequences from the ATGCs and the reference EPSPS of *Vibrio cholerae serotype O1* (vcEPSPS; Class I (Light *et al.*, 2016; Firdous *et al.*, 2018); Supplementary - Table 5) with the program Muscle (Edgar, 2004). The vcEPSPS sequence was used to determine the location of residues in the active site that interact directly with PEP (Schönbrunn *et al.*, 2001; Healy-Fried *et al.*, 2007). Glyphosate inhibits EPSPS by competing with PEP (Schönbrunn *et al.*, 2001). We counted the number of alternative amino acids in each position of the alignment to determine the degree of amino acid substitution in the active site and the rest of the residues. We determined an active site region, i.e. a wider area around the active site residues (± 5 amino acids) that interacts with PEP.

Statistical analysis

The tested Spearman correlation coefficients between sensitivity and total genome dynamics (as the sum of gains, losses, expansions and reductions; GLER), genome number (genome no.), genome size, protein-coding genes, gene families, median gene cluster content, median guanine+cytosine (median GC), median dN/dS, synteny distance (dY) and percentage of aromatic amino acids were not statistically significant (p -value > 0.05) (Supplementary Table 4). We performed a chi-squared test to determine associations between sensitivity to glyphosate and lifestyle (FL vs FHA) and taxonomic group (Actinobacteria, Firmicutes and Proteobacteria). A Kruskal–Wallis test was

used to analyse conservation of the amino acids in the EPSPS. The importance of each variable (ATGC, taxonomic groups, %GC, genome size and lifestyle) to determine potential sensitivity to glyphosate was performed with a MCA and a RF classification model. Input variables are available in the Supplementary Table 11.

Acknowledgements

Pere Puigbò's research is supported by funds from the Turku Collegium for Science and Medicine (Turku, Finland). Miia Rainio was funded by the Academy of Finland (grant no. 311077 to Marjo Helander).

References

- Abraham W. (2020) US7771736B2 - Glyphosate formulations and their use for the inhibition of 5-enolpyruvylshikimate-3-phosphate synthase - Google Patents [Internet]. [cited 2020 Apr 24]. URL <https://patents.google.com/patent/US7771736B2/en>
- Barry GF, Kishore GM, Padgett SR, Stallings WC. (1997) US5627061A - Glyphosate-tolerant 5-enolpyruvylshikimate-3-phosphate synthases, USA.
- Bentley, R. (1990). The shikimate pathway - a metabolic tree with many branches. *Crit Rev Biochem Mol Biol* **25**: 307–384.
- Bote, K., Pöppe, J., Merle, R., Makarova, O., and Roesler, U. (2019) Minimum inhibitory concentration of glyphosate and of a glyphosate-containing herbicide formulation for *Escherichia coli* isolates - differences between Pathogenic and non-pathogenic isolates and between host species. *Front Microbiol* **10**: 932.
- Carozzi N, Carr B, Hammer P. (2006) US20060253921A1 - Identification of a new class of EPSP synthases, USA.
- Edgar, R.C. (2004) MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Res* **32**: 1792–1797.
- Firdous, S., Iqbal, S., Anwar, S., and Jabeen, H. (2018) Identification and analysis of 5-enolpyruvylshikimate-3-phosphate synthase (EPSPS) gene from glyphosate-resistant *Ochrobactrum intermedium* Sq20. *Pest Manag Sci* **74**: 1184–1196.
- Funke, T., Han, H., Healy-Fried, M.L., Fischer, M., and Schönbrunn, E. (2006) Molecular basis for the herbicide resistance of roundup ready crops. *Proc Natl Acad Sci U S A* **103**: 13010–13015.
- Galperin, M.Y., Kristensen, D.M., Makarova, K.S., Wolf, Y.I., and Koonin, E.V. (2019) Microbial genome analysis: the COG approach. *Brief Bioinformatics* **20**: 1063–1070.
- Galperin, M.Y., Makarova, K.S., Wolf, Y.I., and Koonin, E.V. (2015) Expanded microbial genome coverage and improved protein family annotation in the COG database. *Nucleic Acids Res* **43**: D261–D269.
- Gómez-Gallego, C., Rainio, M.J., Collado, M.C., Mantziari, A., Salminen, S., Saikkonen, K., *et al.* (2020) Glyphosate-based herbicide affects the composition of microbes associated with Colorado potato beetle (*Leptinotarsa decemlineata*). *FEMS Microbiol Lett* **367**: fnaa050.
- Healy-Fried, M.L., Funke, T., Priestman, M.A., Han, H., and Schönbrunn, E. (2007) Structural basis of glyphosate

- tolerance resulting from mutations of Pro101 in *Escherichia coli* 5-enolpyruvylshikimate-3-phosphate synthase. *J Biol Chem* **282**: 32949–32955.
- Helander, M., Saloniemi, I., Omacini, M., Druille, M., Salminen, J.-P., and Saikkonen, K. (2018) Glyphosate decreases mycorrhizal colonization and affects plant-soil feedback. *Sci Total Environ* **642**: 285–291.
- Helander, M., Saloniemi, I., and Saikkonen, K. (2012) Glyphosate in northern ecosystems. *Trends Plant Sci* **17**: 569–574.
- Kurenbach, B., Hill, A.M., Godsoe, W., van Hamelsveld, S., and Heinemann, J.A. (2018) Agrichemicals and antibiotics in combination increase antibiotic resistance evolution. *PeerJ* **6**: e5801.
- Kurenbach, B., Marjoshi, D., Amábile-Cuevas, C.F., Ferguson, G.C., Godsoe, W., Gibson, P., et al. (2015) Sublethal exposure to commercial formulations of the herbicides dicamba, 2,4-dichlorophenoxyacetic acid, and glyphosate cause changes in antibiotic susceptibility in *Escherichia coli* and *Salmonella enterica* serovar typhimurium. *MBio* **6**: e00009–15.
- Leino, L., Tall, T., Helander, M., Saloniemi, I., Saikkonen, K., Ruuskanen, S., et al. (2020) Classification of the glyphosate target enzyme (5-enolpyruvylshikimate-3-phosphate synthase) for assessing sensitivity of organisms to the herbicide. *J Hazard Mater*: 124556.
- Light, S.H., Krishna, S.N., Minasov, G., and Anderson, W.F. (2016) An unusual cation-binding site and distinct domain-domain interactions distinguish class II Enolpyruvylshikimate-3-phosphate synthases. *Biochemistry* **55**: 1239–1245.
- Lira JM, Cicchillo RM, Nair SK. (2013) US20130217577A1 - Novel class of glyphosate resistance genes, USA.
- Liu, J., Gefen, O., Ronin, I., Bar-Meir, M., and Balaban, N.Q. (2020) Effect of tolerance on the evolution of antibiotic resistance under drug combinations. *Science* **367**: 200–204.
- McConkey, G.A. (1999) Targeting the shikimate pathway in the malaria parasite *Plasmodium falciparum*. *Antimicrob Agents Chemother* **43**: 175–177.
- Meseguer, M.A., Alvarez, A., Rejas, M.T., Sánchez, C., Pérez-Díaz, J.C., and Baquero, F. (2003) Mycoplasma pneumoniae: a reduced-genome intracellular bacterial pathogen. *Infect Genet Evol* **3**: 47–55.
- Mesnage R, Teixeira M, Mandrioli D, Falcioni L, Ducarmon QR, Zwiittink RD, et al. (2019) Shotgun metagenomics and metabolomics reveal glyphosate alters the gut microbiome of Sprague-Dawley rats by inhibiting the shikimate pathway.
- Motta, E.V.S., Raymann, K., and Moran, N.A. (2018) Glyphosate perturbs the gut microbiota of honey bees. *Proc Natl Acad Sci U S A* **115**: 10305–10310.
- Myers, J.P., Antoniou, M.N., Blumberg, B., Carroll, L., Colborn, T., Everett, L.G., et al. (2016) Concerns over use of glyphosate-based herbicides and risks associated with exposures: a consensus statement. *Environ Health* **15**: 19.
- Novichkov, P.S., Ratnere, I., Wolf, Y.I., Koonin, E.V., and Dubchak, I. (2009) ATGC: a database of orthologous genes from closely related prokaryotic genomes and a research platform for microevolution of prokaryotes. *Nucleic Acids Res* **37**: D448–D454.
- Novick, A., and Doolittle, W.F. (2020) Horizontal persistence and the complexity hypothesis. *Biol Philos* **35**: 2.
- Priestman, M.A., Funke, T., Singh, I.M., Crupper, S.S., and Schönbrunn, E. (2005) 5-Enolpyruvylshikimate-3-phosphate synthase from *Staphylococcus aureus* is insensitive to glyphosate. *FEBS Lett* **579**: 728–732.
- Puigbò, P., Lobkovsky, A.E., Kristensen, D.M., Wolf, Y.I., and Koonin, E.V. (2014) Genomes in turmoil: quantification of genome dynamics in prokaryote supergenomes. *BMC Biol* **12**: 66.
- Richards, T.A., Dacks, J.B., Campbell, S.A., Blanchard, J.L., Foster, P.G., McLeod, R., and Roberts, C.W. (2006) Evolutionary origins of the eukaryotic shikimate pathway: gene fusions, horizontal gene transfer, and endosymbiotic replacements. *Eukaryot Cell* **5**: 1517–1531.
- Ruuskanen, S., Rainio, M.J., Gómez-Gallego, C., Selenius, O., Salminen, S., Collado, M.C., et al. (2020) Glyphosate-based herbicides influence antioxidants, reproductive hormones and gut microbiome but not reproduction: a long-term experiment in an avian model. *Environ Pollut* **266**: 115108.
- Schönbrunn, E., Eschenburg, S., Shuttleworth, W.A., Schloss, J.V., Amrhein, N., Evans, J.N., et al. (2001) Interaction of the herbicide glyphosate with its target enzyme 5-enolpyruvylshikimate 3-phosphate synthase in atomic detail. *Proc Natl Acad Sci U S A* **98**: 1376–1380.
- Shehata, A.A., Schrödl, W., Aldin, A.A., Hafez, H.M., and Krüger, M. (2013) The effect of glyphosate on potential pathogens and beneficial members of poultry microbiota in vitro. *Curr Microbiol* **66**: 350–358.
- Staub, J.M., Brand, L., Tran, M., Kong, Y., and Rogers, S.G. (2012) Bacterial glyphosate resistance conferred by over-expression of an *E. coli* membrane efflux transporter. *J Ind Microbiol Biotechnol* **39**: 641–647.
- Steinrücken, H.C., and Amrhein, N. (1980) The herbicide glyphosate is a potent inhibitor of 5-enolpyruvylshikimate acid-3-phosphate synthase. *Biochem Biophys Res Commun* **94**: 1207–1212.
- Woodburn, A.T. (2000) Glyphosate: production, pricing and use worldwide. *Pest Manag Sci* **56**: 309–312.
- Zomorodipour, A., and Andersson, S.G. (1999) Obligate intracellular parasites: *Rickettsia prowazekii* and *Chlamydia trachomatis*. *FEBS Lett* **452**: 11–15.

Supporting Information

Additional Supporting Information may be found in the online version of this article at the publisher's web-site:

Appendix S1: Supplementary Information