

RESEARCH OUTPUTS / RÉSULTATS DE RECHERCHE

Screening and comparative study of in vitro antioxidant and antimicrobial activities of ethanolic extracts of selected Vietnamese plants

Le Anh Dao, Nguyen; Phu, Tran Minh; Douny, Caroline; Quetin-Leclercq, Joëlle; Hue, Bui Thi Buu; Bach, Le Thi; Quynh Nhu, Truong; Thi Bich Hang, Bui; Thi Thanh Huong, Do; Thanh Phuong, Nguyen; Kestemont, Patrick; Scippo, Marie Louise

Published in:

International Journal of Food Properties

DOI:

[10.1080/10942912.2020.1737541](https://doi.org/10.1080/10942912.2020.1737541)

Publication date:

2020

Document Version

Publisher's PDF, also known as Version of record

[Link to publication](#)

Citation for published version (HARVARD):

Le Anh Dao, N, Phu, TM, Douny, C, Quetin-Leclercq, J, Hue, BTB, Bach, LT, Quynh Nhu, T, Thi Bich Hang, B, Thi Thanh Huong, D, Thanh Phuong, N, Kestemont, P & Scippo, ML 2020, 'Screening and comparative study of in vitro antioxidant and antimicrobial activities of ethanolic extracts of selected Vietnamese plants', *International Journal of Food Properties*, vol. 23, no. 1, pp. 481-496. <https://doi.org/10.1080/10942912.2020.1737541>


General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



Screening and comparative study of *in vitro* antioxidant and antimicrobial activities of ethanolic extracts of selected Vietnamese plants

Nguyen Le Anh Dao, Tran Minh Phu, Caroline Douny, Joëlle Quetin-Leclercq, Bui Thi Buu Hue, Le Thi Bach, Truong Quynh Nhu, Bui Thi Bich Hang, Do Thi Thanh Huong, Nguyen Thanh Phuong, Patrick Kestemont & Marie-Louise Scippo

To cite this article: Nguyen Le Anh Dao, Tran Minh Phu, Caroline Douny, Joëlle Quetin-Leclercq, Bui Thi Buu Hue, Le Thi Bach, Truong Quynh Nhu, Bui Thi Bich Hang, Do Thi Thanh Huong, Nguyen Thanh Phuong, Patrick Kestemont & Marie-Louise Scippo (2020) Screening and comparative study of *in vitro* antioxidant and antimicrobial activities of ethanolic extracts of selected Vietnamese plants, International Journal of Food Properties, 23:1, 481-496, DOI: [10.1080/10942912.2020.1737541](https://doi.org/10.1080/10942912.2020.1737541)

To link to this article: <https://doi.org/10.1080/10942912.2020.1737541>



© 2020 Nguyen Le Anh Dao, Tran Minh Phu, Caroline Douny, Joëlle Quetin-Leclercq, Bui Thi Buu Hue, Le Thi Bach, Truong Quynh Nhu, Bui Thi Bich Hang, Do Thi Thanh Huong, Nguyen Thanh Phuong, Patrick Kestemont and Marie-Louise Scippo. Published with license by Taylor & Francis Group, LLC.



Published online: 12 Mar 2020.



Submit your article to this journal [↗](#)



Article views: 2527



View related articles [↗](#)





View Crossmark data [↗](#)



Citing articles: 8 View citing articles [↗](#)

Screening and comparative study of *in vitro* antioxidant and antimicrobial activities of ethanolic extracts of selected Vietnamese plants

Nguyen Le Anh Dao ^{a,b}, Tran Minh Phu^a, Caroline Douny^b, Joëlle Quetin-Leclercq^c, Bui Thi Buu Hue^d, Le Thi Bach^d, Truong Quynh Nhu^a, Bui Thi Bich Hang^a, Do Thi Thanh Huong^a, Nguyen Thanh Phuong^a, Patrick Kestemont^e, and Marie-Louise Scippo ^b

^aCollege of Aquaculture and Fisheries, Can Tho University, Can Tho City, Vietnam; ^bFARAH – Veterinary Public Health, Laboratory of Food Analysis, University of Liège, Liege, Belgium; ^cLouvain Drug Research Institute, Pharmacognosy Research Group, Université Catholique De Louvain, Brussels, Belgium; ^dCollege of Natural Sciences, Can Tho University, Can Tho City, Vietnam; ^eResearch Unit in Environmental and Evolutionary Biology, Research Institute of Life, Earth & Environment (ILEE), University of Namur, Namur, Belgium

ABSTRACT

This study aimed to screen the *in vitro* antioxidant and antimicrobial activities of ethanolic extracts from 20 plants and three herbal commercial products empirically used for aquaculture improvement in Vietnam. The results of 2,2-diphenyl-1-picrylhydrazyl (DPPH) radical scavenging assays showed that *Phyllanthus amarus* extract was the strongest antioxidant, followed by four extracts in the subsequent order: *Piper betle* > *Psidium guajava* > *Euphorbia hirta* > *Mimosa pudica*. These five plant extracts were very active in a DPPH radical scavenging assay with concentrations needed to scavenge half of the DPPH (IC₅₀) below 30 µg/mL. Seven plant extracts showed an IC₅₀ ranging from 31.9 to 59.7 µg/mL, while eleven extracts showed an IC₅₀ above 70 µg/mL. A positive association was found between phenolic content (expressed as gallic acid equivalents) and antioxidant activity of the plant extracts. Concerning *in vitro* antimicrobial activities, *P. amarus* extract showed the highest activity against two different strains of *Aeromonas hydrophila* as demonstrated by its low minimal inhibitory concentration (MIC) of 156 and 625 µg/mL, respectively; whereas, *P. betle* displayed a moderate activity against *Edwardsiella ictaluri* with a MIC value of 625 µg/mL. Tannins were observed as significant factors contributing to antioxidant and antimicrobial properties of the plant extracts tested.

ARTICLE HISTORY

Received 18 November 2019
Revised 22 February 2020
Accepted 28 February 2020



KEYWORDS

Phyllanthus amarus;
Aeromonas hydrophila;
Edwardsiella ictaluri; tannins;
Vietnam

Introduction

Medicinal therapies using plant extracts expanded popularly in the late 1990 s.^[1] Beside the Chinese herbs, during the last decades, Vietnamese medicinal plants have also received interest as novel sources of alternative medication. It is estimated that approximately 2500 species of the Southeast Asian tropical exotic herbals have been used in folk medicine for their biological/therapeutic properties,^[2,3] such as diuretic,^[4] antioxidant,^[5] cytotoxicity,^[6,7] and antimicrobial.^[8] Moreover, a variety of herbs and plants have been receiving enormous interest as alternatives to synthetic additives or preservatives in the food industry in general and in aquaculture products in particular.^[9]

Aquaculture in Vietnam has been growing remarkably in recent years, producing 3.84 million tons aquaculture products in 2017.^[10] Mekong Delta is the main fish production area in the

CONTACT Marie-Louise Scippo  mlscippo@uliege.be  Department of Food Sciences, FARAH-VPH (Fundamental and Applied Research for Animal & Health, Veterinary Public Health), (M-L. Scippo) University of Liège, Quartier Vallée 2, Avenue De Cureghem 10 (B43b), liege 4000, Belgium

© 2020 Nguyen Le Anh Dao, Tran Minh Phu, Caroline Douny, Joëlle Quetin-Leclercq, Bui Thi Buu Hue, Le Thi Bach, Truong Quynh Nhu, Bui Thi Bich Hang, Do Thi Thanh Huong, Nguyen Thanh Phuong, Patrick Kestemont and Marie-Louise Scippo. Published with license by Taylor & Francis Group, LLC. This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Southern part of Vietnam with a contribution of 70% to the national aquaculture production. Striped catfish (*Pangasianodon hypophthalmus*) is the dominating cultured species for export with an annual production of 1.25 million tons.^[11] However, *Aeromonas hydrophila* and *Edwardsiella ictaluri* are pathogenic bacteria causing major diseases in the striped catfish industry, including “bacillary necrosis of pangasius” (BNP)^[12] or motile aeromonad septicemia (MAS),^[13] which is often related to stressed or immunocompromised hosts.^[14]

Chemicals including antimicrobials have been extensively applied to control pathogens and water quality management.^[15,16] The imprudent use of antimicrobials to control bacterial disease in fish farming may lead to development of antibiotic resistant bacteria, residues of antimicrobials in fish products, environmental impacts, etc. The access to export markets is certainly warned by the risk of noncompliance to international food safety regulations and quality standards. The final outcome could be a decline of profits made by the aquaculture sector, which is one of the most important activity in Vietnam (especially in the Mekong Delta).

Nowadays, more environment friendly prophylactic and protective solutions are claimed and natural bio-active products are examined for enhancing the immune system and health status of cultivated animals.^[17,18] However, in spite of a great variety of wild plants allocated in the various eco-regions of Vietnam and the concern of aquaculture farmers in using alternatives to antibiotics, the use of natural products in aquaculture is not yet popular in the country.

In this study, 20 plants, which are locally available, inexpensive and have been empirically used by fish farmers in the Mekong Delta regions, have been selected for a literature review about their antioxidant and antimicrobial activities (in particular against *A. hydrophila* and *E. ictaluri*). Ethanolic extracts of these 20 selected plants have been tested *in vitro* testing for these activities (Table 1). Besides these 20 plants, three commercial products have been added (Table 2), coming from Vietnamese companies, which are supposed to be efficient in treating white feces syndrome, enteritis and hepatic disease in shrimp, as well as in regenerating liver tissue in fish.

Materials and methods

Chemicals and media

DPPH (2,2-diphenyl-1-picrylhydrazyl), (\pm)- α -tocopherol, Folin-Ciocalteu's reagent, gallic acid, resazurin and gentamicin were purchased from Sigma-Aldrich Chemical Co. (St. Louis, MO, USA). Brain Heart Infusion media (BHI), isosensitest media, buffer pepton water (BPW) and plate count agar (PCA) were obtained from Oxoid (Basingstoke, UK). All solvents and reagents used in the analysis were of analytical grade.

Plant extracts preparation

Twenty fresh plants (Table 1) were collected from various areas in Mekong Delta, Vietnam. The plants were authenticated by the Department of Biology, College of Natural Science, Can Tho University and compared to literature. All collected parts of plants were then washed to remove mud and dust; the rotten and damaged parts were also discarded. Samples were air dried in shade for some days and then put in an oven at about 60°C until well-dried. After that, they were ground into a fine powder with a blender and stored in sealed containers in a dry and cool place. The dried-powder (100 g) was soaked in ethanol 96% (800 mL) for at least 24 hours at room temperature with frequent agitation. The solvent-containing extracts were then decanted and filtered. The ground samples were further extracted 4 times with ethanol 96%. The filtrates from each extraction were combined and the solvent was evaporated under reduced pressure using a rotary evaporator to give crude ethanolic extracts.

Three commercial products, named A, B, C were purchased from purchasers willing to stay anonymous. However, they gave us as a personal communication the plant composition of their



Table 1. List of 20 selected plants used for the screening of their antioxidant and antimicrobial activities.

| Identification number* | Scientific name | Common name | Ecology | Distributed in Vietnam | Level of threats in the wild | | Status of species | Part used | Reference |
|------------------------|--|--|---|--|------------------------------|-------------|-------------------------------|------------|-----------|
| | | | | | Low | Invasive | | | |
| CTU1731 | <i>Ageratum conyzoides</i> | Billygoat-weed, chick weed | Grow and develop on different soils; from lowland to highland 1800 m | Widely grown in Vietnam | Low | Invasive | The whole plant | [19–21] | |
| CTU1922 | <i>Allium sativum</i> | Garlic | Spice herb | Widely grown in Vietnam | No | Native | Bulb | [19–21] | |
| CTU1716 | <i>Alternanthera sessilis</i> | Sessile joyweed and dwarf copperleaf | Uncultivated growth in the garden, paths at edge of ricefield, edge of canal | Widely grown in Vietnam (at height 1000 m) | Low | Native | Aerial parts (leaves + twigs) | [19–21] | |
| CTU1720 | <i>Annona reticulata</i> | Custard apple; bullock's heart; raamphal plant | Aquatic growth, edge of canal (alum tolerant tree) | Widely grown in Mekong Delta | Low | Native | Leaves | [20–22] | |
| CTU1608 | <i>Azadirachta indica</i> | Neem | Grow in the forest, leaves harvested as vegetable | Forest: Ninh Thuận, An Giang, Kiên Giang; cultivated in the South of Vietnam | Low | Native | Leaves, flower and stem bark | [19–21] | |
| CTU17161 | <i>Cayratia trifolia</i> | Fox grape | Scattered growth in secondary forest, shrub, garden | Widely grown in Vietnam | Low | Native | The whole plant | [20,21,23] | |
| CTU1823 | <i>Centella asiatica</i> | Asiatic pennywort, Indian pennywort | Grow in clusters at wild land, paths at edge of ricefield, roadside | Widely grown in Vietnam | Low | Native | The whole plant | [20,21,23] | |
| CTU1836 | <i>Eclipta prostrata/ Eclipta alba</i> | False daisy | Grow in paths at edge of ricefield, roadside, from lowland to highland 1800 m | Widely grown in Vietnam | Low | Native | The whole plant | [19–21] | |
| CTU1874 | <i>Euphorbia hirta</i> | Asthma-plant | Photophilic herb, grow in wild land, paths at edge of ricefield | Widely grown in Vietnam | Low | Native | The whole plant | [20,21,23] | |
| CTU17174 | <i>Houttuynia cordata</i> | Fish mint | Grow in clusters at wild land, paths at edge of ricefield, edge of canal | Uncultivated and widely grown in Vietnam | Low | Native | Leaves + stem | [21,22,24] | |
| CTU17113 | <i>Mimosa pudica</i> | Sensitive plant, sleepy plant, shy plant | Grow in roadside, wild land, sea wall | Widely grown in Vietnam (exotic species) | Medium | Introduced* | The whole plant | [21,22,24] | |
| CTU1871 | <i>Momordica charantia</i> L. | Bitter melon, bitter squash | Uncultivated growth in secondary forest or cultivated in farms | Widely grown in Vietnam (as vegetable) | Low | Native | Leaves, stem | [21,22,24] | |
| CTU1896 | <i>Ocimum basilicum</i> | Basil | Photophilic herb | Widely grown in Vietnam (as spice herb) | No | Native | Aerial parts (leaves + twigs) | [21,23,24] | |
| CTU1899 | <i>Perilla frutescen</i> | Perilla mint | Photophilic herb | Widely grown in Vietnam (as spice herb) | No | Native | Leaves | [21,23,24] | |
| CTU1778 | <i>Phyllanthus amarus</i> | Chamber bitter, gripe-weed, shatterstone, stone-breaker or leaf-flower | Uncultivated growth in paths of grass, paths at edge of ricefield or mountain field, roadside | Widely grown in Vietnam | Low | Native | Aerial parts (leaves + twigs) | [21,23,24] | |
| CTU1623 | <i>Piper betle</i> | Betel | Planted in home gardens | Widely grown in Vietnam | No | Native | Leaves | [21,22,24] | |

(Continued)



Table 1. (Continued).

| Identification number* | Scientific name | Common name | Ecology | Distributed in Vietnam | Level of threats in the wild | Status of species | Part used | Reference |
|------------------------|----------------------------------|-------------|--|--|------------------------------|-------------------|-----------------|------------|
| CTU17137 | <i>Portulaca oleracea</i> | Purslane | Scattered growth or in clusters at wild land, edges of ricefield | Widely grown in Vietnam | Low | Native | The whole plant | [19,21,24] |
| CTU17125 | <i>Psidium guajava</i> | Apple guava | Planted in farms | Widely grown in Vietnam | Low | Introduced* | Leaves | [21,23,24] |
| CTU1644 | <i>Wedelia chinensis</i> | - | Uncultivated grown in paths of grass | Widely grown in the Central and North of Vietnam | Low | Native | Leaves | [19,21,24] |
| CTU1898 | <i>Zingiber officinale</i> Rosc. | Ginger | Uncultivated grown in forest | Widely grown in Vietnam | Low invasion | Native | Rhizome | [19,21,24] |

*Species identification was done in/by Can Tho University (CTU), Vietnam.

Introduced*: this introduced species was perennially planted in Mekong Delta, Vietnam.

Table 2. List of 3 commercial products used for the screening of their antioxidant and antimicrobial activities.

| Product name | Composition* | Form |
|--------------|---|--------|
| A | <i>Blumea balsamifera</i> | Solid |
| B | <i>Cynara cardunculus</i> var. <i>scolymus</i> <i>Phyllanthus emblica</i> | Liquid |
| C | <i>Terminalia arjuna</i> <i>Phyllanthus urinaria</i> <i>Eclipta prostrata</i> | Solid |

*Personal communication from the purchasers who asked to remain anonymous

product (Table 2). For both commercial powder products, a 1 g dry sample was continuously shaken in 30 mL ethanol (96%) for 24 hours. After filtration, the filtrates obtained were evaporated to dryness as explained above. The liquid commercial product was lyophilized. All ethanolic extracts were finally lyophilized until dryness to remove any trace of water and stored in a dry place at room temperature before use.

Total phenolic compounds determination

The amount of total phenolic compounds (TPC) was determined using Folin-Ciocalteu reagent.^[25] This assay was performed in triplicate for all plant extracts, dissolved in methanol. A standard curve was established using gallic acid as standard, in the range of 0–10 µg/mL (in methanol). The amount of total phenolic compounds in plant extracts was calculated as gallic acid equivalents (GAE) in mg of GAE per 100 mg (%) of freeze-dried plant material.

Antioxidant capacity

Antioxidant capacity of plant extracts was measured through DPPH (2,2'-diphenyl-1-picrylhydrazyl) radical scavenging assay.^[26] The assay was performed in triplicate for each plant extract sample and α -tocopherol was used as a positive control. Plant extracts dissolved in methanol were tested for their DPPH scavenging activity in concentrations ranging from 1 to 125 µg/mL. Plant extract concentrations were plotted against percentages of remaining DPPH after 30 minutes reaction. A sigmoidal curve was then fitted, allowing the determination of the IC₅₀, i.e. the plant extract concentration (in µg/mL) needed to scavenge 50% of the DPPH initially introduced.

In vitro investigation of antimicrobial activities of plant extracts

Resazurin method was performed as described by Sarker et al.^[27] with some adaptations for minimum inhibitory concentration (MIC) determination of plant extracts against *A. hydrophila* and *E. ictaluri*. Stock solutions of the plant extracts were prepared at a concentration of 10 mg/mL in sterilized normal saline containing 25% dimethyl sulfoxide (DMSO) (when testing *A. hydrophila*) or 12.5% DMSO (when testing *E. ictaluri*). The resazurin and the antibiotic gentamicin solutions were prepared in sterile distilled water at the concentration of 10 mg/mL and 200 µg/mL, respectively. Two strains of *Aeromonas hydrophila* (1 and 2) were isolated from red tilapia (*Oreochromis* sp.), whereas, *Edwardsella ictaluri* was isolated from striped catfish (*P. hypophthalmus*) in Mekong Delta, Vietnam. These strains were subcultured in BHI broth, incubated for 24 h at 37°C (*A. hydrophila*) and for 48 h at 30°C (*E. ictaluri*), and then pelleted in 20 mL of sterile normal saline solution. The approximate number of bacteria were estimated by the optical density at 600 nm, reaching a value of 4×10^8 and 1×10^8 cfu/mL for *A. hydrophila* and *E. ictaluri*, respectively according to the turbidity of McFarland Standards (e.g., Dalynn Biologicals, cat no. TM50-TM60). Further dilutions were performed to use final concentrations of 2×10^4 in double strength isosensitest broth and 0.5×10^4

cfu/mL in BHI broth for *A. hydrophila* and *E. ictaluri*, respectively. One mg of resazurin dissolved in water was then added into 10 mL of the liquid inoculum.

The MIC determination was performed in a 96-well plate, using concentrations of plant extracts ranging from 5 µg/mL to 2500 µg/mL and 100 µL of working inoculum solution per well. The plates were prepared in triplicate and gentamicin was used as positive control in each plate. After incubation at 37°C for 18 h (*A. hydrophila*) or 48 h (*E. ictaluri*), plates were visually observed to determine the lowest concentration at which color changed from purple to pink, which was taken as the MIC value.^[28]

Tannins quantification and removal from plant extracts

The extracts of *P. amarus* and *E. hirta* were prepared at 10 mg/mL in methanol. Tannins were quantified according to the European pharmacopoeia method and the results were expressed in pyrogallol equivalents (g/100 g extract).^[29] Tannins were excluded from methanolic plant extracts by using a polyamide column (Macherey – Nagel, Germany), as described by Houghton and Raman.^[30] Ten grams of polyamide were soaked in 100 mL of water during one night. A column was packed with the gel. Ten milligrams of plant extract dissolved in 1 mL of methanol were loaded onto the column. A volume of 200 mL of methanol was gradually poured into the column for the elution. An empty dried flask was prepared to collect the eluates, which were evaporated to dryness. Tannins were retained on the column while the dried residue contained the extract free from tannins.

Statistical analysis

Data of total phenolic content and IC₅₀ values were expressed as mean ± standard deviation by Microsoft Excel software. Analysis of variance (one-way ANOVA) was performed by using SPSS 16.0 (SPSS Inc, Chicago, IL, USA).

Results and discussion

Total phenolic content and antioxidant activity of 23 plant extracts

The content of phenolic compounds determined in the 23 plant extracts samples ranged between 0.5 and 18.8 mg gallic acid equivalent/100 mg plant extract (Table 3). Among the first five plant extracts mentioned in Table 3, considered as having a “high” antioxidant activity (see below), *P. amarus* showed the significantly ($p < .05$) highest total phenolic content. Interestingly, the commercial products contained lower levels of phenolic compounds than these 5 plant extracts.

The antioxidant activity measured using the DPPH assay was expressed as an IC₅₀, corresponding to the concentration of plant extract needed to scavenge half of the DPPH. According to Thiangthum et al.^[26], the antioxidant activity of samples can be classified as high (IC₅₀ < 30 µg/mL), intermediate (30 < IC₅₀ < 50 µg/mL), low (50 < IC₅₀ < 70 µg/mL) or absent (IC₅₀ > 70 µg/mL). In our assay, the IC₅₀ of the (±)- α -tocopherol, the reference antioxidant, was 12 µg/mL. Eleven out of the 23 extracts displayed an antioxidant activity, with concentrations able to inhibit half of the maximum response (IC₅₀) ranging from 5.83 to 49.5 µg/mL (Table 3). *P. amarus* extract showed the apparent strongest radical scavenging effect (IC₅₀ = 5.83 µg/mL) but no significant difference ($p > .05$) was observed with the antioxidant activities of *P. betle* and *P. guajava* extracts. The remaining 2 samples showing a high antioxidant activity (i.e. IC₅₀ < 30 µg/mL) were *E. hirta* and *M. pudica*. A group of six samples, including *Z. officinale*, commercial product B, *E. prostrata*, commercial product A, *A. reticulata*, *H. cordata*, showed an intermediate antioxidant capacity (i.e. 30 µg/mL < IC₅₀ < 50 µg/mL).

For the 23 selected plant extracts, the antioxidant activity was related with their phenolic content with a correlation coefficient $R^2 = 0.9137$ (Figure 1). The antioxidant properties of plant extracts might be influenced by the number of specific chemical groups from phenolic compounds, such as

Table 3. Total phenolic content and results from the DPPH radical scavenging assay (IC_{50}) of 23 plant extracts (the plants are shown by decreasing antioxidant capacity, i.e. increasing IC_{50}), and comparison with data from the literature.

| Scientific name | This study | | Literature studies | | |
|-------------------------------|-------------------------------|--------------------------------|-----------------------|--------------------------------|-----------|
| | Phenolic content (%)* | IC_{50} ($\mu\text{g/mL}$) | Phenolic content (%)* | IC_{50} ($\mu\text{g/mL}$) | Reference |
| <i>Phyllanthus amarus</i> | 18.8 ^j \pm 0.75 | 5.83 ^a \pm 0.50 | 17 | 11 | [26,31] |
| <i>Piper betle</i> | 16.4 ⁱ \pm 0.76 | 8.32 ^a \pm 0.90 | 15.4 | 11 | [32] |
| <i>Psidium guajava</i> | 14.5 ^h \pm 0.79 | 8.55 ^a \pm 0.53 | 15.6 | 1.56 | [33] |
| <i>Euphorbia hirta</i> | 10.3 ^g \pm 0.91 | 17.8 ^b \pm 1.28 | 29.1 | 2.81 | [34] |
| <i>Mimosa pudica</i> | 7.46 ^f \pm 0.29 | 18.0 ^b \pm 0.96 | 4.3 | 127 | [35] |
| <i>Zingiber officinale</i> | 6.92 ^e \pm 0.31 | 31.9 ^c \pm 2.48 | 1 | 46.5 | [36] |
| Commercial product B | 0.53 ^a \pm 0.04 | 33.7 ^{cd} \pm 0.34 | - | - | - |
| <i>Eclipta prostrata</i> | 4.57 ^d \pm 0.25 | 36.6 ^{cd} \pm 0.86 | 13.2 | 42 | [37] |
| Commercial product A | 5.65 ^e \pm 0.20 | 38.1 ^d \pm 2.55 | - | - | - |
| <i>Annona reticulata</i> | 5.01 ^{de} \pm 0.24 | 45.1 ^e \pm 4.58 | 13.6 | 51 | [38] |
| <i>Houttuynia cordata</i> | 5.45 ^e \pm 0.28 | 49.5 ^e \pm 5.83 | 12.6 | 73 | [39] |
| <i>Cayratia trifolia</i> | 3.31 ^c \pm 0.22 | 59.7 ^f \pm 5.11 | 7.3 | 74 | [40,41] |
| <i>Perilla frutescens</i> | 3.61 ^c \pm 0.21 | 73.7 ^g \pm 2.09 | 11.6 | 7.97 | [42] |
| <i>Azadirachta indica</i> | 4.33 ^d \pm 0.39 | 79.2 ^h \pm 3.93 | 10.1 | 60.6 | [43] |
| Commercial product C | 2.04 ^b \pm 0.17 | 85.9 ⁱ \pm 1.95 | - | - | - |
| <i>Ageratum conyzoides</i> | 2.05 ^b \pm 0.07 | 118 ^j \pm 1.92 | 0.85 | 214 | [44] |
| <i>Portulaca oleracea</i> | ND | > 125 | 0.43 | 2950 | [45] |
| <i>Allium sativum</i> | ND | > 125 | 0.005 | 600 | [46] |
| <i>Ocimum basilicum</i> | 2.25 ^b \pm 0.26 | > 125 | 0.07 | 350 | [47] |
| <i>Centella asiatica</i> | 2.27 ^b \pm 0.09 | > 125 | 2.4 | 45 | [48] |
| <i>Wedelia chinensis</i> | 1.66 ^b \pm 0.02 | > 125 | - | 45 | [49] |
| <i>Momordica charantia</i> | ND | > 125 | 1.02 | 307 | [50] |
| <i>Alternanthera sessilis</i> | 1.73 ^b \pm 0.11 | > 125 | 3.7 | 946 | [51,52] |

*mg gallic acid equivalent/100 mg dry plant extract; ND = Not determined; "-": no information available.

High antioxidant activity: $IC_{50} < 30 \mu\text{g/mL}$, Intermediate antioxidant activity: $30 < IC_{50} < 50 \mu\text{g/mL}$, No activity: $IC_{50} > 70 \mu\text{g/mL}$. Values are mean \pm SD (n = 3). Mean values within a column with the same letter are not significantly different ($p > 0.05$).




Figure 1. Correlation between phenolic compounds contents (expressed as mg GAE/100 mg extract) and antioxidant activity measured in the DPPH assay (expressed as $1/IC_{50}$, μM^{-1}) of the 23 selected extracts.

hydroxyl or methoxy groups, keto or free carboxylic groups, as well as by other antioxidant secondary metabolites, such as vitamins, volatile oils and carotenoids.^[53] In the present study, the highest antioxidant activity was found for the *P. amarus* extract. This result agrees with the previous study of Thiangthum et al.^[26] reporting that *P. amarus* extracts showed high antioxidant activity as

shown by low IC_{50} values in a DPPH assay, ranging from 10 to 16 $\mu\text{g/mL}$. Regi Raphael et al.^[54] also reported that the extract of *P. amarus* was found as a remarkable antioxidant throughout its inhibition capacity in scavenging free radicals *in vitro*, while other studies confirmed that the high content of phenolic compounds in this plant was positively correlated with its radical scavenging potential.^[55]

Many investigations were conducted concerning the chemical components of this *Phyllanthus* species and bioactive constituents. According to Igwe et al.^[56], extracts from leaves of *P. amarus* (which were the part of the plant used in this study) contained high level of saponins and tannins: 24 and 17%, respectively. In the study of Sharma et al.^[57], the leaves of *P. amarus* were shown to contain a high quantity of lignans (phyllanthin (0.7%) and hypophyllanthin (0.3%)) in comparison with other parts of the plant. These active compounds might be responsible for the major part of the antioxidant activity.

The good relationship found in this study between total phenolic contents and DPPH measurements is also in agreement with a study of Dudonne et al.^[58] who showed a high positive correlation between the free radical scavenging and total phenolic content of 30 plant extracts. For both total phenolic content and IC_{50} found in the DPPH assay, Table 3 compares the results of this study and data found in the literature. For most of the tested plants, the results are roughly similar except for two plants giving opposite results. In this study, the ethanolic extract from the whole *Mimosa pudica* displayed a high antioxidant activity (IC_{50} of 21 $\mu\text{g/mL}$) while Das et al.^[35] reported an IC_{50} of 127 $\mu\text{g/mL}$ for leaves methanolic extract of the same plant, which would mean no antioxidant capacity (Table 3). On the contrary, for *Perilla frutescens*, the ethanolic extract from its leaves showed a low antioxidant capacity in this study (IC_{50} of 70 $\mu\text{g/mL}$), but a high one in Lin et al.^[42] (IC_{50} of 8 $\mu\text{g/mL}$).

Surprisingly, the antioxidant capacity determined in two (A and B) out of the three tested commercial products was shown to be intermediate only and was lower than the one of the 5 plant extracts mentioned here above, while the third commercial product (C) was qualified as non active. Plants used as the main ingredients in each product are herbs that have been acknowledged since ancient times for their valuable and therapeutic efficiency. Extracts of *Blumea balsamifera* (contained in the commercial product A), *Cynara cardunculus* var. *scolymus*, *Phyllanthus emblica*, *Terminalia arjuna* (contained in the commercial product B), or *Phyllanthus urinaria* and *Eclipta prostrata* (contained in the commercial product C) revealed different activities in the DPPH assay with IC_{50} values of 72 $\mu\text{g/mL}$,^[59] 23 $\mu\text{g/mL}$,^[60] 11 $\mu\text{g/mL}$,^[61] 8 $\mu\text{g/mL}$,^[62] 17 $\mu\text{g/mL}$,^[63] and 42 $\mu\text{g/mL}$,^[37] respectively. However, it is well known that many factors of manufacture (such as the materials, the extraction procedure of plant extracts, the ingredients, etc.) can affect the product's activity. Moreover, Bruno and Munro^[64] determined that the formulation of commercial products frequently include constituents other than the active ingredient(s), which can be stabilizers, carriers or diluent agents.^[65]

Antimicrobial activities of plant extracts against *A. hydrophila* and *E. ictaluri*

The *in vitro* antimicrobial activities of the 23 samples against two *A. hydrophila* strains isolated from red tilapia (*Oreochromis* sp.) in Vietnam are presented in Table 4, which also presents the antibacterial activities of the selected plants reported in the literature. According to Kuete^[82], antimicrobial activities can be classified as weak or absent if MIC are above 1250 or 2500 $\mu\text{g/mL}$, respectively. For MIC between 156 and 625 $\mu\text{g/mL}$, the antimicrobial activity was qualified as moderate. Four plant extracts (*P. amarus*, *P. betle*, *P. guajava* and *E. hirta*) and one commercial product (A) showed moderate antimicrobial activities against the first strain of *A. hydrophila* (1) (Table 4), with MICs values ranging between 156 and 625 $\mu\text{g/mL}$, while the rest of the samples showed weak or no antibacterial activity (MIC \geq 1250 or 2500 $\mu\text{g/mL}$, respectively). The growth of the first strain of *A. hydrophila* (1) was also inhibited by *P. amarus*, *P. betle* and commercial product A, but at higher concentrations of plant extracts (MIC = 625 $\mu\text{g/mL}$). *P. guajava* and *E. hirta*, which



Table 4. *In vitro* antibacterial activity of 23 plant extracts against two strains of *A. hydrophila* (1 and 2) and *E. ictaluri*, and other antibacterial activities from literature studies.

| Scientific name | MIC ($\mu\text{g/ml}$) | | | MIC against other bacteria, from literature |
|-------------------------------|--------------------------|------------------------|--------------------|---|
| | <i>A. hydrophila</i> 1 | <i>A. hydrophila</i> 2 | <i>E. ictaluri</i> | |
| <i>Phyllanthus amarus</i> | 156 | 625 | >2500 | <i>Staphylococcus aureus</i> (100 $\mu\text{g/ml}$), <i>Streptococcus pneumoniae</i> (400 mg/ml), <i>Shigella</i> spp. (25 $\mu\text{g/ml}$), <i>E. coli</i> (50 $\mu\text{g/ml}$) ^[66] |
| <i>Piper betle</i> | 156 | 625 | 625 | <i>S. aureus</i> (1000 $\mu\text{g/ml}$), <i>Propionibacterium acnes</i> (4000 $\mu\text{g/ml}$) ^[67] |
| <i>Psidium guajava</i> | 312 | 1250 | >2500 | <i>S. mutans</i> (250 $\mu\text{g/ml}$), <i>S. mitis</i> (250 $\mu\text{g/ml}$), <i>S. oralis</i> (250 $\mu\text{g/ml}$) ^[68] |
| Commercial product A | 312 | 625 | 625 | - |
| <i>Euphorbia hirta</i> | 625 | 1250 | >2500 | <i>S. aureus</i> (25 mg/ml), <i>Candida albicans</i> (12.5 mg/ml) ^[69] |
| <i>Mimosa pudica</i> | 1250 | 2500 | >2500 | <i>Escherichia coli</i> (250 mg/ml), <i>S. aureus</i> (250 mg/ml), <i>Bacillus subtilis</i> (200 mg/ml) ^[70] |
| <i>Eclipta prostrata</i> | 1250 | 1250 | >2500 | <i>E. coli</i> (12.5 mg/ml), <i>S. aureus</i> (3.125 mg/ml), <i>B. subtilis</i> (6.25 mg/ml), <i>B. cereus</i> (1.56 mg/ml) ^[71] |
| Commercial product B | >2500 | >2500 | >2500 | - |
| <i>Zingiber officinale</i> | 2500 | 2500 | >2500 | <i>Pseudomonas aeruginosa</i> (40 mg/ml), <i>E. coli</i> (40 mg/ml), <i>S. aureus</i> (20 mg/ml) ^[72] |
| <i>Annona reticulata</i> | 2500 | 2500 | 2500 | <i>E. coli</i> (30 $\mu\text{g/ml}$), <i>S. aureus</i> (40 $\mu\text{g/ml}$), <i>B. subtilis</i> (10 $\mu\text{g/ml}$) ^[73] |
| <i>Houttuynia cordata</i> | 2500 | 2500 | >2500 | <i>Bacillus dysenteriae</i> (0.08 mg/ml) ^[74] |
| <i>Cayratia trifolia</i> | 2500 | 2500 | >2500 | - |
| <i>Perilla frutescens</i> | >2500 | >2500 | >2500 | - |
| <i>Azadirachta indica</i> | >2500 | >2500 | >2500 | <i>E. coli</i> (0.781 mg/ml), <i>K. pneumoniae</i> (1.562 mg/ml), <i>E. faecalis</i> (3.125 mg/ml), <i>S. aureus</i> (1.562 mg/ml), <i>P. aeruginosa</i> (1.562 mg/ml) ^[43] |
| Commercial product C | 2500 | 2500 | >2500 | - |
| <i>Ageratum conyzoides</i> | >2500 | >2500 | >2500 | <i>E. coli</i> (100 $\mu\text{g/ml}$), <i>S. aureus</i> (200 $\mu\text{g/ml}$) ^[75] |
| <i>Portulaca oleracea</i> | >2500 | >2500 | >2500 | <i>S. aureus</i> (12.5 mg/ml), <i>Streptococcus pyogenes</i> (12.5 mg/ml), <i>P. aeruginosa</i> (50 mg/ml), <i>E. coli</i> (50 mg/ml) ^[76] |
| <i>Allium sativum</i> | 2500 | 2500 | >2500 | <i>E. coli</i> (150 $\mu\text{g/ml}$), <i>Klebsiella pneumoniae</i> (150 $\mu\text{g/ml}$), <i>B. subtilis</i> (100 $\mu\text{g/ml}$) ^[77] |
| <i>Ocimum basilicum</i> | >2500 | >2500 | >2500 | <i>Bacillus cereus</i> (62.5 $\mu\text{g/ml}$), <i>B. subtilis</i> (125 $\mu\text{g/ml}$), <i>S. aureus</i> (62.5 mg/ml), <i>E. coli</i> (125 $\mu\text{g/ml}$), <i>Salmonella typhi</i> (500 $\mu\text{g/ml}$) ^[78] |
| <i>Centella asiatica</i> | 2500 | 2500 | >2500 | <i>S. aureus</i> (8 mg/ml) ^[79] |
| <i>Wedelia chinensis</i> | >2500 | >2500 | >2500 | <i>B. subtilis</i> (6.25 mg/ml), <i>S. aureus</i> (6.25 mg/ml), <i>E. coli</i> (25 mg/ml) ^[80] |
| <i>Momordica charantia</i> | >2500 | >2500 | >2500 | <i>Enterococcus faecalis</i> (1.25 mg/ml), <i>E. coli</i> (5 mg/ml), <i>K. pneumoniae</i> (5 mg/ml) ^[81] |
| <i>Alternanthera sessilis</i> | >2500 | >2500 | >2500 | - |
| Gentamicin | 6.25 | 6.25 | 12.5 | - |

Significant activity: MIC < 100 $\mu\text{g/ml}$

Moderate activity: 100 < MIC \leq 625 $\mu\text{g/ml}$

Weak activity: MIC \geq 625 $\mu\text{g/ml}$

inhibited the growth of the first strain of *A. hydrophila* (1) at a concentration of 312 and 625 µg/mL respectively, needed a concentration of 1250 µg/mL to inhibit the second strain (2) (Table 4). The results showed that both strains of *A. hydrophila* display the same pattern of sensitivity toward tested plant extracts, but the first strain was more sensitive than the second one. The MIC of the gentamycin antibiotic, used as positive control, was 6.25 µg/mL for both strains (Table 4).

The antibacterial activities of the 23 samples against *Edwardsiella ictaluri* strain are summarized in Table 4. Results show that most plant extracts do not possess antimicrobial activities (MIC ≥ 2500 µg/mL). *Piper betle* and commercial product A showed a moderate activity, with MIC values of 625 µg/mL. However, the strain of *E. ictaluri* used in this work was not very sensitive to gentamicin (MIC = 12.5 µg/mL). In another study, where 64 different strains of *E. ictaluri* were tested for their sensitivity to antibiotics, gentamicin showed MIC of 2 µg/mL for 50% of the tested strains and 1 µg/mL (25%) or 4 µg/mL (25%) for the remaining strains.^[83]

Limited information is available about previous studies concerning the antimicrobial capacity of *P. amarus* and *P. betle* against *Aeromonas* spp. and *Edwardsiella* spp. For example, a methanolic extract of *P. amarus* showed an antibacterial activity against *A. hydrophila* with a MIC of 128 µg/mL according to De Britto et al.^[84], while, for the same bacteria, a MIC of 25 µg/mL was found by Caruso et al.^[85] for the ethanolic extract from leaves of *Piper betle*. The capacity of *P. amarus* to inhibit bacterial growth showed a concentration-dependent antibacterial activity particularly against gram-negative microbes,^[86,87] while Kaveti et al.^[88] showed that ethanolic extracts of *P. betle* leaves were active against several strains of gram positive and negative bacteria.

The findings of this study regarding a significant antibacterial activity of two plant extracts (*P. amarus* and *P. betle*) against two strains of the pathogenic bacteria *A. hydrophila* could be useful for the initial selection of natural alternative to antibiotics as well as to prevent bacterial growth in fish products during storage.

Impact of tannin removal on the antimicrobial and antioxidant activities of *P. amarus* and *E. hirta* ethanolic extracts

The medicinal usefulness of the *P. amarus* and *E. hirta* have been the subject of numerous chemical and microbiology studies. Guha et al.^[89] demonstrated that the extract of *P. amarus* is active in inhibiting lipid peroxidation and in scavenging hydroxyl and superoxide radicals. Moreover, Sheikhlar et al.^[83] investigated the antibacterial activity of several extracts of *E. hirta*, *Citrus lemon* and *Trigonella foenum-graecum*. Results showed that the methanolic extract of *E. hirta* exhibited the strongest antimicrobial activity with the lowest MIC (70 µg/mL) against *A. hydrophila* and their potential to be a beneficial dietary supplement for enhancing the resistance of *Clarius gariepinus* to *A. hydrophila* contamination. That is why *P. amarus* and *E. hirta* were chosen as the representative plants for further investigations. Previous screenings of bioactive compounds in *P. amarus* and *E. hirta* revealed that tannins may be responsible for their antioxidant and antimicrobial properties.^[90,91]

Tannins constitute a group of secondary metabolites which are widely distributed among vegetal species.^[92] They are found in approximately 80% of wooden and 15% of herbal dicotyledonous species. Tannins are known to be responsible for general antimicrobial and antioxidant activities.^[93,94] Up to now, investigations on the effects of tannin removals from crude plant extracts are limited. Additionally, the prediction of which class of compounds (alkaloids, anthraquinones, flavonoids, saponins and tannins) possess antioxidant and antimicrobial activities is a challenge.^[1,95,96] Hence, the comparison of the activities of two selected extracts before and after tannins removal on a polyamide column was performed to clarify whether tannins are the main contributors to their antioxidant and antimicrobial activities.

The results showed that, after tannin removal, both *P. amarus* and *E. hirta* extracts were less active against both strains of *A. hydrophila* (Table 5), while their inhibiting activity against *E. ictaluri* remained as weak as before tannin removal (Table 5). As expected, for both plant extracts, the total

Table 5. Antimicrobial activity of *Phyllanthus amarus* and *Euphorbia hirta* before and after tannins removal.

| | MIC ($\mu\text{g/mL}$) | | | |
|---------------------------|-------------------------------|----------|------------------------|----------|
| | Before removing tannins | | After removing tannins | |
| | <i>Aeromonas hydrophila</i> 1 | | | |
| <i>Phyllanthus amarus</i> | 156 | Moderate | 625 | Moderate |
| <i>Euphorbia hirta</i> | 625 | Moderate | 2500 | Weak |
| | <i>Aeromonas hydrophila</i> 2 | | | |
| <i>Phyllanthus amarus</i> | 625 | Moderate | >1250 | Weak |
| <i>Euphorbia hirta</i> | 1250 | Weak | 2500 | Weak |
| | <i>Edwardsiella ictaluri</i> | | | |
| <i>Phyllanthus amarus</i> | >2500 | Weak | >2500 | Weak |
| <i>Euphorbia hirta</i> | >2500 | Weak | >2500 | Weak |

Table 6. Antioxidant activities of *Phyllanthus amarus* and *Euphorbia hirta* before and after tannins removal.

| Name | Before removing tannins | | After removing tannins | |
|---------------------------|---|--------------------------|---|--------------------------|
| | DPPH assay IC ₅₀ ($\mu\text{g/mL}$) | Phenolic content (%*) | DPPH assay IC ₅₀ ($\mu\text{g/mL}$) | Phenolic content (%*) |
| <i>Phyllanthus amarus</i> | 5.83 \pm 0.50 | 18.8 \pm 0.75 | 83.6 \pm 5.38 | 2.68 \pm 0.20 |
| <i>Euphorbia hirta</i> | 17.8 \pm 1.28 | 10.3 \pm 0.91 | 38.3 \pm 2.40 | 5.02 \pm 0.54 |

Values are mean \pm SD (n = 3); * Expressed as mg GAE per 100 mg plant extract

phenolic content decreased after tannin removal while their IC₅₀ in the DPPH assay increased, meaning less antioxidant activity (Table 6). About half of the initial phenolic content remains for *E. hirta* extracts and only the seventh for *P. amarus*, showing that a high amount of tannins was present in these extracts. This apparent higher proportion of tannins in *P. amarus* was confirmed after quantification of tannins in both extracts according to the European pharmacopoeia method, which gave 5.6% for *P. amarus* and 1.79% for *E. hirta* (expressed in pyrogallol). Hydrolysable tannins appear to be the main polyphenol constituents found in *P. amarus*, with geraniin being the most abundant.^[97–99] This could explain the fact that *P. amarus* displayed high antioxidant and antimicrobial activities. This is in agreement with Catteau et al.^[100], reporting that tannins were at least in part responsible for the activities of several plant methanolic extract effects, but other compounds may also explain a part of the activity.

Patel et al.^[101] reported that high content of tannins could be isolated from *P. amarus* which are associated with some health importance and antimicrobial activity. According to Bensky et al.^[90], hydrolysable tannins are among the main secondary metabolites, which have been discovered to prevent viral DNA polymerase and reverse transcriptase in HIV infection, and to act on angiotensin-converting enzymes in diabetes complication. Moreover, antimicrobial activity of *E. hirta* was attributed to tannins and other bioactive components. Yoshida et al.^[91] and Yoshida et al.^[102,103] reported the isolation of hydrolysable dimeric ellagitannins (euphorbin A, B, C and E) from the leaves of the plant.

Conclusion

In our study, twenty three ethanolic extracts from 20 plant and 3 commercial products obtained in Mekong Delta (Vietnam) were screened for their *in vitro* antioxidant and antimicrobial activities. The results from DPPH free radical scavenging assays revealed that the *P. amarus* extract showed the highest antioxidant activity. Four other plant extracts showed a high antioxidant activity in the following descending order: *P. betle* > *P. guajava* > *E. hirta* > *M. pudica*. There was a positive correlation between total phenolic content and antioxidant activity of the 23 extracts. Beside its high antioxidant activity, *P. amarus* extract also showed antibacterial activity against two different strains

of *A. hydrophila* and one strain of *E. ictaluri*. Tannins were shown to be mainly responsible of these activities, as after tannins removal from *P. amarus* and *E. hirta* extracts, both antioxidant and antimicrobial activities decreased. Both *P. amarus* and *E. hirta* appeared to be promising active plants for further studies. The extracts of these two plants could be valuable natural antibiotic alternatives or natural additives to improve seafood and aquaculture product preservation.

Acknowledgments

The authors are grateful to Ir. Guy Degand and François Brose, who worked in the Laboratory of Food Analysis (University of Liège), for their scientific assistance and valuable support in this project. The authors declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

This work was supported by the Académie de Recherche et d'Enseignement supérieur - Commission de la Coopération au Développement (ARES-CCD) through AquaBioActive (Natural bio-active plant products for environmental friendly aquaculture production in the Mekong Delta) project.

ORCID

Nguyen Le Anh Dao  <http://orcid.org/0000-0001-7534-6768>

Marie-Louise Scippo  <http://orcid.org/0000-0002-8463-3264>

References

- [1] Cowan, M. M. Plant Products as Antimicrobial Agents. *Clin Microbiol Rev.* **1999**, *12*, 564–582. DOI: [10.1128/CMR.12.4.564](https://doi.org/10.1128/CMR.12.4.564).
- [2] Banskota, A. H.; Tezuka, Y.; Tran, Q.; Kadota, S. Chemical Constituents and Biological Activities of Vietnamese Medicinal Plants. *Curr. Top. Med. Chem.* **2003**, *3*, 227–248. DOI: [10.2174/1568026033392516](https://doi.org/10.2174/1568026033392516).
- [3] Chi, V. V. *Dictionary of Medicinal Plants in Vietnam*; Medical Publishing House: Hanoi, **1997**; pp 210–215.
- [4] Doan, D. D.; Nguyen, N.; Doan, H.; Nguyen, T.; Phan, T.; Grabe, M.; Johansson, R.; Lindgren, G.; Stjernström, N. Studies on the Individual and Combined Diuretic Effects of Four Vietnamese Traditional Herbal Remedies (*Zea Mays*, *Imperata Cylindrica*, *Plantago Major* and *Orthosiphon Stamineus*). *J. Ethnopharmacol.* **1992**, *36*, 225–231. DOI: [10.1016/0378-8741\(92\)90048-V](https://doi.org/10.1016/0378-8741(92)90048-V).
- [5] Thuong, P. T.; Na, M.-K.; Dang, N. H.; Hung, T. M.; Ky, P. T.; Thanh, T. V.; Nam, N. H.; Thuan, N. D.; Sok, D.-E.; Bae, K.-H. Antioxidant Activities of Vietnamese Medicinal Plants. *Nat. Prod. Sci.* **2006**, *12*, 29–37.
- [6] Namhui, Y.; Bae, K. Screening of Vietnamese Medicinal Plants for Cytotoxic Activity. *Nat. Prod. Sci.* **2010**, *16*, 1.
- [7] Nguyen, A. T.; Fontaine, J.; Malonne, H.; Vanhaelen, M.; Dubois, J.; Ky Pham, T.; Duez, P. Cytotoxicity of Five Plants Used as Anticancer Remedies in Vietnamese Traditional Medicine. In *Recent Progress in Medicinal Plants*; Singh, V. K., Singh, A. V. K., Govil, J. N., Eds.; Studium-Press: New Delhi, India, **2005**; pp 145–155.
- [8] Hue Ngan, D.; Hoai, H. T. C.; Mai Huong, L.; Hansen, P. E.; Vang, O. Bioactivities and Chemical Constituents of a Vietnamese Medicinal Plant Che Vang, *Jasminum Subtripplinerve* Blume (Oleaceae). *Nat. Prod. Res.* **2008**, *22*, 942–949. DOI: [10.1080/14786410701647119](https://doi.org/10.1080/14786410701647119).
- [9] Madsen, H. L.; Bertelsen, G. Spices as Antioxidants. *Trends Food Sci. Technol.* **1995**, *6*, 271–277. DOI: [10.1016/S0924-2244\(00\)89112-8](https://doi.org/10.1016/S0924-2244(00)89112-8).
- [10] GSO. **2018**. http://www.gso.gov.vn/default_en.aspx?tabid=778
- [11] VASEP. **2018**. <http://mseafod.vasep.com.vn/685/onecontent/fishery-profile.htm/>
- [12] Crumlish, M.; Dung, T.; Turnbull, J.; Ngoc, N.; Ferguson, H. Identification of *Edwardsiella Ictaluri* from Diseased Freshwater Catfish, *Pangasius Hypophthalmus* (Sauvage), Cultured in the Mekong Delta, Vietnam. *J. Fish Dis.* **2002**, *25*, 733–736. DOI: [10.1046/j.1365-2761.2002.00412.x](https://doi.org/10.1046/j.1365-2761.2002.00412.x).
- [13] Subagja, J.; Slembrouck, J.; Hung, L. T.; Legendre, M. Larval Rearing of an Asian Catfish *Pangasius Hypophthalmus* (Siluroidei, Pangasiidae): Analysis of Precocious Mortality and Proposition of Appropriate Treatments. *Aquat. Living Res.* **1999**, *12*, 37–44. DOI: [10.1016/S0990-7440\(99\)80013-8](https://doi.org/10.1016/S0990-7440(99)80013-8).
- [14] Roberts, R. J.; Motile Aeromonad Septicemia. In *Bacterial Diseases of Fish*; Inglis, V., Roberts, R. J., Bromage, N. R., Eds.; John Wiley and Sons: New York, **1993**; pp 143–156.

- [15] Phu, T. M.; Phuong, N. T.; Dung, T. T.; Hai, D. M.; Son, V. N.; Rico, A.; Clausen, J. H.; Madsen, H.; Murray, F.; Dalsgaard, A. An Evaluation of Fish Health-management Practices and Occupational Health Hazards Associated with *Pangasius Catfish (Pangasianodon Hypophthalmus)* Aquaculture in the Mekong Delta, Vietnam. *Aquac. Res.* **2016**, *47*, 2778–2794. DOI: [10.1111/are.12728](https://doi.org/10.1111/are.12728).
- [16] Rico, A.; Phu, T. M.; Satapornvanit, K.; Min, J.; Shahabuddin, A.; Henriksson, P. J.; Murray, F. J.; Little, D. C.; Dalsgaard, A.; Van den Brink, P. J. Use of Veterinary Medicines, Feed Additives and Probiotics in Four Major Internationally Traded Aquaculture Species Farmed in Asia. *Aquaculture.* **2013**, *412*, 231–243. DOI: [10.1111/are.12728](https://doi.org/10.1111/are.12728).
- [17] Nhu, T. Q.; Hang, B. T. B.; Hue, B. T. B.; Quetin-Leclercq, J.; Scippo, M.-L.; Phuong, N. T.; Kestemont, P. Plant Extract-based Diets Differently Modulate Immune Responses and Resistance to Bacterial Infection in Striped Catfish (*Pangasianodon Hypophthalmus*). *Fish Shellfish Immunol.* **2019**, *92*, 913–924. DOI: [10.1016/j.fsi.2019.07.064](https://doi.org/10.1016/j.fsi.2019.07.064).
- [18] Nhu, T. Q.; Hang, B. T. B.; Vinikas, A.; Hue, B. T. B.; Quetin-Leclercq, J.; Scippo, M.-L.; Phuong, N. T.; Kestemont, P. Screening of Immuno-modulatory Potential of Different Herbal Plant Extracts Using Striped Catfish (*Pangasianodon Hypophthalmus*) Leukocyte-based *in Vitro* Tests. *Fish Shellfish Immunol.* **2019**, *93*, 296–307. DOI: [10.1016/j.fsi.2019.07.025](https://doi.org/10.1016/j.fsi.2019.07.025).
- [19] Ho, P. H. *An Illustrated Flora of Vietnam*; Tre Publishing House: Ho Chi Minh City, Vietnam, **2003**; Vol. 3.
- [20] Chi, V. V. *Dictionary of Common Plants in Vietnam*; Science and Technic Publishing House: Ha Noi, **2003**; Vol. 1.
- [21] Ban, N. T. *Checklist of Plant Species of Vietnam*; Agriculture Publishing House: Ha Noi, **2005**; Vol. 3.
- [22] Ho, P. H. *An Illustrated Flora of Vietnam*; Tre Publishing House: Ho Chi Minh City, **2003**; Vol. 1.
- [23] Ho, P. H. *An Illustrated Flora of Vietnam*; Tre Publishing House: Ho Chi Minh City, **2003**; Vol. 2.
- [24] Chi, V. V. *Dictionary of Common Plants in Vietnam*; Science and Technic Publishing House: Ha Noi, **2004**; Vol. 2.
- [25] Singleton, V. L.; Rossi, J. A. Colorimetry of Total Phenolics with Phosphomolybdic-phosphotungstic Acid Reagents. *Am. J. Enol. Vitic.* **1965**, *16*, 144–158.
- [26] Thiangthum, S.; Dejaegher, B.; Goodarzi, M.; Tistaert, C.; Gordien, A.; Hoai, N. N.; Van, M. C.; Quetin-Leclercq, J.; Suntornsuk, L.; Vander Heyden, Y. Potentially Antioxidant Compounds Indicated from *Mallotus* and *Phyllanthus* Species Fingerprints. *J. Chromatogr. B.* **2012**, *910*, 114–121. DOI: [10.1016/j.jchromb.2012.06.025](https://doi.org/10.1016/j.jchromb.2012.06.025).
- [27] Sarker, S. D.; Nahar, L.; Kumarasamy, Y. Microtitre Plate-based Antibacterial Assay Incorporating Resazurin as an Indicator of Cell Growth, and Its Application in the *in Vitro* Antibacterial Screening of Phytochemicals. *Methods.* **2007**, *42*, 321–324. DOI: [10.1016/j.ymeth.2007.01.006](https://doi.org/10.1016/j.ymeth.2007.01.006).
- [28] Drummond, A. J.; Waigh, R. D. The Development of Microbiological Methods for Phytochemical Screening. *Rec. Res. Dev. Phytochem.* **2000**, *4*, 143–152.
- [29] European Pharmacopoeia Commission. Tannins in Herbal Drugs (Method 2. 8.14). In *European Pharmacopoeia*, 9th ed.; European Directorate for the Quality of Medicines (EDQM): Strasbourg, **2018**; pp 275–276.
- [30] Houghton, P.; Raman, A. Methods for Extraction and Sample Clean-up. In *Laboratory Handbook for the Fractionation of Natural Extracts*, Pharmacognosy Research Laboratories, Department of Pharmacy, King's College London, UK; Chapman and Hall: London, UK, **1998**; pp 199.
- [31] Kumaran, A.; Karunakaran, R. J. *In Vitro* Antioxidant Activities of Methanol Extracts of Five *Phyllanthus* Species from India. *LWT-Food Sci. Technol.* **2007**, *40*, 344–352. DOI: [10.1016/j.lwt.2005.09.011](https://doi.org/10.1016/j.lwt.2005.09.011).
- [32] Atiya, A.; Sinha, B. N.; Ranjan Lal, U. New Chemical Constituents from the *Piper Betle* Linn. (Piperaceae). *Nat. Prod. Res.* **2018**, *32*, 1080–1087. DOI: [10.1080/14786419.2017.1380018](https://doi.org/10.1080/14786419.2017.1380018).
- [33] Ekalu, U.; Ikpeme, E.; Ekerette, E.; Chukwu, C. *In Vitro* Antioxidant and Free Radical Activity of Some Nigerian Medicinal Plants: Bitter Leaf (*Vernonia Amygdalina* L.) And Guava (*Psidium Guajava* DeL.). *Res. Res. J. Med. Plant.* **2015**, *9*, 215–226. DOI: [10.3923/rjmp.2015.215.226](https://doi.org/10.3923/rjmp.2015.215.226).
- [34] Teeli, R. A.; Ganie, S. A.; Dar, M. S.; Raja, W.; Yadav, S. S. Antioxidant Activity of *Euphorbia Hirta* L. Leaf Extracts. *Res. J. Pharm. Technol.* **2018**, *11*, 199–202. DOI: [10.5958/0974-360X.2018.00037.9](https://doi.org/10.5958/0974-360X.2018.00037.9).
- [35] Das, K.; Yasin, M.; Mahbub, N. U.; Islam, M. S.; Mahbuba, N. Evaluation of Antioxidant and Cytotoxic Activity of Methanolic Extract of *Mimosa Pudica* Leaves. *Pharma Innov.* **2014**, *3*, 32.
- [36] Mojani, M.; Ghasemzadeh, A.; Rahmat, A.; Loh, S.; Ramasamy, R. Assessment of Bioactive Compounds, Nutritional Composition and Antioxidant Activity of Malaysian Young Ginger (*Zingiber Officinale* Roscoe). *Int. Food Res. J.* **2014**, *21*, 1931.
- [37] Le, Q. U.; Lay, H. L.; Wu, M. C. Antioxidant Activities and HepG2 Cells Growth Inhibitory Capacity of Whole Plant Ethanol Extracts (*Eclipta Alba* Hassk and *Mesona Procumbens* Hems). *J. Food Biochem.* **2018**, *42*, e12454. DOI: [10.1111/jfbc.12454](https://doi.org/10.1111/jfbc.12454).
- [38] Subba, B.; Aryal, P. Study of Biological Activity and Chemical Constituent of *Annona Reticulata*. *J. Inst. Sci. Techn.* **2016**, *21*, 157–163. DOI: [10.3126/JIST.V21I1.16068](https://doi.org/10.3126/JIST.V21I1.16068).
- [39] Wang, K.-H.; Lin, R.-D.; Hsu, F.-L.; Huang, Y.-H.; Chang, H.-C.; Huang, C.-Y.; Lee, M.-H. Cosmetic Applications of Selected Traditional Chinese Herbal Medicines. *J. Ethnopharmacol.* **2006**, *106*, 353–359. DOI: [10.1016/j.jep.2006.01.010](https://doi.org/10.1016/j.jep.2006.01.010).
- [40] Rabeta, M.; Lin, S. Effects of Different Drying Methods on the Antioxidant Activities of Leaves and Berries of *Cayratia Trifolia*. *Sains Malays.* **2015**, *44*, 275–280. DOI: [10.17576/jsm-2015-4402-16](https://doi.org/10.17576/jsm-2015-4402-16).

- [41] Perumal, P. C.; Sophia, D.; Raj, C. A.; Ragavendran, P.; Starlin, T.; Gopalakrishnan, V. K. *In Vitro* Antioxidant Activities and HPTLC Analysis of Ethanolic Extract of *Cayratia Trifolia* (L.). *Asian Pac. J. Trop. Dis.* **2012**, *2*, S952–S956. DOI: [10.1016/S2222-1808\(12\)60299-0](https://doi.org/10.1016/S2222-1808(12)60299-0).
- [42] Lin, E.-S.; Chou, H.-J.; Kuo, P.-L.; Huang, Y.-C. Antioxidant and Antiproliferative Activities of Methanolic Extracts of *Perilla Frutescens*. *J. Med. Plants Res.* **2010**, *4*, 477–483. DOI: [10.5897/JMPR10.035](https://doi.org/10.5897/JMPR10.035).
- [43] Narendhirakannan, R. T.; Nirmala, J. G.; Caroline, A.; Lincy, S.; Saj, M.; Durai, D. Evaluation of Antibacterial, Antioxidant and Wound Healing Properties of Seven Traditional Medicinal Plants from India in Experimental Animals. *Asian Pac. J. Trop. Biomed.* **2012**, *2*, S1245–S1253. DOI: [10.1016/S2221-1691\(12\)60394-3](https://doi.org/10.1016/S2221-1691(12)60394-3).
- [44] Neelabh, C.; Nahid, A.; Kumar, N. Study on Methanolic Extract of *Ageratum Conyzoides* for Its Ability to Act as an Antioxidant and to Suppress the Microbial Growth. *Pharma Innov.* **2017**, *6*, 170.
- [45] Alam, M.; Juraimi, A. S.; Rafii, M.; Abdul Hamid, A.; Aslani, F.; Hasan, M.; Zainudin, M.; Asraf, M.; Uddin, M. Evaluation of Antioxidant Compounds, Antioxidant Activities, and Mineral Composition of 13 Collected Purslane (*Portulaca Oleracea* L.) Accessions. *Biomed Res. Int.* **2014**, *2014*, 1–10. DOI: [10.1155/2014/296063](https://doi.org/10.1155/2014/296063).
- [46] Chekki, R. Z.; Snoussi, A.; Hamrouni, I.; Bouzouita, N. Chemical Composition, Antibacterial and Antioxidant Activities of Tunisian Garlic (*Allium Sativum*) Essential Oil and Ethanol Extract. *Mediterranean J. Chem.* **2014**, *3*, 947–956. DOI: [10.13171/mjc.3.4.2014.09.07.11](https://doi.org/10.13171/mjc.3.4.2014.09.07.11).
- [47] Aydemir, T.; Becerik, S. Phenolic Content and Antioxidant Activity of Different Extracts from *Ocimum Basilicum*, *Apium Graveolens* and *Lepidium Sativum* Seeds. *J. Food Biochem.* **2011**, *35*, 62–79. DOI: [10.1111/j.1745-4514.2010.00366.x](https://doi.org/10.1111/j.1745-4514.2010.00366.x).
- [48] Dewi, R. T.; Maryani, F. Antioxidant and α -glucosidase Inhibitory Compounds of *Centella Asiatica*. *Procedia Chem.* **2015**, *17*, 147–152. DOI: [10.1016/j.proche.2015.12.130](https://doi.org/10.1016/j.proche.2015.12.130).
- [49] Gurusamy, K.; Saranya, P. *In Vitro* Antioxidant Potential of Ethanolic Contents of *Eclipta Alba* and *Wedelia Chinensis*. *J. Pharm. Res.* **2010**, *3*, 2825–2827.
- [50] Rezaeizadeh, A.; Zuki, A.; Abdollahi, M.; Goh, Y.; Noordin, M.; Hamid, M.; Azmi, T. Determination of Antioxidant Activity in Methanolic and Chloroformic Extracts of *Momordica Charantia*. *Afr. J. Biotechnol.* **2011**, *10*, 4932–4940. DOI: [10.5897/AJB10.1972](https://doi.org/10.5897/AJB10.1972).
- [51] Ho, Y.-L.; Huang, S.-S.; Deng, J.-S.; Lin, Y.-H.; Chang, Y.-S.; Huang, G.-J. *In Vitro* Antioxidant Properties and Total Phenolic Contents of Wetland Medicinal Plants in Taiwan. *Bot. Stud.* **2012**, *53*, 55–66.
- [52] Othman, A.; Ismail, A.; Hassan, F. A.; Yusof, B. N. M.; Khatib, A. Comparative Evaluation of Nutritional Compositions, Antioxidant Capacities, and Phenolic Compounds of Red and Green Sessile Joyweed (*Alternanthera Sessilis*). *J. Funct. Foods.* **2016**, *21*, 263–271. DOI: [10.1016/j.jff.2015.12.014](https://doi.org/10.1016/j.jff.2015.12.014).
- [53] Pratt, D. E.; Hudson, B. J. Natural Antioxidants Not Exploited Commercially. In *Food Antioxidants*; Springer, **1990**; pp 171–191. DOI: [10.1007/978-94-009-0753-9_5](https://doi.org/10.1007/978-94-009-0753-9_5).
- [54] Regi Raphael, K.; Ajith, T.; Joseph, S.; Kuttan, R. Anti-mutagenic Activity of *Phyllanthus Amarus* Schum & Thonn *in Vitro* as Well as *in Vivo*. *Teratog. Carcinog. Mutagen.* **2002**, *22*, 285–291. DOI: [10.1002/tcm.10021](https://doi.org/10.1002/tcm.10021).
- [55] Guha, G.; Rajkumar, V.; Kumar, R. A.; Mathew, L. Aqueous Extract of *Phyllanthus Amarus* Inhibits Chromium (Vi)-induced Toxicity in MDA-MB-435S Cells. *Food Chem. Toxicol.* **2010**, *48*, 396–401. DOI: [10.1016/j.fct.2009.10.028](https://doi.org/10.1016/j.fct.2009.10.028).
- [56] Igwe, C. U.; Nwaogu, L. A.; Ujuwundu, C. O. Assessment of the Hepatic Effects, Phytochemical and Proximate Compositions of *Phyllanthus Amarus*. *Afr. J. Biotechnol.* **2007**, *6*, 728–731.
- [57] Sharma, A.; Singh, R. T.; Handa, S. S. Estimation of Phyllanthin and Hypophyllanthin by High Performance Liquid Chromatography in *Phyllanthus Amarus*. *Phytochem. Anal.* **1993**, *4*, 226–229. DOI: [10.1002/pca.2800040507](https://doi.org/10.1002/pca.2800040507).
- [58] Dudonne, S.; Vitrac, X.; Coutiere, P.; Woillez, M.; Merillon, J.-M. Comparative Study of Antioxidant Properties and Total Phenolic Content of 30 Plant Extracts of Industrial Interest Using DPPH, ABTS, FRAP, SOD, and ORAC Assays. *J. Agric. Food Chem.* **1768-1774**, 2009(57). DOI: [10.1021/jf803011r](https://doi.org/10.1021/jf803011r).
- [59] Shyur, L.-F.; Tsung, J.-H.; Chen, J.-H.; Chiu, C.-Y.; Lo, C.-P. Antioxidant Properties of Extracts from Medicinal Plants Popularly Used in Taiwan. *Int. J. Appl. Sci. Eng.* **2005**, *3*, 195–202.
- [60] Soumaya, K.; Chaouachi, F.; Ksouri, R.; El Gazzah, M. Polyphenolic Composition in Different Organs of Tunisia Populations of *Cynara Cardunculus* L. And Their Antioxidant Activity. *J. Food Nutr. Res.* **2013**, *1*, 1–6. DOI: [10.12691/jfnr-1-1-1](https://doi.org/10.12691/jfnr-1-1-1).
- [61] Liu, X.; Zhao, M.; Wang, J.; Yang, B.; Jiang, Y. Antioxidant Activity of Methanolic Extract of Emblica Fruit (*Phyllanthus Emblica* L.) From Six Regions in China. *J. Food Compos. Anal.* **2008**, *21*, 219–228. DOI: [10.1016/j.jfca.2007.10.001](https://doi.org/10.1016/j.jfca.2007.10.001).
- [62] Shahriar, M.; Akhter, S.; Hossain, M. I.; Haque, M. A.; Bhuiyan, M. A. Evaluation of *in Vitro* Antioxidant Activity of Bark Extracts of *Terminalia Arjuna*. *J. Med. Plants Res.* **2012**, *6*, 5286–5298. DOI: [10.5897/JMPR12.580](https://doi.org/10.5897/JMPR12.580).
- [63] Eldeen, I.; Seow, E.; Abdullah, R.; Sulaiman, S. *In Vitro* Antibacterial, Antioxidant, Total Phenolic Contents and anti-HIV-1 Reverse Transcriptase Activities of Extracts of Seven *Phyllanthus* Sp. *S. Afr. J. Bot.* **2011**, *77*, 75–79. DOI: [10.1016/j.sajb.2010.05.009](https://doi.org/10.1016/j.sajb.2010.05.009).

- [64] Bruno, D.; Munro, A. *The Calculation of Various Treatment Dose Rates in Fish Farming*; Scottish Office Agriculture and Fisheries Department, Marine Laboratory: Aberdeen, Scotland, **1991**; pp 10.
- [65] Rodgers, C.; Furones, M. Antimicrobial Agents in Aquaculture: Practice, Needs and Issues. *Options Médit.* **2009**, *86*, 41–59.
- [66] Mazumder, A.; Mahato, A.; Mazumder, R. Antimicrobial Potentiality of *Phyllanthus Amarus* against Drug Resistant Pathogens. *Nat. Prod. Res.* **2006**, *20*, 323–326. DOI: [10.1080/14786410600650404](https://doi.org/10.1080/14786410600650404).
- [67] Taukoora, U.; Lall, N.; Mahomoodally, F. *Piper Betle* L. (Betel Quid) Shows Bacteriostatic, Additive, and Synergistic Antimicrobial Action When Combined with Conventional Antibiotics. *S. Afr. J. Bot.* **2016**, *105*, 133–140. DOI: [10.1016/j.sajb.2016.01.006](https://doi.org/10.1016/j.sajb.2016.01.006).
- [68] Braga, T. V.; Das Dores, R. G. R.; Ramos, C. S.; Evangelista, F. C. G.; da Silva Tinoco, L. M.; de Pilla Varotti, F.; Das Graças Carvalho, M.; de Paula Sabino, A. Antioxidant, Antibacterial and Antitumor Activity of Ethanolic Extract of the *Psidium Guajava* Leaves. *Am. J. Plant Sci.* **2014**, *5*, 3492. DOI: [10.4236/ajps.2014.523365](https://doi.org/10.4236/ajps.2014.523365).
- [69] Gupta, D.; Kumar, M.; Gupta, V. An *in Vitro* Investigation of Antimicrobial Efficacy of *Euphorbia Hirta* and *Murraya Koenigii* against Selected Pathogenic Microorganisms. *Asian J. Pharm. Clin. Res.* **2018**, *11*, 359–363. DOI: [10.22159/ajpcr.2018.v11i5.24578](https://doi.org/10.22159/ajpcr.2018.v11i5.24578).
- [70] Tomar, R. S.; Shrivastava, V.; Kaushik, S. *In Vitro* Efficacy of Methanolic Extract of *Mimosa Pudica* against Selected Micro-organisms for Its Broad Spectrum Antimicrobial Activity. *Int. J. Curr. Microbiol. Appl. Sci.* **2014**, *3*, 780–784.
- [71] Borkatak, M.; Kakoty, B.; Saikia, L. Proximate Analysis and Antimicrobial Activity of *Eclipta Alba* (L.) Hassk. —a Traditionally Used Herb. *Int. J. Pharm. Pharm. Sci.* **2013**, *5*, 149–154.
- [72] Aghazadeh, M.; Bialvaei, A. Z.; Aghazadeh, M.; Kabiri, F.; Saliyani, N.; Yousefi, M.; Eslami, H.; Kafil, H. S. Survey of the Antibiofilm and Antimicrobial Effects of *Zingiber Officinale* (*In Vitro* Study). *Jundishapur J. Microbiol.* **2016**, *9*. DOI: [10.5812/jjm.30167](https://doi.org/10.5812/jjm.30167).
- [73] Jamkhande, P. G.; Wattamwar, A. S.; Kankudte, A. D.; Tidke, P. S.; Kalaskar, M. G. Assessment of *Annona Reticulata* Linn. Leaves Fractions for *in Vitro* Antioxidative Effect and Antimicrobial Potential against Standard Human Pathogenic Strains. *Alexandria J. Med.* **2016**, *52*, 19–25. DOI: [10.1016/j.ajme.2014.12.007](https://doi.org/10.1016/j.ajme.2014.12.007).
- [74] Zhou, G.; Yu, H.; Yu, Y. Study on Bacteriostatic Action in Different Parts of *Houttuynia Cordata*. *Human Forest Sci. Technol.* **2006**, *33*, 38–40.
- [75] Prajapati, R.; Roy, S.; Mishra, S.; Raza, S.; Thakur, L. Formulation Development, Standardization and Antimicrobial Activity of *Ageratum Conyzoides* Extracts and Their Formulation. *Int. J. Pharm. Pharm. Sci.* **2014**, *6*, 369–374.
- [76] Agyare, C.; Baiden, E.; Apenteng, J. A.; Boakye, Y. D.; Adu-Amoah, L. Anti-infective and Anti-inflammatory Properties of *Portulaca Oleracea* L. *Donnish. J. Med. Plant Res.* **2015**, *2*, 1–6.
- [77] Meriga, B.; Mopuri, R.; MuraliKrishna, T. Insecticidal, Antimicrobial and Antioxidant Activities of Bulb Extracts of *Allium Sativum*. *Asian Pac. J. Trop. Med.* **2012**, *5*, 391–395. DOI: [10.1016/S1995-7645\(12\)60065-0](https://doi.org/10.1016/S1995-7645(12)60065-0).
- [78] Hossain, M. A.; Kabir, M.; Salehuddin, S.; Rahman, S. M.; Das, A.; Singha, S. K.; Alam, M. K.; Rahman, A. Antibacterial Properties of Essential Oils and Methanol Extracts of Sweet Basil *Ocimum Basilicum* Occurring in Bangladesh. *Pharm. Biol.* **2010**, *48*, 504–511. DOI: [10.3109/13880200903190977](https://doi.org/10.3109/13880200903190977).
- [79] Taemchua, D.; Rukkamsuk, T.; Sakpuaram, T.; Ruangwises, N. Antibacterial Activity of Crude Extracts of *Centella Asiatica* against *Staphylococcus Aureus* in Bovine Mastitis. *Kasetsart Vet.* **2009**, *19*, 119–128.
- [80] Darah, I.; Lim, S.; Nithianantham, K. Effects of Methanol Extract of *Wedelia Chinensis* Osbeck (Asteraceae) Leaves against Pathogenic Bacteria with Emphasis on *Bacillus Cereus*. *Indian J. Pharm. Sci.* **2013**, *75*, 533–539.
- [81] Svobodova, B.; Barros, L.; Calhelha, R. C.; Heleno, S.; Alves, M. J.; Walcott, S.; Bittova, M.; Kuban, V.; Ferreira, I. C. Bioactive Properties and Phenolic Profile of *Momordica Charantia* L. Medicinal Plant Growing Wild in Trinidad and Tobago. *Ind. Crops Prod.* **2017**, *95*, 365–373. DOI: [10.1016/j.indcrop.2016.10.046](https://doi.org/10.1016/j.indcrop.2016.10.046).
- [82] Kuete, V. Potential of Cameroonian Plants and Derived Products against Microbial Infections: A Review. *Planta Med.* **2010**, *76*, 1479–1491. DOI: [10.1055/s-0030-1250027](https://doi.org/10.1055/s-0030-1250027).
- [83] Dung, T. T.; Haesebrouck, F.; Tuan, N. A.; Sorgeloos, P.; Baele, M.; Decostere, A. Antimicrobial Susceptibility Pattern of *Edwardsiella Ictaluri* Isolates from Natural Outbreaks of Bacillary Necrosis of *Pangasianodon Hypophthalmus* in Vietnam. *Microb. Drug Resist.* **2008**, *14*, 311–316. DOI: [10.1089/mdr.2008.0848](https://doi.org/10.1089/mdr.2008.0848).
- [84] De Britto, A. J.; Gracelin, D. H. S.; Sebastian, S. R. Antibacterial Activity of a Few Medicinal Plants against *Xanthomonas Campestris* and *Aeromonas Hydrophila*. *J. Biopesticides.* **2011**, *4*, 57.
- [85] Caruso, D.; Lusiatusti, A. M.; Taukhid, T.; Avarre, J. C.; Yuhana, M.; Sarter, S. Ethnobotanical Uses and Antimicrobial Properties of Plants in Small-scale Tropical Fish Farms: The Case of Indonesian Fish Farmers in Java (Indonesia). *J. World Aquac. Soc.* **2017**, *48*, 83–92. DOI: [10.1111/jwas.12345](https://doi.org/10.1111/jwas.12345).
- [86] Saranraj, P.; Sivasakthivelan, P. Screening of Antibacterial Activity of the Medicinal Plant *Phyllanthus Amarus* against Urinary Tract Infection Causing Bacterial Pathogens. *Appl. J. Hyg.* **2012**, *1*, 19–24. DOI: [10.5829/idosi.ajh.2012.1.3.71111](https://doi.org/10.5829/idosi.ajh.2012.1.3.71111).
- [87] Ushie, O.; Neji, P.; Etim, E.; Nsor, G. Phytochemical Screening and Antimicrobial Activities of *Phyllanthus Amarus* Stem Bark Extracts. *Int. J. Mod. Biol. Med.* **2013**, *3*, 101–112.

- [88] Kaveti, B.; Tan, L.; Sarnnia, K. T.; Baig, M. Antibacterial Activity of *Piper Betle* Leaves. *Int. J. Pharm. Teach. Pract.* **2011**, *2*, 129–132.
- [89] Sheikhlari, A.; Meng, G. Y.; Alimon, R.; Romano, N.; Ebrahimi, M. Dietary *Euphorbia Hirta* Extract Improved the Resistance of Sharptooth Catfish *Clarias Gariepinus* to *Aeromonas Hydrophila*. *J. Aquat. Anim. Health.* **2017**, *29*, 225–235. DOI: [10.1080/08997659.2017.1374310](https://doi.org/10.1080/08997659.2017.1374310).
- [90] Bensky, D.; Gamble, A.; Kaptchuk, T. *Chinese Herbal Medicine Materia Medica*; Eastland Press: Seattle, USA, **1993**.
- [91] Yoshida, T.; Chen, L.; Shingu, T.; Okuda, T. Tannins and Related Polyphenols of Euphorbiaceous Plants. IV.: Euphorbins A and B, Novel Dimeric Dehydroellagitannins from *Euphorbia Hirta* L. *Chem. Pharm. Bull.* **1988**, *36*, 2940–2949. DOI: [10.1248/cpb.36.2940](https://doi.org/10.1248/cpb.36.2940).
- [92] Haslam, E., *Plant Polyphenols: Vegetable Tannins Revisited*; Cambridge University Press: Cambridge, UK, **1989**.
- [93] Chung, K.-T.; Wong, T. Y.; Wei, C.-I.; Huang, Y.-W.; Lin, Y. Tannins and Human Health: A Review. *Crit. Rev. Food Sci. Nutr.* **1998**, *38*, 421–464. DOI: [10.1080/10408699891274273](https://doi.org/10.1080/10408699891274273).
- [94] Gu, H.-F.; Li, C.-M.; Xu, Y.-J.; Hu, W.-F.; Chen, M.-H.; Wan, Q.-H. Structural Features and Antioxidant Activity of Tannin from Persimmon Pulp. *Food Res. Int.* **2008**, *41*, 208–217. DOI: [10.1016/j.foodres.2007.11.011](https://doi.org/10.1016/j.foodres.2007.11.011).
- [95] Cushnie, T. T.; Lamb, A. J. Antimicrobial Activity of Flavonoids. *Int. J. Antimicrob. Agents.* **2005**, *26*, 343–356. DOI: [10.1016/j.ijantimicag.2005.09.002](https://doi.org/10.1016/j.ijantimicag.2005.09.002).
- [96] Karou, D.; Savadogo, A.; Canini, A.; Yameogo, S.; Montesano, C.; Simpore, J.; Colizzi, V.; Traore, A. S. Antibacterial Activity of Alkaloids from *Sida Acuta*. *Afr. J. Biotechnol.* **2006**, *5*, 195–200.
- [97] Foo, L. Y. Amariin, a Di-dehydrohexahydroxydiphenoyl Hydrolysable Tannin from *Phyllanthus Amarus*. *Phytochemistry.* **1993**, *33*, 487–491. DOI: [10.1016/0031-9422\(93\)85545-3](https://doi.org/10.1016/0031-9422(93)85545-3).
- [98] Foo, L. Y. Amariin Acid and Related Ellagitannins from *Phyllanthus Amarus*. *Phytochemistry.* **1995**, *39*, 217–224. DOI: [10.1016/0031-9422\(94\)00836-I](https://doi.org/10.1016/0031-9422(94)00836-I).
- [99] Foo, L. Y.; Wong, H. Phyllanthusiin D, an Unusual Hydrolysable Tannin from *Phyllanthus Amarus*. *Phytochemistry.* **1992**, *31*, 711–713. DOI: [10.1016/0031-9422\(92\)90071-W](https://doi.org/10.1016/0031-9422(92)90071-W).
- [100] Catteau, L.; Van Bambeke, F.; Quetin-Leclercq, J. Preliminary Evidences of the Direct and Indirect Antimicrobial Activity of 12 Plants Used in Traditional Medicine in Africa. *Phytochem. Rev.* **2015**, *14*, 975–991. DOI: [10.1007/s11101-015-9437-x](https://doi.org/10.1007/s11101-015-9437-x).
- [101] Patel, J. R.; Tripathi, P.; Sharma, V.; Chauhan, N. S.; Dixit, V. K. *Phyllanthus Amarus*: Ethnomedicinal Uses, Phytochemistry and Pharmacology: A Review. *J. Ethnopharmacol.* **2011**, *138*, 286–313. DOI: [10.1016/j.jep.2011.09.040](https://doi.org/10.1016/j.jep.2011.09.040).
- [102] Yoshida, T.; Namba, O.; Chen, L.; Okuda, T. Euphorbin E, a Hydrolyzable Tannin Dimer of Highly Oxidized Structure, from *Euphorbia Hirta*. *Chem. Pharm. Bull.* **1990**, *38*, 1113–1115. DOI: [10.1248/cpb.38.1113](https://doi.org/10.1248/cpb.38.1113).
- [103] Yoshida, T.; Namba, O.; Chen, L.; Okuda, T. Tannins and Related Polyphenols of Euphorbiaceous Plants. V.: Euphorbin C, an Equilibrated Dimeric Dehydroellagitannin Having a New Tetrameric Galloyl Group. *Chem. Pharm. Bull.* **1990**, *38*, 86–93. DOI: [10.1248/cpb.38.86](https://doi.org/10.1248/cpb.38.86).