

## Visualizing Chlorophyll a Fluorescence: A Practical Demonstration

Ali Ahmad – University of Granada


Santiago Atero Calvo – University of Granada


Begoña Blasco León – University of Granada

Safa Selmi – Association of Safeguard of Matmata - University of Gabes


Alessandro Candiani – DNAPhone SRL


Vanessa Martos Núñez – University of Granada

 0000-0001-5530-7374

 0000-0001-8446-5515

 0000-0001-8061-5141

 0000-0002-3389-6806

 0000-0002-6200-7705

 0000-0001-6442-7968

Recepción: 14.11.2022 | Aceptado: 18.11.2022

Correspondencia a través de **ORCID**: Ali Ahmad

 **0000-0001-5530-7374**

Citar: Ahmad, A, Atero Calvo, S, Blasco León, B, Selmi, S, Candiani, A, & Martos, V (2022). Visualizing Chlorophyll a Fluorescence: A Practical Demonstration. *REIDOCREA*, 11(63), 713-718.

Financiación: This work was supported by the projects: “VIRTUOUS” funded from the European Union’s Horizon 2020 Project H2020-MSCA-RISE-2019. Ref. 872181, “SUSTAINABLE” funded from the European Union’s Horizon 2020 Project H2020-MSCA-RISE-2020. Ref. 101007702, and the “Project of Excellence” from FEDER (Fondo Europeo de Desarrollo Regional)- Junta de Andalucía 2018. Ref. P18-H0-4700.

Área o categoría del conocimiento: Fisiología Vegetal – Docencia

**Abstract:** Chlorophylls are the principal components of plants for light harvesting. They utilize the energy retrieved from solar radiations to carry out the process of photosynthesis and produce reduced organic compounds such as carbohydrates. However, all of the incident light is not used in photosynthesis process, it confronts two other fates. A part of it is dissipated as heat, whereas the other is emitted as fluorescence. These processes occur simultaneously and whether one or the other occurs to a greater or lesser extent will depend both on the physiological status of the plant and the environmental conditions it faces. Chlorophyll (Chl) fluorescence is inversely proportional to the yield of photosynthesis and therefore is of prime importance in plant physiology. Furthermore, there are a lot of studies where Chl a fluorescence has been used as a probe for estimating photosynthetic yield, drought, salinity, vigor, and environmental effects on crop production yield. Therefore, this study was undertaken to demonstrate the visualization of the light emitted from Chl, commonly known as Chl a fluorescence. Plant leaves were dark adapted for 20 minutes before their exposure to ultraviolet (UV) light. Red glasses were used to visualize the emitted red light (fluorescence) from the leaves. This study may instill further interest in the plant physiology students to deepen and expand their learning by undertaking simple demonstrations like this.

**Keyword:** Chlorophyll a Fluorescence

### **Visualización de la fluorescencia de clorofila a: Una demostración práctica**


**Resumen:** Las clorofilas son los principales componentes de las plantas para recolectar la luz. Utilizan la energía procedente de la radiación solar para llevar a cabo el proceso de fotosíntesis y producir moléculas orgánicas reducidas como los hidratos de carbono. Sin embargo, el total de la utilización de la luz incidente no toda es aprovechada en el proceso de fotosíntesis, ya que esta se enfrenta a otros dos destinos. Una parte se disipa como calor, mientras que la otra se emite como fluorescencia. Estos procesos ocurren de forma simultánea y que se de uno u otro proceso en mayor o menor medida, va a depender tanto del estatus fisiológico de la planta como de las condiciones ambientales a las que se enfrenta. La fluorescencia de la clorofila (Chl) es inversamente proporcional al rendimiento o la tasa de la fotosíntesis y, por lo tanto, tiene una importancia fundamental en los estudios de fisiología vegetal. Asimismo, hay muchos estudios en los que la fluorescencia de Chl a se ha utilizado como sonda para estimar el rendimiento fotosintético, la sequía, la salinidad, el vigor y los efectos ambientales en la producción y el rendimiento de los cultivos. Por lo tanto, este estudio se realizó para demostrar la visualización de la luz emitida por la Chl, comúnmente conocida como fluorescencia de Chl a. Las hojas de las plantas fueron adoptadas en la oscuridad durante 20 minutos antes de su exposición a la luz ultravioleta (UV). Se utilizaron gafas rojas para visualizar la luz roja emitida (fluorescencia) de las hojas. Este estudio puede infundir más interés en los estudiantes de fisiología vegetal para que profundicen y amplíen su aprendizaje realizando demostraciones sencillas como esta.

**Palabra clave:** Fluorescencia de clorofila a

## Introduction

Chlorophylls are the principal components of plants for light harvesting. They utilize the harvested light to produce sugars through the process of photosynthesis. However, all the incident light is not absorbed by chlorophyll (Chl) and a part of it is dissipated (Maxwell & Johnson, 2000). More precisely, the incident light on Chl encounters the following three fates (Figure 1):

1. Absorbance for photosynthesis process (also known as photochemistry)
2. Dissipation in the form of heat
3. Re-emission in the form of light (known as Chl fluorescence)



**Figure 1.** The incident light on plant leaf (chlorophyll) confronts three fates. A part of it is used in photosynthesis process, while a part of it is emitted as heat and other part is emitted as fluorescence.

Photosynthesis, Chl *a* fluorescence, and heat dissipation all of these processes occur simultaneously. The solar energy is absorbed by electron acceptor of photosystem II (PSII), namely plastoquinone A ( $Q_A$ ), which deliver it to photosynthetic apparatus. Once  $Q_A$  has accepted an electron it cannot accept another electron unless it transfers that to another electron acceptor. Reaction center which accepts electron is referred to as “closed” when  $Q_A$  cannot accept another electron. Consequently, light is emitted in the form of fluorescence and heat; thereby resulting in lower rate of photosynthesis. Therefore, estimating the yield of one process could provide useful information about the other processes (Cosgrove & Borowitzka, 2010). For instance, Chl fluorescence yield would suggest the yield of photochemistry and heat emission. Consequently, this would indicate the efficiency of  $CO_2$  assimilation in photosynthesis process. The intensity or yield of Chl fluorescence is inversely proportional to yield of photosynthesis and dissipative heat energy (Duysens, 1963; Krause & Weis, 1991). Hence, it is of paramount importance in plant physiology studies.

It has been reported that Chl fluorescence accounts for only a very small portion of absorbed light i.e., 0.5 to 10% (Barber et al., 1989; Brody & Rabinowitch, 1957; Latimer et al., 1956; Porcar-Castell et al., 2014). Similarly, almost all of the Chl fluorescence derives from PSII accounting for nearly 90—95% (Cosgrove & Borowitzka, 2010). The spectrum of Chl fluorescence is different from light absorption spectrum i.e., the energy emitted as Chl fluorescence has a higher wavelength than absorbed (Maxwell &

Johnson, 2000). Kautsky and co-authors were the first to report Chl fluorescence changes in 1960 (Kautsky et al., 1960). They reported an increase in the Chl fluorescence over a short period of time when dark adapted plants were exposed to light. The present study is meant to practically demonstrate this process of increased Chl a fluorescence for dark adapted leaf.

### **Importance of Chl a Fluorescence**

The implication of studying Chl a fluorescence is multifold. Its yield provides useful information about various photosynthetic parameters including non-photochemical quenching (NPQ), JIP test, dark-adaptation kinetics of OJIP transients, statistic aspects of the measurements of parameters, the actinic light wavelength dependence of photosynthesis, rapid light curves (RLCs), Chl a fluorescence and 820nm transmission, electron transport rate (ETR), Chl a fluorescence and delayed fluorescence, flash-induced fluorescence, imaging, Chl a fluorescence and photoacoustic spectroscopy, and quenching analysis (Bukhov et al., 2001; Bukhov et al., 1997; Buschmann & Kocsányi, 1989; Bussotti et al., 2011; de Wijn & Van Gorkom, 2001; Goltsev et al., 2012; Gorbe & Calatayud, 2012; Hideg & Schreiber, 2007; Horton & Hague, 1988; Ioannidis et al., 2000; Kalaji et al., 2012; Kalaji et al., 2017; Klughammer & Schreiber, 1994; Krall & Edwards, 1990; Lichtenthaler et al., 2007; Nedbal & Whitmarsh, 2004; Ralph & Gademann, 2005; Schansker et al., 2003; Schansker et al., 2005; Schreiber et al., 2012; Schreiber et al., 1986; Snel et al., 1990; Strasser et al., 2004).

Plant Chl content is highly depictive of its vigor and is prone to changing environment (Ahmad, del Moral Garrido, et al., 2022). Therefore, any change affecting Chl would result in an alteration in Chl a fluorescence. Consequently, it could be implied to draw valuable information about various plant biological processes. In this regard, a summary of Chl a fluorescence with respect to various physiological parameters and its eco-physiological applications is provided in Table 1.

**Table 1.** A summary of few representative research articles on the application of chlorophyll a fluorescence in various agronomic and environmental studies.

<b>Assay or Condition</b>	<b>Reference</b>
Fungicide stress	(Ahmad, Navarro-León, et al., 2022)
Heat stress	(Brestic et al., 2012)
UV stress	(Guidi et al., 2011)
Salt stress	(Ahmad, Blasco, et al., 2022)
Photoinhibition	(Matsubara et al., 2011)
Drought stress	(Banks, 2018)
Urban tree conditions	(Swoczyna et al., 2010)
Environmental pollution	(Oláh et al., 2021)
Sulfur-deprivation	(Duan et al., 2019)
Water quality	(Sang et al., 2020)

## **Methodology**

### **Plant material and growth conditions**

The experiment was conducted on a laboratory grown lettuce (*Lactuca sativa*) plants that were 40 days old. Plants were grown in 13cm x 13cm x 12.5 cm plastic pot containing peat and vermiculite. They were irrigated twice a week with tap water. The growth chamber had a relative humidity of 60-80%, photosynthetically active radiation (PAR) of 350  $\mu\text{mol m}^{-2} \text{s}^{-1}$ , and a photoperiod of 14-10 hrs with the subsequent temperature of 22 °C at night and 18 °C during the day.

## Resources

An ultraviolet (UV) torch light, red color glasses, and a piece of paper to block light was used for this experiment. Similarly, Nikon-D5300 digital camera was used to record the video.

## Visualization of Chl a fluorescence


Plants were dark adapted for 15-20 minutes (Figure 2) before their exposure to UV light, in order to record the high Chl a fluorescence (Kautsky et al., 1960). Subsequently, a small piece of paper was placed on a leaf along, whereas the rest of the leaf was continuously exposed to UV. After, few seconds the piece of paper was removed and a high fluorescence in the form of red light was recorded.



**Figure 2.** Dark adapted plant for 15-20 minutes, where all processes of energy entrapment and emission halt.

## Results and Discussion

The dark adapted leaves when exposed to UV light emitted a strong signal of Chl a fluorescence, which was observed with the help of red glasses (Figure 3).



**Figure 3.** An illustration to observe chlorophyll a fluorescence under UV light with the help of red glasses (A). Visualization of red fluorescence from dark adapted leaf under UV light without red glasses (B) and with red glasses (C). Images captured through Nikon D-5300 digital camera.

This increase in Chl a fluorescence over a short period of time when dark adapted leaves were exposed to light had been previously documented by Kautsky and co-authors (Kautsky et al., 1960). The explanation of this lies in the fact that whenever a dark adapted leaf is brought to light, plastoquinone (electron acceptor of PSII) captures an

electron for its further transport. At this stage, the reaction center of Chl *a* is termed as “closed”, which means no further electron can be accepted by plastoquinone. Consequently, the incident light is emitted back in the form of fluorescence and heat, leading to lower photosynthetic yield.

## Conclusion

The incident light on plant leaves is emitted as fluorescence. This Chl *a* fluorescence can be visualized using a simple demonstration. The present study demonstrates the basic process of Chl *a* fluorescence using UV light and red glasses. Red glasses assist in visualizing the emitted red signal of fluorescence when the reaction centers are closed. This study may instill further interest in the plant physiology students to deepen and expand their learning by undertaking simple demonstrations like this.

## References

- Ahmad, A, Blasco, B, & Martos, V (2022). Combating Salinity Through Natural Plant Extracts Based Biostimulants: A Review. *Frontiers in Plant Science*, 1665.
- Ahmad, A, del Moral Garrido, MBG, & Martos, V (2022). Learning about chlorophyll and anthocyanins as potential indicators of plant physiological state. *ReiDoCrea: Revista electrónica de investigación y docencia creativa*(11), 171-176.
- Ahmad, A, Navarro-León, E, Izquierdo-Ramos, MJ, Rios, JJ, Blasco, B, Navarro-Morillo, I, & Ruiz, JM (2022). Analysis of RAZORMIN® as a Biostimulant and Its Effect on the Phytotoxicity Mitigation Caused by Fungicide Azoxystrobin in Pepper. *Agronomy*, 12(6), 1418.
- Banks, JM (2018). Chlorophyll fluorescence as a tool to identify drought stress in *Acer* genotypes. *Environmental and Experimental Botany*, 155, 118-127.
- Barber, J, Malkin, S, & Telfer, A (1989). The origin of chlorophyll fluorescence in vivo and its quenching by the photosystem II reaction centre. *Philosophical Transactions of the Royal Society of London. B, Biological Sciences*, 323(1216), 227-239.
- Brestic, M, Zivcak, M, Kalaji, HM, Carpentier, R, & Allakhverdiev, SI (2012). Photosystem II thermostability in situ: environmentally induced acclimation and genotype-specific reactions in *Triticum aestivum* L. *Plant Physiology and Biochemistry*, 57, 93-105.
- Brody, SS, & Rabinowitch, E (1957). Excitation lifetime of photosynthetic pigments in vitro and in vivo. *Science*, 125(3247), 555-555.
- Bukhov, N, Egorova, E, Krendeleva, T, Rubin, A, Wiese, C, & Heber, U (2001). Relaxation of variable chlorophyll fluorescence after illumination of dark-adapted barley leaves as influenced by the redox states of electron carriers. *Photosynthesis Research*, 70(2), 155-166.
- Bukhov, NG, Boucher, N, & Carpentier, R (1997). The correlation between the induction kinetics of the photoacoustic signal and chlorophyll fluorescence in barley leaves is governed by changes in the redox state of the photosystem II acceptor side. A study under atmospheric and high CO<sub>2</sub> concentrations. *Canadian journal of botany*, 75(9), 1399-1406.
- Buschmann, C, & Kocsányi, L (1989). Light-induced heat production correlated with fluorescence and its quenching mechanisms. *Photosynthesis Research*, 21(2), 129-136.
- Bussotti, F, Desotgiu, R, Cascio, C, Pollastrini, M, Gravano, E, Gerosa, G, Marzuoli, R, Nali, C, Lorenzini, G, & Salvatori, E (2011). Ozone stress in woody plants assessed with chlorophyll *a* fluorescence. A critical reassessment of existing data. *Environmental and Experimental Botany*, 73, 19-30.
- Cosgrove, J, & Borowitzka, MA (2010). Chlorophyll fluorescence terminology: an introduction. In *Chlorophyll *a* fluorescence in aquatic sciences: methods and applications* (pp. 1-17). Springer.
- de Wijn, R, & Van Gorkom, HJ (2001). Kinetics of electron transfer from QA to QB in photosystem II. *Biochemistry*, 40(39), 11912-11922.
- Duan, J, Fu, B, Kang, H, Song, Z, Jia, M, Cao, D, & Wei, A (2019). Response of gas-exchange characteristics and chlorophyll fluorescence to acute sulfur dioxide exposure in landscape plants. *Ecotoxicology and Environmental Safety*, 171, 122-129.
- Duysens, L (1963). Mechanism of the two photochemical reactions in algae as studied by means of fluorescence. *Studies on microalgae and photosynthetic bacteria*, 353-372.
- Goltsev, V, Zaharieva, I, Chemev, P, Kouzmanova, M, Kalaji, HM, Yordanov, I, Krasteva, V, Alexandrov, V, Stefanov, D, & Allakhverdiev, SI (2012). Drought-induced modifications of photosynthetic electron transport in intact leaves: analysis and use of neural networks as a tool for a rapid non-invasive estimation. *Biochimica et Biophysica Acta (BBA)-Bioenergetics*, 1817(8), 1490-1498.
- Goebel, E, & Calatayud, A (2012). Applications of chlorophyll fluorescence imaging technique in horticultural research: A review. *Scientia Horticulturae*, 138, 24-35.
- Guidi, L, Degl'Innocenti, E, Remorini, D, Bircolli, S, Fini, A, Ferrini, F, Nicese, FP, & Tattini, M (2011). The impact of UV-radiation on the physiology and biochemistry of *Ligustrum vulgare* exposed to different visible-light irradiance. *Environmental and Experimental Botany*, 70(2-3), 88-95.

- Hideg, E, & Schreiber, U (2007). Parallel assessment of ROS formation and photosynthesis in leaves by fluorescence imaging. *Photosynthesis Research*, 92(1), 103-108.
- Horton, P, & Hague, A (1988). Studies on the induction of chlorophyll fluorescence in isolated barley protoplasts. IV. Resolution of non-photochemical quenching. *Biochimica et Biophysica Acta (BBA)-Bioenergetics*, 932, 107-115.
- Ioannidis, N, Schansker, G, Barynin, VV, & Petrouleas, V (2000). Interaction of nitric oxide with the oxygen evolving complex of photosystem II and manganese catalase: a comparative study. *Journal of Biological Inorganic Chemistry*, 5(3), 354-363.
- Kalaji, HM, Carpentier, R, Allakhverdiev, SI, & Bosa, K (2012). Fluorescence parameters as early indicators of light stress in barley. *Journal of Photochemistry and Photobiology B: Biology*, 112, 1-6.
- Kalaji, HM, Schansker, G, Brestic, M, Bussotti, F, Calatayud, A, Ferroni, L, Goltsev, V, Guidi, L, Jajoo, A, & Li, P (2017). Frequently asked questions about chlorophyll fluorescence, the sequel. *Photosynthesis Research*, 132(1), 13-66.
- Kautsky, H, Appel, W, & Amann, H (1960). Chlorophyll-fluorescenz und Kohlensäureassimilation. XIII. Mitteilung. Die Fluorescenzkurve und die Photochemie der Pflanze. *Biochem. Z.*, 332, 277-292.
- Klughammer, C, & Schreiber, U (1994). An improved method, using saturating light pulses, for the determination of photosystem I quantum yield via P700<sup>+</sup>-absorbance changes at 830 nm. *Planta*, 192(2), 261-268.
- Krall, J, & Edwards, GE (1990). Quantum yields of photosystem II electron transport and carbon dioxide fixation in C4 plants. *Functional Plant Biology*, 17(5), 579-588.
- Krause, GH, & Weis, E (1991). Chlorophyll fluorescence and photosynthesis: the basics. *Annual review of plant biology*, 42(1), 313-349.
- Latimer, P, Bannister, T, & Rabinowitch, E (1956). Quantum yields of fluorescence of plant pigments. *Science*, 124(3222), 585-586.
- Lichtenthaler, HK, Ač, A, Marek, MV, Kalina, J, & Urban, O (2007). Differences in pigment composition, photosynthetic rates and chlorophyll fluorescence images of sun and shade leaves of four tree species. *Plant Physiology and Biochemistry*, 45(8), 577-588.
- Matsubara, S, Chen, Y, Caliandro, R, & Clegg, RM (2011). Photosystem II fluorescence lifetime imaging in avocado leaves: contributions of the lutein-epoxide and violaxanthin cycles to fluorescence quenching. *Journal of Photochemistry and Photobiology B: Biology*, 104(1-2), 271-284.
- Maxwell, K, & Johnson, GN (2000). Chlorophyll fluorescence—a practical guide. *Journal of experimental botany*, 51(345), 659-668.
- Nedbal, L, & Whitmarsh, J (2004). Chlorophyll fluorescence imaging of leaves and fruits. In *Chlorophyll a Fluorescence* (pp. 389-407). Springer.
- Oláh, V, Hepp, A, Irfan, M, & Mészáros, I (2021). Chlorophyll fluorescence imaging-based duckweed phenotyping to assess acute phytotoxic effects. *Plants*, 10(12), 2763.
- Porcar-Castell, A, Tyystjärvi, E, Atherton, J, Van der Tol, C, Flexas, J, Pfündel, EE, Moreno, J, Frankenberg, C, & Berry, JA (2014). Linking chlorophyll a fluorescence to photosynthesis for remote sensing applications: mechanisms and challenges. *Journal of experimental botany*, 65(15), 4065-4095.
- Ralph, PJ, & Gademann, R (2005). Rapid light curves: a powerful tool to assess photosynthetic activity. *Aquatic botany*, 82(3), 222-237.
- Sang, H, Guo, W, Gao, Y, Jiao, X, & Pan, X (2020). Effects of Alternating Fresh and Saline Water Irrigation on Soil Salinity and Chlorophyll Fluorescence of Summer Maize. *Water*, 12(11), 3054.
- Schansker, G, Srivastava, A, & Strasser, RJ (2003). Characterization of the 820-nm transmission signal paralleling the chlorophyll a fluorescence rise (OJIP) in pea leaves. *Functional Plant Biology*, 30(7), 785-796.
- Schansker, G, Tóth, SZ, & Strasser, RJ (2005). Methylviologen and dibromothymoquinone treatments of pea leaves reveal the role of photosystem I in the Chl a fluorescence rise OJIP. *Biochimica et Biophysica Acta (BBA)-Bioenergetics*, 1706(3), 250-261.
- Schreiber, U, Klughammer, C, & Kolbowski, J (2012). Assessment of wavelength-dependent parameters of photosynthetic electron transport with a new type of multi-color PAM chlorophyll fluorometer. *Photosynthesis Research*, 113(1), 127-144.
- Schreiber, U, Schliwa, U, & Bilger, W (1986). Continuous recording of photochemical and non-photochemical chlorophyll fluorescence quenching with a new type of modulation fluorometer. *Photosynthesis Research*, 10(1), 51-62.
- Snel, JF, Kooijman, M, & Vredenberg, WJ (1990). Correlation between chlorophyll fluorescence and photoacoustic signal transients in spinach leaves. *Photosynthesis Research*, 25(3), 259-268.
- Strasser, RJ, Tsimilli-Michael, M, & Srivastava, A (2004). Analysis of the chlorophyll a fluorescence transient. In *Chlorophyll a fluorescence* (321-362). Springer.
- Swoczyna, T, Kalaji, HM, Pietkiewicz, S, Borowski, J, & Zaráś-Januskiewicz, E (2010). Monitoring young urban trees tolerance to roadside conditions by application of chlorophyll fluorescence. *Zesz. Probl. Postepow Nauk Roln*, 545, 303-309.