

FEATURED ARTICLE

Detecting amyloid positivity in early Alzheimer's disease using combinations of plasma A β 42/A β 40 and p-tau

Shorena Janelidze¹ | Sebastian Palmqvist^{1,2} | Antoine Leuzy¹ | Erik Stomrud^{1,2} |
Inge M.W. Verberk³ | Henrik Zetterberg^{4,5,6,7} | Nicholas J. Ashton^{4,8,9,10} |
Pedro Pesini¹¹ | Leticia Sarasa¹¹ | José Antonio Allué¹¹ | Charlotte E. Teunissen³ |
Jeffrey L. Dage¹² | Kaj Blennow^{4,5} | Niklas Mattsson-Carlgren^{1,13,14} | Oskar Hansson^{1,2}

¹ Clinical Memory Research Unit, Department of Clinical Sciences Malmö, Lund University, Lund, Sweden

² Skåne University Hospital, Malmö, Sweden

³ Neurochemistry Laboratory, Department of Clinical Chemistry, Amsterdam Neuroscience, Vrije Universiteit Amsterdam, Amsterdam UMC, Amsterdam, the Netherlands

⁴ Department of Psychiatry and Neurochemistry, Institute of Neuroscience & Physiology, the Sahlgrenska Academy at the University of Gothenburg, Mölndal, Sweden

⁵ Clinical Neurochemistry Laboratory, Sahlgrenska University Hospital, Mölndal, Sweden

⁶ Department of Neurodegenerative Disease, UCL Institute of Neurology, London, UK

⁷ UK Dementia Research Institute at UCL, London, UK

⁸ Wallenberg Centre for Molecular and Translational Medicine, University of Gothenburg, Gothenburg, Sweden

⁹ King's College London, Institute of Psychiatry, Psychology and Neuroscience, Maurice Wohl Institute Clinical Neuroscience Institute, London, UK

¹⁰ NIHR Biomedical Research Centre for Mental Health and Biomedical Research Unit for Dementia at South London and Maudsley NHS Foundation, London, UK

¹¹ Araclon Biotech Ltd., Zaragoza, Spain

¹² Eli Lilly and Company, Indianapolis, Indiana, USA

¹³ Department of Neurology, Skåne University Hospital, Lund, Sweden

¹⁴ Wallenberg Center for Molecular Medicine, Lund University, Lund, Sweden

Correspondence

Oskar Hansson, Memory Clinic, Skåne University Hospital, S:t Johannesgatan 8, SE-20502 Malmö, Sweden.

E-mail: Oskar.Hansson@med.lu.se

Funding information

Swedish Research Council, Grant/Award Number: 2016-00906; Knut and Alice Wallenberg Foundation, Grant/Award Number: 2017-0383; Marianne and Marcus Wallenberg Foundation, Grant/Award Number: 2015.0125; Strategic Research Area Multi-Park (Multidisciplinary Research in Parkinson's disease) at Lund University; Swedish Alzheimer Foundation, Grant/Award Numbers: AF-745911, AF-846521; Swedish Brain Foundation, Grant/Award Number: FO2019-0326; Parkinson Foundation of Sweden, Grant/Award

Abstract

Introduction: We studied usefulness of combining blood amyloid beta (A β)42/A β 40, phosphorylated tau (p-tau)217, and neurofilament light (NfL) to detect abnormal brain A β deposition in different stages of early Alzheimer's disease (AD).

Methods: Plasma biomarkers were measured using mass spectrometry (A β 42/A β 40) and immunoassays (p-tau217 and NfL) in cognitively unimpaired individuals (CU, N = 591) and patients with mild cognitive impairment (MCI, N = 304) from two independent cohorts (BioFINDER-1, BioFINDER-2).

Results: In CU, a combination of plasma A β 42/A β 40 and p-tau217 detected abnormal brain A β status with area under the curve (AUC) of 0.83 to 0.86. In MCI, the models including p-tau217 alone or A β 42/A β 40 and p-tau217 had similar AUCs (0.86–0.88); however, the latter showed improved model fit. The models were implemented

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial](https://creativecommons.org/licenses/by-nc/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.

© 2021 The Authors. *Alzheimer's & Dementia* published by Wiley Periodicals LLC on behalf of Alzheimer's Association.

Number: 1280/20; Skåne University Hospital Foundation, Grant/Award Number: 2020-O000028; Regionalt Forskningsstöd, Grant/Award Number: 2020-0314; Bundy Academy; Konung Gustaf V:s och Drottning Victorias Frimurarestiftelse; Swedish federal government under the ALF agreement, Grant/Award Number: 2018-Projekt0279; European Commission (Marie Curie International Training Network, JPND); Health Holland; Dutch Research Council (ZonMW); Alzheimer Drug Discovery Foundation; Selfridges Group Foundation; Alzheimer Netherlands; Alzheimer Association

in an online application providing individualized risk assessments (<https://brainapps.shinyapps.io/PredictABplasma/>).

Discussion: A combination of plasma A β 42/A β 40 and p-tau217 discriminated A β status with relatively high accuracy, whereas p-tau217 showed strongest associations with A β pathology in MCI but not in CU.

KEYWORDS

Alzheimer's disease, amyloid, A β 42/A β 40, blood biomarkers, neurofilament light, p-tau217

1 | BACKGROUND

In Alzheimer's disease (AD), the underlying brain amyloid beta (A β) and tau pathologies and ensuing neurodegeneration can be reliably detected and monitored using cerebrospinal fluid (CSF) and imaging biomarkers. CSF concentrations of A β 42 (alone or as a ratio with A β 40) and phosphorylated tau (p-tau) reflect AD-related changes in A β and tau metabolism in the brain.¹ A β and tau positron emission tomography (PET) are used to measure A β plaque load and insoluble paired helical filament (PHF) tau aggregates.^{2,3} Biomarkers of neurodegeneration (due to AD or other causes) include fluorodeoxyglucose (FDG) PET, magnetic resonance imaging (MRI), and CSF neurofilament light (NfL).⁴⁻⁷ The CSF and imaging biomarkers have been successfully used in research settings and in specialized clinics in some countries and were recently incorporated into an ATN (amyloid, tau, neurodegeneration) classification system, a research framework proposed by the National Institute on Aging for the diagnosis of AD.⁸ More accessible and inexpensive methods, like blood tests, however, are needed for widespread applicability in clinical trials as well as for future implementation in routine clinical care. AD biomarker concentrations in blood are low, making their quantification in the presence of other high abundance proteins challenging. However, recent technological advances in mass spectrometry and immunodetection have led to the development of novel methods that have allowed for reliable assessment of A β 42/A β 40, p-tau, and NfL in blood.⁹

Blood A β 42/A β 40 correlates with CSF A β 42/A β 40 and A β -PET and can identify with relatively high precision individuals with abnormal brain A β burden or those at high risk of future conversion to A β -PET positivity.¹⁰⁻¹⁵ When measured in plasma, tau phosphorylated at threonine 217 and 181 (p-tau217 and p-tau181) accurately detect amyloid and tau pathology assessed by PET, differentiate AD from non-AD neurodegenerative disorders, and predict future progression to AD dementia.¹⁶⁻²¹ Plasma levels of p-tau217 have been shown to increase in very early preclinical stages of AD and continue to increase over time in patients with preclinical and prodromal AD.^{22,23} Elevated levels of blood NfL have also been reported in mild cognitive impairment (MCI) and AD dementia stages in sporadic disease and already in pre-symptomatic phases in familial AD,²⁴⁻²⁶ increasing over time in parallel with other signs of neurodegeneration.²⁷ However, NfL is not specific to neurodegeneration in AD but is increased (in both CSF and blood) in many other disorders of the central nervous system, including,

for example, frontotemporal dementia, progressive supranuclear palsy, corticobasal syndrome, amyotrophic lateral sclerosis, and Creutzfeldt-Jakob disease.²⁸⁻³²

Plasma biomarkers have the potential to greatly accelerate the development of effective disease-modifying treatments in AD by facilitating the identification of individuals at the earliest disease stages (i.e., subjects with preclinical or prodromal AD), when the treatments are most likely to be successful. Although plasma AD biomarkers have shown relatively fair accuracy for detecting A β pathology, it remains to be established if a blood test combining these biomarkers could offer improved performance. For example, decrease in plasma A β levels in AD is very modest (15%–20% at most), while plasma p-tau is a more dynamic biomarker that could better mirror progressive increases in brain A β burden. At the same time, because AD biomarkers follow different trajectories with A β 42/A β 40 starting to change first followed next by p-tau and then by NfL,^{33,34} it is very likely that blood A β 42/A β 40 would be the most accurate in the very early disease stages. In the present study, we measured plasma A β 42/A β 40 (Araclon mass spectrometry assay), plasma p-tau217 (Lilly-developed immunoassay), and plasma NfL (Simoa-based immunoassay) in two independent cohorts of cognitively unimpaired (CU) participants (n = 591) and patients with MCI (n = 304). We first assessed the accuracy of individual biomarkers and then different combinations of biomarkers to detect abnormal brain A β status (defined using either CSF A β 42/A β 40 or A β -PET) in the CU subjects and patients with MCI, separately. Finally, we tested whether the accuracy was improved by including information on apolipoprotein E (APOE) ϵ 4 status of the examined individuals.

2 | METHODS

2.1 | Study participants

The study was approved by the Regional Ethics Committee in Lund, Sweden. All participants provided written informed consent. They were recruited in southern Sweden (Skåne University Hospital and the Hospital of Ängelholm) as previously reported.^{16,20} Further details on study design and recruitment procedures are given in the supporting information. From BioFINDER-1 (clinical trial no. NCT01208675), we included 123 cognitively healthy controls, 118 patients with subjective

cognitive decline (SCD), and 140 MCI patients recruited between 2010 and 2015. The BioFINDER-2 cohort (clinical trial no. NCT03174938) comprised 235 cognitively healthy controls, 115 SCD, and 164 MCI patients recruited between 2017 and 2019. In accordance with the research framework by the National Institute on Aging-Alzheimer's Association study patients with SCD and cognitively healthy controls were considered the CU group (BioFINDER-1 N = 241, BioFINDER-2 N = 350).⁸ From both cohorts, we included all participants with available plasma A β 42/A β 40, plasma p-tau217, plasma NfL, and APOE genotype data.

2.2 | Plasma and CSF sampling and analysis

Ethylenediaminetetraacetic acid (EDTA)-plasma and CSF samples were collected and handled as previously described.^{12,16} Plasma p-tau217 concentration was measured according to the published protocols using immunoassay on a Mesoscale Discovery platform developed by Lilly Research Laboratories.^{20,22} BioFINDER-1 and BioFINDER-2 samples were analyzed at the Clinical Memory Research Unit, Lund University (Sweden) and at Lilly Research Laboratories, respectively. Briefly, biotinylated-IBA493 was used as a capture antibody and SULFO-TAG-4G10-E2 (anti-tau) as the detector and samples were diluted 1:2. In BioFINDER-2, the assay was calibrated with a recombinant tau (4R2N) protein that was phosphorylated *in vitro* using a reaction with glycogen synthase kinase-3 and characterized by mass spectrometry, while in BioFINDER-1, we used a synthetic p-tau217 peptide.

Given that prior studies have suggested a potentially better performance of blood A β 42/A β 40 measured with mass spectrometry compared to immunoassays,^{10–12,14,15} we used mass spectrometry to quantify plasma levels of A β 42 and A β 40 in the present study. Further details of the assays are described in the supporting information.

In BioFINDER-1, plasma NfL concentration was measured using Simoa N4PE kit (Quanterix) at the Neurochemistry Laboratory of the Amsterdam UMC location VUmc (Netherlands). In BioFINDER-2, plasma NfL concentration was measured at the Clinical Neurochemistry Laboratory in Gothenburg using a Simoa kit (Quanterix).

CSF A β 42 and A β 40 concentrations in BioFINDER-1 were quantified at Euroimmun using enzyme-linked immunosorbent assay (ELISA) kits (Euroimmun). CSF A β 42 and A β 40 concentrations in BioFINDER-2 were measured with Meso Scale Discovery immunoassays (MSD) at the Clinical Neurochemistry Laboratory in Gothenburg. CSF A β 42/A β 40 measured using either Euroimmun or MSD assays perform equally well when predicting A β -PET assessment outcome.³⁵ CSF A β 42/A β 40 data were binarized using previously published cutoffs (< 0.091 in BioFINDER-1 and < 0.0752 in BioFINDER-2^{20,36}).

2.3 | A β -PET imaging

A β imaging was performed using [¹⁸F]flutemetamol PET, as described in the supporting information. Briefly, standardized uptake value ratio (SUVR) images were created using dynamic (list-mode) 90- to

RESEARCH IN CONTEXT

- 1. Systematic review:** We searched and reviewed the literature on Alzheimer's disease (AD), blood biomarkers of amyloid and tau pathologies, and neurodegeneration using PubMed. While prior publications suggested that blood amyloid beta (A β)42/A β 40, phosphorylated tau (p-tau), and neurofilament light (NfL) become abnormal very early in the disease course, no studies have investigated which combination of these biomarkers most accurately predicts brain A β pathology in early AD.
- 2. Interpretation:** The main findings from two independent cohorts were that in preclinical and prodromal AD, a combination of plasma A β 42/A β 40 and plasma p-tau217 detected with high precision abnormal brain A β status.
- 3. Future directions:** Blood biomarkers have the potential to greatly accelerate the development of effective disease-modifying treatments in AD by facilitating identification of individuals at the earliest disease stages when the treatments are most likely to be successful. Replication in other cohorts will be needed before the findings of the present study could be implemented in clinical trials.

HIGHLIGHTS

- Brain A β pathology in preclinical and prodromal Alzheimer's disease (AD) was accurately detected by a combination of plasma A β 42/A β 40 and P-tau217.
- In prodromal AD, plasma P-tau217 by itself exhibited high predictive value for A β status.

100-minute post-injection data and the whole cerebellum, pons/brainstem, and eroded cortical white matter as reference region.³⁷ A β -PET status (abnormal/normal) was determined by applying a Gaussian mixture modeling-based cutoff to neocortical SUVR values.

2.4 | Statistical analysis

All analyses were performed using SPSS version 26 (IBM). Differences in baseline demographic and clinical data and biomarker levels were tested with Chi-square and Mann-Whitney tests. Plasma biomarker data were transformed to z-scores based on the distribution in the A β -CU sample. Discrimination accuracies of biomarkers were determined with logistic regression models and receiver operating characteristic (ROC) curve analysis. APOE risk allele status was modeled as one variable coded for the presence of ϵ 4 allele (1 for ϵ 4 carriers and 0 for noncarriers). Improvements in model fit were estimated using Akaike

TABLE 1 Demographic and clinical characteristics

	BioFINDER-2			BioFINDER-1		
	CU	MCI	P-value	CU	MCI	P-value
N	350	164		241	140	
Age, years	64 (53–75)	71 (66–76)	<.0001	72 (68–75)	71 (67–76)	.47
Female no., (%)	183 (52.3)	79 (48.2)	.38	134 (55.6)	50 (35.7)	.0002
Duration of education, years*	12 (10–15)	12 (9–15)	.54	12 (9–14)	11 (9–13)	0.006
MMSE	29 (28–30)	27 (25–29)	<.0001	29 (28–30)	27 (26–29)	<.0001
APOE ε4 positivity No., %	155 (44.3)	87 (53)	.06	90 (37.3)	73 (52.1)	.005
CSF Aβ42/Aβ0†	1.00 (0.78–1.12)	0.69 (0.50–1.04)	<.0001	0.12 (0.08–0.14)	0.07 (0.05–0.13)	<.0001
CSF Aβ42/Aβ0 positivity, No., (%)	81 (23.1)	89 (54.3)	<.0001	78 (32.4)	86 (61.4)	<.0001
Aβ-PET, [¹⁸ F]Flutemetamol SUVR neocortical meta-ROI*	0.62 (0.59–0.66)	0.72 (0.61–0.72)	<.0001	0.67 (0.64–0.77)	0.93 (0.70–1.09)	<.0001
Aβ-PET positivity No., %	75 (22.0)	83 (52.9)	<.0001	673 (29.1)	86 (66.2)	<.0001
Plasma Aβ42/Aβ0	0.22 (0.20–0.25)	0.21 (0.19–0.24)	.044	0.31 (0.29–0.34)	0.29 (0.27–0.32)	.0002
Plasma p-tau217 pg/mL‡	0.94 (0.31–1.78)	1.66 (0.43–3.70)	<.0001	0.15 (0.07–0.24)	0.27 (0.13–0.39)	<.0001
Plasma NfL pg/mL‡	12.13 (8.30–16.96)	16.35 (11.60–22.95)	<.0001	27.58 (21.61–36.45)	35.00 (26.87–45.32)	<.0001

Notes: Data are shown median (interquartile range) unless otherwise specified; P values are from Chi-square (sex, APOE ε4 status, CSF Aβ42/Aβ0 status, Aβ-PET status), Mann-Whitney tests (age, duration of education, MMSE) and univariate analysis of variance adjusted for age and sex (plasma biomarkers, CSF Aβ42/Aβ0, [¹⁸F]Flutemetamol SUVR).

*In BioFINDER-2, education was missing for 1 MCI patient and Aβ-PET was not available for 16 participants; in BioFINDER-1, education was missing for 3 MCI patients and Aβ-PET was not available for 21 participants.

†Samples from BioFINDER-1 and BioFINDER-2 were analyzed using different assays (as described in the Materials and Methods section) and biomarker concentrations are therefore not comparable across the cohorts.

Abbreviations: Aβ, amyloid beta; APOE, apolipoprotein E; CSF, cerebrospinal fluid; CU, cognitively unimpaired; MCI, mild cognitive impairment; MMSE, Mini-Mental State Examination; NfL, neurofilament light; p-tau, phosphorylated tau; ROI, region of interest; SUVR, standardized uptake value ratio

information criterion (AIC) with a decrease of 2 or more in AIC indicating better model fit.³⁸ The AIC values were transformed to the Akaike weights that can be interpreted as the probability that the respective model is the most correct among the candidate models in the given sample.³⁹ The performance of plasma biomarkers was assessed separately in BioFINDER-1 and BioFINDER-2. We also performed external validation across cohorts by testing the model estimates derived in BioFINDER-2 on BioFINDER-1 and vice versa. The fitted models were used to create an online application that calculates individualized probability for Aβ positivity. Area under the curve (AUC) of two ROC curves were compared to DeLong test with adjustment for multiple comparisons using the Benjamini-Hochberg method and a false discovery rate (FDR) of 5%. The FDR correction was applied separately for the analysis in the BioFINDER-1 and BioFINDER-2 cohorts. Statistical significance was set at $P < .05$. Out of 895 study participants, 334 had plasma p-tau217 levels below the detection limit of the assay. When it was not possible to interpolate plasma p-tau217 concentrations from the standard curve due to the very low signal, the values were imputed to the lowest measurable value. Out of 334 samples below the detection limit, plasma p-tau217 values were only imputed for 95 cases (10% of the whole study population). Almost all imputed data (97%, $N = 92$) were in the Aβ- group. Furthermore, the large majority of

the samples below the detection limit (87%, $N = 289$) were also in the Aβ- group. Given that data below the detection limit were confined to the Aβ- group, these values were considered to represent truly very low p-tau217 concentrations and were included in all statistical analyses. Results from the larger BioFINDER-2 cohort are presented first.

3 | RESULTS

3.1 | Participant characteristics

The BioFINDER-2 cohort included 350 CU participants (269 CSF Aβ-; 81 CSF Aβ+) and 164 patients with MCI (75 CSF Aβ-; 89 CSF Aβ+). The BioFINDER-1 cohort included 241 CU participants (163 CSF Aβ-; 78 CSF Aβ+) and 140 patients with MCI (54 CSF Aβ-; 86 CSF Aβ+). The majority of participants in BioFINDER-2 ($n = 498$) and BioFINDER-1 ($n = 360$) also underwent Aβ-PET. Baseline demographic and clinical characteristics of both cohorts are summarized in Table 1 and Tables S1 and S2 in supporting information. In BioFINDER-2, CU participants were on average younger than MCI participants. In BioFINDER-1, the CU group included more men than the MCI group. In both cohorts,




FIGURE 1 Receiver operating characteristic (ROC) curve analyses for discriminating cerebrospinal fluid (CSF) amyloid beta ($A\beta$)42/ $A\beta$ 40 status. ROC curves are shown for plasma $A\beta$ 42/ $A\beta$ 40, plasma phosphorylated tau (p-tau)217, plasma neurofilament light (NfL), a combination of plasma $A\beta$ 42/ $A\beta$ 40 and plasma p-tau 217, and the full model including all three plasma biomarkers (plasma $A\beta$ 42/ $A\beta$ 40, plasma P-tau 217, and plasma NfL). AUC, area under the curve; CU, cognitively unimpaired; MCI, mild cognitive impairment

the CSF and plasma biomarkers as well as $A\beta$ -PET measures were more abnormal in MCI patients compared to CU individuals (Table 1). Plasma biomarker concentrations are shown in Figure S1 in supporting information.

3.2 | Detecting abnormal CSF $A\beta$ status in the BioFINDER-2 cohort

3.2.1 | CU participants

In CU participants, univariate associations with abnormal CSF $A\beta$ 42/ $A\beta$ 40 status were stronger for plasma $A\beta$ 42/ $A\beta$ 40 (AUC = 0.79 [confidence interval (CI) 0.73–0.84]; odds ratio [OR] = 0.26, $P < .0001$) and plasma p-tau217 (AUC = 0.81 [CI 0.75–0.86]; OR = 2.49, $P < .0001$) than for plasma NfL (AUC = 0.70 [CI 0.65–0.76], OR = 1.71, $P < .0001$; Table 2). The model combining plasma p-tau217 and plasma $A\beta$ 42/ $A\beta$ 40 showed high discriminative accuracy with an

AUC of 0.86 (CI 0.82–0.91) and was not significantly different from the model including all three plasma biomarkers (Δ AUC = 0.005, $P = .61$), although the AIC indicated somewhat better fit for the three-biomarker model (Table 2, Figure 1A).

3.2.2 | MCI patients

In MCI patients, plasma p-tau217 (AUC = 0.88 [CI 0.83–0.94]; OR = 2.29, $P < .0001$) outperformed plasma $A\beta$ 42/ $A\beta$ 40 (AUC = 0.70 [CI 0.62–0.78]; OR = 0.48, $P = .0001$), while plasma NfL was not significant (AUC = 0.54, CI [0.45–0.63]; OR = 0.97, $P = .68$) to detect $A\beta$ positivity (Table 3). The plasma p-tau217 model was noninferior to the model with all three plasma biomarkers (Δ AUC = 0.00, $P = .95$) and to the model including plasma p-tau217 and plasma $A\beta$ 42/ $A\beta$ 40 (Δ AUC = 0.00, $P = .95$), even though the AIC suggested that the model combining plasma p-tau217 and plasma $A\beta$ 42/ $A\beta$ 40 fit the data better than all the other models (Table 3 and Figure 1B).

TABLE 2 Associations with CSF A β 42/A β 40 status in CU individuals in BioFINDER-2 and BioFINDER-1

Model	Odds ratio (P-value)				P-value vs. full plasma model*	AIC (Δ AIC) vs. full plasma model	wAIC
	A β 42/A β 40	p-tau217	NfL	AUC (95% CI)			
BioFINDER-2[†]							
A β 42/A β 40, p-tau217, NfL	0.28 ($P < .0001$)	2.23 ($P < .0001$)	1.48 ($p = 0.008$)	0.868 [0.822, 0.914]	NA	283 (ref)	0.92
A β 42/A β 40, p-tau217	0.27 ($P < .0001$)	2.42 ($P < .0001$)	NA	0.863 [0.817, 0.910]	.61	288 (5)	0.08
A β 42/A β 40	0.26 ($P < .0001$)	NA	NA	0.786 [0.732, 0.841]	<.0001	314 (31)	1.7e-07
p-tau217	NA	2.49 ($P < .0001$)	NA	0.805 [0.748, 0.862]	.030	343 (60)	8.6e-14
NfL	NA	NA	1.71 ($p < 0.0001$)	0.704 [0.646, 0.762]	<.0001	363 (80)	3.9e-18
BioFINDER-1[‡]							
A β 42/A β 40, p-tau217, NfL	0.34 ($P < .0001$)	2.11 ($P < .0001$)	1.29 ($P = .094$)	0.837 [0.782, 0.891]	NA	228 (ref)	0.62
A β 42/A β 40, p-tau217	0.33 ($P < .0001$)	2.20 ($P < .0001$)	NA	0.833 [0.778, 0.888]	.60	229 (1)	0.38
A β 42/A β 40	0.33 ($P < .0001$)	NA	NA	0.790 [0.730, 0.851]	.08	260 (31)	7.0e-08
p-tau217	NA	2.10 ($P < .0001$)	NA	0.731 [0.664, 0.798]	.001	270 (42)	4.7e-10
NfL	NA	NA	1.50 ($P = .0011$)	0.639 [0.565, 0.713]	<.0001	295 (66)	1.8e-15

Notes: Data are from logistic regression models with binarized CSF A β 42/A β 40 status as outcome. For plasma biomarkers, odds ratios represent increased risk of CSF A β 42/A β 40 positivity for each SD change in biomarker value. Δ AIC, the difference between the AIC values of the reference model and other models; wAIC, the Akaike weight for a given model calculated from Δ AIC.

*P-values (adjusted for multiple comparisons) are for comparisons of AUCs (using DeLong test) between the full model (A β 42/A β 40, p-tau217, NfL) and other models.

[†]Out of 350 CU participants, 269 were classified as CSF A β 42/A β 40 negative and 81 were classified as CSF A β 42/A β 40 positive.

[‡]Out of 241 CU participants, 163 were classified as CSF A β 42/A β 40 negative and 78 were classified as CSF A β 42/A β 40 positive.

Abbreviations: A β , amyloid beta; AIC, Akaike information criterion; APOE, apolipoprotein E; AUC, area under the curve; CI, confidence interval; CSF, cerebrospinal fluid; CU, cognitively unimpaired; NfL, neurofilament light; p-tau, phosphorylated tau; SD, standard deviation

3.3 | Validation in the BioFINDER-1 cohort and across the cohorts

3.3.1 | CU participants in BioFINDER-1

In CU participants, plasma A β 42/A β 40 (AUC = 0.79 [CI 0.73–0.85], OR = 0.33, $P < .0001$) and plasma p-tau217 (AUC = 0.73 [CI 0.66–0.80]; OR = 2.10, $P < .0001$) were more strongly associated with abnormal CSF A β 42/A β 40 status than plasma NfL (AUC = 0.64 [CI 0.56–0.71], OR = 1.50, $P = .001$; Table 2). Just as in BioFINDER-2, the model combining plasma A β 42/A β 40 and plasma p-tau217 showed high discriminative accuracy (AUC = 0.83 [CI 0.78–0.89]), which was non-inferior to the model including all three plasma biomarkers in terms of both AUC or AIC (Δ AUC = 0.004, $P = .60$; Δ AIC = 1; Table 2 and Figure 1C).

3.3.2 | MCI patients in BioFINDER-1

In MCI patients, plasma p-tau217 (AUC 0.86 [CI 0.80–0.92]; OR = 3.28, $P < .0001$) outperformed plasma A β 42/A β 40 (AUC 0.71 [CI 0.63–0.80]; OR = 0.44, $P = .0003$), while plasma NfL was not significant (AUC 0.60, CI [0.50–0.71]; OR = 1.00, $P = .98$; Table 3). Similar to what was seen in BioFINDER-2, the AUC for the model including plasma p-tau217 was noninferior to the models including all three plasma biomarkers (Δ AUC = 0.014, $P = .31$) or plasma A β 42/A β 40 and plasma p-tau217

(Δ AUC = 0.014, $P = .31$) with somewhat better model fit for the last two models (Table 3 and Figure 1D).

3.3.3 | Validation across cohorts

To assess the generalizability of the findings of the present study to wider samples we performed external cross-validation by testing the model fit from BioFINDER-2 in BioFINDER-1 and vice versa. Using this approach, we again observed that a combination of plasma A β 42/A β 40 and plasma p-tau217 could discriminate CSF A β 42/A β 40 status in CU with AUCs of 0.83 to 0.87 and in MCI with AUCs of 0.87 to 0.89 and no further improvement was seen when adding plasma NfL to the models (Table 4).

3.4 | Added value of APOE in the BioFINDER-1 and 2 cohorts

Adding APOE ϵ 4 status improved model fit as determined using AIC for all tested plasma biomarkers and biomarker combinations (Tables S3–6 in supporting information). However, in both cohorts, there were no significant differences in AUCs when adding APOE ϵ 4 to the combinations of plasma A β 42/A β 40 and plasma p-tau217 in CU participants and in MCI patients (Tables S3–6).

TABLE 3 Associations with CSF Aβ42/Aβ40 status in MCI patients in BioFINDER-2 and BioFINDER-1

Model	Odds ratio (P-value)			AUC (95% CI)	P-value vs. full plasma model*	AIC (ΔAIC) vs. full plasma model	wAIC
	Aβ42/Aβ40	p-tau217	NfL				
BioFINDER-2†							
Aβ42/Aβ40, p-tau217, NfL	0.42 (P = .0001)	2.32 (P < .0001)	0.99 (P = .89)	0.879 [0.827, 0.931]	NA	182 (ref)	0.27
Aβ42/Aβ40, p-tau217	0.42 (P = .0001)	2.32 (P < .0001)	NA	0.880 [0.828, 0.932]	.61	180 (-2)	0.73
Aβ42/Aβ40	0.48 (P = .0001)	NA	NA	0.703 [0.621, 0.784]	<.0001	213 (31)	5.0e-08
p-tau217	NA	2.29 (P < .0001)	NA	0.882 [0.828, 0.936]	.95	195 (13)	0.0004
NfL	NA	NA	0.97 (P = .68)	0.538 [0.448, 0.628]	<.0001	230 (48)	1.0e-11
BioFINDER-1‡							
Aβ42/Aβ40, p-tau217, NfL	0.48 (P = .0072)	3.20 (P < .0001)	0.88 (P = .45)	0.877 [0.821, 0.934]	NA	124 (ref)	0.36
Aβ42/Aβ40, p-tau217	0.50 (P = .0096)	3.19 (P < .0001)	NA	0.877 [0.820, 0.934]	.86	123 (-1)	0.59
Aβ42/Aβ40	0.44 (P = .0003)	NA	NA	0.714 [0.628, 0.801]	.0005	176 (51)	1.8e-12
p-tau217	NA	3.28 (P < .0001)	NA	0.863 [0.803, 0.924]	.31	128 (4)	0.05
NfL	NA	NA	1.00 (P = .98)	0.603 [0.500, 0.705]	<.0001	191 (67)	1.0e-15

Notes: Data are from logistic regression models with binarized CSF Aβ42/Aβ40 status as outcome. For plasma biomarkers, odds ratios represent increased risk of CSF Aβ42/Aβ40 positivity for each SD change in biomarker value. ΔAIC, the difference between the AIC values of the reference model and other models; wAIC, the Akaike weight for a given model calculated from ΔAIC.

*P-values (adjusted for multiple comparisons) are for comparisons of AUCs (using DeLong test) between the full model (Aβ42/Aβ40, p-tau217, NfL) and other models.

†Out of 164 MCI patients, 75 were classified as CSF Aβ42/Aβ40 negative and 89 were classified as CSF Aβ42/Aβ40 positive.

‡Out of 140 MCI patients, 54 were classified as CSF Aβ42/Aβ40 negative and 86 were classified as CSF Aβ42/Aβ40 positive.

Abbreviations: Aβ, amyloid beta; AIC, Akaike information criterion; APOE, apolipoprotein E; AUC, area under the curve; CI, confidence interval; CSF, cerebrospinal fluid; CU, cognitively unimpaired; MCI, mild cognitive impairment; NfL, neurofilament light; p-tau, phosphorylated tau; SD, standard deviation.

TABLE 4 Associations with CSF Aβ42/Aβ40 status, external validation across cohorts

Model	BioFINDER-1*		BioFINDER-2†	
	AUC (95% CI)	P-value vs. full plasma model‡	AUC (95% CI)	P-value vs. full plasma model‡
CU				
Aβ42/Aβ40, p-tau217, NfL	0.837 [0.783, 0.892]	NA	0.870 [0.824, 0.916]	NA
Aβ42/Aβ40, p-tau217	0.834 [0.779, 0.889]	.73	0.866 [0.819, 0.912]	.61
Aβ42/Aβ40	0.790 [0.730, 0.851]	.08	0.786 [0.732, 0.841]	.0001
p-tau217	0.731 [0.664, 0.798]	.018	0.805 [0.748, 0.862]	.026
NfL	0.639 [0.565, 0.713]	<.0001	0.704 [0.646, 0.762]	<.0001
MCI				
Aβ42/Aβ40, p-tau217, NfL	0.872 [0.814, 0.930]	NA	0.890 [0.840, 0.940]	NA
Aβ42/Aβ40, p-tau217	0.872 [0.814, 0.930]	.86	0.895 [0.846, 0.943]	.61
Aβ42/Aβ40	0.714 [0.628, 0.801]	.0001	0.703 [0.621, 0.784]	<.0001
p-tau217	0.863 [0.803, 0.924]	.70	0.882 [0.828, 0.936]	.79
NfL	0.603 [0.500, 0.705]	<.0001	0.538 [0.448, 0.629]	<.0001

Notes: Data are from logistic regression models with binarized CSF Aβ42/Aβ40 status as outcome.

*Regression estimates from the models fit with the data from BioFINDER-2 were tested in BioFINDER-1.

†Regression estimates from the models fit with the data from BioFINDER-1 were tested in BioFINDER-2.

‡P-values (adjusted for multiple comparisons) are for comparisons of AUCs (using DeLong test) between the full model (Aβ42/Aβ40, p-tau217, NfL) and other models.

Abbreviations: Aβ, amyloid beta; AIC, Akaike information criterion; APOE, apolipoprotein E; AUC, area under the curve; CI, confidence interval; CSF, cerebrospinal fluid; CU, cognitively unimpaired; MCI, mild cognitive impairment; NfL, neurofilament light; p-tau, phosphorylated tau.

Subject information

Cognitive status
Cognitively unimpaired

APOE ϵ 4
Not available


Plasma P-tau (z) (higher is worse)
-0.6 3.6 5

Plasma A β 42/A β 40 (z) (lower is worse)
-3 -1.2 3

Plot options
 95% prediction interval (graphic)
 95% prediction interval (numeric)
 Mean estimates (numeric)

RESULTS

INFO



Subject information

Cognitive status
Cognitively unimpaired

APOE ϵ 4
Not available

Plasma P-tau (z) (higher is worse)
-0.6 0.3 5

Plasma A β 42/A β 40 (z) (lower is worse)
-3 0.9 3

Plot options
 95% prediction interval (graphic)
 95% prediction interval (numeric)
 Mean estimates (numeric)

RESULTS

INFO




FIGURE 2 Individualized probability for A β positivity. Implementation of logistic regression model from BioFINDER-2 at <https://brainapps.shinyapps.io/PredictABplasma/>. The online application allows user to enter diagnosis (CU, MCI), plasma A β 42/A β 40 and p-tau217 values (z-scores) and APOE ϵ 4 (1 for ϵ 4 carriers, 0 for noncarriers, NA for not available) and based on these data calculates individualized probability for A β positivity. For example, a CU individual with no APOE ϵ 4 status available and z-score values of -1.2 and 3.6 for plasma A β 42/A β 40 and p-tau217, respectively, has 93% probability of being CSF A β 42/A β 40 positive; whereas a CU individual with no APOE ϵ 4 status available and z-score values of 0.9 and 0.3 for plasma A β 42/A β 40 and p-tau217, respectively, has 5% probability of being CSF A β 42/A β 40 positive. A β , amyloid beta; APOE, apolipoprotein E; CU, cognitively unimpaired; MCI, mild cognitive impairment; NfL, neurofilament light; p-tau, phosphorylated tau

3.5 | Estimating individualized probability for A β positivity

Individual predicted probabilities from logistic regression models in BioFINDER-2 and BioFINDER-1 are shown in Figures S2 and S3 in supporting information. Given that a combination of plasma A β 42/A β 40 and plasma p-tau217 showed the best performance with respect to AUC and/or AIC, we implemented the two biomarker fitted model

(with or without APOE ϵ 4 status) from BioFINDER-2 as an online tool (Figure 2, <https://brainapps.shinyapps.io/PredictABplasma/>). The online application allows the user to enter diagnosis (CU, MCI), plasma A β 42/A β 40 and p-tau217 values (z-scores), and APOE ϵ 4 (1 for ϵ 4 carriers, 0 for noncarriers, NA for not available) and based on these data calculates individualized probability for CSF A β positivity. For example, a CU individual with no APOE ϵ 4 status available and z-score values of -1.2 and 3.6 for plasma A β 42/A β 40 and p-tau217, respectively, has

93% probability of being CSF A β 42/A β 40 positive; whereas a CU individual with no APOE ϵ 4 status available and z-score values of 0.9 and 0.3 for plasma A β 42/A β 40 and p-tau217, respectively, has 5% probability of being CSF A β 42/A β 40 positive (Figure 2). We added an option of entering APOE ϵ 4 status because the models including APOE ϵ 4 fit the data better compared to the models without APOE ϵ 4 and because previous studies have shown better performance of plasma A β 42/A β 40 when combined with APOE ϵ 4 status.^{12,13}

3.6 | Associations with A β -PET in the BioFINDER-1 and 2 cohorts

The performances of the plasma biomarkers were very similar using A β -PET status as outcome instead of CSF A β 42/A β 40. A combination of plasma A β 42/A β 40 and plasma p-tau217 could predict A β -PET status in CU with AUCs of 0.82 to 0.84 and in MCI with AUCs of 0.86 to 0.91 (Tables S7–S10 in supporting information). The AUCs for these models were not statistically different from the AUCs of the full three-biomarker models in the same study sample (Tables S7–S10). In both cohorts, there were no significant changes in AUCs when adding APOE ϵ 4 to the most models including plasma A β 42/A β 40 and plasma p-tau217 as predictors (Tables S7–S10).

Voxel-based analysis revealed strong associations of A β -PET retention in especially medial frontoparietal regions with the combination of plasma A β 42/A β 40 and plasma p-tau217 in CU participants and with plasma p-tau217 in MCI patients (Figure S4 in supporting information).

4 | DISCUSSION

In this study including two independent cohorts, we explored the utility of currently available plasma biomarkers to detect A β pathology at different disease stages. We show that in CU individuals, plasma A β 42/A β 40 and plasma p-tau217 discriminated A β status more accurately than plasma NfL and that the best performing model included plasma A β 42/A β 40 and plasma p-tau217 with no added value of plasma NfL. In patients with MCI, plasma p-tau217 was superior to plasma A β 42/A β 40 and plasma NfL. Adding plasma A β 42/A β 40 and plasma NfL did not significantly improve the performance of p-tau217 in terms of AUC. However, the models combining plasma A β 42/A β 40 and p-tau217 fit the data better than the model including plasma p-tau217 by itself or all three biomarkers. Adding APOE ϵ 4 status did not result in significantly better discriminative accuracy, even though the model fits were improved. The findings were consistent using either CSF A β 42/A β 40 or A β -PET status as outcome.

Previous research suggested that plasma A β 42/A β 40 can be used to detect pathological CSF A β 42/A β 40 and A β -PET scans across different disease stages, especially when combined with APOE genotype.^{10–12,14,15} Recent data have also indicated that plasma p-tau can accurately discriminate abnormal versus normal A β -PET status.^{16,18,20,21} Here we show that when measured in the same cohorts, plasma A β 42/A β 40 and plasma p-tau217 identified CU indi-

viduals with abnormal A β status with similar precision (A β 42/A β 40, AUC 0.79; p-tau217, AUC 0.73–0.81). Furthermore, this is the first study to demonstrate in CU participants that the combination of A β 42/A β 40 and plasma p-tau217 improved discriminative accuracy with AUCs reaching 0.83 to 0.86 in the two independent cohorts. In contrast, in MCI patients, plasma p-tau217 (AUCs 0.86–0.88) outperformed plasma A β 42/A β 40 (AUCs 0.70–0.71) and there was no further increase in AUC when combining the two biomarkers. These findings are not surprising given prior data suggesting differences in the biomarker dynamics in AD. While in both CSF and blood, A β 42/A β 40 start to change before p-tau,^{34,40,41} p-tau levels continue to increase over the course of AD^{23,42} and the magnitude of this increase (especially for p-tau217) is considerably larger compared to the drop in A β biomarkers levels.^{10,12,20} Thus, it is likely that plasma A β 42/A β 40 captures CU individuals in the earliest stages of the disease leading to an improved performance of combined plasma A β 42/A β 40 and plasma p-tau217 measures in preclinical AD.

The discriminative accuracy of plasma A β 42/A β 40 (quantified using mass spectrometry) was somewhat lower compared to some of the previous mass spectrometry findings,^{10,11} but not others.⁴³ It remains to be seen whether differences in the performance of plasma A β 42/A β 40 between the present and other studies and between the CU and MCI groups that we report here are cohort specific. For this, head-to-head comparisons of available blood A β assays across different diagnostic groups would be needed.

We did not find any improvements in AUC combining plasma A β 42/A β 40 and plasma p-tau217 with plasma NfL. In CU participants in BioFINDER-2, the models including all three biomarkers fit the data better compared to a combination of plasma A β 42/A β 40 and plasma p-tau217. However, in the same group in BioFINDER-1, plasma NfL was not a significant predictor of CSF A β status in the models including all three biomarkers. These results indicate that while the performance of plasma A β 42/A β 40 and plasma p-tau217 was consistent across the cohorts, that was not the case for NfL and that the effects of NfL were small (if any). NfL is a biomarker of axonal injury and neuronal loss and higher plasma levels of this biomarker are associated with faster rates of atrophy on MRI and cognitive deterioration.^{25,27} When combined, plasma p-tau and plasma NfL were reported to better predict longitudinal changes in Mini-Mental State Examination and conversion to AD dementia in patients with MCI.⁴⁴ Thus, while useful in the evaluation of other neurodegenerative and acute brain disorders, in patients with suspected early AD, plasma NfL might be more suitable to predict disease progression rather than as a biomarker linked to A β pathology.

In line with published data,^{12,13} we found that the models combining plasma biomarkers and APOE ϵ 4 status fit the data better than the corresponding models without APOE ϵ 4 status when using AIC. However, adding APOE ϵ 4 did not significantly improve the AUCs of the best performing models. While information on APOE genotype might potentially improve performance of the plasma AD biomarkers, it is important to consider that APOE ϵ 4 does not reflect A β status but merely indicates disease risk and that its use for patient screening and selection in clinical trials may lead to biased inclusion of APOE ϵ 4 carriers.

This study has several limitations. Although the performance of the plasma biomarkers was validated across independent BioFINDER-1 and BioFINDER-2 cohorts, these are specialized cohorts that could have characteristics distinguishing them from other specialized cohorts. Therefore, it is important that our findings are replicated in a more heterogeneous population-based sample and within the intended population in primary care.^{45,46} Plasma levels of p-tau217 were measured using a research grade assay, which could be one explanation for the slight difference in AUCs between the BioFINDER-1 and 2 cohorts. For some cases, p-tau217 concentrations were below the detection limit of the assay and a more sensitive assay on a fully automated platform is needed to reliably measure plasma p-tau217 at low concentration. Nonetheless, as long as individuals with levels below the detection limit represent truly low values (as previously shown for this assay²⁰), this insufficient sensitivity will not strongly affect the accuracy for detecting A β pathology. Finally, implementation of blood-based biomarkers would require standardization of pre-analytical and analytical procedures and development of the certified reference materials. However, to provide an example of potential clinical utility of plasma biomarkers we built an online application in which the user could obtain individualized probability of A β positivity after entering diagnosis, plasma A β 42/A β 40 and p-tau217 values, and optionally APOE ϵ 4 status.

To conclude, we show that the presence of A β pathology in early AD could be effectively detected by combining plasma measurements of A β 42/A β 40 and p-tau217. In patients with MCI, plasma p-tau217 exhibited the highest predictive value for A β status compared to other biomarkers. These findings will aid the implementation of plasma biomarkers in clinical practice and drug trials.

ACKNOWLEDGMENTS

The study was supported by the Swedish Research Council (2016-00906); the Knut and Alice Wallenberg Foundation (2017-0383); the Marianne and Marcus Wallenberg Foundation (2015.0125); the Strategic Research Area MultiPark (Multidisciplinary Research in Parkinson's disease) at Lund University, the Swedish Alzheimer Foundation (AF-745911, AF-846521); the Swedish Brain Foundation (FO2019-0326); The Parkinson Foundation of Sweden (1280/20); the Skåne University Hospital Foundation (2020-O000028); Regionalt Forskningsstöd (2020-0314); The Bundy Academy; the Konung Gustaf V:s och Drottning Victorias Frimurarestiftelse; and the Swedish federal government under the ALF agreement (2018-Projekt0279), the European Commission (Marie Curie International Training Network, JPND), Health Holland, the Dutch Research Council (ZonMW), Alzheimer Drug Discovery Foundation, The Selfridges Group Foundation, Alzheimer Netherlands, Alzheimer Association. The funding sources had no role in the design and conduct of the study; in the collection, analysis, interpretation of the data; or in the preparation, review, or approval of the manuscript. The precursor of [¹⁸F]flutemetamol was provided by GE Healthcare. Analysis of plasma p-tau217 and A β 42/A β 40 were performed at Eli Lilly and Company and Araclon Biotech Ltd, respectively. Analysis of CSF A β 42/A β 40 was performed at Euroimmun.

CONFLICT OF INTEREST

SJ, SP, AL, NMC, ES, IMVV, NJA report no conflicts of interest. HZ has served on scientific advisory boards for Denali, Roche Diagnostics, Wave, Samumed, Siemens Healthineers, Pinteon Therapeutics, and CogRx; has given lectures in symposia sponsored by Fujirebio, Alzecure, and Biogen; and is a co-founder of Brain Biomarker Solutions in Gothenburg AB (BBS), which is a part of the GU Ventures Incubator Program. KB has served as a consultant, on advisory boards, or on data monitoring committees for Abcam, Axon, Biogen, JOMDD/Shimadzu, Julius Clinical, Lilly, MagQu, Novartis, Roche Diagnostics, and Siemens Healthineers, and is a co-founder of Brain Biomarker Solutions in Gothenburg AB (BBS), which is a part of the GU Ventures Incubator Program. PP, LS, and JAA are full-time employees at Araclon Biotech-Grifols. CET has a collaboration contract with ADx Neurosciences and Quanterix, performed contract research or received grants from AC-Immune, Axon Neurosciences, Biogen, Brainstorm Therapeutics, Celgene, EIP Pharma, Eisai, Roche, Toyama, Vivoryon. JLD is a full-time employee of Eli Lilly and Company. OH has acquired research support (for the institution) from AVID Radiopharmaceuticals, Biogen, Eli Lilly, Eisai, GE Healthcare, Pfizer, and Roche. In the past 2 years, he has received consultancy/speaker fees from AC Immune, Alzpath, Biogen, Cerveau, and Roche.

REFERENCES

1. Blennow K, Hampel H, Weiner M, Zetterberg H. Cerebrospinal fluid and plasma biomarkers in Alzheimer disease. *Nat Rev Neurol*. 2010;6:131-144.
2. Chetelat G, Arbizu J, Barthel H, et al. Amyloid-PET and (18)F-FDG-PET in the diagnostic investigation of Alzheimer's disease and other dementias. *Lancet Neurol*. 2020;19:951-962.
3. Ossenkoppele R, Rabinovici GD, Smith R, et al. Discriminative accuracy of [¹⁸F]flortaucipir positron emission tomography for Alzheimer disease vs other neurodegenerative disorders. *JAMA*. 2018;320:1151-1162.
4. Dickerson BC, Bakkour A, Salat DH, et al. The cortical signature of Alzheimer's disease: regionally specific cortical thinning relates to symptom severity in very mild to mild AD dementia and is detectable in asymptomatic amyloid-positive individuals. *Cereb Cortex*. 2009;19:497-510.
5. Fox NC, Crum WR, Scahill RI, Stevens JM, Janssen JC, Rossor MN. Imaging of onset and progression of Alzheimer's disease with voxel-compression mapping of serial magnetic resonance images. *Lancet*. 2001;358:201-205.
6. Landau SM, Harvey D, Madison CM, et al. Associations between cognitive, functional, and FDG-PET measures of decline in AD and MCI. *Neurobiol Aging*. 2011;32:1207-1218.
7. Zetterberg H, Skillback T, Mattsson N, et al. Association of cerebrospinal fluid neurofilament light concentration with Alzheimer disease progression. *JAMA Neurol*. 2016;73:60-67.
8. Jack CR Jr, Bennett DA, Blennow K, et al. NIA-AA research framework: toward a biological definition of Alzheimer's disease. *Alzheimers Dement*. 2018;14:535-562.
9. Simren J, Ashton NJ, Blennow K, Zetterberg H. An update on fluid biomarkers for neurodegenerative diseases: recent success and challenges ahead. *Curr Opin Neurobiol*. 2020;61:29-39.
10. Nakamura A, Kaneko N, Villemagne VL, et al. High performance plasma amyloid-beta biomarkers for Alzheimer's disease. *Nature*. 2018;554:249-254.
11. Ovod V, Ramsey KN, Mawuenyega KG, et al. Amyloid beta concentrations and stable isotope labeling kinetics of human plasma specific to

- central nervous system amyloidosis. *Alzheimers Dement*. 2017;13:841-849.
12. Palmqvist S, Janelidze S, Stomrud E, et al. Performance of fully automated plasma assays as screening tests for Alzheimer disease-related beta-amyloid status. *JAMA Neurol*. 2019;76(9):1060-1069.
 13. Schindler SE, Bollinger JG, Ovod V, et al. High-precision plasma beta-amyloid 42/40 predicts current and future brain amyloidosis. *Neurology*. 2019;93:e1647-e59.
 14. Verberk IMW, Slot RE, Verfaillie SCJ, et al. Plasma amyloid as prescreener for the earliest Alzheimer pathological changes. *Ann Neurol*. 2018;84:648-658.
 15. Vergallo A, Megret L, Lista S, et al. Plasma amyloid beta 40/42 ratio predicts cerebral amyloidosis in cognitively normal individuals at risk for Alzheimer's disease. *Alzheimers Dement*. 2019;15:764-775.
 16. Janelidze S, Mattsson N, Palmqvist S, et al. Plasma P-tau181 in Alzheimer's disease: relationship to other biomarkers, differential diagnosis, neuropathology and longitudinal progression to Alzheimer's dementia. *Nat Med*. 2020;26:379-386.
 17. Karikari TK, Benedet AL, Ashton NJ, et al. Diagnostic performance and prediction of clinical progression of plasma phospho-tau181 in the Alzheimer's disease neuroimaging initiative. *Mol Psychiatry*. 2020.
 18. Karikari TK, Pascoal TA, Ashton NJ, et al. Blood phosphorylated tau 181 as a biomarker for Alzheimer's disease: a diagnostic performance and prediction modelling study using data from four prospective cohorts. *Lancet Neurol*. 2020;19:422-433.
 19. Lantero Rodriguez J, Karikari TK, Suarez-Calvet M, et al. Plasma p-tau181 accurately predicts Alzheimer's disease pathology at least 8 years prior to post-mortem and improves the clinical characterisation of cognitive decline. *Acta Neuropathol*. 2020;140:267-278.
 20. Palmqvist S, Janelidze S, Quiroz YT, et al. Discriminative accuracy of plasma phospho-tau217 for Alzheimer disease vs other neurodegenerative disorders. *JAMA*. 2020;24(8):772-781.
 21. Thijssen EH, La Joie R, Wolf A, et al. Diagnostic value of plasma phosphorylated tau181 in Alzheimer's disease and frontotemporal lobar degeneration. *Nat Med*. 2020;26:387-397.
 22. Janelidze S, Berron D, Smith R, et al. Associations of plasma phospho-tau217 levels with tau positron emission tomography in early Alzheimer disease. *JAMA Neurol*. 2020.
 23. Mattsson-Carlgrén N, Janelidze S, Palmqvist S, et al. Longitudinal plasma p-tau217 is increased in early stages of Alzheimer's disease. *Brain*. 2020.
 24. Mattsson N, Andreasson U, Zetterberg H, Blennow K. Alzheimer's disease neuroimaging I, association of plasma neurofilament light with neurodegeneration in patients with Alzheimer disease. *JAMA Neurol*. 2017;74:557-566.
 25. Weston PSJ, Poole T, Ryan NS, et al. Serum neurofilament light in familial Alzheimer disease: a marker of early neurodegeneration. *Neurology*. 2017;89:2167-2175.
 26. Preische O, Schultz SA, Apel A, et al. Serum neurofilament dynamics predicts neurodegeneration and clinical progression in presymptomatic Alzheimer's disease. *Nat Med*. 2019;25:277-283.
 27. Mattsson N, Cullen NC, Andreasson U, Zetterberg H, Blennow K. Association between longitudinal plasma neurofilament light and neurodegeneration in patients with Alzheimer disease. *JAMA Neurol*. 2019;76:791-799.
 28. Benatar M, Zhang L, Wang L, et al. Validation of serum neurofilaments as prognostic and potential pharmacodynamic biomarkers for ALS. *Neurology*. 2020;95:e59-e69.
 29. Bridel C, van Wieringen WN, Zetterberg H, et al. Diagnostic value of cerebrospinal fluid neurofilament light protein in neurology: a systematic review and meta-analysis. *JAMA Neurol*. 2019.
 30. Hansson O, Janelidze S, Hall S, et al. Blood-based NfL: a biomarker for differential diagnosis of parkinsonian disorder. *Neurology*. 2017;88:930-937.
 31. Rohrer JD, Woollacott IO, Dick KM, et al. Serum neurofilament light chain protein is a measure of disease intensity in frontotemporal dementia. *Neurology*. 2016;87:1329-1336.
 32. Thompson AGB, Luk C, Heslegrave AJ, et al. Neurofilament light chain and tau concentrations are markedly increased in the serum of patients with sporadic Creutzfeldt-Jakob disease, and tau correlates with rate of disease progression. *J Neurol Neurosurg Psychiatry*. 2018;89:955-961.
 33. Jack CR Jr, Knopman DS, Jagust WJ, et al. Tracking pathophysiological processes in Alzheimer's disease: an updated hypothetical model of dynamic biomarkers. *Lancet Neurol*. 2013;12:207-216.
 34. Palmqvist S, Insel PS, Stomrud E, et al. Cerebrospinal fluid and plasma biomarker trajectories with increasing amyloid deposition in Alzheimer's disease. *EMBO Mol Med*. 2019;11:e11170.
 35. Janelidze S, Pannee J, Mikulskis A, et al. Concordance between different amyloid immunoassays and visual amyloid positron emission tomographic assessment. *JAMA Neurol*. 2017;74:1492-1501.
 36. Minta K, Brinkmalm G, Janelidze S, et al. Quantification of total apolipoprotein E and its isoforms in cerebrospinal fluid from patients with neurodegenerative diseases. *Alzheimers Res Ther*. 2020;12:19.
 37. Mattsson-Carlgrén N, Leuzy A, Janelidze S, et al. The implications of different approaches to define AT(N) in Alzheimer disease. *Neurology*. 2020.
 38. Olofson E, Dahan A. Using Akaike's information theoretic criterion in mixed-effects modeling of pharmacokinetic data: a simulation study. *F1000Res*. 2013;2:71.
 39. Wagenmakers EJ, Farrell S. AIC model selection using Akaike weights. *Psychon Bull Rev*. 2004;11:192-196.
 40. McDade E, Wang G, Gordon BA, et al. Longitudinal cognitive and biomarker changes in dominantly inherited Alzheimer disease. *Neurology*. 2018;91:e1295-e306.
 41. Schindler SE, Li Y, Todd KW, et al. Emerging cerebrospinal fluid biomarkers in autosomal dominant Alzheimer's disease. *Alzheimers Dement*. 2019;15:655-665.
 42. Mattsson-Carlgrén N, Andersson E, Janelidze S, et al. A β deposition is associated with increases in soluble and phosphorylated tau that precede a positive Tau PET in Alzheimer's disease. *Science Advances*. 2020;6:eaaz2387.
 43. Keshavan A, Pannee J, Karikari TK, et al. Population-based blood screening for preclinical Alzheimer's disease in a British birth cohort at age 70. *Brain*. 2021.
 44. Cullen NC, Leuzy A, Palmqvist S, et al. Individualized prognosis of longitudinal cognitive decline and dementia in mild cognitive impairment based on plasma biomarker combinations. *Nature Aging*. 2020.
 45. O'Bryant SE, Mielke MM, Rissman RA, et al. Blood-based biomarkers in Alzheimer disease: current state of the science and a novel collaborative paradigm for advancing from discovery to clinic. *Alzheimers Dement*. 2017;13:45-58.
 46. OECD. Emerging trends in biomedicine and health technology innovation: addressing the global challenge of Alzheimer's. 2013.

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section at the end of the article.

How to cite this article: Janelidze S, Palmqvist S, Leuzy A, et al. Detecting amyloid positivity in early Alzheimer's disease using combinations of plasma A β 42/A β 40 and p-tau. *Alzheimer's Dement*. 2022;18:283-293. <https://doi.org/10.1002/alz.12395>