

El Niño as a predictor of round sardinella distribution along the northwest African coast

Jorge **López-Parages**^{1,2}, Pierre-Amaël Auger^{3,4}, Belén Rodríguez-Fonseca^{2,5}, Noel Keenlyside⁶, Carlo Gaetan⁷, Angelo Rubino⁷, Maeregu Woldeyes Arisido⁸, Timothée Brochier^{9,10}

¹Physical Oceanography group, University of Málaga, Málaga, Spain. E-mail: parages@uma.es

²Departamento de Física de la Tierra y Astrofísica, Universidad Complutense de Madrid, Madrid, Spain

³Instituto Milenio de Oceanografía and Escuela de Ciencias del Mar, Pontificia Universidad Católica de Valparaíso, Valparaíso, Chile

⁴Université de Brest, CNRS, IRD, Ifremer, Laboratoire d'Océanographie Physique et Spatiale, IUEM, Brest, France

⁵Instituto de Geociencias IGEO UCM-CSIC, Madrid, Spain

⁶Geophysical Institute, University of Bergen and Bjerknes Centre for Climate Research, Bergen, Norway

⁷Dipartimento di Scienze Ambientali, Informatica e Statistica - Ca' Foscari University of Venice, Venice, Italy

⁸Department of Sociology and Social Research, University of Milano-Bicocca, Milan, Italy

⁹Institut de Recherche pour le Développement (IRD), Sorbonne Université, UMMISCO, Bondy, France

¹⁰Université Cheikh Anta Diop de Dakar, Ecole Supérieure Polytechnique, UMMISCO, Dakar, Senegal

The El Niño Southern Oscillation (ENSO) produces global marine environment conditions that can cause changes in abundance and distribution of distant fish populations worldwide. Understanding mechanisms acting locally on fish population dynamics is crucial to develop forecast skill useful for fisheries management. The present work addresses the role played by ENSO on the round sardinella population biomass and distribution in the central-southern portion of the Canary Current Upwelling System (CCUS). A combined physical-biogeochemical framework is used to understand the climate influence on the hydrodynamical conditions in the study area. Then, an evolutionary individual-based model is used to simulate the round sardinella spatio-temporal biomass variability. According to model experiments, anomalous oceanographic conditions forced by El Niño along the African coast cause anomalies in the latitudinal migration pattern of the species. A robust anomalous increase and decrease of the simulated round sardinella biomass is identified in winter off the Cape Blanc and the Saharan coast region, respectively, in response to El Niño variations. The resultant anomalous pattern is an alteration of the normal migration between the Saharan and the Mauritanian waters. It is primarily explained by the modulating role that El Niño exerts on the currents off Cape Blanc, modifying therefore the normal migration of round sardinella in the search of acceptable temperature conditions. This climate signature can be potentially predicted up to six months in advance based on El Niño conditions in the Pacific.