



LJMU Research Online

Sekar, D, Tusubira, D and Ross, K

TDP-43 and NEAT long non-coding RNA: Roles in neurodegenerative disease

<http://researchonline.ljmu.ac.uk/id/eprint/18176/>

Article

Citation (please note it is advisable to refer to the publisher's version if you intend to cite from this work)

Sekar, D, Tusubira, D and Ross, K (2022) TDP-43 and NEAT long non-coding RNA: Roles in neurodegenerative disease. *Frontiers in Cellular Neuroscience*, 16.

LJMU has developed **LJMU Research Online** for users to access the research output of the University more effectively. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LJMU Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain.

The version presented here may differ from the published version or from the version of the record. Please see the repository URL above for details on accessing the published version and note that access may require a subscription.

For more information please contact researchonline@ljmu.ac.uk

<http://researchonline.ljmu.ac.uk/>



OPEN ACCESS

EDITED BY

Adriana Magalska,
Nencki Institute of Experimental
Biology (PAS), Poland

REVIEWED BY

Anna Konopka,
Macquarie University, Australia
Kunikazu Tanji,
Hirosaki University, Japan
Claudia Fallini,
University of Rhode Island,
United States

*CORRESPONDENCE

Deusedit Tusubira
dtusubira@must.ac.ug
Kehinde Ross
o.k.ross@ijmu.ac.uk

SPECIALTY SECTION

This article was submitted to
Cellular Neurophysiology,
a section of the journal
Frontiers in Cellular Neuroscience

RECEIVED 27 May 2022

ACCEPTED 11 October 2022

PUBLISHED 26 October 2022



CITATION

Sekar D, Tusubira D and Ross K (2022)
TDP-43 and NEAT long non-coding
RNA: Roles in neurodegenerative
disease.
Front. Cell. Neurosci. 16:954912.
doi: 10.3389/fncel.2022.954912

COPYRIGHT

© 2022 Sekar, Tusubira and Ross. This
is an open-access article distributed
under the terms of the [Creative
Commons Attribution License \(CC BY\)](https://creativecommons.org/licenses/by/4.0/).
The use, distribution or reproduction in
other forums is permitted, provided
the original author(s) and the copyright
owner(s) are credited and that the
original publication in this journal is
cited, in accordance with accepted
academic practice. No use, distribution
or reproduction is permitted which
does not comply with these terms.

TDP-43 and NEAT long non-coding RNA: Roles in neurodegenerative disease

Durairaj Sekar¹, Deusedit Tusubira ^{2*} and
Kehinde Ross ^{3,4*}

¹Centre for Cellular and Molecular Research, Saveetha Dental College and Hospitals, Saveetha Institute of Medical and Technical Sciences (SIMATS), Saveetha University, Chennai, India, ²Department of Biochemistry, Mbarara University of Science and Technology, Mbarara, Uganda, ³School of Pharmacy and Biomolecular Sciences, Liverpool John Moores University, Liverpool, United Kingdom, ⁴Institute for Health Research, Liverpool John Moores University, Liverpool, United Kingdom

Understanding and ameliorating neurodegenerative diseases represents a key challenge for supporting the health span of the aging population. Diverse protein aggregates have been implicated in such neurodegenerative disorders, including amyloid- β , α -synuclein, tau, fused in sarcoma (FUS), and transactivation response element (TAR) DNA-binding protein 43 (TDP-43). Recent years have seen significant growth in our mechanistic knowledge of relationships between these proteins and some of the membrane-less nuclear structures that fulfill key roles in the cell function. These include the nucleolus, nuclear speckles, and paraspeckles. The ability of macromolecular protein:RNA complexes to partition these nuclear condensates through biophysical processes that involve liquid-liquid phase separation (LLPS) has also gained attention recently. The paraspeckle, which is scaffolded by the architectural long-non-coding RNA nuclear enriched abundant transcript 1 (NEAT1) plays central roles in RNA processing and metabolism and has been linked dynamically to TDP-43. In this mini-review, we outline essential early and recent insights in relation to TDP-43 proteinopathies. We then appraise the relationships between TDP-43 and NEAT1 in the context of neuronal paraspeckles and neuronal stress. We highlight key areas for investigation based on recent advances in our understanding of how TDP-43 affects neuronal function, especially in relation to messenger ribosomal nucleic acid (mRNA) splicing. Finally, we offer perspectives that should be considered for translational pipelines in order to improve health outcomes for the management of neurodegenerative diseases.

KEYWORDS

TDP-43, long non-coding RNA, NEAT1, neurons, paraspeckles, TAR DNA-binding protein 43, nucleic acid therapies, swimming microrobots

Abbreviations: AD, Alzheimer's disease; ALS, amyotrophic lateral sclerosis; ASO, antisense oligonucleotide; CB, Cajal body; DMD, Duchenne muscular dystrophy; FTL, frontotemporal lobar degeneration; FUS, fused in sarcoma; HD, Huntington's disease; NEAT, nuclear enriched abundant transcript 1; NONO, non-POU domain-containing octamer-binding protein; PD, Parkinson's disease; PSPs, paraspeckle proteins; SFPQ, splicing factor proline and glutamine rich; snRNPs, small nuclear ribonucleoproteins; STMN2, stathmin 2; TDP-43, transactivation response element (TAR) binding protein.

Introduction

Neurodegenerative diseases are a diverse set of conditions characterized by the progressive degradation of the structure and function of the central or peripheral nervous systems (Dugger and Dickson, 2017). Major neurodegenerative disorders include Alzheimer's disease (AD), amyloid lateral sclerosis (ALS; also known as Lou Gehrig's disease), frontotemporal lobar degeneration (FTLD), Huntington's disease (HD), and Parkinson's disease (PD). Collectively, such neurodegenerative disorders are projected to become the second leading cause of death after cardiovascular disease by 2040 (Gammon, 2014). In contrast to rapid neuronal loss caused by metabolic or toxic illnesses, neurodegenerative disorders are characterized by the gradual loss of selectively vulnerable populations of neurons, thus diminishing the functional properties of the brain and peripheral organs (Dugger and Dickson, 2017). Cell death and shrinkage in a specific region of the brain represent a major pathogenic trait in several neurodegenerative illnesses (Dugger and Dickson, 2017).

Pathological deposition of intracellular or extracellular protein aggregates is associated with several neurodegenerative disorders. Depending on the specific protein, these conditions can be grouped into amyloidosis (amyloid- β), α -synucleinopathies (α -synuclein), tauopathies (tau), FUSopathies [Fused in sarcoma (FUS)], and TDP-43 proteinopathies [transactivation response element (TAR) DNA-binding protein 43 (TDP-43)] (Kovacs, 2017). Such protein aggregates, and/or their oligomeric species frequently stress neurons, resulting in neuronal death (Chi et al., 2018).

The TDP-43 proteinopathies include ALS, FTLD-TDP, primary lateral sclerosis, and progressive muscular atrophy and are defined by TDP-43-related pathology (Dugger and Dickson, 2017). The hallmarks of TDP-43 proteinopathies include cytoplasmic accumulation, phosphorylation, ubiquitination, and truncation of TDP-43 into neurotoxic C-terminal fragments (Chhangani et al., 2021). Most cases of ALS are sporadic, though 10% of ALS cases are considered to be familial (Taylor et al., 2016). In addition to TDP-43, a further 15 genes harboring mutations have been unequivocally associated with ALS by strong genetic and functional data, as reviewed by Taylor et al. (2016). These include Cu-Zn superoxide dismutase 1 (SOD1), FUS, optineurin (OPTN), TANK binding kinase 1 (TBK1), and C9ORF72 (Taylor et al., 2016). More recently, 12 additional genomic loci have been implicated in ALS by genome-wide association studies (GWAS), including KIF5A, GPX3, TNIP1, and UNC13A (van Rheenen et al., 2016, 2021; Benyamin et al., 2017; Nicolas et al., 2018). A machine learning approach that integrated epigenetic features of motor neurons with ALS GWAS data also revealed a role for KANK1 in ALS pathology, from 690 ALS-associated genes that were identified (Zhang et al., 2022).

Transactivation response element (TAR) DNA-binding protein 43 is a highly conserved heterogeneous ribonucleoprotein (hnRNP) multi-domain protein first identified as a 43-kDa protein that bound the TAR in human immunodeficiency virus (Ou et al., 1995). Under normal physiological conditions, TDP-43 is subjected to nucleocytoplasmic shuttling while residing predominantly in the nucleus (Ayala et al., 2008). This localization to both nuclear and cytosolic compartments is reflected in the processes regulated by TDP-43, which span messenger ribosomal nucleic acid (mRNA) transcription, splicing, maturation, and mRNA transport as well as the formation of stress granules and the regulation of miRNA processing, as reviewed recently by Prasad et al. (2019). Unsurprisingly, therefore, mutations that increase TDP-43 aggregation, increase TDP-43 half-life, or alter TDP-43 interactions with other proteins are thought to contribute to disease pathology in TDP-43 proteinopathies, and over 52 TDP-43 mutations have been linked to disease (Buratti, 2015).

There is growing recognition that TDP-43 undergoes liquid-liquid phase separation, (LLPS) a biophysical process that gives rise to biomolecular condensates reminiscent of an oil-in-vinegar mixture (Conicella et al., 2020; Ditlev, 2021; Schmit et al., 2021; Cascella et al., 2022). Driven by multivalent intermolecular interactions involving proteins and nucleic acids, biophysical LLPS is thought to support thermodynamically favorable formation of membraneless organelles endowed with specific functions within the nucleus and cytoplasm (Banani et al., 2017; Peng et al., 2021). Several such membraneless compartments have been defined in the nucleus, include the nucleolus, nuclear speckle, Cajal body, histone locus body, and paraspeckles which are broadly related by various aspects of RNA processing (Mao et al., 2011), as well as Polycomb bodies, promyelocytic leukemia (PML) bodies and nuclear stress bodies (Courchaine et al., 2016; Sawyer et al., 2019; Corpet et al., 2020; Wang et al., 2021). In this review, we appraise some of these nuclear bodies briefly then comment on recent developments in our understanding of TDP-43 in neurodegenerative disorders, highlighting areas for further research and therapeutic strategies that may help ameliorate disease.

Nuclear bodies and liquid-liquid phase separation

The nucleolus was established as the site for ribosome biogenesis in the 1960s and through a variety of technological innovations, we now recognize that key nucleolar proteins involved in rRNA transcription shuttle rapidly between the nucleolus and nucleoplasm (Chen and Huang, 2001). Further, the nucleolus consists of three phase-separated subcompartments: multiple copies of the fibrillar center (FC) and the dense fibrillar component (DFC) embedded within the granular component (GC) (Boisvert et al., 2007;

Lafontaine et al., 2021). Recent studies indicate the assembly of the DFC is driven by phase-separation mediated by nascent pre-rRNA sorting (Yao et al., 2019), while binding of the abundant nucleolar protein nucleophosmin (NPM1) to proteins and RNA drives the formation of phase separated multicomponent liquid-like droplets in the GC matrix (Mitrea et al., 2016). Importantly, the GC also functions as a quality control center into which misfolded proteins are sequestered reversibly during periods of cell stress in order to prevent their aggregation, allowing subsequent heat shock protein 70-dependent refolding during recovery from the stress (Frottin et al., 2019). Interestingly, nucleolin, another abundant nucleolar protein, was recently shown to alleviate TDP-43-related toxicity, at least in yeast and HEK293 cell models, apparently by facilitating the nuclear retention of TDP-43, thus preventing its accumulation as toxic cytoplasmic condensates (Peggion et al., 2021).

Nuclear speckles (interchromatin granule clusters) are highly dynamic structures enriched for transcription factors, chromatin remodeling factors and other proteins that enable nuclear speckles to function as a hub for coordinated regulation of the various steps associated with gene expression (Galganski et al., 2017). These include gene positioning, chromosome localization, chromatin modification, transcription, splicing, 3' end processing, and mRNA modification (Galganski et al., 2017; Faber et al., 2022). Notably, in an attempt to clarify exactly what should be called a nuclear speckle and based on recent findings, Ilik and Aktas, 2021 proposed to define as nuclear speckles by LLPS and the presence of two proteins SON and serine/arginine repetitive matrix protein 2 (SRRM2), that are rich in intrinsically disordered regions.

Cajal bodies (CB) are multifunctional biomolecular condensates that support the maturation of small nuclear ribonucleoproteins (snRNPs). The snRNPs are multimeric protein-RNA complexes that form the spliceosome and it is thought that snRNPs are concentrated in CB in order to promote the assembly of the individual monomers into higher order complexes (Klingauf et al., 2006; Morris, 2008; Shaw et al., 2008). Cajal bodies are defined by the presence of coilin, although other proteins such as survival motor neuron protein (SMN; the spinal muscular atrophy disease gene product), Geminin, Nopp140, and fibrillarin, as reviewed in Lafarga et al. (2017). Loss of coilin is semi-lethal, with about half of coilin knockout mice dying late in gestation (Walker et al., 2009). The reproductive fecundity of the mice was reduced, with smaller litter sizes and fewer litters overall (Walker et al., 2009). Thus, it appears that CB function has evolved to be somewhat robust such that in the absence of coilin, the "residual" CB can maintain levels of function that support viability. Interestingly, the localization of small Cajal body RNAs (scaRNAs) was recently shown to be regulated

by TDP-43 and this in turn regulated the site-specific 2'-O-methylation of U1 and U2 small nuclear RNAs (Izumikawa et al., 2019).

Interestingly, early work showed CB interactions with replication-dependent histone (RDH) loci (Shopland et al., 2001). These histone locus bodies (HLBs) have now been linked to the formation of a ternary complex with CB and RNA polymerase II, facilitating 3'-end processing of RDH genes (Imada et al., 2021; Suzuki et al., 2022). The dependence of these HLBs on histone pre-mRNA transcripts has led to the histone pre-mRNA transcripts being classified as architectural or scaffolding RNAs (Shevtsov and Dundr, 2011; Chujo and Hirose, 2017).

Three long non-coding RNAs (lncRNAs) have been also been categorized as scaffolding or architectural RNAs associated with nuclear bodies in human cells (Chujo and Hirose, 2017). These are (a) mammalian nuclear enriched abundant transcript 1 isoform 2 (NEAT1_2) lncRNA in the paraspeckle, (b) ribosomal intergenic spacer (IGS) lncRNA that drives amyloid body assembly for local nuclear protein synthesis (Theodoridis et al., 2021), and (c) human satellite III (SatIII) lncRNA in the nuclear stress body (Valgardsdottir et al., 2005, 2008). In addition, *Drosophila* heat shock RNA (Hsr) omega in the omega speckle (Mallik and Lakhota, 2009) and the fission yeast lncRNA meiRNA in the Mei2 dot (Yamashita et al., 1998) also meet the criteria to be considered architectural RNAs (Chujo and Hirose, 2017).

In healthy cells, TDP-43 resides in the nucleus and regulates RNA processing and translation, functions that are lost following of mis-localization of TDP-43 as cytoplasmic inclusions (Bjork et al., 2022). Both the loss of function associated with nuclear depletion and the gain of toxicity due to cytoplasmic accumulation are implicated in the pathogenic effects of TDP-43 (Chhangani et al., 2021). Recent studies have started to uncover complex relationships between TDP-43 and the lncRNA NEAT1 in TDP-43 proteinopathies, and we now turn to this theme.

Transactivation response element DNA-binding protein 43 proteinopathies: A brief overview

Landmark proteomics studies on post-mortem brain extracts from FTL and ALS patients in 2006 uncovered TDP-43 as a key protein associated with FTL, specifically ubiquitin-positive but tau- and alpha-synuclein negative FTL (Arai et al., 2006; Neumann et al., 2006). This predominantly nuclear protein that was known to bind the TAR element of the human immunodeficiency virus has since been implicated in the pathogenesis of multiple neurodegenerative diseases collectively known as TDP-43 proteinopathies (de Boer et al., 2020; Huang et al., 2020; Carlos and Josephs, 2022). Early

studies observed TDP-43 cytoplasmic inclusions in ALS and AD (Arai et al., 2006; Igaz et al., 2008). Further, a transgenic mouse model with moderate human TDP-43 overexpression showed intranuclear and cytoplasmic phosphorylated TDP-43 aggregates in neurons. Importantly, these mice showed axonal and myelin degeneration which were reflected in gait abnormalities and early mortality (Xu et al., 2010). Some TDP-43 aggregates localize to mitochondria and, crucially, inhibition of TDP-43 mitochondrial localization suppresses TDP-43 neuronal toxicity (Mori et al., 2008; Wang et al., 2016). Recent studies have now revealed the structural basis for these observations in ALS as aggregated TDP-43 sequesters selected microRNAs (miRNAs) and proteins, leading to altered expression of nuclear-genome-encoded mitochondrial proteins (Zuo et al., 2021). This in turn generates a global mitochondrial imbalance that drives oxidative stress (Zuo et al., 2021). Interestingly, TDP-43 were shown to form amyloid-like filaments in the brains of patients who had ALS with FTLD (Arseni et al., 2022) so it will be intriguing to determine how the filaments drive protein and miRNA sequestration. In addition, very recent work indicated that brain samples from TDP-43 proteinopathies as well as tauopathies and synucleinopathies are characterized by amyloid fibrils consisting of a 135-amino acid C-terminal fragment of transmembrane protein 106B (TMEM106B) (Chang et al., 2022; Jiang et al., 2022; Schweighauser et al., 2022). However, the relationships of these TMEM106B fibrils to disease pathology remains to be established.

In an early study of protein:RNA interactions in the post-mortem brain tissue by Tollervey et al. (2011), most TDP-43 binding was found on introns, long ncRNAs and intergenic RNAs. Similar results were observed in cultured cells. In addition, the authors found that TDP-43 regulates pre-mRNA splicing, generating alternative mRNA isoforms for several proteins involved in neuronal development or disease (Tollervey et al., 2011). What has become clear more recently is that TDP-43 suppresses the incorporation of poorly conserved cryptic exons into mRNA, and TDP-43 deficiency in the nucleus is associated with the generation of mis-spliced transcripts that are targeted for nonsense-mediated decay (Ling et al., 2015; Humphrey et al., 2017; Jeong et al., 2017; Sun et al., 2017).

Transactivation response element DNA-binding protein 43 and Alzheimer's disease

In addition to ALS and FTLD, roles for TDP-43 in the pathology of AD have also been established. Several studies have observed TDP-43 immunoreactivity in a subset of AD cases with neuroanatomical distribution patterns that were broadly distinct from those observed in FTLD (Amador-Ortiz et al., 2007; Higashi et al., 2007; Uryu et al., 2008; Arai et al., 2009;

TABLE 1 Transactivation response element (TAR) DNA-binding protein 43 (TDP-43) positive immunoreactivity in Alzheimer's disease (AD) patients.

TDP-43 positive patients/Total number of patients	Percentage	References	Comments
9/85	11	Davidson et al., 2011	Early onset AD
19/93	20	Amador-Ortiz et al., 2007	
47/182	26	Uryu et al., 2008	
24/84	29	Davidson et al., 2011	Late onset AD
5/16	31	Kadokura et al., 2009	
5/15	33	Higashi et al., 2007	
19/53	36	Arai et al., 2009	
14/25	56	Arai et al., 2009	
34/46	74	McAleese et al., 2017	
6/8	75	King et al., 2010	

Kadokura et al., 2009; King et al., 2010; Davidson et al., 2011; McAleese et al., 2017). The findings are summarized in Table 1. Subsequent studies have linked TDP-43 with lower cognitive function in AD patients in which hippocampal sclerosis has been reported (Nag et al., 2015). Importantly, even when subjects had comparable levels of AD pathology based on Braak staging, TDP-43 positive cohorts were much more likely to have cognitive malfunction compared to their TDP-negative counterparts, and the effects of TDP-43 appeared to be independent of hippocampal sclerosis (Josephs et al., 2014). Of note, TDP-43 deposition appears to be part of the normal physiological response during aging with neuronal cytoplasmic inclusions observed in 36% in the brains of cognitively normal subjects aged 71–100 years (Arnold et al., 2013).

Early work revealed that the amygdala was particularly vulnerable to TDP-43 in advanced AD region (Hu et al., 2008). Further, detailed analysis of TDP-43 distribution across 14 brain regions led to the six-stage scheme in which TDP-43 immunoreactivity is limited to the amygdala at stage 1 and extends progressively to the entorhinal cortex and subiculum (stage 2), the dentate gyrus of the hippocampus and occipitotemporal cortex (stage 3), insular cortex, ventral striatum, basal forebrain, and inferior temporal cortex (stage 4), substantia nigra and midbrain tectum (stage 5), and basal ganglia and middle frontal cortex (stage 6) (Josephs et al., 2016). Interestingly, the distinct distribution patterns of TDP-43 have been observed in relation to these stages: stages 4–6 were linked to “typical” TDP-43 immunoreactive inclusions defined by neuronal cytoplasmic inclusions, neuronal intranuclear inclusions, and dystrophic neurites, while stages 1–3 associated with a “type-β” deposition in which TDP-43 immunoreactivity occurs adjacent to tau-positive neurofibrillary tangles in the

same neuron (Josephs et al., 2019). However, both types were associated with smaller amygdala and hippocampal volumes and such volumetric differences may be predictive of TDP-43 status while patients are alive. For further details on the relationships between TDP-43 and classical neuropathological features of AD such as amyloid- β and tau oligomers, the reader is referred to an excellent recent review by Meneses et al. (2021) that also includes a helpful depiction of the TDP-43 distribution pattern in AD.

Transactivation response element DNA-binding protein 43 clearance and altered splicing

One mechanism of disease recently implicated in ALS and FTLT-DTP links the loss of nuclear TDP-43 to mis-splicing and depletion of stathmin 2 (*STMN2*), a regulator of the microtubule cytoskeleton (Klim et al., 2019; Melamed et al., 2019; Prudencio et al., 2020; Pickles et al., 2022). Given the importance of microtubules for normal neuron function, it is noteworthy that two independent studies have now demonstrated that stathmin deficiency leads to motor dysfunction in mouse models (Guerra San Juan et al., 2022; Krus et al., 2022).

Loss of nuclear TDP-43 has also been linked to mis-splicing events that decrease the expression of *UNC13A* in neurons (Brown et al., 2022; Ma et al., 2022). The *UNC13A* protein is critical for normal synaptic function and altered *UNC13A* splicing introduces a cryptic exon that leads to premature stop codons (Brown et al., 2022; Ma et al., 2022). The relationships between TDP-43 and cryptic exon splicing leading to neuronal dysfunction are summarized in **Figure 1**. Importantly, single-nucleotide polymorphisms (SNPs) associated with FTLT and ALS risk were found in the region containing the cryptic exon (Brown et al., 2022; Ma et al., 2022). Genetic variants associated with disease risk correlated with elevated levels of *UNC13A* cryptic exon inclusion. Further, the SNPs influenced TDP-43 binding to *UNC13A* pre-mRNA and enhanced cryptic exon inclusion when *UNC13A* minigenes were transcribed (Brown et al., 2022; Ma et al., 2022). The conclusions reached in these studies was not so much that the *UNC13A* risk variants play a causative role in disease but rather, against a backdrop of TDP-43 clearance from the nucleus, insufficient TDP-43 binding to the cryptic exon permits incorporation of cryptic exon into pre-mRNA.


The study by Tollervey et al. (2011) also detected substantially more TDP-43 binding to lncRNA NEAT1 in FTLT-DTP samples, possibly due to increased NEAT1 expression. Given that NEAT1 serves as an architectural lncRNA for paraspeckle formation, we now turn to the relationships between NEAT1, TDP-43, and paraspeckles in the neuronal nucleus.

Nuclear enriched abundant transcript 1 and transactivation response element DNA-binding protein 43 proteinopathy in the neuronal paraspeckle

Early work by Hutchinson et al. (2007) led to the identification of NEAT1 and NEAT2 (also known as MALAT-1). Subsequent independent work from Hirose and Spector groups defined two NEAT1 isoforms transcribed from the multiple endocrine neoplasia I (MEN I) genomic locus, a short isoform of 3.7 kb (initially MEN ϵ , now known as NEAT1_1) and a much longer transcript of 23 kb (initially MEN β , now NEAT1_2) (Sasaki et al., 2009; Sunwoo et al., 2009). These studies also provided initial evidence linking NEAT1 lncRNAs to paraspeckles (Sasaki et al., 2009; Sunwoo et al., 2009). Further work confirmed the involvement of NEAT1 in paraspeckle assembly, with NEAT1_2 in particular serving as a scaffold or architectural lncRNA to support paraspeckle formation (Clemson et al., 2009; Souquere et al., 2010).

In neurons, upregulation of NEAT1 was linked to neuronal and oligodendrocyte maturation (Mercer et al., 2010) but it was work by Tollervey et al. (2011) that established a possible connection to FTLT by showing that NEAT1 and MALAT-1 bound TDP-43 in post-mortem human brain samples. Further evidence for NEAT1 involvement in neurodegenerative disorders came from studies by Nishimoto et al. (2013) whose analysis of 633 human spinal motor neurons derived from six ALS patients revealed elevated expression of NEAT1_2 in the early stages of the disease. The investigators also confirmed the ability of TDP-43 to bind NEAT1_2 and colocalize with NEAT1_2 in paraspeckles (Nishimoto et al., 2013). These findings together gave rise to the notion of NEAT1_2 lncRNA as a scaffold for the assembly of macromolecular protein and RNA complexes in the nuclei of ALS motor neurons in the early phase of the disease. Importantly, elevated expression of NEAT1 has also been observed in other neurodegenerative diseases, including frontotemporal dementia, AD, HD, and PD, as reviewed in An et al. (2018).

These early studies thus revealed a clear axis of neurodysregulation associated with NEAT1_2 and TDP-43. What was less clear was whether NEAT1_2 roles were pathogenic or a compensatory effect to stave off neuronal death. Two main concepts have now begun to emerge, however. First, that LLPS appears to be the driving force for paraspeckle assembly, with the middle domain in particular recruiting the essential paraspeckle proteins (PSPs) non-POU domain-containing octamer-binding protein (NONO) and SFPQ (splicing factor proline and glutamine rich), with NONO forming dimers that support oligomerization with other PSPs (Yamazaki et al., 2018). From a biophysical perspective, the spatial arrangement of NEAT1 and the PSPs



appears to mimic amphiphatic block copolymer micelles in which connected hydrophilic and hydrophobic polymers self-assemble into spherical and cylindrical nanostructures

in water (Yamazaki et al., 2021). Secondly, that upregulation of NEAT1 and of paraspeckles exerts a protective effect on neurons, to which we now turn.


Nuclear enriched abundant transcript 1: Transactivation response element DNA-binding protein 43 as protective partners in the neuronal stress response

Early work from Nakagawa et al. (2011) found widespread constitutive expression of NEAT1_1, whereas NEAT1_2 expression was limited to specific cells in the stomach and intestine. There was no obvious phenotype to *NEAT1*-deficient mice, beyond the loss of paraspeckles although subsequent work revealed a high expression of NEAT1 in the corpus luteum of mouse ovaries, impaired formation of luteal tissue, low serum progesterone, and a stochastic inability to become pregnant (Nakagawa et al., 2014). In contrast, no essential physiological roles were identified from studies on mice specifically deficient in NEAT1_1 (Adriaens et al., 2019). It would therefore be interesting to explore the effects of NEAT1_1 depletion in other mammalian orders, especially in primates or primate tissues, to uncover potential functions in these organisms.

Notably, as illustrated in Figure 2 and discussed below, NEAT1 nonetheless appears to exert cytoprotective effects in diverse models. Neurons do not express NEAT1_2 under basal conditions and are devoid of paraspeckles at least *in vitro* (Shelkovnikova et al., 2018). Unsurprisingly, therefore, mice lacking NEAT1 show no evidence of neuronal loss, neuroinflammation, or gross synaptic dysfunction (Kukharsky et al., 2020). On the other hand, Shelkovnikova et al. (2018) recently observed paraspeckles in a small cohort of ALS patients (two with TDP-43 pathology, four with C9ORF72 pathology, and seven with sporadic ALS), with ~30% of spinal cord neurons displaying paraspeckles compared to no paraspeckles in neurons from healthy control patients. An associated increased NEAT1 expression was also reported (Shelkovnikova et al., 2018). Given that key features of TDP-43 proteinopathy include clearance of the protein from the nucleus with concomitant accumulation and aggregation in the cytoplasm

(Arai et al., 2006; Neumann et al., 2006), it was notable that silencing of TDP-43 in SH-SY5Y neuroblastoma cells raised the number of paraspeckles per nucleus and expression of the paraspeckle-specific NEAT1 isoform, NEAT1_2 (Shelkovnikova et al., 2018). More importantly, because TDP-43 can enhance miRNA biogenesis (Buratti et al., 2010; Kawahara and Mieda-Sato, 2012), the formation of paraspeckles in TDP-43-deficient neuroblastoma cells appears to serve a protective role by marshaling the pri-miRNA processing machinery to sustain the generation of miRNAs that target pro-apoptotic genes (Bottini et al., 2017; Jiang et al., 2017; Shelkovnikova et al., 2018). However, although some TDP-43-binding mRNAs were altered in the serum and cerebrospinal fluid (CSF) of sporadic ALS patients (Freischmidt et al., 2013), the relationships between TDP-43 and motor neuron miRNA expression has not been fully established.

The idea that the upregulation of NEAT1_2 and paraspeckle abundance exerts a neuroprotective role is further supported by recent work from Wang et al. (2020). Distinct triggers of cell stress, including arsenite and leptomycin B (an inhibitor of nuclear export), induced TDP-43 to form dynamic, reversible liquid-droplet-like nuclear bodies in mammalian cells, including mouse neurons (Wang et al., 2020). Importantly, studies on fly models of ALS-related function suggested that the TDP-43 nuclear bodies were neuroprotective *in vivo*. Stress also raised the levels of NEAT1 RNA in neurons, which in turn supported the condensation of TDP-43 through mechanisms that exploited LLPS. One particular ALS-associated TDP-43 mutation D169G, but not other mutations examined, reduced NEAT1 mediated TDP-43 LLPS. Importantly, the TDP-43 nuclear bodies appear to exert a cytoprotective effect in human embryonic kidney (HEK293T) cells as well as *Drosophila* neurons (Wang et al., 2020). Clearly, it will be crucial to determine whether NEAT1-induced TDP-43 nuclear bodies formed in response to neuronal stress suppresses the incorporation of cryptic, neurodegenerative *STMN2* and *UNC13A* exons (Klim et al., 2019; Melamed et al., 2019; Prudencio et al., 2020; Brown et al., 2022; Ma et al., 2022).



Depletion of TDP-43 drives paraspeckle assembly through elevation of NEAT1_2 expression that is mediated by type 1 interferon (Shelkovnikova et al., 2018). This upregulation of NEAT1_2 expression appears to result from the accumulation of miRNA species (Shelkovnikova et al., 2018). On the other hand, more recent work has suggested that overexpression of full length TDP-43 that localizes to the nucleus (but not TDP-43 lacking the nuclear localization signal) can elevate NEAT1_1 expression in SH-SY5Y neuroblastoma cells and in the central nervous systems (CNS) cortex of a mouse model (Matsukawa et al., 2021). Even though paraspeckles are not observed in cells that express only NEAT1_1, which instead distributes in a diffuse “microspeckle” pattern that does not co-localize with paraspeckle markers in the nucleoplasm (Nakagawa et al., 2011; Li R. et al., 2017), NEAT1_1 alleviated TDP-43-dependent toxicity in yeast and *Drosophila* models of TDP-43 proteinopathy (Matsukawa et al., 2021). One idea proposed for the ability of NEAT1_1 to ameliorate TDP-43 toxicity is that NEAT1_1 binding to TDP-43 may block unwanted interactions of TDP-43 with other nuclear RNAs (Matsukawa et al., 2021). Hence, it will be interesting to determine whether NEAT1_1 can also prevent TDP-43-dependent toxicity in non-human primate models (Uchida et al., 2012).

Work from Mann et al. (2019) recently indicated that a 34-nt TDP-43-binding oligonucleotide abrogated the phase transitions and neurotoxicity of pathologically-relevant TDP-43 inclusions, at least as monitored using optogenetic (i.e., light-dependent) oligomerization strategies. Strikingly, the oligonucleotide apparently helped maintain optogenetic TDP-43 in the nucleus and prevent localization to the cytoplasm, though the mechanisms of this potentially therapeutic axis requires further investigation. Further work is also required to establish whether the action of such oligonucleotides is associated with the induction of paraspeckle formation in neurons. Beyond these studies, there is clearly an urgent need to establish whether TDP-43-binding oligonucleotides can restore the ability of TDP-43 to suppress splicing of the cryptic *STMN2* and *UNC13A* exons implicated in neurodegeneration (Klim et al., 2019; Melamed et al., 2019; Prudencio et al., 2020; Brown et al., 2022; Ma et al., 2022).

The above considerations together suggest complex effects of TDP-43 in supporting neuroprotective responses *via* elevation of NEAT1_2 and NEAT1_1. Further studies are required to validate the therapeutic potential of paraspeckle induction and NEAT1_1 elevation in TDP-43 neurodegenerative disorders. These will need to span zebrafish, mouse and primate models of TDP-43 proteinopathy as have been used in recent studies (Lissouba et al., 2018; Yin et al., 2019; Huang et al., 2021). One important question to address will be whether NEAT1_2 elevation (or more generally, enhancement of paraspeckle numbers) in parallel with increased expression of NEAT1_1 has an additive or synergistic effect on neuroprotection from TDP-43 toxicity.

Therapeutic horizons: Oligonucleotide approaches

Interestingly, *de novo* paraspeckle assembly appears to be a key feature of ALS TDP-43 proteinopathy (Shelkovnikova et al., 2018). Exposure of SH-SY5Y neuroblastoma cells to enoxacin, a small molecule that promotes miRNA biogenesis, boosted NEAT1_2 and paraspeckle abundance, adding a lncRNA:paraspeckle axis to the mode of action of enoxacin (Shan et al., 2008; Melo et al., 2011; Emde et al., 2015; Shelkovnikova et al., 2018). These paraspeckles appear to serve a protective role in neuroblastoma cells. However, neither enoxacin nor other agents (such as histone deacetylase inhibitors) that induced NEAT1_2 and *de novo* paraspeckles in neuroblastoma cells were able to evoke paraspeckle assembly in human embryonic cell-derived motor neurons (Shelkovnikova et al., 2018). Nonetheless, it is noteworthy that in mouse embryonic stem cells, a 20-mer phosphorothioate-modified (PS) antisense oligonucleotide (ASOs) drove the assembly of PSPs into paraspeckle-like condensates lacking NEAT1 but containing the paraspeckle protein P54nrb (Shen et al., 2014). Whether such ASO-dependent paraspeckles can assemble in motor neurons needs to be established, along with evaluating the impact of such induced paraspeckles on nuclear retention and splicing functions of TDP-43. With recent regulatory approvals of several ASOs, including nusinersen (Spinraza[®]) for spinal muscular atrophy (Finkel et al., 2016; Corey, 2017), casimersen for Duchenne muscular dystrophy (DMD) (Shirley, 2021; Wagner et al., 2021) and volanesorsen for familial chylomicronemia syndrome (Gouni-Berthold et al., 2021), the prospects for therapeutic ASOs seem strong, so it will be important to exploit their potential for alleviating TDP-proteinopathies. Casimersen, is an exon-skipping ASO which functions by facilitating the expression of an internally truncated but functional dystrophin protein in DMD patients with a mutation that is amenable to exon 45 skipping (Shirley, 2021; Wagner et al., 2021). It will be interesting to determine whether similar exon skipping approaches can restore efficacious *STMN2* and *UNC13A* expression in ALS and FTLN neurons. The challenge will be specific delivery of such exon-skipping ASO to the relevant regions of the brain. Alternatively, a PS-modified version of the 34-nt TDP-43-binding oligonucleotide (bait oligonucleotide) shown by Mann et al. (2019) to enhance nuclear retention of TDP-43 may also have translational potential once targeted safe delivery to specific anatomical locations of the brain is achieved. Conceivably, computational modeling of wild type and mutant TDP-43, for instance using AlphaFold or RosettaFold algorithms (Baek et al., 2021; Jumper et al., 2021) will enable the selection of PS-rich oligonucleotides that can target TDP-43 mutants for functional retention in the nucleus. There are at least 52 missense mutations associated with ALS (Ratti and Buratti, 2016), so the ability to rapidly determine

the effects of each mutation on the predicted structure of TDP-43 will provide new insights to improve the bait oligonucleotide design. A 12-nucleotide GU-repeat RNA sequence recently shown to associate with TDP-43 and prevent aggregation and localization to the cytoplasm may also hold translational potential (Rengifo-Gonzalez et al., 2021; Cascella et al., 2022).

It will also be important to determine whether ASOs or bait oligonucleotides ASOs can support the generation of the neuroprotective TDP-43 nuclear bodies that Wang et al. (2020) observed in mammalian cells and fly neurons; see NEAT1:TDP43 as protective partners in the neuronal stress response section above. Given, as mentioned above, the ability of a 20-mer PS-ASO to nucleate the assembly of PSPs into nuclear condensates lacking NEAT1 (Shen et al., 2014), can strategies be designed (for instance *via* machine learning) to uncover similar PS-ASOs that condense NEAT1 and TDP-43? In this regard, it is noteworthy that an attempt to deploy AlphaFold and other computational approaches in expanding the targetable chemical space of TDP-43 has already been reported (Scott et al., 2022). However, the recent early termination of promising ASO clinical trials for HD, due to lack of efficacy (Kwon, 2021), is a reminder of how challenging the effective deployment of ASO in any neurodegenerative disorders is likely to remain for some time.

One obvious limitation of the above approaches, however, is that while they may restore expression of functional *STMN2* or *UNC13A* they do nothing to address any pathology that may be due to cytoplasmic TDP-43 aggregates. Hence such strategies will need to demonstrate that therapeutic restoration of *STMN2* or *UNC13A* protein expression enhances neuronal survival even in the presence of cytoplasmic TDP-43 inclusions. More pragmatically, small molecules that have shown potential as disruptors of TDP-43 aggregates, such as an acridine derivative generated by Prasad et al. (2016) are likely to be required for co-administration with ASO in order to maximize improvements in patient outcomes.

Therapeutic horizons: Inspiration from COVID-19 messenger ribosomal nucleic acid vaccines

The neuroprotective abilities of NEAT1_1 in TDP-43 proteinopathies reported by Matsukawa et al. (2021) aligns with earlier work showing overexpression of NEAT1_1 protected mouse neuroblasts from oxidative damage (Sunwoo et al., 2017). Upregulation of NEAT1_1 in human HD post-mortem brains was thus considered to be a neuroprotective response to limit neuronal death (Sunwoo et al., 2017). The corollary to these observations is that direct delivery of synthetic NEAT1_1 lncRNA or expression vector may help avert neuronal damage and dysfunction. The global pandemic trigger by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) has witnessed the unprecedented deployment of

mRNA in the form of mRNA vaccines, as reviewed recently (Chaudhary et al., 2021). Therefore, it should be possible to exploit similar RNA technologies in the development of NEAT1_1 for targeted delivery to neurons associated with TDP-43 proteinopathies and Huntington's disease. The challenge will center on the development of safe and effective delivery vehicles but promising approaches for brain-targeted nucleic acid therapies (NATs) have started to emerge. These include lipid nanoparticles (Tanaka et al., 2018), cationic liposomes (Dhaliwal et al., 2020), both of which were used to deliver mRNA, and focused ultrasound (Mathew et al., 2021), which was used to deliver an expression plasmid. Clearly, these technologies can also be developed for the delivery of TDP-43 binding oligonucleotides to neurons, and it will be interesting to see which delivery system best supports the formation of neuroprotective TDP-43 nuclear bodies.

Future horizons: Can tiny machines target transactivation response element DNA-binding protein 43 proteinopathies?

Taking inspiration from the motility of single-cell biological entities such as sperm and *Escherichia coli*, a whole new field of “swimming microrobots,” “microswimmers,” or “miniature medical robots” has emerged in recent years for a wide range of biomedical applications (Peyer et al., 2013; King, 2022). The premise of these approaches is that such untethered micromachines can deliver therapeutic payloads in a precise manner to regions of the body that are normally difficult to access and also perform minimally invasive surgical procedures (Li J. et al., 2017). Magnetic fields are arguably the most widely tested actuation methods to drive the motion of swimming microrobots (Yang and Zhang, 2020; Shao et al., 2021). However, other approaches such as ultrasound or light have also been explored, along with self-navigating methods based on chemotaxis, phototaxis, magnetotaxis, gravitaxis, and rheotaxis (Yu et al., 2021). Interestingly, swimming microrobots have become an area of intense study for targeted drug delivery to neurons (Dong et al., 2020; Kim et al., 2020). We are therefore likely to see application of these micromachines to neurodegenerative disorders in the coming years.

Conclusion

Much progress has been made in deciphering how TDP-43 mislocalization and dysfunction contributes to neurodegeneration. As argued above, it seems that the induction of paraspeckles in the neuronal nucleus or the retention of

TDP-43 as nuclear bodies can support neural viability. However, further studies are needed to clarify the relationships between induced paraspeckles and TDP-43 localization and function, especially in relation to mRNA splicing.

The emergence of NATs spanning ASO and other modalities such as short-interfering RNA (Kulkarni et al., 2021) offer a tantalizing prospect for NAT-based strategies based on oligonucleotides delivered to the brain in order to bind TDP-43 and enhance nuclear retention and restore nuclear function of TDP-43 if the promising findings from Mann et al. (2019) are reproduced in large animal models of TDP-proteinopathy. Therefore, approaches that combine TDP-43 binding oligonucleotides with studies of swimming microrobots on non-human primate models of TDP-43-mediated neurodegeneration are likely to have a transformative impact on our understanding of how to modulate the neuronal nucleus to ameliorate TDP proteinopathies.

Author contributions

KR, DT, and DS conceptualized the idea. KR wrote the first draft. All authors read the manuscript and approved the submitted version.

References

- Adriaens, C., Rambow, F., Bervoets, G., Silla, T., Mito, M., Chiba, T., et al. (2019). The long noncoding RNA NEAT1_1 is seemingly dispensable for normal tissue homeostasis and cancer cell growth. *RNA* 25, 1681–1695. doi: 10.1261/rna.071456.119
- Amador-Ortiz, C., Lin, W. L., Ahmed, Z., Personett, D., Davies, P., Duara, R., et al. (2007). TDP-43 immunoreactivity in hippocampal sclerosis and Alzheimer's disease. *Ann. Neurol.* 61, 435–445. doi: 10.1002/ana.21154
- An, H., Williams, N. G., and Shelkovernikova, T. A. (2018). NEAT1 and paraspeckles in neurodegenerative diseases: A missing lnc found? *Noncoding RNA Res.* 3, 243–252. doi: 10.1016/j.ncrna.2018.11.003
- Arai, T., Hasegawa, M., Akiyama, H., Ikeda, K., Nonaka, T., Mori, H., et al. (2006). TDP-43 is a component of ubiquitin-positive tau-negative inclusions in frontotemporal lobar degeneration and amyotrophic lateral sclerosis. *Biochem. Biophys. Res. Commun.* 351, 602–611. doi: 10.1016/j.bbrc.2006.10.093
- Arai, T., Mackenzie, I. R., Hasegawa, M., Nonaka, T., Niizato, K., Tsuchiya, K., et al. (2009). Phosphorylated TDP-43 in Alzheimer's disease and dementia with Lewy bodies. *Acta Neuropathol.* 117, 125–136. doi: 10.1007/s00401-008-0480-1
- Arnold, S. J., Dugger, B. N., and Beach, T. G. (2013). TDP-43 deposition in prospectively followed, cognitively normal elderly individuals: Correlation with argyrophilic grains but not other concomitant pathologies. *Acta Neuropathol.* 126, 51–57. doi: 10.1007/s00401-013-1110-0
- Arseni, D., Hasegawa, M., Murzin, A. G., Kametani, F., Arai, M., Yoshida, M., et al. (2022). Structure of pathological TDP-43 filaments from ALS with FTL. *Nature* 601, 139–143. doi: 10.1038/s41586-021-04199-3
- Ayala, Y. M., Zago, P., D'Ambrogio, A., Xu, Y. F., Petrucelli, L., Buratti, E., et al. (2008). Structural determinants of the cellular localization and shuttling of TDP-43. *J. Cell Sci.* 121, 3778–3785. doi: 10.1242/jcs.038950
- Baek, M., DiMaio, F., Anishchenko, I., Dauparas, J., Ovchinnikov, S., and Lee, G. R. (2021). Accurate prediction of protein structures and interactions using a three-track neural network. *Science* 373, 871–876. doi: 10.1126/science.abj8754
- Banani, S. F., Lee, H. O., Hyman, A. A., and Rosen, M. K. (2017). Biomolecular condensates: Organizers of cellular biochemistry. *Nat. Rev. Mol. Cell Biol.* 18, 285–298. doi: 10.1038/nrm.2017.7
- Benyamin, B., He, J., Zhao, Q., Gratten, J., Garton, F., and Leo, P. J. (2017). Cross-ethnic meta-analysis identifies association of the GPX3-TNIP1 locus with amyotrophic lateral sclerosis. *Nat. Commun.* 8:611. doi: 10.1038/s41467-017-00471-1
- Bjork, R. T., Mortimore, N. P., Loganathan, S., and Zarnescu, D. C. (2022). Dysregulation of Translation in TDP-43 Proteinopathies: Deficits in the RNA Supply Chain and Local Protein Production. *Front. Neurosci.* 16:840357. doi: 10.3389/fnins.2022.840357
- Boisvert, F. M., van Koningsbruggen, S., Navasques, J., and Lamond, A. I. (2007). The multifunctional nucleolus. *Nat. Rev. Mol. Cell Biol.* 8, 574–585. doi: 10.1038/nrm2184
- Bottini, S., Hamouda-Tekaya, N., Mategot, R., Zaragosi, L. E., Audebert, S., Pisano, S., et al. (2017). Post-transcriptional gene silencing mediated by microRNAs is controlled by nucleoplasmic Sfpq. *Nat. Commun.* 8:1189. doi: 10.1038/s41467-017-01126-x
- Brown, A. L., Wilkins, O. G., Keuss, M. J., Hill, S. E., Zanovello, M., Lee, W. C., et al. (2022). TDP-43 loss and ALS-risk SNPs drive mis-splicing and depletion of UNC13A. *Nature* 603, 131–137. doi: 10.1038/s41586-022-04436-3
- Buratti, E. (2015). Functional Significance of TDP-43 Mutations in Disease. *Adv. Genet.* 91, 1–53. doi: 10.1016/bs.adgen.2015.07.001
- Buratti, E., De Conti, L., Stuardi, C., Romano, M., Baralle, M., and Baralle, F. (2010). Nuclear factor TDP-43 can affect selected microRNA levels. *FEBS J.* 277, 2268–2281. doi: 10.1111/j.1742-4658.2010.07643.x
- Carlos, A. F., and Josephs, K. A. (2022). Frontotemporal lobar degeneration with TAR DNA-binding protein 43 (TDP-43): Its journey of more than 100 years. *J. Neurol.* 269, 4030–4054. doi: 10.1007/s00415-022-11073-3
- Cascella, R., Bigi, A., Riffert, D. G., Gagliani, M. C., Ermini, E., Moretti, M., et al. (2022). A quantitative biology approach correlates neuronal toxicity with

Acknowledgments

We thank reviewers for helpful feedback that significantly improved the quality of the manuscript. We apologize to the many excellent researchers whose work has not been cited on this occasion.

Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

- the largest inclusions of TDP-43. *Sci. Adv.* 8:eabm6376. doi: 10.1126/sciadv.abm6376
- Chang, A., Xiang, X., Wang, J., Lee, C., Arakhamia, T., Simjanoska, M., et al. (2022). Homotypic fibrillation of TMEM106B across diverse neurodegenerative diseases. *Cell* 185:e1315. doi: 10.1016/j.cell.2022.02.026
- Chaudhary, N., Weissman, D., and Whitehead, K. A. (2021). mRNA vaccines for infectious diseases: Principles, delivery and clinical translation. *Nat. Rev.* 20, 817–838. doi: 10.1038/s41573-021-00283-5
- Chen, D., and Huang, S. (2001). Nucleolar components involved in ribosome biogenesis cycle between the nucleolus and nucleoplasm in interphase cells. *J. Cell Biol.* 153, 169–176. doi: 10.1083/jcb.153.1.169
- Chhangani, D., Martin-Pena, A., and Rincon-Limas, D. E. (2021). Molecular, functional, and pathological aspects of TDP-43 fragmentation. *iScience* 24:102459. doi: 10.1016/j.isci.2021.102459
- Chi, H., Chang, H. Y., and Sang, T. K. (2018). Neuronal Cell Death Mechanisms in Major Neurodegenerative Diseases. *Int. J. Mol. Sci.* 19:3082. doi: 10.3390/ijms19103082
- Chujo, T., and Hirose, T. (2017). Nuclear Bodies Built on Architectural Long Noncoding RNAs: Unifying Principles of Their Construction and Function. *Mol. Cells* 40, 889–896.
- Clemson, C. M., Hutchinson, J. N., Sara, S. A., Ensminger, A. W., Fox, A. H., Chess, A., et al. (2009). An architectural role for a nuclear noncoding RNA: NEAT1 RNA is essential for the structure of paraspeckles. *Mol. Cell* 33, 717–726. doi: 10.1016/j.molcel.2009.01.026
- Conicella, A. E., Dignon, G. L., Zerze, G. H., Schmidt, H. B., D'Ordine, A. M., Kim, Y. C., et al. (2020). TDP-43 alpha-helical structure tunes liquid-liquid phase separation and function. *Proc. Natl. Acad. Sci. U S A.* 117, 5883–5894. doi: 10.1073/pnas.1912055117
- Corey, D. R. (2017). Nusinersen, an antisense oligonucleotide drug for spinal muscular atrophy. *Nat. Neurosci.* 20, 497–499. doi: 10.1038/nn.4508
- Corpet, A., Kleijwegt, C., Roubille, S., Juillard, F., Jacquet, K., Texier, P., et al. (2020). PML nuclear bodies and chromatin dynamics: Catch me if you can! *Nucleic Acids Res.* 48, 11890–11912. doi: 10.1093/nar/gkaa828
- Courchaine, E. M., Lu, A., and Neugebauer, K. M. (2016). Droplet organelles? *EMBO J.* 35, 1603–1612. doi: 10.15252/embj.201593517
- Davidson, Y. S., Raby, S., Foulds, P. G., Robinson, A., Thompson, J. C., Sikkink, S., et al. (2011). TDP-43 pathological changes in early onset familial and sporadic Alzheimer's disease, late onset Alzheimer's disease and Down's syndrome: Association with age, hippocampal sclerosis and clinical phenotype. *Acta Neuropathol.* 122, 703–713. doi: 10.1007/s00401-011-0879-y
- de Boer, E. M. J., Orié, V. K., Williams, T., Baker, M. R., De Oliveira, H. M., and Polvikoski, T. (2020). TDP-43 proteinopathies: A new wave of neurodegenerative diseases. *J. Neurol. Neurosurg. Psychiatry* 92, 86–95. doi: 10.1136/jnnp-2020-322983
- Dhaliwal, H. K., Fan, Y., Kim, J., and Amiji, M. M. (2020). Intranasal Delivery and Transfection of mRNA Therapeutics in the Brain Using Cationic Liposomes. *Mol. Pharm.* 17, 1996–2005. doi: 10.1021/acs.molpharmaceut.0c00170
- Ditlev, J. A. (2021). Membrane-associated phase separation: Organization and function emerge from a two-dimensional milieu. *J. Mol. Cell Biol.* 13, 319–324. doi: 10.1093/jmcb/mjab010
- Dong, M., Wang, X. P., Chen, X. Z., Mushtaq, F., Deng, S. Y., Zhu, C. H., et al. (2020). 3D-Printed Soft Magnetolectric Microswimmers for Delivery and Differentiation of Neuron-Like Cells. *Adv. Funct. Mater.* 30:1910323. doi: 10.1002/adfm.201910323
- Dugger, B. N., and Dickson, D. W. (2017). Pathology of Neurodegenerative Diseases. *Cold Spring Harb. Perspect. Biol.* 9:a028035. doi: 10.1101/cshperspect.a028035
- Emde, A., Eitan, C., Liou, L. L., Libby, R. T., Rivkin, N., Magen, I., et al. (2015). Dysregulated miRNA biogenesis downstream of cellular stress and ALS-causing mutations: A new mechanism for ALS. *EMBO J.* 34, 2633–2651. doi: 10.15252/embj.201490493
- Faber, G. P., Nadav-Eliyahu, S., and Shav-Tal, Y. (2022). Nuclear speckles - a driving force in gene expression. *J. Cell Sci.* 135:jcs259594. doi: 10.1242/jcs.259594
- Finkel, R. S., Chiriboga, C. A., Vajsar, J., Day, J. W., Montes, J., De Vivo, D. C., et al. (2016). Treatment of infantile-onset spinal muscular atrophy with nusinersen: A phase 2, open-label, dose-escalation study. *Lancet* 388, 3017–3026. doi: 10.1016/S0140-6736(16)31408-8
- Freischmidt, A., Müller, K., Ludolph, A. C., and Weishaupt, J. H. (2013). Systemic dysregulation of TDP-43 binding microRNAs in amyotrophic lateral sclerosis. *Acta Neuropathol. Commun.* 1:42. doi: 10.1186/2051-5960-1-42
- Frottin, F., Schueder, F., Tiwary, S., Gupta, R., Korner, R., Schlichthaerle, T., et al. (2019). The nucleolus functions as a phase-separated protein quality control compartment. *Science* 365, 342–347. doi: 10.1126/science.aaw9157
- Galganski, L., Urbaneck, M. O., and Krzyzosiak, W. J. (2017). Nuclear speckles: Molecular organization, biological function and role in disease. *Nucleic Acids Res.* 45, 10350–10368. doi: 10.1093/nar/gkx759
- Gammon, K. (2014). Neurodegenerative disease: Brain windfall. *Nature* 515, 299–300. doi: 10.1038/nj7526-299a
- Gouni-Berthold, I., Alexander, V. J., Yang, Q., Huh, E., Steinhagen-Thiessen, E., and Moriarty, P. M. (2021). Efficacy and safety of volanesorsen in patients with multifactorial chylomicronaemia (COMPASS): A multicentre, double-blind, randomised, placebo-controlled, phase 3 trial. *Lancet Diabetes Endocrinol.* 9, 264–275. doi: 10.1016/S2213-8587(21)00046-2
- Guerra San Juan, I., Nash, L. A., Smith, K. S., Leyton-Jaimes, M. F., Qian, M., Klim, J. R., et al. (2022). Loss of mouse Stmn2 function causes motor neuropathy. *Neuron* 110:e1676. doi: 10.1016/j.neuron.2022.02.011
- Higashi, S., Iseki, E., Yamamoto, R., Minegishi, M., Hino, H., Fujisawa, K., et al. (2007). Concurrence of TDP-43, tau and alpha-synuclein pathology in brains of Alzheimer's disease and dementia with Lewy bodies. *Brain Res.* 1184, 284–294.
- Hu, W. T., Josephs, K. A., Knopman, D. S., Boeve, B. F., Dickson, D. W., Petersen, R. C., et al. (2008). Temporal lobar predominance of TDP-43 neuronal cytoplasmic inclusions in Alzheimer disease. *Acta Neuropathol.* 116, 215–220.
- Huang, C., Li, J., Zhang, G., Lin, Y., Li, C., Zheng, X., et al. (2021). TBN improves motor function and prolongs survival in a TDP-43M337V mouse model of ALS. *Hum. Mol. Genet.* 30, 1484–1496. doi: 10.1093/hmg/ddab101
- Huang, W., Zhou, Y., Tu, L., Ba, Z., Huang, J., Huang, N., et al. (2020). TDP-43: From Alzheimer's Disease to Limbic-Predominant Age-Related TDP-43 Encephalopathy. *Front. Mol. Neurosci.* 13:26. doi: 10.3389/fnmol.2020.00026
- Humphrey, J., Emmett, W., Fratta, P., Isaacs, A. M., and Plagnol, V. (2017). Quantitative analysis of cryptic splicing associated with TDP-43 depletion. *BMC Med. Genomics* 10:38. doi: 10.1186/s12920-017-0274-1
- Hutchinson, J. N., Ensminger, A. W., Clemson, C. M., Lynch, C. R., Lawrence, J. B., and Chess, A. (2007). A screen for nuclear transcripts identifies two linked noncoding RNAs associated with SC35 splicing domains. *BMC Genomics* 8:39. doi: 10.1186/1471-2164-8-39
- Igaz, L. M., Kwong, L. K., Xu, Y., Truax, A. C., Uryu, K., Neumann, M., et al. (2008). Enrichment of C-terminal fragments in TAR DNA-binding protein-43 cytoplasmic inclusions in brain but not in spinal cord of frontotemporal lobar degeneration and amyotrophic lateral sclerosis. *Am. J. Pathol.* 173, 182–194. doi: 10.2353/ajpath.2008.080003
- Ilik, I. A., and Aktas, T. (2021). Nuclear speckles: Dynamic hubs of gene expression regulation. *FEBS J.* doi: 10.1111/febs.16117
- Imada, T., Shimi, T., Kaiho, A., Saeki, Y., and Kimura, H. (2021). RNA polymerase II condensate formation and association with Cajal and histone locus bodies in living human cells. *Genes Cells* 26, 298–312. doi: 10.1111/gtc.12840
- Izumikawa, K., Nobe, Y., Ishikawa, H., Yamauchi, Y., Taoka, M., Sato, K., et al. (2019). TDP-43 regulates site-specific 2'-O-methylation of U1 and U2 snRNAs via controlling the Cajal body localization of a subset of C/D scaRNAs. *Nucleic Acids Res.* 47, 2487–2505. doi: 10.1093/nar/gkz086
- Jeong, Y. H., Ling, J. P., Lin, S. Z., Donde, A. N., Braunstein, K. E., Majounie, E., et al. (2017). Tdp-43 cryptic exons are highly variable between cell types. *Mol. Neurodegener.* 12:13. doi: 10.1186/s13024-016-0144-x
- Jiang, L., Shao, C., Wu, Q. J., Chen, G., Zhou, J., Yang, B., et al. (2017). NEAT1 scaffolds RNA-binding proteins and the Microprocessor to globally enhance pri-miRNA processing. *Nat. Struct. Mol. Biol.* 24, 816–824. doi: 10.1038/nsmb.3455
- Jiang, Y. X., Cao, Q., Sawaya, M. R., Abskharon, R., Ge, P., DeTure, M., et al. (2022). Amyloid fibrils in FTLTDP are composed of TMEM106B and not TDP-43. *Nature* 605, 304–309. doi: 10.1038/s41586-022-04670-9
- Josephs, K. A., Murray, M. E., Tosakulwong, N., Weigand, S. D., Serie, A. M., Perkerson, R. B., et al. (2019). Pathological, imaging and genetic characteristics support the existence of distinct TDP-43 types in non-FTLD brains. *Acta Neuropathol.* 137, 227–238. doi: 10.1007/s00401-018-1951-7
- Josephs, K. A., Murray, M. E., Whitwell, J. L., Tosakulwong, N., Weigand, S. D., Petrucelli, L., et al. (2016). Updated TDP-43 in Alzheimer's disease staging scheme. *Acta Neuropathol.* 131, 571–585. doi: 10.1007/s00401-016-1537-1

- Josephs, K. A., Whitwell, J. L., Weigand, S. D., Murray, M. E., Tosakulwong, N., Liesinger, A. M., et al. (2014). TDP-43 is a key player in the clinical features associated with Alzheimer's disease. *Acta Neuropathol.* 127, 811–824. doi: 10.1007/s00401-014-1269-z
- Jumper, J., Evans, R., Pritzel, A., Green, T., Figurnov, M., Ronneberger, O., et al. (2021). Highly accurate protein structure prediction with AlphaFold. *Nature* 596, 583–589. doi: 10.1038/s41586-021-03819-2
- Kadokura, A., Yamazaki, T., Lemere, C. A., Takatama, M., and Okamoto, K. (2009). Regional distribution of TDP-43 inclusions in Alzheimer disease (AD) brains: Their relation to AD common pathology. *Neuropathology* 29, 566–573. doi: 10.1111/j.1440-1789.2009.01017.x
- Kawahara, Y., and Mieda-Sato, A. (2012). TDP-43 promotes microRNA biogenesis as a component of the Drosha and Dicer complexes. *Proc. Natl. Acad. Sci. U S A.* 109, 3347–3352. doi: 10.1073/pnas.1112427109
- Kim, E., Jeon, S., An, H.-K., Kianpour, M., Yu, S.-W., Kim, J.-Y., et al. (2020). A magnetically actuated microrobot for targeted neural cell delivery and selective connection of neural networks. *Sci. Adv.* 6:eabb5696. doi: 10.1126/sciadv.abb5696
- King, A. (2022). Miniature medical robots step out from sci-fi. *Nature* 2022:9. doi: 10.1038/d41586-022-00859-0
- King, A., Sweeney, F., Bodi, I., Troakes, C., Maekawa, S., and Al-Sarraj, S. (2010). Abnormal TDP-43 expression is identified in the neocortex in cases of dementia pugilistica, but is mainly confined to the limbic system when identified in high and moderate stages of Alzheimer's disease. *Neuropathology* 30, 408–419. doi: 10.1111/j.1440-1789.2009.01085.x
- Klim, J. R., Williams, L. A., Limone, F., Guerra San Juan, I., Davis-Dusenbery, B. N., Mordes, D. A., et al. (2019). ALS-implicated protein TDP-43 sustains levels of STMN2, a mediator of motor neuron growth and repair. *Nat. Neurosci.* 22, 167–179. doi: 10.1038/s41593-018-0300-4
- Klingauf, M., Stanek, D., and Neugebauer, K. M. (2006). Enhancement of U4/U6 small nuclear ribonucleoprotein particle association in Cajal bodies predicted by mathematical modeling. *Mol. Biol. Cell* 17, 4972–4981. doi: 10.1091/mbc.e06-06-0513
- Kovacs, G. G. (2017). Concepts and classification of neurodegenerative diseases. *Handb. Clin. Neurol.* 145, 301–307. doi: 10.1016/B978-0-12-802395-2.00021-3
- Krus, K. L., Strickland, A., Yamada, Y., Devault, L., Schmidt, R. E., Bloom, A. J., et al. (2022). Loss of Stathmin-2, a hallmark of TDP-43-associated ALS, causes motor neuropathy. *Cell Rep.* 39:111001. doi: 10.1016/j.celrep.2022.111001
- Kukharsky, M. S., Ninkina, N. N., An, H., Telezhkin, V., Wei, W., Meritens, C. R., et al. (2020). Long non-coding RNA Neat1 regulates adaptive behavioural response to stress in mice. *Transl. Psychiatry* 10:171. doi: 10.1038/s41398-020-0854-2
- Kulkarni, J. A., Witzigmann, D., Thomson, S. B., Chen, S., Leavitt, B. R., Cullis, P. R., et al. (2021). The current landscape of nucleic acid therapeutics. *Nat. Nanotechnol.* 16, 630–643. doi: 10.1038/s41565-021-00898-0
- Kwon, D. (2021). Failure of genetic therapies for Huntington's devastates community. *Nature* 593:180. doi: 10.1038/d41586-021-01177-7
- Lafarga, M., Tapia, O., Romero, A. M., and Berciano, M. T. (2017). Cajal bodies in neurons. *RNA Biol.* 14, 712–725. doi: 10.1080/15476286.2016.1231360
- Lafontaine, D. L. J., Riback, J. A., Bascetin, R., and Brangwynne, C. P. (2021). The nucleolus as a multiphase liquid condensate. *Nat. Rev. Mol. Cell Biol.* 22, 165–182. doi: 10.1038/s41580-020-0272-6
- Li, J., Esteban-Fernandez de Avila, B., Gao, W., Zhang, L., and Wang, J. (2017). Micro/Nanorobots for Biomedicine: Delivery, Surgery, Sensing, and Detoxification. *Sci. Robot.* 2:eaa6431. doi: 10.1126/scirobotics.aam6431
- Li, R., Harvey, A. R., Hodgetts, S. I., and Fox, A. H. (2017). Functional dissection of NEAT1 using genome editing reveals substantial localization of the NEAT1_1 isoform outside paraspeckles. *RNA* 23, 872–881. doi: 10.1261/rna.059477.116
- Ling, J. P., Pletnikova, O., Troncoso, J. C., and Wong, P. C. (2015). TDP-43 repression of nonconserved cryptic exons is compromised in ALS-FTD. *Science* 349, 650–655. doi: 10.1126/science.aab0983
- Lissouba, A., Liao, M., Kabashi, E., and Drapeau, P. (2018). Transcriptomic Analysis of Zebrafish TDP-43 Transgenic Lines. *Front. Mol. Neurosci.* 11:463. doi: 10.3389/fnmol.2018.00463
- Ma, X. R., Prudencio, M., Koike, Y., Vatsavayai, S. C., Kim, G., Harbinski, F., et al. (2022). TDP-43 represses cryptic exon inclusion in the FTD-ALS gene UNC13A. *Nature* 603, 124–130. doi: 10.1038/s41586-022-04424-7
- Mallik, M., and Lakhotia, S. C. (2009). RNAi for the large non-coding hsr omega transcripts suppresses polyglutamine pathogenesis in Drosophila models. *RNA Biol.* 6, 464–478. doi: 10.4161/rna.6.4.9268
- Mann, J. R., Gleixner, A. M., Mauna, J. C., Gomes, E., DeChellis-Marks, M. R., Needham, P. G., et al. (2019). RNA Binding Antagonizes Neurotoxic Phase Transitions of TDP-43. *Neuron* 102:e328. doi: 10.1016/j.neuron.2019.01.048
- Mao, Y. S., Zhang, B., and Spector, D. L. (2011). Biogenesis and function of nuclear bodies. *Trends Genet.* 27, 295–306. doi: 10.1016/j.tig.2011.05.006
- Mathew, A. S., Gorick, C. M., and Price, R. J. (2021). Single-cell mapping of focused ultrasound-transfected brain. *Gene Ther.* [Preprint]. doi: 10.1038/s41434-021-00226-0
- Matsukawa, K., Kukharsky, M. S., Park, S. K., Park, S., Watanabe, N., Iwatsubo, T., et al. (2021). Long non-coding RNA NEAT1_1 ameliorates TDP-43 toxicity in vivo models of TDP-43 proteinopathy. *RNA Biol.* 18, 1546–1554. doi: 10.1080/15476286.2020.1860580
- McAleese, K. E., Walker, L., Erskine, D., Thomas, A. J., McKeith, I. G., and Attems, J. (2019). TDP-43 pathology in Alzheimer's disease, dementia with Lewy bodies and ageing. *Brain Pathol.* 27, 472–479. doi: 10.1111/bpa.12424
- Melamed, Z., Lopez-Erauskin, J., Baughn, M. W., Zhang, O., Drenner, K., Sun, Y., et al. (2019). Premature polyadenylation-mediated loss of stathmin-2 is a hallmark of TDP-43-dependent neurodegeneration. *Nat. Neurosci.* 22, 180–190. doi: 10.1038/s41593-018-0293-z
- Melo, S., Villanueva, A., Moutinho, C., Davalos, V., Spizzo, R., Ivan, C., et al. (2011). Small molecule enoxacin is a cancer-specific growth inhibitor that acts by enhancing TAR RNA-binding protein 2-mediated microRNA processing. *Proc. Natl. Acad. Sci. U S A.* 108, 4394–4399. doi: 10.1073/pnas.1014720108
- Meneses, A., Koga, S., O'Leary, J., Dickson, D. W., Bu, G., and Zhao, N. (2021). TDP-43 Pathology in Alzheimer's Disease. *Mol. Neurodegener.* 16:84. doi: 10.1186/s13024-021-00503-x
- Mercer, T. R., Qureshi, I. A., Gokhan, S., Dinger, M. E., Li, G., Mattick, J. S., et al. (2010). Long noncoding RNAs in neuronal-glia fate specification and oligodendrocyte lineage maturation. *BMC Neurosci.* 11:14. doi: 10.1186/1471-2202-11-14
- Mitrea, D. M., Cika, J. A., Guy, C. S., Ban, D., Banerjee, P. R., Stanley, C. B., et al. (2016). Nucleophosmin integrates within the nucleolus via multimodal interactions with proteins displaying R-rich linear motifs and rRNA. *eLife* 5:e13571. doi: 10.7554/eLife.13571
- Mori, F., Tanji, K., Zhang, H. X., Nishihira, Y., Tan, C. F., Takahashi, H., et al. (2008). Maturation process of TDP-43-positive neuronal cytoplasmic inclusions in amyotrophic lateral sclerosis with and without dementia. *Acta Neuropathol.* 116, 193–203. doi: 10.1007/s00401-008-0396-9
- Morris, G. E. (2008). The Cajal body. *Biochim. Biophys. Acta* 1783, 2108–2115. doi: 10.1016/j.bbamcr.2008.07.016
- Nag, S., Yu, L., Capuano, A. W., Wilson, R. S., Leurgans, S. E., Bennett, D. A., et al. (2015). Hippocampal sclerosis and TDP-43 pathology in aging and Alzheimer disease. *Ann. Neurol.* 77, 942–952. doi: 10.1002/ana.24388
- Nakagawa, S., Naganuma, T., Shioi, G., and Hirose, T. (2011). Paraspeckles are subpopulation-specific nuclear bodies that are not essential in mice. *J. Cell Biol.* 193, 31–39. doi: 10.1083/jcb.201011110
- Nakagawa, S., Shimada, M., Yanaka, K., Mito, M., Arai, T., Takahashi, E., et al. (2014). The lncRNA Neat1 is required for corpus luteum formation and the establishment of pregnancy in a subpopulation of mice. *Development* 141, 4618–4627. doi: 10.1242/dev.110544
- Neumann, M., Sampathu, D. M., Kwong, L. K., Truax, A. C., Micsenyi, M. C., Chou, T. T., et al. (2006). Ubiquitinated TDP-43 in frontotemporal lobar degeneration and amyotrophic lateral sclerosis. *Science* 314, 130–133. doi: 10.1126/science.1134108
- Nicolas, A., Kenna, K. P., Renton, A. E., Ticozzi, N., Faghri, F., and Chia, R. (2018). Genome-wide Analyses Identify KIF5A as a Novel ALS Gene. *Neuron* 97:e1266.
- Nishimoto, Y., Nakagawa, S., Hirose, T., Okano, H. J., Takao, M., Shibata, S., et al. (2013). The long non-coding RNA nuclear-enriched abundant transcript 1_2 induces paraspeckle formation in the motor neuron during the early phase of amyotrophic lateral sclerosis. *Mol. Brain* 6:31. doi: 10.1186/1756-6606-6-31
- Ou, S. H., Wu, F., Harrich, D., Garcia-Martinez, L. F., and Gaynor, R. B. (1995). Cloning and characterization of a novel cellular protein, TDP-43, that binds to human immunodeficiency virus type 1 TAR DNA sequence motifs. *J. Virol.* 69, 3584–3596. doi: 10.1128/jvi.69.6.3584-3596.1995
- Peggion, C., Massimino, M. L., Stella, R., Bortolotto, R., Agostini, J., Maldì, A., et al. (2021). Nucleolin Rescues TDP-43 Toxicity in Yeast and Human Cell Models. *Front. Cell. Neurosci.* 15:625665. doi: 10.3389/fncel.2021.625665
- Peng, P. H., Hsu, K. W., and Wu, K. J. (2021). Liquid-liquid phase separation (LLPS) in cellular physiology and tumor biology. *Am. J. Cancer Res.* 11, 3766–3776.

- Peyer, K. E., Zhang, L., and Nelson, B. J. (2013). Bio-inspired magnetic swimming microrobots for biomedical applications. *Nanoscale* 5, 1259–1272. doi: 10.1039/C2NR32554C
- Pickles, S., Gendron, T. F., Koike, Y., Yue, M., Song, Y., Kachergus, J. M., et al. (2022). Evidence of cerebellar TDP-43 loss of function in FTLD-TDP. *Acta Neuropathol. Commun.* 10:107. doi: 10.1186/s40478-022-01408-6
- Prasad, A., Bharathi, V., Sivalingam, V., Girdhar, A., and Patel, B. K. (2019). Molecular Mechanisms of TDP-43 Misfolding and Pathology in Amyotrophic Lateral Sclerosis. *Front. Mol. Neurosci.* 12:25. doi: 10.3389/fnmol.2019.00025
- Prasad, A., Raju, G., Sivalingam, V., Girdhar, A., Verma, M., Vats, A., et al. (2016). An acridine derivative, [4,5-bis{(N-carboxy methyl imidazolium)methyl}acridine] dibromide, shows anti-TDP-43 aggregation effect in ALS disease models. *Sci. Rep.* 6:39490. doi: 10.1038/srep39490
- Prudencio, M., Humphrey, J., Pickles, S., Brown, A. L., Hill, S. E., Kachergus, J. M., et al. (2020). Truncated stathmin-2 is a marker of TDP-43 pathology in frontotemporal dementia. *J. Clin. Invest.* 130, 6080–6092. doi: 10.1172/JCI139741
- Ratti, A., and Buratti, E. (2016). Physiological functions and pathobiology of TDP-43 and FUS/TLS proteins. *J. Neurochem.* 138, 95–111. doi: 10.1111/jnc.13625
- Rengifo-Gonzalez, J. C., El Hage, K., Clement, M. J., Steiner, E., Joshi, V., Craveur, P., et al. (2021). The cooperative binding of TDP-43 to GU-rich RNA repeats antagonizes TDP-43 aggregation. *eLife* 10:e67605. doi: 10.7554/eLife.67605
- Sasaki, Y. T., Ideue, T., Sano, M., Mituyama, T., and Hirose, T. (2009). MENepsilon/beta noncoding RNAs are essential for structural integrity of nuclear paraspeckles. *Proc. Natl. Acad. Sci. U S A.* 106, 2525–2530. doi: 10.1073/pnas.0807899106
- Sawyer, I. A., Bartek, J., and Dundr, M. (2019). Phase separated microenvironments inside the cell nucleus are linked to disease and regulate epigenetic state, transcription and RNA processing. *Semin. Cell Dev. Biol.* 90, 94–103. doi: 10.1016/j.semcdb.2018.07.001
- Schmit, J. D., Feric, M., and Dundr, M. (2021). How Hierarchical Interactions Make Membraneless Organelles Tick Like Clockwork. *Trends Biochem. Sci.* 46, 525–534. doi: 10.1016/j.tibs.2020.12.011
- Schweighauser, M., Arseni, D., Bacioglu, M., Huang, M., Lovestam, S., Shi, Y., et al. (2022). Age-dependent formation of TMEM106B amyloid filaments in human brains. *Nature* 605, 310–314. doi: 10.1038/s41586-022-04650-z
- Scott, D. D., Mowrey, D., Nagarajan, K., François-Moutal, L., Nair, A., and Khanna, M. (2022). Molecular Dynamics simulation of TDP-43 RRM in the presence and absence of RNA. *bioRxiv* [Preprint]. doi: 10.1101/2022.03.15.484514
- Shan, G., Li, Y., Zhang, J., Li, W., Szulwach, K. E., Duan, R., et al. (2008). A small molecule enhances RNA interference and promotes microRNA processing. *Nat. Biotechnol.* 26, 933–940. doi: 10.1038/nbt.1481
- Shao, Y., Fahmy, A., Li, M., Li, C., Zhao, W., and Sieng, J. (2021). Study on Magnetic Control Systems of Micro-Robots. *Front. Neurosci.* 15:736730. doi: 10.3389/fnins.2021.736730
- Shaw, D. J., Eggleton, P., and Young, P. J. (2008). Joining the dots: Production, processing and targeting of U snRNP to nuclear bodies. *Biochim. Biophys. Acta* 1783, 2137–2144. doi: 10.1016/j.bbamcr.2008.07.025
- Shelkovnikova, T. A., Kukharsky, M. S., An, H., Dimasi, P., Alexeeva, S., Shabir, O., et al. (2018). Protective paraspeckle hyper-assembly downstream of TDP-43 loss of function in amyotrophic lateral sclerosis. *Mol. Neurodegener.* 13:30. doi: 10.1186/s13024-018-0263-7
- Shen, W., Liang, X. H., and Crooke, S. T. (2014). Phosphorothioate oligonucleotides can displace NEAT1 RNA and form nuclear paraspeckle-like structures. *Nucleic Acids Res.* 42, 8648–8662. doi: 10.1093/nar/gku579
- Shevtsov, S. P., and Dundr, M. (2011). Nucleation of nuclear bodies by RNA. *Nat. Cell Biol.* 13, 167–173. doi: 10.1038/ncb2157
- Shirley, M. (2021). Casimersen: First Approval. *Drugs* 81, 875–879. doi: 10.1007/s40265-021-01512-2
- Shopland, L. S., Byron, M., Stein, J. L., Lian, J. B., Stein, G. S., and Lawrence, J. B. (2001). Replication-dependent histone gene expression is related to Cajal body (CB) association but does not require sustained CB contact. *Mol. Biol. Cell* 12, 565–576. doi: 10.1091/mbc.12.3.565
- Souquere, S., Beauclair, G., Harper, F., Fox, A., and Pierron, G. (2010). Highly ordered spatial organization of the structural long noncoding NEAT1 RNAs within paraspeckle nuclear bodies. *Mol. Biol. Cell* 21, 4020–4027. doi: 10.1091/mbc.e10-08-0690
- Sun, M., Bell, W., LaClair, K. D., Ling, J. P., Han, H., Kageyama, Y., et al. (2017). Cryptic exon incorporation occurs in Alzheimer's brain lacking TDP-43 inclusion but exhibiting nuclear clearance of TDP-43. *Acta Neuropathol.* 133, 923–931. doi: 10.1007/s00401-017-1701-2
- Sunwoo, H., Dinger, M. E., Wilusz, J. E., Amaral, P. P., Mattick, J. S., and Spector, D. L. (2009). MEN epsilon/beta nuclear-retained non-coding RNAs are up-regulated upon muscle differentiation and are essential components of paraspeckles. *Genome Res.* 19, 347–359. doi: 10.1101/gr.087775.108
- Sunwoo, J. S., Lee, S. T., Im, W., Lee, M., Byun, J. I., Jung, K. H., et al. (2017). Altered Expression of the Long Noncoding RNA NEAT1 in Huntington's Disease. *Mol. Neurobiol.* 54, 1577–1586. doi: 10.1007/s12035-016-9928-9
- Suzuki, H., Abe, R., Shimada, M., Hirose, T., Hirose, H., Noguchi, K., et al. (2012). The 3' Pol II pausing at replication-dependent histone genes is regulated by Mediator through Cajal bodies' association with histone locus bodies. *Nat. Commun.* 13:2905. doi: 10.1038/s41467-022-30632-w
- Tanaka, H., Nakatani, T., Furihata, T., Tange, K., Nakai, Y., Yoshioka, H., et al. (2018). In Vivo Introduction of mRNA Encapsulated in Lipid Nanoparticles to Brain Neuronal Cells and Astrocytes via Intracerebroventricular Administration. *Mol. Pharm.* 15, 2060–2067. doi: 10.1021/acs.molpharmaceut.7b01084
- Taylor, J. P., Brown, R. H. Jr., and Cleveland, D. W. (2016). Decoding ALS: From genes to mechanism. *Nature* 539, 197–206. doi: 10.1038/nature20413
- Theodoridis, P. R., Bokros, M., Marijan, D., Balukoff, N. C., Wang, D., Kirk, C. C., et al. (2021). Local translation in nuclear condensate amyloid bodies. *Proc. Natl. Acad. Sci. U S A.* 118:e2014457118. doi: 10.1073/pnas.2014457118
- Tollervey, J. R., Curk, T., Rogelj, B., Briese, M., Cereda, M., Kayikci, M., et al. (2011). Characterizing the RNA targets and position-dependent splicing regulation by TDP-43. *Nat. Neurosci.* 14, 452–458. doi: 10.1038/nn.2778
- Uchida, A., Sasaguri, H., Kimura, N., Tajiri, M., Ohkubo, T., Ono, F., et al. (2018). Non-human primate model of amyotrophic lateral sclerosis with cytoplasmic mislocalization of TDP-43. *Brain* 135, 833–846. doi: 10.1093/brain/awr348
- Uryu, K., Nakashima-Yasuda, H., Forman, M. S., Kwong, L. K., Clark, C. M., Grossman, M., et al. (2008). Concomitant TAR-DNA-binding protein 43 pathology is present in Alzheimer disease and corticobasal degeneration but not in other tauopathies. *J. Neuropathol. Exp. Neurol.* 67, 555–564. doi: 10.1097/NEN.0b013e31817713b5
- Valgardsdottir, R., Chiodi, I., Giordano, M., Cobianchi, F., Riva, S., and Biamonti, G. (2005). Structural and functional characterization of noncoding repetitive RNAs transcribed in stressed human cells. *Mol. Biol. Cell* 16, 2597–2604. doi: 10.1091/mbc.e04-12-1078
- Valgardsdottir, R., Chiodi, I., Giordano, M., Rossi, A., Bazzini, S., Ghigna, C., et al. (2008). Transcription of Satellite III non-coding RNAs is a general stress response in human cells. *Nucleic Acids Res.* 36, 423–434. doi: 10.1093/nar/gkm1056
- van Rheenen, W., Shatunov, A., Dekker, A. M., McLaughlin, R. L., Diekstra, F. P., and Pulit, S. L. (2016). Genome-wide association analyses identify new risk variants and the genetic architecture of amyotrophic lateral sclerosis. *Nat. Genet.* 48, 1043–1048. doi: 10.1038/ng.3622
- van Rheenen, W., van der Spek, R. A. A., Bakker, M. K., van Vugt, J., Hop, P. J., and Zwaborn, R. A. J. (2021). Common and rare variant association analyses in amyotrophic lateral sclerosis identify 15 risk loci with distinct genetic architectures and neuron-specific biology. *Nat. Genet.* 53, 1636–1648. doi: 10.1038/s41588-021-00973-1
- Wagner, K. R., Kuntz, N. L., Koenig, E., East, L., Upadhyay, S., Han, B., et al. (2021). Safety, tolerability, and pharmacokinetics of casimersen in patients with Duchenne muscular dystrophy amenable to exon 45 skipping: A randomized, double-blind, placebo-controlled, dose-titration trial. *Muscle Nerve* 64, 285–292. doi: 10.1002/mus.27347
- Walker, M. P., Tian, L., and Matera, A. G. (2009). Reduced viability, fertility and fecundity in mice lacking the cajal body marker protein, coilin. *PLoS One* 4:e6171. doi: 10.1371/journal.pone.0006171
- Wang, B., Zhang, L., Dai, T., Qin, Z., Lu, H., Zhang, L., et al. (2021). Liquid-liquid phase separation in human health and diseases. *Signal Transduct. Target. Ther.* 6:290. doi: 10.1038/s41392-021-00678-1
- Wang, C., Duan, Y., Duan, G., Wang, Q., Zhang, K., Deng, X., et al. (2020). Stress Induces Dynamic, Cytotoxicity-Antagonizing TDP-43 Nuclear Bodies via Paraspeckle LncRNA NEAT1-Mediated Liquid-Liquid Phase Separation. *Mol. Cell* 79:e447. doi: 10.1101/802058
- Wang, W., Wang, L., Lu, J., Siedlak, S. L., Fujioka, H., Liang, J., et al. (2016). The inhibition of TDP-43 mitochondrial localization blocks its neuronal toxicity. *Nat. Med.* 22, 869–878. doi: 10.1038/nm.4130
- Xu, Y. F., Gendron, T. F., Zhang, Y. J., Lin, W. L., D'Alton, S., Sheng, H., et al. (2010). Wild-type human TDP-43 expression causes TDP-43 phosphorylation,

mitochondrial aggregation, motor deficits, and early mortality in transgenic mice. *J. Neurosci.* 30, 10851–10859. doi: 10.1523/JNEUROSCI.1630-10.2010

Yamashita, A., Watanabe, Y., Nukina, N., and Yamamoto, M. (1998). RNA-assisted nuclear transport of the meiotic regulator Mei2p in fission yeast. *Cell* 95, 115–123. doi: 10.1016/S0092-8674(00)81787-0

Yamazaki, T., Souquere, S., Chujo, T., Kobelke, S., Chong, Y. S., Fox, A. H., et al. (2018). Functional Domains of NEAT1 Architectural lncRNA Induce Paraspeckle Assembly through Phase Separation. *Mol. Cell* 70:e1037. doi: 10.1016/j.molcel.2018.05.019

Yamazaki, T., Yamamoto, T., Yoshino, H., Souquere, S., Nakagawa, S., Pierron, G., et al. (2021). Paraspeckles are constructed as block copolymer micelles. *EMBO J.* 40:e107270. doi: 10.15252/embj.2020107270

Yang, Z., and Zhang, L. (2020). Magnetic Actuation Systems for Miniature Robots: A Review. *Adv. Intel. Syst.* 2:2000082. doi: 10.1002/aisy.202000082

Yao, R. W., Xu, G., Wang, Y., Shan, L., Luan, P. F., Wang, Y., et al. (2019). Nascent Pre-rRNA Sorting via Phase Separation Drives the Assembly of Dense Fibrillar Components in the Human Nucleolus. *Mol. Cell* 76:e711.

Yin, P., Guo, X., Yang, W., Yan, S., Yang, S., Zhao, T., et al. (2019). Caspase-4 mediates cytoplasmic accumulation of TDP-43 in the primate brains. *Acta Neuropathol.* 137, 919–937. doi: 10.1007/s00401-019-01979-0

Yu, S., Cai, Y., Wu, Z., and He, Q. (2021). Recent progress on motion control of swimming micro/nanorobots. *View* 2:20200113. doi: 10.1002/VIW.20200113

Zhang, S., Cooper-Knock, J., Weimer, A. K., Shi, M., Moll, T., and Marshall, J. N. G. (2022). Genome-wide identification of the genetic basis of amyotrophic lateral sclerosis. *Neuron* 110:e1011.

Zuo, X., Zhou, J., Li, Y., Wu, K., Chen, Z., Luo, Z., et al. (2021). TDP-43 aggregation induced by oxidative stress causes global mitochondrial imbalance in ALS. *Nat. Struct. Mol. Biol.* 28, 132–142. doi: 10.1038/s41594-020-00537-7