

Review

# Hyperglycaemia-Linked Diabetic Foot Complications and Their Management Using Conventional and Alternative Therapies

Prawej Ansari <sup>1,2,\*</sup>, Samia Akther <sup>1</sup>, Joyeeta T. Khan <sup>1</sup>, Sara S. Islam <sup>1</sup>, Md. Samim R. Masud <sup>1</sup>, Anisur Rahman <sup>1</sup>, Veronique Seidel <sup>3</sup> and Yasser H. A. Abdel-Wahab <sup>2</sup>

<sup>1</sup> Department of Pharmacy, School of Pharmacy and Public Health, Independent University, Bangladesh (IUB), Dhaka 1229, Bangladesh

<sup>2</sup> School of Biomedical Sciences, Ulster University, Coleraine BT52 1SA, UK

<sup>3</sup> Natural Products Research Laboratory, Strathclyde Institute of Pharmacy and Biomedical Sciences, University of Strathclyde, Glasgow G4 0RE, UK

\* Correspondence: pr.ansari@iub.edu.bd; Tel.: +880-132-387-9720

**Citation:** Ansari, P.; Akther, S.; Khan, J.T.; Islam, S.S.; Masud, M.S.R.; Rahman, A.; Seidel, V.; Abdel-Wahab, Y.H.A. Hyperglycaemia-Linked Diabetic Foot Complications and Their Management using Conventional and Alternative Therapies. *Appl. Sci.* **2022**, *12*, 11777. <https://doi.org/10.3390/app122211777>

Academic Editor: Marco G. Alves

Received: 18 October 2022

Accepted: 15 November 2022

Published: 19 November 2022

**Publisher's Note:** MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Copyright:** © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

**Abstract:** Diabetes mellitus, a major cause of mortality around the globe, can result in several secondary complications, including diabetic foot syndrome, which is brought on by diabetic neuropathy and ischemia. Approximately 15% of diabetic patients suffer from diabetic foot complications, and among them 25% are at risk of lower limb amputations. Diabetic foot ulcers are characterized as skin lesions, gangrene, or necrosis, and may develop due to several reasons, including hyperglycemia and slower wound healing in diabetic patients. A management protocol involving wound cleaning, oral antibiotics, skin ointments, and removing dead tissue is currently followed to treat diabetic foot ulcers. In severe cases, amputation is performed to prevent the infection from spreading further. The existing therapy can be costly and present adverse side effects. Combined with a lack of vascular surgeons, this ultimately results in disability, especially in developing nations. There is a growing interest in the use of alternative therapies, such as medicinal plants, to discover more efficient and affordable treatments for diabetic foot syndrome. It has been observed that treatment with numerous plants, including *Carica papaya*, *Annona squamosa*, *Catharanthus roseus*, and *Centella asiatica*, promotes wound healing, reduces inflammation, and may decrease the number of amputations. However, little information is currently available on the prevention and management of diabetic foot ulcers, and additional research is necessary to completely understand the role of alternative therapies in the treatment of diabetic foot complications.

**Keywords:** hyperglycemia; diabetic foot; infection; glucose; alternative medicine

## 1. Introduction

Diabetes mellitus (DM) is considered a worldwide epidemic, being one of the top ten diseases causing mortality globally and affecting more than 10.5% of the adult population [1,2]. It is a chronic metabolic disorder manifested by persistent high blood glucose levels occurring due to deficiencies in insulin production, insulin resistance, or both [3]. According to its etiology and pathogenesis, diabetes mellitus can be broadly classified into two major types: type 1 diabetes mellitus (T1DM) and type 2 diabetes mellitus (T2DM) [4]. T1DM is a result of the autoimmune destruction of the pancreatic  $\beta$ -cells of the islets of Langerhans, leading to significantly diminished insulin secretion, whereas T2DM, often associated with obesity and an unhealthy lifestyle, is the outcome of either impaired insulin secretion, decreased insulin sensitivity to cells, or both [5,6]. The long-term secondary complications of diabetes include coronary heart disease, peripheral

artery disease, cardiomyopathy, stroke, and cerebrovascular disease, as well as diabetic retinopathy (which causes visual impairment), diabetic nephropathy (which leads to kidney failure), and diabetic neuropathy (which increases the risk of foot ulcers) [7–9]. Individuals with long-term DM are prone to these complications, which may lead to increased morbidity and mortality [10].

Diabetic foot is a major complication of DM that affects approximately 15% of diabetic patients, with 25% of them facing the possibility of lower limb amputation [11–13]. Diabetic patients are often hospitalized due to the consequences of foot-related conditions such as infections, ulcers, and gangrene or foot necrosis [14]. The ulceration is the starting point for a severe stage of the disease, and if left untreated, this can lead to amputations. Diabetic neuropathy, neuro-ischemia, and infections all have an impact on whether lesions heal or deteriorate, and amputations in diabetic patients are often precipitated by a foot ulcer that progresses to serious gangrene or infection [12].

To accelerate the healing process of diabetic foot ulcerations, various interventions such as wound cleansing, revascularization, antibiotic therapy, dead tissue removal, and in extreme situations surgery have been implemented [15]. The conventional therapy for treating diabetic foot is often impracticable, particularly in underdeveloped countries, because of the high costs, adverse side effects, and the unavailability of vascular surgeons. In recent years, there has been a surge of interest in the search for alternative medications derived from natural resources, especially medicinal plants, for the prevention and treatment of diabetic foot [16,17].

Since ancient times, herbal remedies have been used as effective therapies to treat a wide range of ailments, including diabetic foot [16]. Recent findings have revealed that biologically-active phytochemicals present in plants demonstrate various pharmacological effects that help to prevent different forms of cell damage, including chronic wounds [17]. Numerous medicinal plants, including *Aloe vera*, *Annona squamosa*, *Azadirachta indica*, *Carica papaya*, *Catharanthus roseus*, *Centella asiatica*, *Curcuma longa*, *Hylocereus undatus*, and *Punica granatum*, have exhibited potential wound healing, anti-inflammatory, antioxidative, antibacterial, and other antidiabetic properties with little to no side effects, and might be considered as possible effective treatments for diabetic injuries [18–20]. Several bioactive phytochemicals such as quercetin, kaempferol, isoquercitrin, apigenin, tangeretin, naringenin, luteolin, catechin, gallic acid, methyl gallate, and rutin present in medicinal plants including *Allium cepa*, *Beta vulgaris*, *Citrus sinensis*, *Schinus polygamus*, and *Sorghum bicolor* exert their antidiabetic action via different mechanistic pathways to improve  $\beta$ -cell function and insulin secretion; enhance insulin sensitivity; increase GLUT-4 expression; decrease gluconeogenesis; and inhibit  $\alpha$ -amylase,  $\alpha$ -glucosidase, and DPP-IV activity and the formation of advanced glycation end products. Such phytochemicals not only ameliorate diabetic foot wounds but also contribute to the overall antidiabetic action of the medicinal plants to prevent and manage other diabetes-related complications [21,22]. Thus, due to the abundance of bioactive compounds in plants, researchers have directed their focus toward examining the significance of medicinal plants and isolating their phytoconstituents to assess their prospective wound-healing properties [18]. This review article primarily aims to explore the role of traditional medicinal plants and their isolated phytochemicals in the prevention and management of diabetic foot syndrome, as well as their future scope as antidiabetic medications.

## 2. Epidemiology

Nearly 537 million people are currently suffering from diabetes mellitus worldwide. This is projected to rise to 783 million by the year 2045 [2]. Diabetes mellitus is the primary cause of non-traumatic lower limb amputations across the globe [23]. This growing incidence of DM can lead to the increased prevalence of diabetic foot complications, which have become a serious medical, social, and economic concern of global importance [24]. Diabetic foot ulcers are foot lesions that damage the skin, soft tissues, and bones in the

legs and feet, generating an aggravated infection in diabetic patients and potentially leading to lower limb amputations. Recent reports showed that around 60 to 80% of these ulcers can heal, 10 to 15% can stay active, and 5 to 24% eventually result in limb amputation (Figure 1) [23].

More than 85% of all amputations in diabetic foot patients are preceded by ulceration leading to severe gangrene or infection [12,25]. On average, about 6.4% of the worldwide population suffers from diabetic foot, with 2 to 5% yearly occurrence rates of ulcer or necrosis [26]. It is predicted that 19 to 34% of diabetic individuals will encounter foot disease at some stage in their lives [23]. A systemic meta-analytical review has shown that diabetic foot ulcers (DFU) are more common in men (4.5%) than in women (3.5%) and more prevalent in type 2 diabetic patients (6.4%) compared to type 1 individuals (5.5%) (Figure 1). Despite the rapidly growing prevalence of diabetic foot, only a few studies have been performed regarding its epidemiology. Thus, a detailed epidemiological study on the global update of diabetic foot is crucial in order for improved health and to minimize the financial strain on diabetic patients by preventing, controlling, and properly treating diabetic foot ulcerations [27].




Figure 1. Incidence rates of diabetic foot ulcers in T1DM and T2DM and their conditions.

### 3. Etiology


Diabetic foot ulcers are associated with predisposing factors such as a history of foot ulcers, hyperglycemia, calluses, foot deformity, inappropriate footwear, trauma, dry skin, underlying peripheral arterial disease, and peripheral neuropathy [28,29]. Initially, patients fail to identify the foot ulcer due to the presence of neuropathy and arterial disease, which along with the predisposing factors can ultimately progress to partial or complete limb amputation [30–33].

#### 3.1. Diabetic Peripheral Neuropathy

Diabetic peripheral nephropathy (DPN) is one of the most severe chronic complications of DM characterized by peripheral nerve malfunction. The dysfunction of autonomic, sensory, and motor nerves in DPN inhibits sweating, pain, temperature, touch, and pressure sensations, as well as increases inflammation, dry skin, cracking, bone deformities, tissue necrosis, and weakness in patients [31,32,34]. When an individual loses their protective sensation, thermal trauma arises and the body's ability to sense the foot's position disappears. Simultaneously, motor neuropathy causes the protrusion of abnormal bones, which changes the normal structure of the foot bones and gives rise to foot deformities such as hallux rigidus and hammer toe. Finally, autonomic nerve malfunction causes dry skin, decreased sweating, and increased secondary arterial capillary refill, leading to skin fissures and crusts and diabetic foot ulcers [33].

#### 3.2. Peripheral Arterial Disease

Hyperglycemia-induced peripheral arterial diseases are attributed to increased oxidative stress and atherosclerosis. In DM, blood sugar accumulation causes blockage, weakening, and narrowing of the arteries. Moreover, elevated oxidative stress prolongs inflammation in the microcirculation and affects the elasticity of capillaries, contributing to ischemia [31]. Additionally, atherosclerosis affects the blood vessels associated with the femoral artery and knee, contributing to developing foot ulcers two times more frequently in diabetic patients compared to non-diabetic patients [33]. The presence of atherosclerosis in the lower limbs has been documented to be 20 times more frequent in T2DM patients than in normal individuals, indicating a higher chance of developing DFU [35]. Many studies indicate that about 90% of DFU cases are attributed to neuropathy and the remaining 10% are associated with ischemia and other contributing factors [36]. A summary of the etiology of diabetic foot ulcers is illustrated in Figure 2.



**Figure 2.** Diagrammatic synopsis of the etiology of diabetic foot ulcers involving prolonged hyperglycemia, diabetic peripheral neuropathy, and peripheral arterial diseases. The sustained hyperglycemia impairs the wound-healing process via collagen crosslinking disorder, cell defense disorders, decreased inflammatory response, and disrupted angiogenesis; peripheral neuropathy affects the sensory, motor, and autonomic nervous systems, causing protective sensation loss, bone deformation, dry skin, skin fissures, and infections; peripheral arterial diseases such as ischemia, inflammation in lower limb microcirculation, and high plantar pressure in the foot can occur from atherosclerosis and elevated oxidative stress.

### 3.3. Other Contributive Factors

Recent studies have highlighted several other important factors leading to DFU, such as prolonged hyperglycemia (>10 years), gender (male), older age (around 65), obesity, retinopathy, peripheral neuropathy, a higher glycated hemoglobin level, nicotine intake, a previously attempted amputation, a history of ulcers, elevated plantar pressure, infection, and inappropriate foot care [37–41].

In DM, the wound-healing process is hampered via collagen crosslinking disorders, immune disorders, tinea and onychomycosis infections, metalloproteinase matrix functional dysfunction, cellular defense disorders, inflammation, and neovascularization disorders. Hyperglycemia weakens the acute inflammatory responses and disrupts angiogenesis, impairing the wound-healing process, and is considered the main culprit in the formation of DFU [42].

DFU is associated with severe infections ranging from uncomplicated cellulitis to life-threatening necrotizing fasciitis. The immunological dysfunction mediated by hyperglycemia mainly induces infection, which further leads to poor perfusion in soft tissues of the skin, damaged skin, and bacterial infections caused by bacteria such as *Escherichia coli* and *Staphylococcus aureus*, and this ultimately leads to sepsis. The risk of amputation increases if infection, sepsis, and resistant bacterial strains are present [14]. In summary, diabetic foot ulcer is a multifactorial life-threatening disease attributed to poor glycemic control.

#### 4. Types of Diabetic Foot Complications

Diabetic foot ulcers can be classified depending on the degree of tissue loss, size, perfusion, infection site, depth, area, and sensation [43]. The changes in these parameters are based on an individual's age, sex, medical conditions, and existing comorbidities, including a loss of peripheral sensation (LOPS) and peripheral arterial diseases (PAD), which ultimately categorize diabetic foot ulcers into different grades [44].

According to Meggitt and Wagner, diabetic foot ulcers can be classified into 6 grades (Figure 3):

**Grade zero:** No lesion on the skin, potential bone deformity or hyperkeratotic lesion, pain;


**Grade one:** Superficial viable or necrotic ulcers; subcutaneous tissue loss; potential bone deformity; no penetration into the deeper layers of the skin;

**Grade two:** Deeper penetrates including bones, tendons, ligaments, or deep fascia; bone deformity prominent in some aspects; absence of abscess or osteomyelitis;

**Grade Three:** Presence of osteomyelitis, deep abscess, or tendinitis; severe infection symptoms (e.g., redness, heat, and swelling);

**Grade four:** Gangrene (dry, wet, infected, or non-infected) in toe or forefoot; surgical ablation of the foot required with minimal blood supply for below-knee amputations;

**Grade five:** Gangrene spreads to the whole foot; no healing signs; amputation below the knee is strongly recommended [28,45,46].



**Figure 3.** Diabetic foot ulcers are categorized using the Wagner–Meggitt classification. Grade 0 means no break in the skin. Grade 1 indicates a superficial ulcer. Grade 2 indicates a deep ulcer. Grade 3 shows the presence of osteomyelitis. Grade 4 is identified as forefoot gangrene. Grade 5 is recognized as complete foot exposure to gangrene.

Another scientist, named Brodsky, later discovered that the grade 4 and grade 5 foot ulcers in Wagner–Meggitt's classification were ischemic, and revised the classification in the following manner (Figure 4):


**Grade 0:** Intact skin; no sign of ulceration but the foot is at risk;

**Grade 1:** Superficial ulcer; no sign of osteomyelitis or exposed bones; no deep ulceration;

**Grade 2:** Deep ulceration; deeper penetration towards bones; bone deformity may be present to some extent;

**Grade 3:** Presence of osteomyelitis or abscess; severe infection and redness; no gangrene exposure;

**Grade A:** Not gangrenous; no ischemia;  
**Grade B:** Presence of ischemia but no gangrene;  
**Grade C:** Presence of ischemia and partial foot gangrene;  
**Grade D:** Presence of ischemia and complete foot gangrene [47].



**Figure 4.** Diabetic foot ulcers based on Brodsky's classification. Grade 0 has no sign of ulceration but persistent pain is present. Grade 1 is a superficial ulcer. Grade 2 indicates deep ulceration towards the bones. Grade 3 means exposure to severe infection (osteomyelitis). Grade A has no ischemia. Grade B indicates ischemia but no gangrene. Grades C and D are partial and complete gangrene infections with ischemia, respectively.

## 5. Pathophysiology

Diabetic foot ulcers generally develop in three stages. Initially, neuropathy induces skin inflammation and forms a callus. Then, the involvement of the motor and autonomic nervous system causes bone deformation, trauma, and abnormal skin conditions. Finally, a subcutaneous hemorrhage mediated by frequent trauma erodes and forms the ulcer [48]. In addition, atherosclerosis in the microcirculation causes neovascularization and inflammation, contributing to the delayed healing of wounds, necrosis, and gangrene [28]. The stages of DFU formation and impaired wound healing are, however, associated with certain dysfunctional mechanisms in diabetic individuals [49].


The neuropathic foot is generally warm, dry, numb, and lacks pain sensations, involving neuropathic ulcers, the Charcot joint, and neuropathic edema. On the contrary, the ischemic foot is usually cold and has localized pressure necrosis and gangrene with the absence of pulses [49]. The key features of diabetes mellitus (i.e., hyperglycemia, insulin resistance, hyperlipidemia, and increased oxidative stress) are considered the main causes of endothelial dysfunction, cellular defense disorders, and other diabetes-related complications. In hyperglycemia, the endothelial nitric oxide production is suppressed by the inhibition of nitric oxide synthase, and as a result the high level of reactive oxygen species (ROS), particularly superoxide radicals, increases the hydrogen peroxide levels. This, in turn, causes the formation of highly reactive hydroxyl radicals and damages the cells. Nitric oxide and superoxide together produce peroxynitrite, which affects the endothelial vasodilation and mediates lipid peroxidation. The concentrations of low-density lipoproteins are increased, followed by atherosclerosis in the microcirculation, increased inflammation, abnormal intimal growth, platelet aggregation,

and thrombosis [50,51]. Moreover, the impairment of the autonomic nervous system is linked to degeneration of the postganglionic unmyelinated sudomotor axons, reducing sweat production. This, in turn, triggers dry skin, thick plaques, and callus formation [49]. Thus, diabetic neuropathy can be regarded as one of the key contributing factors of DFU, which proceed from a foot deformity and callus formation to increased local pressure and repeated injury, tissue necrosis, and finally ulceration [31,52].

Neuropathy, defined as a long-term painless degenerative arthropathy or Charcot neuroarthropathy, is also characterized by disrupted a sensory innervation of the foot joint. The affected autonomic nervous system increases the local and resting blood flow, causing massive calcium dissolution, leading to subluxation, joint dislocation, bone deformation, osteolysis, and soft tissue edema [53]. Studies have also reported that elevated inflammatory cytokines such as tumor necrosis factor- $\alpha$  and interleukin-1 $\beta$  activate the nuclear factor NF- $\kappa$ b, leading to osteoclast maturation, ultimately contributing to bone deformities [54].

In peripheral arteries, the endothelial dysfunction elevates the vasoconstrictor thromboxane A<sub>2</sub>, which can aggregate platelets and increase the risk of plasma hypercoagulation. This can lead to ischemia in the lower limb microcirculation and eventually results in ulceration. The ulcer may become exposed and develop into gangrene and infection [55,56]. Moreover, the inhibition of nitric oxide, which is known as an anticoagulant, also contributes to the propensity of atherosclerosis, constricting the blood vessels in the microcirculation and ultimately leading to ischemia. Clinical reports suggest that DFU patients with ischemia and vascular insufficiency experience intense pain, limb hair loss, thinner skin, and a lack of peripheral pulses [54,55].

DFU gets worse over time due to the impaired healing ability existing in diabetic patients. When skin tissues, blood vessels, nerves, and other associated tissues become damaged, even controlled blood glucose levels fail to improve the condition. The slow wound-healing process deteriorates the condition and leads to life-threatening infections including cellulitis, osteomyelitis, abscesses, gangrene, and sepsis [56,57]. Additionally, the immune system of diabetic patients becomes weaker than normal and the hyperglycemic state increases the number of pro-inflammatory cytokines, affecting cell defenses such as phagocytosis, intracellular killing, chemotaxis, and leukocyte activity. The loss of leukocyte activity, decreased chemotaxis, negative nitrogen balance, and increased gluconeogenesis, as well as the impaired synthesis of fibroblasts, protein, and collagen, alter the normal wound-healing process and lead to a prolonged inflammatory state [55,58,59]. In summary, hyperglycemia, neuropathy, vascular insufficiency, arterial diseases, neuroarthropathy, and impaired immunology all contribute to the pathophysiology of diabetic foot ulcers (Figure 5) [54].



**Figure 5.** A schematic diagram representing the pathophysiology of diabetic foot ulcers via different mechanistic pathways. Endothelial dysfunction, ischemia, neuroarthropathy, and impaired immunology contribute to the pathophysiology of DFU.

## 6. Current Therapy for Diabetic Foot Complications

In order to treat and manage the diabetic foot, it is essential to properly diagnose its stage and severity. Alongside the maintenance of blood sugar levels, current therapeutic strategies involve targeting wound healing, controlling the spread of infection, relieving pressure, and improving blood flow [29]. Analgesic medications such as tapentadol, pregabalin, tramadol, duloxetine, acetaminophen, and some opioids (e.g., oxycodone) are employed to alleviate diabetic peripheral neuropathy (DPN)-associated pain. Although these drugs are effective in reducing mild to moderate pain, their frequent use causes nausea, constipation, drowsiness, and confusion [29]. Many antibiotics including nafcillin, flucloxacillin, dicloxacillin, ceftazidime, cefazolin, ceftriaxone, dalbavancin, oritavancin, telavancin, doxycycline, sulfamethoxazole, or trimethoprim have been effectively used to treat diabetic foot infections, even though no prospective comparative trials have been conducted [60]. It should be noted that several factors such as wound age, the host's immunological state, polymicrobial infection, sanitary conditions, and former antimicrobial therapy may contribute to the development of antimicrobial resistance [61]. Furthermore, the cost of antimicrobial agents recommended for the treatment of infected diabetic foot burdens a large number of diabetic patients, particularly in underdeveloped nations. This results in advanced grades of diabetic foot ulcers that are untreatable in a resource-constrained healthcare system [60,61]. The initial and most crucial step to heal wounds and lower the likelihood of amputations in patients with DFU is debridement. This involves the removal of necrotic and senescent tissues along with foreign and infectious materials from the wound. Debridement tends to lower the bacterial count while increasing the generation of local growth factors, as well as relieving pressure and facilitating wound drainage [36]. The primary purpose of surgical debridement is to transform a chronic ulcer into an acute ulcer. This has been demonstrated to be more successful in treating DFU than mechanical, biological, enzymatic, or autolytic debridement methods [36]. The use of appropriate wound dressing, revascularization for ischemia, pressure relief, laser therapy, and surgery also play an important role in the treatment and control of diabetic foot [29,36,60]. More research is needed, however, to assess the patient demographics for whom these treatments are promising, as well as their cost-to-benefit ratios. [29]. Table 1 summarizes some of the available conventional therapies and their advantages and disadvantages in the management of diabetic foot.



**Table 1.** Current therapeutic strategies for the treatment of diabetic foot.

Conventional Therapies for Diabetic Foot Management		Examples	Advantages	Disadvantages	References
1.	Analgesics	Tapentadol, pregabalin, tramadol, duloxetine, acetaminophen, oxycodone	Alleviates diabetic peripheral neuropathy-induced pain	Constipation, nausea, drowsiness, confusion, drug abuse (opioids)	[29]
2.	Antibiotics	Nafcillin, ceftazidime, cefazolin, clindamycin, dalbavancin, sulfamethoxazole/tri methoprim	The most common treatment for diabetic foot infections	Antibiotic resistance, high cost, unavailability	[29,61]
3.	Debridement	Surgical, autolytic, mechanical, enzymatic, maggot debridement, polysaccharide beads/paste	Aids in a complete assessment of the wound, enhances breakdown of necrotic tissue, speeds up ulcer healing	High cost, time-consuming, patient reluctance, surgical debridement may increase wound size	[62–64]
4.	Wound dressings	Hydrogels, hydrocolloids	Prevent infections, easy to use, effective, provides thermal insulation and mechanical protection	Costly, unavailability	[29,62,64]
5.	Revascularization	-	Heals diabetic foot ulcers	Costly	[65]
6.	Pressure relief	Half-shoes, rigid-soled post-operative shoes, total contact casts, accommodative dressings	Heals diabetic foot ulcers, significantly reduces pressure, high acceptability	Costly, patient compliance, ineffective against inflammation	[29,66]
7.	Laser therapy	Low-level laser therapy	Effective in healing wounds, reduces inflammatory phase	High cost, discontinuous in efficacy	[29]
8.	Surgery	-	Heals amputation or non-healing ulcers, one of the most effective methods to treat foot ulcers	Costly, lack of skilled surgeons	[67]

### 7. Alternative Therapy for Diabetic Foot

The global burden of DFU has expanded surprisingly over the last few decades with the increased incidence of DM. DFU treatment has been estimated to be as expensive as one-third of the total diabetes mellitus management alone is [68]. The overall expenditure and the amputation of lower limbs affect the quality of life of patients, as well as causing social stigma among individuals. To mitigate this public health crisis, alternative therapies can be recommended to DFU patients as adjunctive treatments with fewer side effects and promising health benefits [69].

Synthetic medicines, either alone or in conjunction with insulin, are commonly used as first-line treatments to manage DM and its complications. Such medications may occasionally fail to perform their intended effect properly and may have adverse side effects on the patient's health. As a result, there has been an increase in the pursuit of alternative sources of antidiabetic medications based on natural resources such as plants

and animals [16]. Plants are a good source of medicines and have traditionally been used as antidiabetic therapies for centuries. Several drugs currently used in the treatment of DM also derive directly or indirectly from medicinal plants [70]. The use of plants solely, or in combination with existing antidiabetic agents, has been reported to minimize the problems associated with foot conditions in diabetic individuals. Many of these herbs and their biologically active components are currently being studied to elucidate their mode of action [70].

The major challenge with DFU is the impaired and delayed wound-healing process [69,71]. Medicinal plants have long been acknowledged as a major source of therapies that are utilized both traditionally and conventionally [16]. The traditional remedies possess various therapeutic properties, such as wound healing, antioxidant, antimicrobial, anti-inflammatory, and antihyperglycemic activities. The majority of recent studies indicate that natural remedies are generally used for wound dressing, with few employed as oral drugs to treat DFU [18,71]. Among the various kinds of traditional healthcare practices, the medicinal plants used in *Ayurveda*, *Unani*, traditional Chinese medicine, and other traditional health systems have significantly contributed to the healing of wounds in diabetic foot syndrome. In Western nations, these medical systems, which integrate medicinal herbs, acupuncture, dietary therapy, massage, and therapeutic exercise for treating and preventing diseases, are regarded as a form of complementary and alternative medicine [72]. Chinese herbal medicine, the main part of traditional Chinese medicine, is extensively employed in clinical practice to treat chronic wounds of diabetic foot ulcers. Such an approach can be used either alone or in combination with conventional medicines with the intent to promote wound healing [73]. Some studies have also demonstrated that the combination of insulin with *Ayurveda* medicine has not only improved patients' treatment satisfaction but has also had a positive impact on their quality of life [74].

The numerous active phytoconstituents of medicinal herbs act via different mechanistic pathways to accelerate wound healing. This includes stimulating fibroblasts, stimulating the early expression of growth factors, displaying free radical scavenging or antioxidant activity, preventing wound bleeding, inhibiting microbial growth, stimulating collagen synthesis and improving collagen strength, improving blood circulation to wounds, preventing cell damage, promoting DNA synthesis, improving wound contraction and epithelialization, as well as increasing the production and migration of keratinocytes to the wound site [18,75–78]. Such attributes, together with various other beneficial effects, have contributed to the use of medicinal plants in the treatment of diabetic foot complications, as well as their future potential as alternative antidiabetic medicines [16]. Many phytomedicines that have been marketed and used all around the world have been developed from traditionally used medicinal plants [16]. The formulation of phytomedicines may be either single or poly-herbal. Several marketed poly-herbal formulations including Diabeta [79], Angipars [80], WinVivo [81], Jathyadi Thailam, Jatyadi Ghritam [82], and others have been employed to heal diabetic foot wounds. These products are prepared from individual ingredients from various antidiabetic herbs and are often sold with instructions regarding diet, exercise, and rest. These adjustments ultimately help the patients to re-establish a healthy lifestyle [79]. The most common medicinal plants and their isolated phytochemicals, traditional uses, and modes of action responsible for their healing effect on diabetic foot are summarized in Table 2.

**Table 2.** Traditional uses, isolated active phytochemicals, and mechanism of action of medicinal plants used to treat diabetic foot conditions.

Medicinal Plants	Parts Used	Isolated Phytochemicals	Traditional Uses	Extract Form	Mechanism of Action	References
1. <i>Acacia nilotica</i>	Leaves, roots, bark, seeds	Gallic acid, ellagic acid, isoquercitrin, leucocianidolum, kaempferol	Diabetes, wounds, burns, stomachache, cough, malaria, pneumonia	Aqueous extract	Decreases oxidative stress; suppresses TNF- $\alpha$ and IL-1 $\beta$ activity; enhances cellular proliferation, re-epithelization of wounds, and dermal tissue regeneration	[83,84]
2. <i>Acalypha langiana</i>	Leaves	Acalyphine, triacetoneamine	Diabetes, wounds, diarrhea, urinary infection, asthma, tuberculosis	Aqueous extract	Minimizes wound area, faster tissue regeneration in granulation section, improves congestion and edema	[18,85–87]
3. <i>Actinidia deliciosa</i>	Fruit, stem, roots	Ursolic acid, oleanolic acid, epicatechin, quercetin, emodin	Diabetes, wound healing, fever, diarrhea, hypertension, hyperlipidemia	Pure normal concentrated kiwifruit extract	Enhances angiogenesis and vascularization, increases collagen and granulation tissues, minimizes ulcer size, initiates wound closure, inhibits bacterial infection	[88,89]
4. <i>Aloe vera</i>	Leaves	Emodin, aloin, aloesin, acemannan	Diabetes, wound healing, dermatitis, constipation, infection, worm infestation	Aqueous extract	Enhances collagen cross-linking, increases DNA and glycosaminoglycans, reduces wound area, and accelerates healing rate of ulcer	[90,91]
5. <i>Alternanthera sessilis</i>	Stem, leaves	Sebacic acid, bis(2-ethylhexyl) ester, neophytadiene, hexadecanoic acid	Diabetes, ulcer, wounds, lesion, fever, skin rash, burn, indigestion	Ethanol extract	Initiates wound closure, improves dermal fibroblast and keratinocyte proliferation	[92,93]
6. <i>Annona squamosa</i>	Leaves	Squamosin, squamostatin, rutin, linalool, $\beta$ -caryophyllene	Diabetes, ulcer, hypertension, wounds, hemorrhage, epilepsy	Ethanol extract	Increases DNA synthesis, protein and collagen contents; decreases lipid peroxidation and wound size, restores epidermis; enhances tensile strength, proliferation,	[94–96]

						epithelialization, and migration of keratinocytes	
7.	Anogeissus leiocarpus	Bark, leaves	Ellagic acid, gallic acid, anogeissinin, rutin, quercetin	Diabetes, infection, wounds, diabetes, tuberculosis, hyperlipidemia	Ethanol extract	Reduces oxidative stress, lipid peroxidation, and wound area; enhances vascularity, wound contraction, and rate of epithelialization; prevents onset of cell necrosis and microbial infection	[97–99]
8.	Arnebia euchroma	Roots	Naphthoquinone, shikonin, teracrylshikonin $\beta$ -hydroxyisovalerylalkannin	Diabetes, eye infection, cuts, wound, toothache, earache	Aqueous, ethanol extract	Enhances granulation tissue, wound contraction, re-epithelization; restores epidermis; minimizes wound size; accelerates wound healing	[100–102]
9.	Azadirachta indica	Leaves	Rutin, nimbin, deacetylnimbin, quercetin, azadiradione	Diabetes, ulcer, lesion, wounds, piles, intestinal worms, dermatitis, asthma, cough	Ethanol extract	Increases the rate of wound contraction, improves DNA content and nitric oxide level, enhances tissue regeneration, increases hydroxyproline and protein content	[16,87,103,104]
10.	Balanites aegyptiaca	Bark, fruit, seed, leaves	Furanocoumarin, quercetin, diosgenin, linoleic acid, palmitic acid	Diabetes, malaria, wound healing, diarrhea, constipation, asthma	Aqueous extract	Prevents microbial infection, suppresses lipid peroxidation, enhances wound contraction, and accelerates its healing	[16,105,106]
11.	Camellia sinensis	Leaves	(-) Epigallocatechin-3-gallate, quercitrin, rutin	Diabetes, wound healing, anxiety, flatulence, cardiovascular diseases	Hot water extract	Lowers blood sugar levels, increases insulin secretion, stimulates proliferation and differentiation of keratinocytes and fibroblasts, increases collagen synthesis and angiogenesis	[16,107,108]
12.	Carica papaya	Leaves, fruits	Apigenin, hesperitin, kaempferol, rutin, naringenin	Diabetes, dengue, malaria, ulcer, hypertension, wound, psoriasis	Aqueous, ethanol, methanol extract	Improves granulation tissue weight, collagen and hydroxyproline content, wound size, and contraction; epithelizes	[18,109,110]

					faster and prevents microbial infection		
13.	<i>Catharanthus roseus</i>	Flower, roots, leaves	Catharanthine, vindolinine, isorhamnetin, quercetin, kaempferol, O-feruloylquinic acids	Diabetes, sepsis, wound healing, blood dysentery, blood purification	Ethanol extract	Enhances wound contraction, tensile strength, angiogenesis, epithelialization, vascularization, granulation tissue weight, and hydroxyproline content and protects against microbial infection	[111,112]
14.	<i>Cecropia peltata</i>	Leaves	Orientin, vitexin, rutin, chlorogenic acid, catechin	Diabetes, respiratory tract infection, wounds, kidney diseases, hypertension, sleep disorder	Aqueous, ethanol extract	Minimizes wound area; enhances hydroxyproline, protein, and hexosamine contents; accelerates wound healing	[113,114]
15.	<i>Centella asiatica</i>	Leaves	Kaempferol, hydrochotine, linoleic, oleic, stearic, palmitic acid	Diabetes, ulcer, wounds, eczema, urinary tract infection, amenorrhea, diarrhea	Aqueous, ethanol extract	Improves angiogenesis, collagen content, fibroblast growth factor, and vascular endothelial growth factor; inhibits TNF- $\alpha$ , IL-6, IL-1 $\beta$ , COX-2, prostaglandin E2, and lipoxygenase; initiates re-epithelialization	[115,116]
16.	<i>Citrus x paradisi</i>	Peel, fruit	Narirutin, hesperidin, naringin, neohesperidin, limoline	Diabetes, hypertension, gastrointestinal problems, wound healing, infection, cold, cough	Aqueous peel extract	Reduces wound size; increases collagen synthesis, hydroxyproline and protein contents, and regeneration of tissue; initiates wound closure and healing	[117–190]
17.	<i>Citrus x sinensis</i>	Peel, fruit	Kaempferol, naringin, hesperetin, limocitrin, naringenin	Diabetes, bronchitis, cough, obesity, hypertension, tuberculosis	Aqueous extract	Protects from oxidative damage, inflammatory cytokines, and hydrogen-peroxide-induced cellular damage; enhances cell proliferation, blood flow in tissues, angiogenesis, keratinocyte and fibroblast migration, wound healing	[120–122]

18.	Citrus limon	Peel, fruit, leaves	D-limonene, naringin, hesperetin, hesperidin, apigenin, quercetin, orientin	Diabetes, cough, hypertension, sore throat, scurvy, kidney stones	Aqueous, acetone, ethanol extract	Inhibits gluconeogenesis; enhances collagen synthesis, protein and hydroxyproline contents; increases tissue growth rate; reduces wound area and healing time	[123,124]
19.	Curcuma longa	Rhizome	Curcumin, curcuminoids, turmerin, vanillic acid, $\beta$ -sitosterol, zingiberene	Diabetes, liver disorders, cough, inflammation, ulcer, wounds, pain, infections	Ethanol extract	Improves ulcers, inflammation, wound contraction, proliferation, fibroblast formation, collagen synthesis, granulation tissue and epithelialization; helps in tissue remodeling and wound size minimization; suppresses inflammatory cytokines	[16,76,125–127]
20.	Eucalyptus alba	Leaves	Chlorogenic acid, gallic acid, caffeic acid, vanillic acid, ferulic acid	Diabetes, sinusitis, bronchitis, cold, cough, wounds, sores,	Ethanol, methanol, acetone extract	Inhibits TNF- $\alpha$ , IL-6, and $\alpha$ -glucosidase enzyme; reduces oxidative stress; decreases oxidative free radicals; increases cell proliferation, collagen synthesis, and blood flow to the wound; prevents microbial attack	[128–130]
21.	Euphorbia hirta	Leaves, stem, flower	Heptacosane, camphol, leucocyanidin, kaempferol, gallic acid	Diabetes, cold, asthma, dysentery, wound healing, dermatitis	Ethanol extract	Initiates wound closure and healing; restores epidermis, collagen tissue, and extensive fibrosis; improves granulation tissue weight, inflammatory cells, cellular damage, and re-epithelialization; decreases lipid peroxidation	[16,87,131,132]
22.	Guiera senegalensis	Roots, leaves	Epigallocatechin, epicatechin, gallic acid, quercetin	Diabetes, wound healing, gastrointestinal disorders, cough, malaria, enteritis	Aqueous, methanol extract	Exhibits antimicrobial action, inhibits lipid peroxidation and inflammatory cytokines, minimizes wound size, improves healing time	[133–136]

23.	<i>Gymnema sylvestre</i>	Leaves	Gymnemic acid, gurmardin, lupeol, stigmasterol, kaempferol	Diabetes, anemia, constipation, indigestion, infections, cardiopathy	Hot water, ethanol, methanol extract	Decreases blood glucose levels, bacterial growth, and inflammation; decreases formation of reactive oxidants; improves wound contraction; increases fibroblasts and enhances tissue regeneration	[137,138]
24.	<i>Hibiscus rosa-sinensis</i>	Flower, leaves, roots	Quercetin, stigmasterol, $\beta$ -sitosterol, cyaniding, taraxeryl acetate	Diabetes, wounds, stomach ulcer, hypertension, hypercholesterolemia	Ethanol extract, n-butyl alcohol extract	Increases DNA, collagen, and protein contents of granulation tissues; improves rates of epithelialization and wound contraction, closure rate, granuloma weight, tensile strength	[139–141]
25.	<i>Hylocereus undatus</i>	Peel, fruit	Gallic acid, betacyanin, lycopene, ascorbic acid	Diabetes, asthma, intestinal disease, wounds, ulcer, allergy, hypertension	Aqueous extract	Increases total protein, collagen, hydroxyproline, and DNA contents; enhances rate of epithelialization, wound size, tensile strength, and wound closure	[142,143]
26.	<i>Hypericum perforatum</i>	Leaves, flowers	Amentoflavone, hypericin, hyperforin	Diabetes, depression, wounds healing, mycobacterium and viral infections, GIT disorders, eczema	Aqueous, ethanol extract	Improves glycemic levels, decreases inflammation, inhibits extracellular matrix degradation by matrix metalloproteinase-2 and urokinase, increases epithelial regeneration and revascularization	[103,144]
27.	<i>Kaempferia galanga</i>	Rhizome	Ethyl p-methoxy cinnamate, 4-methoxy cinnamic acid, kaempferol, kaempferide, luteolin	Diabetes, hypertension, asthma, urticaria, rheumatism, wound healing, ulcers	Ethanol extract	Lowers blood glucose levels, exhibits anti-inflammatory and antioxidant effects, enhances wound contraction, increases rate of epithelialization	[19,145–147]
28.	<i>Lawsonia alba</i>	Leaves	Kaempferol, quercetin, luteolin,	Diabetes, wound healing, bacterial and fungal infections, jaundice, skin problems	Aqueous, ethanol extract	Exhibits antibacterial activity, improves wound healing in fissures and cracks in diabetic foot, prevents decubitus ulcers, improves	[19,148]

						fibroblasts and collagen synthesis, increases granulation tissue	
29.	<i>Linum usitatissimum</i>	Seeds	$\alpha$ -linolenic acid, omega-3 fatty acids, p-coumaric acid	Diabetes, diarrhea, skin diseases, wound healing, gastrointestinal infections	Ethanol extract	Improves insulin sensitivity; stimulates fibrin proliferation and collagen synthesis; increases granulation tissue around wounds; promotes vascular contraction, chemotaxis, adhesion, transmigration, and cell activation for tissue repair	[16,149–151]
30.	<i>Lycopodium serratum</i>	Leaves	Lycoposerramine A, serratezomines A, B, C, $\alpha$ -pyridine,	Diabetes, wounds, sores, cuts, and burns, to improve learning and memory efficiency, Alzheimer's disease, otitis media	Ethanol extract	Increases rates of epithelialization and wound contraction, increases content of hydroxyproline at the wound site, and exerts antioxidant and antibacterial effects	[19,152]
31.	<i>Martynia annua</i>	Leaves, flowers	p-hydroxy benzoic acid, chlorogenic acid, luteolin	Diabetes, epilepsy, sore throat, wounds, inflammation	Ethanol extract	Inhibits oxidative stress, exerts antibacterial action, improves angiogenesis of fibroblasts, increases collagen synthesis and hydroxyproline content, enhances epithelialization, increases wound contraction rate	[153,154]
32.	<i>Melilotus officinalis</i>	Flowers, flowering stems	Quercetin, kaempferol, lupeol, oleanolic acid, coumarin compounds	Diabetes, intestinal worms, gastric ulcers, wound treatment, vein problems, hemorrhoids	Methanol, ethanol, hexane, chloroform extract	Stimulates the proliferation of endothelial cells and vascular neoangiogenesis, improves blood circulation, increases wound contraction and collagen synthesis, exerts anti-inflammatory and antioxidant effects	[155,156]
33.	<i>Mimosa tenuiflora</i>	Bark	Lupeol, mimonoside A,B,C, arabinogalactans, campesterol-3-O- $\beta$ -D-	Diabetes, skin burns, ulcers, wounds, psoriasis, inflammation	Aqueous, ethanol extract	Enhances proliferation of dermal fibroblasts, stimulates production of keratinocytes, exerts	[157,158]



		glucopyranosyl, stigmasterol-3-O- $\beta$ -D- glucopyranosyl			antimicrobial activity to improve wound healing		
34.	Momordica charantia	Leaves, fruit	Momocharin, momordicin	Diabetes, wound healing, microbial infections, hypertension, inflammation, skin disease	Aqueous, ethanol, ether extract	Lowers blood glucose levels, increases insulin secretion, rectifies structural abnormalities in peripheral nerve, improves wound contraction, increases granulation tissue and neovascularization	[77,151,159]
35.	Morinda citrifolia	Leaves, fruits	Rutin, quercetin, kaempferol, coumarin compounds	Diabetes, wounds, pain and inflammation, hypertension, constipation, gastric ulcers	Aqueous, ethanol extract	Reduces blood glucose levels, decreases wound size, decreases period of epithelialization, inhibits bacterial growth, reduces inflammation, exerts antioxidant action	[75,160,161]
36.	Myrtus communis	Leaves, fruits	Limonene, $\alpha$ , $\beta$ -pinene, myrtucommulone, gallic acid, ellagic acid, linalool	Diabetes, skin diseases, hemorrhage, peptic ulcers, wounds, conjunctivitis	Aqueous, ethanol extract	Lowers blood glucose levels, stimulates angiogenesis, promotes proliferation and migration of cells to wound sites, reduces inflammation, exerts free radical scavenging activity, prevents microbial infection at the wound site	[162,163]
37.	Nigella sativa	Seeds	Thymoquinone, thymohydroquinone, thymol, oleic acid, linoleic acid	Diabetes, wound healing, microbial infections, rheumatism, dysentery, respiratory tract diseases	Aqueous, ethanol extract	Lowers blood glucose levels; induces angiogenesis, fibroblast proliferation, and collagen synthesis; minimizes bacterial infection; exhibits anti- inflammatory action; decreases neuronal degeneration	[16,151,164,165]
38.	Olea europaea	Leaves	Oleuropein, oleanolic acid, hydroxytyrosol, luteolin	Diabetes, hypertension, constipation, UTI, skin	Ethanol extract	Decreases blood sugar levels; exhibits antioxidant activity; improves blood circulation in	[16,78]

			ulcers, inflammatory wounds		healing tissues; inhibits bacterial growth; enhances epithelialization, collagen synthesis, and development of fibroblasts		
39.	<i>Orthosiphon aristatus</i>	Leaves	Sinensetin, caffeic acid, rosmarinic acid, eupatorin, eugenol, linalool	Diabetes, nephritis, kidney and gallstones, hypertension, arteriosclerosis epilepsy, rheumatism,	Aqueous, ethanol extract	Reduces blood glucose levels; increases insulin secretion and actio; exerts antioxidant and anti-inflammatory action; enhances epithelialization rate; increases wound contraction and collagen synthesis in fibroblasts	[166,167]
40.	<i>Prosopis cineraria</i>	Leaves, stem bark	Pautelin, linolenic acid, luteolin, gallic acid, rutin	Diabetes, rheumatism, asthma, bronchitis, eye infections, wound healing	Ethanol extract	Increases rates of wound healing and tissue repair, enhances wound contraction and epithelialization, increases cell proliferation, exerts antioxidant and antimicrobial actions, increases collagen synthesis, regulates inflammatory markers and oxidative stress	[168,169]
41.	<i>Pterocarpus marsupium</i>	Bark	Marsupin, marsupol, carsupin, epicatechin, pterostilbene	Diabetes, wound healing, skin diseases, ulcer, cough, diarrhea, dysentery	Ethanol extract	Lowers blood glucose levels, improves glucose tolerance, promotes re-epithelialization, exhibits free radical scavenging activity, decrease TNF- $\alpha$ levels, increases expression of angiogenesis-related proteins, reduces bacterial infection in wounds	[16,170]
42.	<i>Punica granatum</i>	Fruit, flowers	Ellagic acid, gallic acid, luteolin, catechin, rutin, quercetin	Diabetes, inflammation, rheumatism, sore throat, leprosy, burns	Aqueous, methanol, ethanol, ethyl acetate extract	Lowers blood sugar levels, increases wound healing rate, enhances cell proliferation and collagen synthesis, reduces inflammation	[171,172]

43.	Radix Rehmanniae	Roots	Catalpol	Diabetes, ulcers, hypertension, liver and kidney diseases, depression	Aqueous extract	Reduces blood sugar levels, heals diabetic foot ulcers, promotes tissue regeneration and angiogenesis, inhibits TNF- $\alpha$ production, increases IL-2 and IFN- $\gamma$ production	[173,174]
44.	Rosmarinus officinalis	Leaves, stems, flowers	Rosmarinic acid, carnosic acid, chlorogenic acid, apigenin, luteolin	Diabetes, anxiety, inflammation, muscle, and joint pain, wound, colds	Ethanol, methanol, chloroform extract	Exhibits free radical scavenging activity, inhibits lipid peroxidation, decreases growth of microorganisms, exerts anti-inflammatory and neuroprotective actions, promotes angiogenesis, improves granulation tissue for faster wound healing	[175,176]
45.	Sesamum indicum	Seeds	Sesamol, sesamin, sesamolin, sesaminol	Diabetes, healing burns and wounds, hypertension, constipation	Ethanol extract	Lowers blood glucose levels, accelerates wound healing, increases wound contraction, decreases wound inflammation, exerts neuroprotection	[19,151,177,178]
46.	Syzygium cumini	Seeds, bark	Quercetin, ellagic acid, gallic acid, kaempferol, iso-coumarin	Diabetes, sore throat, constipation, wounds, skin ulcers, gastritis, constipation	Aqueous extract	Decreases blood sugar levels, enhances insulin secretion and action and re-epithelialization, increases collagen deposition, promotes neovascularization	[16,179,180]
47.	Terminalia chebula	Fruit, leaves	Ellagic acid, gallic acid, tannic acid, chebulagic acid, corilagin	Diabetes, constipation, dementia, wound healing, astringent	Aqueous, methanol, ethyl acetate extract	Lowers blood glucose levels, improves rate of wound contraction, increases proliferation of keratinocytes and fibroblasts, improves free radical scavenging activity	[16,181]
48.	Tinospora cordifolia	Leaves, bark, roots	Tinosporaside, tinosporine, berberine,	Diabetes, asthma, dysentery, diarrhea,	Aqueous, methanol, ethanol extract	Exerts immunomodulatory effects, enhances wound contraction, promotes collagen synthesis,	[16,182]

		jatrorrhizine, $\beta$ -sitosterol, arabinogalactan	jaundice, anemia, allergy, wound healing		exhibits antioxidant and free radical scavenging activity levels, inhibits microbial growth at wound site	
49.	Trigonella-foenum graecum	Seeds	Galactomannan, trigonelline, 4-hydroxy isoleucine	Diabetes, wound healing, indigestion, pneumonia, bronchitis, constipation	Aqueous, ethanol extract	Decreases blood glucose levels, increases insulin secretion, restores function of nerve fibers, improves collagen synthesis and maturation, increases granuloma tissue [19,151,183,184]
50.	Zingiber officinale	Roots, rhizome	Gingerol, paradol, shogaol	Diabetes, wound healing, microbial infections, cough, asthma	Ethanol extract	Decreases blood sugar levels, increases insulin secretion, improves collagen synthesis and maturation, increases wound contraction and epithelialization, inhibits bacterial colonization [185,186]

## 8. Advantages of Alternative Therapies over Conventional Treatment Approaches for Diabetic Foot

Diabetes mellitus and its associated complications including diabetic foot have major detrimental effects on healthcare systems, particularly in poorer nations where both resources and finances are scarce. Natural compounds originating from plants have served, and continue to serve, a pivotal role in pharmaceutical research [16]. Nearly 80% of people living in developing countries rely upon traditional medicines, mostly derived from medicinal plants, to address their basic health needs [18]. Plant-based medicines have become a promising alternative source of medicines due to their ease of access, availability, low cost, and relatively low risk of adverse side effects [16]. The wide range of health benefits possessed by these plants is primarily because of the presence of numerous phytoconstituents (e.g., flavonoids, glycosides, carotenoids, terpenoids, and alkaloids) that have potent antihyperglycemic properties and are generally safe. The blood glucose-lowering effects of plants are mainly attributed to their ability to either improve the  $\beta$ -cell function by increasing insulin production and secretion or decrease intestinal glucose absorption [187]. Many of the studies carried out to date have proved that the bioactive compounds derived from medicinal plants have various pharmacological effects that are useful in preventing cell damage as well as chronic wounds. In addition to being traditionally used as topical formulations to treat burns, infections, inflammation, and skin problems, plants and their associated compounds have also demonstrated wound-healing characteristics *in vitro*, *in vivo*, and in clinical studies [17]. Natural products are increasingly being employed not only as therapeutic agents in their own right but also as templates for the synthesis of novel bioactive molecules. As such, it is important that extensive scientific and clinical research is carried out to validate the medicinal use of plants and phytochemicals. Finally, as plants have proved beneficial for the treatment of diabetes and its associated complications, it is reasonable to assume that the combination of herbal and synthetic medicines may help reduce some of the harmful effects of conventional drugs [175].


## 9. Prevention and Management of Diabetic Foot Complications

Prevention is the primary management strategy to control any lower limb injury linked to DM. Providing basic guidance to patients regarding the choice of suitable footwear and wound treatments, including a specific emphasis on foot care, has been found to help delay the occurrence or re-occurrence of DFU [188]. The use of inappropriate footwear and walking barefoot are the main contributors to foot trauma resulting in ulcers in diabetic patients with sensitive feet. Thus, people with a loss of protective sensation (LOPS) should have unrestricted access to suitable footwear and should be advised to wear them at all times, both indoors and outdoors [53,189].

Some other important aspects involved in the prevention of DFU include identifying the foot likely to be at risk of ulceration and inspecting and examining it on a regular basis, as well as treating pre-ulcerative symptoms [53]. To identify individuals who are at risk of developing foot ulcers, all diabetic patients should get their feet examined at least once annually, and those who have been identified should be checked more frequently. Diabetic patients who show symptoms of pre-ulcers should be treated accordingly. The pre-ulcerative treatments include the removal of excess calluses, the treatment of blisters and ingrown or thickened nails, the administration of antifungal medications for fungal infections, and in case of foot deformities non-surgical techniques such as orthosis may be applicable. These interventions should be carried out by an experienced foot care expert until the pre-ulcerative signs subside and do not reappear [53,189].

To control and manage DFU, it is crucial to heal the wound first. This mostly depends on the ulcer's grade and vascularity, as well as the presence of infection. The affected foot should be given adequate relaxation and pressure relief [32]. Controlling hyperglycemia accelerates wound healing and also helps to avoid negative effects on cellular immunity

and infection [190]. Many antiseptic and antimicrobial agents are being investigated as topical preparations for the treatment and management of DFU. Honey, which has antibacterial properties, has recently gained popularity as an alternative treatment for diabetic foot wounds. It provides a moist environment for wounds, and its high viscosity helps to form a protective barrier against infection. Its immunomodulatory effects are also essential for tissue repair. Much work, including performing well-designed randomized double-blind controlled studies, is still required to properly evaluate the potential of honey, design evidence-based practice guidelines, and assess the risks and benefits of such therapies [191,192]. A summary of the preventive measures and treatment strategies used for diabetic foot conditions is depicted in Figure 6.



**Figure 6.** Prevention and management of diabetic foot conditions.

## 10. Future Strategies Used to Overcome the Burden of Diabetic Foot

To reduce the burden of diabetic foot, it is essential to prevent the occurrence of DFU. However, the prevention of ulcers remains underexplored both in research and clinical settings [193]. Due to the complex etiology of DFU, a multidisciplinary team approach involving diabetologists, dermatologists, primary care doctors and nurses, orthopedic and vascular surgeons, podiatrists, and orthotists is crucial. As discussed earlier, the conventional preventive and management strategies for diabetic foot conditions include local wound healing with the use of dressing, the regular removal of callus and necrotic debris, offloading, and the administration of broad-spectrum antibiotics in the case of foot infections [29,194]. These treatment approaches, however, can be adversely affected by underlying health issues that may impede recovery and be a financial burden on patients. In recent years, the use of traditional herbs as medications has received widespread attention as one of the most effective strategies to treat DFU and infections. Despite the availability of numerous medicinal plants and associated active phytoconstituents, as well as pharmaceutical preparations for the management of diabetes, additional medicinal plants with greater antidiabetic efficacy must be sought. The significance of traditional phytomedicines as one of the treatment strategies for DFU should be seriously considered, particularly in low-income nations [195]. The effects of plant extracts on diabetic wounds should be scientifically investigated in order to develop potential adjunct medications for wound treatment. Large clinical trial studies should also be performed on diabetic and non-diabetic patients to elucidate the mode of action of medicinal plants or constituents on diabetic foot-related diseases [18]. Although there are certain limitations to the extent and scope of using medicinal plants for curing diabetic foot wounds, these offer promising

potential and may even lead to the discovery of novel treatment strategies for wound healing [18].

## 11. Conclusions

Diabetes mellitus and its related complications are on the rise globally, creating an urgent need for effective treatment strategies. The use of conventional synthetic drugs as well as surgical or non-surgical procedures has been proven to be effective in healing diabetic foot syndrome. However, these are accompanied by a number of disadvantages, which include adverse side effects, high costs, and inaccessibility to the vast majority of the global population. As a result, the current research efforts are focused on exploring novel sources of antidiabetic and wound-healing remedies, especially plant-based medicines. Medicinal plants have been documented to exhibit promising effects in treating diabetic foot ulcers and infections. Despite their limitations and the lack of clinical trials, traditional medicine remains the primary and often the only line of therapy for many people, particularly those living in poorer or developing nations, where plants are still used to treat various ailments and infections, including diabetic wounds. Thus, medicinal plants and their bioactive constituents with antidiabetic activity are being extensively studied to discover and develop novel medicines with better potency and fewer side effects. Further research and clinical studies are required to validate the efficacy and safety of herbal medicines, as well as to fully understand the molecular pathways involved in the diabetic wound-healing process.

**Author Contributions:** Conceptualization, P.A.; formal Analysis, P.A., V.S. and S.A.; funding acquisition, Y.H.A.A.-W. and P.A.; investigation, resources, writing, P.A., S.A., J.T.K., S.S.I., M.S.R.M. and A.R.; Visualization, P.A. and J.T.K.; editing, P.A. and V.S.; supervision and reviewing, P.A., V.S. and Y.H.A.A.-W. All authors have read and agreed to the published version of the manuscript.

**Funding:** This study received no external funding.

**Institutional Review Board Statement:** Not applicable.

**Informed Consent Statement:** Not applicable.

**Data Availability Statement:** Not applicable.

**Acknowledgments:** We would like to express our gratitude to Independent University, Bangladesh (IUB), Ulster University, and University Strathclyde, UK, for providing access to library resources in order to collect information for this literature research.

**Conflicts of Interest:** The authors indicate that this manuscript has no conflicts of interest.

## References

1. Oguntibeju, O.O. Type 2 diabetes mellitus, oxidative stress and inflammation: Examining the links. *Int. J. Physiol. Pathophysiol. Pharmacol.* **2019**, *11*, 45.
2. Sun, H.; Saeedi, P.; Karuranga, S.; Pinkepank, M.; Ogurtsova, K.; Duncan, B.B.; Stein, C.; Basit, A.; Chan, J.C.; Mbanya, J.C. IDF Diabetes Atlas: Global, regional and country-level diabetes prevalence estimates for 2021 and projections for 2045. *Diabetes Res. Clin. Pract.* **2022**, *183*, 109119.
3. Alam, U.; Asghar, O.; Azmi, S.; Malik, R.A. General aspects of diabetes mellitus. *Handb. Clin. Neurol.* **2014**, *126*, 211–222.
4. Mayfield, J.A. Diagnosis and classification of diabetes mellitus: New criteria. *Am. Fam. Physician* **1998**, *58*, 1355.
5. Katsarou, A.; Gudbjörnsdóttir, S.; Rawshani, A.; Dabelea, D.; Bonifacio, E.; Anderson, B.J.; Jacobsen, L.M.; Schatz, D.A.; Lernmark, Å. Type 1 diabetes mellitus. *Nat. Rev. Dis. Prim.* **2017**, *3*, 17016.
6. Ohlson, L.-O.; Larsson, B.; Björntorp, P.; Eriksson, H.; Svärdsudd, K.; Welin, L.; Tibblin, G.; Wilhelmsen, L. Risk factors for type 2 (non-insulin-dependent) diabetes mellitus. Thirteen and one-half years of follow-up of the participants in a study of Swedish men born in 1913. *Diabetologia* **1988**, *31*, 798–805.
7. Viigimaa, M.; Sachinidis, A.; Toumpourleka, M.; Koutsampasopoulos, K.; Alliksoo, S.; Titma, T. Macrovascular complications of type 2 diabetes mellitus. *Curr. Vasc. Pharmacol.* **2020**, *18*, 110–116.
8. Fowler, M.J. Microvascular and macrovascular complications of diabetes. *Clin. Diabetes* **2008**, *26*, 77–82.
9. Giurato, L.; Uccioli, L. The diabetic foot: Charcot joint and osteomyelitis. *Nucl. Med. Commun.* **2006**, *27*, 745–749.
10. Nathan, D.M. Long-term complications of diabetes mellitus. *N. Engl. J. Med.* **1993**, *328*, 1676–1685.

11. Volmer-Thole, M.; Lobmann, R. Neuropathy and diabetic foot syndrome. *Int. J. Mol. Sci.* **2016**, *17*, 917.
12. Brocco, E.; Ninkovic, S.; Marin, M.; Whisstock, C.; Bruseghin, M.; Boschetti, G.; Viti, R.; Forlini, W.; Volpe, A. Diabetic foot management: Multidisciplinary approach for advanced lesion rescue. *J. Cardiovasc. Surg.* **2018**, *59*, 670–684.
13. Markakis, K.; Bowling, F.; Boulton, A. The diabetic foot in 2015: An overview. *Diabetes/Metab. Res. Rev.* **2016**, *32*, 169–178.
14. Bandyk, D.F. The diabetic foot: Pathophysiology, evaluation, and treatment. *Semin. Vasc. Surg.* **2018**, *31*, 43–48.
15. Bekele, F.; Chelkeba, L. Amputation rate of diabetic foot ulcer and associated factors in diabetes mellitus patients admitted to Nekemte referral hospital, western Ethiopia: Prospective observational study. *J. Foot Ankle Res.* **2020**, *13*, 65.
16. Ansari, P.; Akther, S.; Hannan, J.M.A.; Seidel, V.; Nujat, N.J.; Abdel-Wahab, Y.H.A. Pharmacologically Active Phytomolecules Isolated from Traditional Antidiabetic Plants and Their Therapeutic Role for the Management of Diabetes Mellitus. *Molecules* **2022**, *27*, 4278.
17. Ahmadian, R.; Bahramsoltani, R.; Marques, A.M.; Rahimi, R.; Farzaei, M.H. Medicinal Plants as Efficacious Agents for Diabetic Foot Ulcers: A Systematic Review of Clinical Studies. *Wounds Compend. Clin. Res. Pract.* **2021**, *33*, 207–218.
18. Oguntibeju, O.O. Medicinal plants and their effects on diabetic wound healing. *Vet. World* **2019**, *12*, 653.
19. Nagori, B.P.; Solanki, R. Role of medicinal plants in wound healing. *Res. J. Med. Plant* **2011**, *5*, 392–405.
20. Sharma, A.; Khanna, S.; Kaur, G.; Singh, I. Medicinal plants and their components for wound healing applications. *Future J. Pharm. Sci.* **2021**, *7*, 53.
21. El-Nashar, H.A.; Mostafa, N.M.; El-Shazly, M.; Eldahshan, O.A. The role of plant-derived compounds in managing diabetes mellitus: A review of literature from 2014 to 2019. *Curr. Med. Chem.* **2021**, *28*, 4694–4730.
22. Abdelghffar, E.A.; Mostafa, N.M.; El-Nashar, H.A.; Eldahshan, O.A.; Singab, A.N.B. Chilean pepper (*Schinus molle*) ameliorates the adverse effects of hyperglycaemia/dyslipidaemia in high fat diet/streptozotocin-induced type 2 diabetic rat model. *Ind. Crops Prod.* **2022**, *183*, 114953.
23. Talukdar, N.; Das, K.; Barman, I. A review on ethanobotanical survey of medicinal plants available in North-East India against microbes involved in diabetic foot ulcer. *J. Diabetol.* **2021**, *12*, 128.
24. Boulton, A.J.; Vileikyte, L.; Ragnarson-Tennvall, G.; Apelqvist, J. The global burden of diabetic foot disease. *Lancet* **2005**, *366*, 1719–1724.
25. Boulton, A.J. The diabetic foot: Grand overview, epidemiology and pathogenesis. *Diabetes/Metab. Res. Rev.* **2008**, *24*, S3–S6.
26. Chun, D.-I.; Kim, S.; Kim, J.; Yang, H.-J.; Kim, J.H.; Cho, J.-H.; Yi, Y.; Kim, W.J.; Won, S.H. Epidemiology and burden of diabetic foot ulcer and peripheral arterial disease in Korea. *J. Clin. Med.* **2019**, *8*, 748.
27. Zhang, P.; Lu, J.; Jing, Y.; Tang, S.; Zhu, D.; Bi, Y. Global epidemiology of diabetic foot ulceration: A systematic review and meta-analysis. *Ann. Med.* **2017**, *49*, 106–116.
28. Oliver, T.I.; Mutluoglu, M. Diabetic foot ulcer. In *StatPearls*; StatPearls Publishing: Treasure Island, FL, USA, 2019.
29. Perez-Favila, A.; Martinez-Fierro, M.L.; Rodriguez-Lazalde, J.G.; Cid-Baez, M.A.; Zamudio-Osuna, M.D.J.; Martinez-Blanco, M.D.R.; Mollinedo-Montaño, F.E.; Rodriguez-Sanchez, I.P.; Castañeda-Miranda, R.; Garza-Veloz, I. Current therapeutic strategies in diabetic foot ulcers. *Medicina* **2019**, *55*, 714.
30. Boulton, A.J. The pathway to foot ulceration in diabetes. *Med. Clin.* **2013**, *97*, 775–790.
31. Alavi, A.; Sibbald, R.G.; Mayer, D.; Goodman, L.; Botros, M.; Armstrong, D.G.; Woo, K.; Boeni, T.; Ayello, E.A.; Kirsner, R.S. Diabetic foot ulcers: Part I. Pathophysiology and prevention. *J. Am. Acad. Dermatol.* **2014**, *70*, e1.
32. Frykberg, R.G. Diabetic foot ulcers: Pathogenesis and management. *Am. Fam. Physician* **2002**, *66*, 1655.
33. Rosyid, F.N. Etiology, pathophysiology, diagnosis and management of diabetics' foot ulcer. *Int. J. Res. Med. Sci.* **2017**, *5*, 4206–4213.
34. Forsythe, R.; Brownrigg, J.; Hinchliffe, R. Peripheral arterial disease and revascularization of the diabetic foot. *Diabetes Obes. Metab.* **2015**, *17*, 435–444.
35. Laing, P. The development and complications of diabetic foot ulcers. *Am. J. Surg.* **1998**, *176*, 115–195.
36. Yazdanpanah, L.; Nasiri, M.; Adarvishi, S. Literature review on the management of diabetic foot ulcer. *World J. Diabetes* **2015**, *6*, 37.
37. Shahbazian, H.; Yazdanpanah, L.; Latifi, S.M. Risk assessment of patients with diabetes for foot ulcers according to risk classification consensus of International Working Group on Diabetic Foot (IWGDF). *Pak. J. Med. Sci.* **2013**, *29*, 730.
38. Iraj, B.; Khorvash, F.; Ebneshahidi, A.; Askari, G. Prevention of diabetic foot ulcer. *Int. J. Prev. Med.* **2013**, *4*, 373.
39. Waaijman, R.; De Haart, M.; Arts, M.L.; Wever, D.; Verlouw, A.J.; Nollet, F.; Bus, S.A. Risk factors for plantar foot ulcer recurrence in neuropathic diabetic patients. *Diabetes Care* **2014**, *37*, 1697–1705.
40. Monteiro-Soares, M.; Boyko, E.; Ribeiro, J.; Ribeiro, I.; Dinis-Ribeiro, M. Predictive factors for diabetic foot ulceration: A systematic review. *Diabetes/Metab. Res. Rev.* **2012**, *28*, 574–600.
41. McEwen, L.N.; Ylitalo, K.R.; Herman, W.H.; Wrobel, J.S. Prevalence and risk factors for diabetes-related foot complications in Translating Research Into Action for Diabetes (TRIAD). *J. Diabetes Its Complicat.* **2013**, *27*, 588–592.
42. Tellechea, A.; Leal, E.; Veves, A.; Carvalho, E. Inflammatory and angiogenic abnormalities in diabetic wound healing: Role of neuropeptides and therapeutic perspectives. *Open Circ. Vasc. J.* **2010**, *3*, 43–55.
43. Schaper, N. Diabetic foot ulcer classification system for research purposes: A progress report on criteria for including patients in research studies. *Diabetes/Metab. Res. Rev.* **2004**, *20*, S90–S95.
44. Jeffcoate, W.J.; Bus, S.A.; Game, F.L.; Hinchliffe, R.J.; Price, P.E.; Schaper, N.C. International Working Group on the Diabetic Foot and the European Wound Management Association. Reporting standards of studies and papers on the prevention and



- management of foot ulcers in diabetes: Required details and markers of good quality. *Lancet Diabetes Endocrinol.* **2016**, *4*, 781–788.
45. Wagner, F.W., Jr. The dysvascular foot: A system for diagnosis and treatment. *Foot Ankle* **1981**, *2*, 64–122.
  46. Abid, A.; Hosseinzadeh, S. Foot Ulcer. In *StatPearls*; StatPearls Publishing: Treasure Island, FL, USA, **2020**.
  47. Robinson, A.; Pasapula, C.; Brodsky, J. Surgical aspects of the diabetic foot. *J. Bone Jt. Surgery. Br. Vol.* **2009**, *91*, 1–7.
  48. Armstrong, D.G.; Boulton, A.J.; Bus, S.A. Diabetic foot ulcers and their recurrence. *N. Engl. J. Med.* **2017**, *376*, 2367–2375.
  49. Edmonds, M. The diabetic foot: Pathophysiology and treatment. *Clin. Endocrinol. Metab.* **1986**, *15*, 889–916.
  50. Apelqvist, J.; Ragnarson-Tennvall, G.; Larsson, J.; Persson, U. Diabetic foot ulcers in a multidisciplinary setting an economic analysis of primary healing and healing with amputation. *J. Intern. Med.* **1994**, *235*, 463–471.
  51. Eslami, M.H.; Zayaruzny, M.; Fitzgerald, G.A. The adverse effects of race, insurance status, and low income on the rate of amputation in patients presenting with lower extremity ischemia. *J. Vasc. Surg.* **2007**, *45*, 55–59.
  52. Galkowska, H.; Wojewodzka, U.; Olszewski, W.L. Low recruitment of immune cells with increased expression of endothelial adhesion molecules in margins of the chronic diabetic foot ulcers. *Wound Repair Regen.* **2005**, *13*, 248–254.
  53. Schaper, N.; Van Netten, J.; Apelqvist, J.; Lipsky, B.; Bakker, K.; Foot, I.W.G.o.t.D. Prevention and management of foot problems in diabetes: A Summary Guidance for Daily Practice 2015, based on the IWGDF Guidance Documents. *Diabetes/Metab. Res. Rev.* **2016**, *32*, 7–15.
  54. Noor, S.; Zubair, M.; Ahmad, J. Diabetic foot ulcer—A review on pathophysiology, classification and microbial etiology. *Diabetes Metab. Syndr. : Clin. Res. Rev.* **2015**, *9*, 192–199.
  55. Clayton, W., Jr.; Elasy, T.A. A review of the pathophysiology, classification, and treatment of foot ulcers in diabetic patients. *Clin. Diabetes* **2009**, *27*, 52–58.
  56. Wild, T.; Rahbarnia, A.; Kellner, M.; Sobotka, L.; Eberlein, T. Basics in nutrition and wound healing. *Nutrition* **2010**, *26*, 862–866.
  57. Sharp, A.; Clark, J. Diabetes and its effects on wound healing. *Nurs. Stand.* **2011**, *25*, 41–47.
  58. Singh, S.; Pai, D.R.; Yuhhui, C. Diabetic foot ulcer—diagnosis and management. *Clin. Res. Foot Ankle* **2013**, *1*, 120.
  59. Syafril, S. Pathophysiology diabetic foot ulcer. In Proceedings of the IOP Conference Series: Earth and Environmental Science, Medan, Indonesia, 15–18 November 2017; IOP Publishing: Bristol, UK, 2018; p. 012161.
  60. Lipsky, B.A.; Aragón-Sánchez, J.; Diggle, M.; Embil, J.; Kono, S.; Lavery, L.; Senneville, É.; Urbančič-Rovan, V.; Van Asten, S.; Peters, E.J. IWGDF guidance on the diagnosis and management of foot infections in persons with diabetes. *Diabetes/Metab. Res. Rev.* **2016**, *32*, 45–74.
  61. Singh, S.; Gupta, B. Choices and challenges of antibiotics therapy in diabetic foot infection. *Indian J. Endocrinol. Metab.* **2017**, *21*, 647.
  62. Ramirez-Acuña, J.M.; Cardenas-Cadena, S.A.; Marquez-Salas, P.A.; Garza-Veloz, I.; Perez-Favila, A.; Cid-Baez, M.A.; Flores-Morales, V.; Martinez-Fierro, M.L. Diabetic foot ulcers: Current advances in antimicrobial therapies and emerging treatments. *Antibiotics* **2019**, *8*, 193.
  63. Edwards, J.; Stapley, S. Debridement of diabetic foot ulcers. *Cochrane Database Syst. Rev.* **2010**, *2010*, CD003556.
  64. Kavitha, K.V.; Tiwari, S.; Purandare, V.B.; Khedkar, S.; Bhosale, S.S.; Unnikrishnan, A.G. Choice of wound care in diabetic foot ulcer: A practical approach. *World J. Diabetes* **2014**, *5*, 546.
  65. Hincliffe, R.; Andros, G.; Apelqvist, J.; Bakker, K.; Fiedrichs, S.; Lammer, J.; Lepantalo, M.; Mills, J.; Reekers, J.; Shearman, C. A systematic review of the effectiveness of revascularization of the ulcerated foot in patients with diabetes and peripheral arterial disease. *Diabetes/Metab. Res. Rev.* **2012**, *28*, 179–217.
  66. Mulder, G.; Alfieri, D. The diabetic foot: Considerations for pressure reduction and off-loading. *Prim. Intent. Aust. J. Wound Manag.* **2007**, *15*, 58–65.
  67. Adams, C.A., Jr.; Deitch, E.A. Diabetic foot infections. In *Surgical Treatment: Evidence-Based and Problem-Oriented*; Holzheimer R.G., Mannick J.A., Eds.; Zuckschwerdt: Munich, Germany, 2001.
  68. Kerr, M.; Barron, E.; Chadwick, P.; Evans, T.; Kong, W.; Rayman, G.; Sutton-Smith, M.; Todd, G.; Young, B.; Jeffcoate, W. The cost of diabetic foot ulcers and amputations to the National Health Service in England. *Diabet. Med.* **2019**, *36*, 995–1002.
  69. Stratman, S.; Schneider, C.; Kirsner, R.S. New Therapies for the Treatment of Diabetic Foot Ulcers: Updated Review of Clinical Trials. *Surg. Technol. Int.* **2020**, *37*, 37–47.
  70. Salehi, B.; Ata, A.V.; Anil Kumar, N.; Sharopov, F.; Ramirez-Alarcon, K.; Ruiz-Ortega, A.; Abdulmajid Ayatollahi, S.; Valere Tsouh Fokou, P.; Kobarfard, F.; Amiruddin Zakaria, Z. Antidiabetic potential of medicinal plants and their active components. *Biomolecules* **2019**, *9*, 551.
  71. Kulprachakarn, K.; Ounjaijean, S.; Wungrath, J.; Mani, R.; Rerkasem, K. Micronutrients and natural compounds status and their effects on wound healing in the diabetic foot ulcer. *Int. J. Low. Extrem. Wounds* **2017**, *16*, 244–250.
  72. Li, S.; Zhao, J.; Liu, J.; Xiang, F.; Lu, D.; Liu, B.; Xu, J.; Zhang, H.; Zhang, Q.; Li, X. Prospective randomized controlled study of a Chinese herbal medicine compound Tangzu Yuyang Ointment for chronic diabetic foot ulcers: A preliminary report. *J. Ethnopharmacol.* **2011**, *133*, 543–550.
  73. Wang, Y.; Cao, H.-J.; Wang, L.-Q.; Lu, C.-L.; Yan, Y.-Q.; Lu, H.; Zhang, K.; Zhang, H.-M.; Liu, J.-P. The effects of Chinese herbal medicines for treating diabetic foot ulcers: A systematic review of 49 randomized controlled trials. *Complement. Ther. Med.* **2019**, *44*, 32–43.

74. Debnath, P.; Prakash, A.; Banerjee, S.; Rao, P.N.; Tripathy, T.B.; Adhikari, A.; Shivakumar. Quality of life and treatment satisfaction observed among Indians with diabetes foot ulcers undergoing ayurvedic adjunct therapy. *J. Evid.-Based Complement. Altern. Med.* **2015**, *20*, 13–19.
75. Alam, G.; Singh, M.P.; Singh, A. Wound healing potential of some medicinal plants. *Int. J. Pharm. Sci. Rev. Res.* **2011**, *9*, 136–145.
76. Tejada, S.; Manayi, A.; Daglia, M.; Nabavi, S.F.; Sureda, A.; Hajheydari, Z.; Gortzi, O.; Pazoki-Toroudi, H.; Nabavi, S.M. Wound healing effects of curcumin: A short review. *Curr. Pharm. Biotechnol.* **2016**, *17*, 1002–1007.
77. Pişkin, A.; Altunkaynak, B.Z.; Tümentemur, G.; Kaplan, S.; Yazıcı, Ö.B.; Hökelek, M. The beneficial effects of *Momordica charantia* (bitter gourd) on wound healing of rabbit skin. *J. Dermatol. Treat.* **2014**, *25*, 350–357.
78. Koca, U.; Süntar, I.; Akkol, E.K.; Yilmazer, D.; Alper, M. Wound repair potential of *Olea europaea* L. leaf extracts revealed by in vivo experimental models and comparative evaluation of the extracts' antioxidant activity. *J. Med. Food* **2011**, *14*, 140–146.
79. Kaur, M.; Valecha, V. Diabetes and antidiabetic herbal formulations: An alternative to Allopathy. *Eur. J. Med.* **2014**, *6*, 226–240.
80. Papanas, N.; Maltezos, E. Polyherbal formulation as a therapeutic option to improve wound healing in the diabetic foot. *Indian J. Med. Res.* **2011**, *134*, 146.
81. Ongarora, B.G. Recent technological advances in the management of chronic wounds: A literature review. *Health Sci. Rep.* **2022**, *5*, e641.
82. Mandrika, I.; Kumar, S.; Zandersone, B.; Eranezhath, S.S.; Petrovska, R.; Liduma, I.; Jezupovs, A.; Pirags, V.; Tracevska, T. Antibacterial and anti-inflammatory potential of polyherbal formulation used in chronic wound healing. *Evid.-Based Complement. Altern. Med.* **2021**, *2021*, 9991454.
83. Roozbeh, N.; Darvish, L. *Acacia nilotica*: New plant for help in pelvic organ prolapse. *J. Menopausal Med.* **2016**, *22*, 129–130.
84. Kankara, S.; Sani, D.; Ibrahim, M.; Mustafa, M.; Go, R. *Acacia nilotica* pods' water extract enhances wound healing in Sprague-Dawley rats by alleviating oxidative stress and suppressing pro-inflammatory cytokines. *Niger. J. Sci. Res.* **2017**, *16*, 202–210.
85. Islam, M.S.; Ara, H.; Ahmad, K.I.; Uddin, M.M. A review on medicinal uses of different plants of Euphorbiaceae family. *UPRA3* **2019**, *4*, 45–49.
86. Gutierrez, R.P. Evaluation of the wound healing properties of *Acalypha langiana* in diabetic rats. *Fitoterapia* **2006**, *77*, 286–289.
87. Shalu, R.; Amanjot, G.S.P.; Kapil, K.; Sukhbir, K. Wound healing potential of medicinal plants with their screening models: A comprehensive review. *J. Drug Deliv. Ther.* **2016**, *6*, 56–66.
88. He, X.; Fang, J.; Chen, X.; Zhao, Z.; Li, Y.; Meng, Y.; Huang, L. *Actinidia chinensis* Planch.: A review of chemistry and pharmacology. *Front. Pharmacol.* **2019**, *10*, 1236.
89. Mohajeri, G.; Safaee, M.; Sanei, M.H. Effects of topical Kiwifruit on healing of neuropathic diabetic foot ulcer. *J. Res. Med. Sci. Off. J. Isfahan Univ. Med. Sci.* **2014**, *19*, 520.
90. Worasakwutiphong, S.; Termwattanaphakdee, T.; Kamolhan, T.; Phimmuan, P.; Sittichokechaiwut, A.; Viyoch, J. Evaluation of the safety and healing potential of a fibroin-aloe gel film for the treatment of diabetic foot ulcers. *J. Wound Care* **2021**, *30*, 1020–1028.
91. Sánchez, M.; González-Burgos, E.; Iglesias, I.; Gómez-Serranillos, M.P. Pharmacological update properties of *Aloe vera* and its major active constituents. *Molecules* **2020**, *25*, 1324.
92. Jalalpure, S.S.; Agrawal, N.; Patil, M.; Chinkode, R.; Tripathi, A. Antimicrobial and wound healing activities of leaves of *Alternanthera sessilis* Linn. *Int. J. Green Pharm. (IJGP)* **2008**, *3*, 141–144.
93. Muniandy, K.; Gothai, S.; Tan, W.S.; Kumar, S.S.; Mohd Esa, N.; Chandramohan, G.; Al-Numair, K.S.; Arulselvan, P. In vitro wound healing potential of stem extract of *Alternanthera sessilis*. *Evid.-Based Complement. Altern. Med.* **2018**, *2018*, 3142073.
94. Kumar, M.; Changan, S.; Tomar, M.; Prajapati, U.; Saurabh, V.; Hasan, M.; Sasi, M.; Maheshwari, C.; Singh, S.; Dhumal, S. Custard apple (*Annona squamosa* L.) leaves: Nutritional composition, phytochemical profile, and health-promoting biological activities. *Biomolecules* **2021**, *11*, 614.
95. Ponrasu, T.; Suguna, L. Efficacy of *Annona squamosa* on wound healing in streptozotocin-induced diabetic rats. *Int. Wound J.* **2012**, *9*, 613–623.
96. Ansari, P.; Flatt, P.R.; Harriott, P.; Abdel-Wahab, Y.H.A. Evaluation of the antidiabetic and insulin releasing effects of *A. squamosa*, including isolation and characterization of active phytochemicals. *Plants* **2020**, *9*, 1348.
97. Singh, D.; Baghel, U.S.; Gautam, A.; Baghel, D.S.; Yadav, D.; Malik, J.; Yadav, R. The genus *Anogeissus*: A review on ethnopharmacology, phytochemistry and pharmacology. *J. Ethnopharmacol.* **2016**, *194*, 30–56.
98. Motto, A.E.; Lawson-Evi, P.; Bakoma, B.; Eklü-Gadegbeku, K.; Aklikokou, K. Antihyperlipidemic and antioxidant properties of hydro-alcoholic extracts from *Anogeissus leiocarpus* (Combretaceae). *Heliyon* **2021**, *7*, e06648.
99. Alhassan, D.A.; Uba, A.I.; Muhammad, A.U.; Muhammad, Y.Y.U. Phytochemical screening and antimicrobial activity of crude stem bark extracts of *Anogeissus leiocarpus*. *Eur. J. Med. Plants* **2016**, *11*, 1–7.
100. Hosseini, A.; Mirzaee, F.; Davoodi, A.; Jouybari, H.B.; Azadbakht, M. The traditional medicine aspects, biological activity and phytochemistry of *Arnebia* spp. *Med. Glas.* **2018**, *15*, 1–9.
101. Nasiri, E.; Hosseinimehr, S.J.; Azadbakht, M.; Akbari, J.; Enayati-Fard, R.; Azizi, S.; Azadbakht, M. The healing effect of *Arnebia euchroma* ointment versus silver sulfadiazine on burn wounds in rat. *World J. Plast. Surg.* **2015**, *4*, 134.
102. Gupta, S.K.; Sharma, O.P.; Raina, N.S.; Sehgal, S. Ethno-botanical study of medicinal plants of Paddar valley of Jammu and Kashmir, India. *Afr. J. Tradit. Complement. Altern. Med.* **2013**, *10*, 59–65.
103. Iabichella, M.L. The use of an extract of *Hypericum perforatum* and *Azadirachta indica* in advanced diabetic foot: An unexpected outcome. *Case Rep.* **2013**, *2013*, bcr2012007299.

104. Maan, P.; Yadav, K.S.; Yadav, N.P. Wound healing activity of *Azadirachta indica* A. juss stem bark in mice. *Pharmacogn. Mag.* **2017**, *13*, S316.
105. Chothani, D.L.; Vaghasiya, H. A review on *Balanites aegyptiaca* Del (desert date): Phytochemical constituents, traditional uses, and pharmacological activity. *Pharmacogn. Rev.* **2011**, *5*, 55.
106. Annan, K.; Dickson, R. Evaluation of wound healing actions of *Hoslundia opposita* vahl, *Anthocleista nobilis* G. Don. and *Balanites aegyptiaca* L. *J. Sci. Technol.* **2008**, *28*, 26–35.
107. Ansari, P.; Flatt, P.R.; Harriott, P.; Abdel-Wahab, Y.H.A. Anti-hyperglycaemic and insulin-releasing effects of *Camellia sinensis* leaves and isolation and characterisation of active compounds. *Br. J. Nutr.* **2021**, *126*, 1149–1163.
108. Shedoeva, A.; Leavesley, D.; Upton, Z.; Fan, C. Wound healing and the use of medicinal plants. *Evid.-Based Complement. Altern. Med.* **2019**, *2019*, 2684108.
109. Hariono, M.; Julianus, J.; Djunarko, I.; Hidayat, I.; Adelya, L.; Indayani, F.; Auw, Z.; Namba, G.; Hariyono, P. The future of *Carica papaya* Leaf extract as an herbal medicine product. *Molecules* **2021**, *26*, 6922.
110. Singh, S.P.; Kumar, S.; Mathan, S.V.; Tomar, M.S.; Singh, R.K.; Verma, P.K.; Kumar, A.; Kumar, S.; Singh, R.P.; Acharya, A. Therapeutic application of *Carica papaya* leaf extract in the management of human diseases. *DARU J. Pharm. Sci.* **2020**, *28*, 735–744.
111. Nayak, B.; Pinto Pereira, L.M. *Catharanthus roseus* flower extract has wound-healing activity in Sprague Dawley rats. *BMC Complement. Altern. Med.* **2006**, *6*, 41.
112. Lee, O.N.; Ak, G.; Zengin, G.; Cziáky, Z.; Jekő, J.; Rengasamy, K.R.; Park, H.Y.; Kim, D.H.; Sivanesan, I. Phytochemical composition, antioxidant capacity, and enzyme inhibitory activity in callus, somaclonal variant, and normal green shoot tissues of *Catharanthus roseus* (L) G. Don. *Molecules* **2020**, *25*, 4945.
113. Rivera-Mondragón, A.; Ortíz, O.O.; Bijttebier, S.; Vlietinck, A.; Apers, S.; Pieters, L.; Caballero-George, C. Selection of chemical markers for the quality control of medicinal plants of the genus *Cecropia*. *Pharm. Biol.* **2017**, *55*, 1500–1512.
114. Nayak, B.S. *Cecropia peltata* L. (Cecropiaceae) has wound-healing potential: A preclinical study in a Sprague Dawley rat model. *Int. J. Low. Extrem. Wounds* **2006**, *5*, 20–26.
115. Arribas-López, E.; Zand, N.; Ojo, O.; Snowden, M.J.; Kochhar, T. A systematic review of the effect of *Centella asiatica* on wound healing. *Int. J. Environ. Res. Public Health* **2022**, *19*, 3266.
116. Gohil, K.J.; Patel, J.A.; Gajjar, A.K. Pharmacological review on *Centella asiatica*: A potential herbal cure-all. *Indian J. Pharm. Sci.* **2010**, *72*, 546.
117. Ahmad, M.; Ansari, M.; Alam, A.; Khan, T. Oral dose of citrus peel extracts promotes wound repair in diabetic rats. *Pak. J. Biol. Sci. PJS* **2013**, *16*, 1086–1094.
118. Lai, S.; Lai, A.; Stange, R.R., Jr.; McCollum, T.G.; Schirra, M. Characterization of the wound-induced material in *Citrus paradisi* fruit peel by carbon-13 CP-MAS solid state NMR spectroscopy. *Phytochemistry* **2003**, *63*, 177–183.
119. Uckoo, R.M.; Jayaprakasha, G.K.; Balasubramaniam, V.; Patil, B.S. Grapefruit (*Citrus paradisi* Macfad) phytochemicals composition is modulated by household processing techniques. *J. Food Sci.* **2012**, *77*, C921–C926.
120. Gupta, D.N.; Rani, R.; Kokane, A.D.; Ghosh, D.K.; Tomar, S.; Sharma, A.K. Characterization of a cytoplasmic 2-Cys peroxiredoxin from *Citrus sinensis* and its potential role in protection from oxidative damage and wound healing. *Int. J. Biol. Macromol.* **2022**, *209*, 1088–1099.
121. Favela-Hernández, J.M.J.; González-Santiago, O.; Ramírez-Cabrera, M.A.; Esquivel-Ferriño, P.C.; Camacho-Corona, M.D.R. Chemistry and Pharmacology of *Citrus sinensis*. *Molecules* **2016**, *21*, 247.
122. Rusak, A.; Rybak, Z. New directions of research related to chronic wound healing. *Polim. Med.* **2013**, *43*, 199–204.
123. Balogun, F.O.; Ashafa, A.O.T. A review of plants used in South African traditional medicine for the management and treatment of hypertension. *Planta Med.* **2019**, *85*, 312–334.
124. Klimek-Szczykutowicz, M.; Szopa, A.; Ekiert, H. *Citrus limon* (Lemon) phenomenon—A review of the chemistry, pharmacological properties, applications in the modern pharmaceutical, food, and cosmetics industries, and biotechnological studies. *Plants* **2020**, *9*, 119.
125. Sami, D.G.; Abdellatif, A.; Azzazy, H.M. Turmeric/oregano formulations for treatment of diabetic ulcer wounds. *Drug Dev. Ind. Pharm.* **2020**, *46*, 1613–1621.
126. Prasad, S.; Aggarwal, B.B. Turmeric, the golden spice. In: *Herbal Medicine: Biomolecular and Clinical Aspects*, 2nd ed.; Benzie I.F.F., Wachtel-Galor S., Eds. CRC Press/Taylor & Francis: Boca Raton, Florida, USA, 2011.
127. Sharifi-Rad, J.; Rayess, Y.E.; Rizk, A.A.; Sadaka, C.; Zgheib, R.; Zam, W.; Sestito, S.; Rapposelli, S.; Neffe-Skocińska, K.; Zielińska, D. Turmeric and its major compound curcumin on health: Bioactive effects and safety profiles for food, pharmaceutical, biotechnological and medicinal applications. *Front. Pharmacol.* **2020**, *11*, 01021.
128. Qabaha, K.; Ras, S.A.; Abbadi, J.; Al-Rimawi, F. Anti-inflammatory of both *Eucalyptus* spp. and *Pistascia lentiscus* were investigated along with their phenolic compounds analysis using HPLC with UV detection. *Afr. J. Tradit. Complement. Altern. Med.* **2016**, *13*, 1–6.
129. Mumtaz, R.; Zubair, M.; Khan, M.A.; Muzammil, S.; Siddique, M.H. Extracts of *Eucalyptus alba* Promote Diabetic Wound Healing by Inhibiting  $\alpha$ -Glucosidase and Stimulating Cell Proliferation. *Evid.-Based Complement. Altern. Med.* **2022**, *2022*, 4953105.
130. Alam, P.; Shakeel, F.; Anwer, M.K.; Foudah, A.I.; Alqarni, M.H. Wound healing study of eucalyptus essential oil containing nanoemulsion in rat model. *J. Oleo Sci.* **2018**, *67*, ess18005.

131. Kumar, S.; Malhotra, R.; Kumar, D. *Euphorbia hirta*: Its chemistry, traditional and medicinal uses, and pharmacological activities. *Pharmacogn. Rev.* **2010**, *4*, 58.
132. Tuhin, R.H.; Begum, M.; Rahman, M.; Karim, R.; Begum, T.; Ahmed, S.U.; Mostofa, R.; Hossain, A.; Abdel-Daim, M.; Begum, R. Wound healing effect of *Euphorbia hirta* linn.(Euphorbiaceae) in alloxan induced diabetic rats. *BMC Complement. Altern. Med.* **2017**, *17*, 423.
133. Bosisio, E.; Mascetti, D.; Verotta, L.; Zani, F.; Mazza, P.; Talbot, M. *Guiera senegalensis* JF Gmelin (Combretaceae): Biological activities and chemical investigation. *Phytomedicine* **1997**, *3*, 339–348.
134. Alshafei, N.K.; Ahmed, S.; Abdelfattah, N. Preliminary observations on the uses of *Guiera senegalensis* as a traditional medicinal plants in Western Kurdofan, Sudan. *J. Plant Biochem. Physiol.* **2016**, *2*, 42–48.
135. Miaffo, D.; Ntchapda, F.; Kamgue, O.G.; Mahamad, A.T.; Kamanyi, A. Glucose-lowering potential of *Guiera senegalensis* roots in a diabetic rat model. *Avicenna J. Phytomedicine* **2020**, *10*, 653.
136. Ifijen, I.; Mamza, A.; Fasina, K.; Omoruyi, J.; Ikhuoria, E. Phytochemical analysis of *Guiera senegalensis* jF Gmel extract and its anti-plasmodial properties on wister albino mice via oral route. *Int. J. Pharmacol. Phytochem. Ethnomed.* **2019**, *13*, 35–44.
137. Daisy, E.A.C.; Rajendran, N.K.; Houreld, N.N.; Marraiki, N.; Elgorban, A.M.; Rajan, M. Curcumin and *Gymnema sylvestre* extract loaded graphene oxide-polyhydroxybutyrate-sodium alginate composite for diabetic wound regeneration. *React. Funct. Polym.* **2020**, *154*, 104671.
138. Tiwari, P.; Mishra, B.; Sangwan, N.S. Phytochemical and pharmacological properties of *Gymnema sylvestre*: An important medicinal plant. *BioMed Res. Int.* **2014**, *2014*, 830285.
139. Bhaskar, A.; Nithya, V. Evaluation of the wound-healing activity of *Hibiscus rosa sinensis* L. (Malvaceae) in Wistar albino rats. *Indian J. Pharmacol.* **2012**, *44*, 694.
140. Shen, H.-M.; Chen, C.; Jiang, J.-Y.; Zheng, Y.-L.; Cai, W.-F.; Wang, B.; Ling, Z.; Tang, L.; Wang, Y.-H.; Shi, G.-G. The N-butyl alcohol extract from *Hibiscus rosa-sinensis* L. flowers enhances healing potential on rat excisional wounds. *J. Ethnopharmacol.* **2017**, *198*, 291–301.
141. Ansari, P.; Azam, S.; Hannan, J.M.A.; Flatt, P.R.; Abdel-Wahab, Y.H.A. Anti-hyperglycaemic activity of *H. rosa-sinensis* leaves is partly mediated by inhibition of carbohydrate digestion and absorption, and enhancement of insulin secretion. *J. Ethnopharmacol.* **2020**, *253*, 112647.
142. Perez G,R.; Vargas S,R.; Ortiz H,Y. Wound healing properties of *Hylocereus undatus* on diabetic rats. *Phytother. Res. Int. J. Devoted Pharmacol. Toxicol. Eval. Nat. Prod. Deriv.* **2005**, *19*, 665–668.
143. Joshi, M.; Prabhakar, B. Phytoconstituents and pharmaco-therapeutic benefits of pitaya: A wonder fruit. *J. Food Biochem.* **2020**, *44*, e13260.
144. Yadollah-Damavandi, S.; Chavoshi-Nejad, M.; Jangholi, E.; Nekouyian, N.; Hosseini, S.; Seifae, A.; Rafiee, S.; Karimi, H.; Ashkani-Esfahani, S.; Parsa, Y. Topical *Hypericum perforatum* improves tissue regeneration in full-thickness excisional wounds in diabetic rat model. *Evid.-Based Complement. Altern. Med.* **2015**, *2015*, 245328.
145. Chowdhury, M.Z.; Al Mahmud, Z.; Ali, M.S.; Bachar, S.C. Phytochemical and pharmacological investigations of rhizome extracts of *Kaempferia galanga*. *Int. J. Pharmacogn.* **2014**, *1*, 185–192.
146. Shanbhag, T.V.; Sharma, C.; Adiga, S.; Bairy, L.K.; Shenoy, S.; Shenoy, G. Wound healing activity of alcoholic extract of *Kaempferia galanga* in Wistar rats. *Indian J. Physiol. Pharmacol.* **2006**, *50*, 384–390.
147. Wang, S.-Y.; Zhao, H.; Xu, H.-T.; Han, X.-D.; Wu, Y.-S.; Xu, F.-F.; Yang, X.-B.; Göransson, U.; Liu, B. *Kaempferia galanga* L.: Progresses in phytochemistry, pharmacology, toxicology and ethnomedicinal uses. *Front. Pharmacol.* **2021**, *12*, 675350.
148. Al-Snafi, A. A review on *Lawsonia inermis*: A potential medicinal plant. *Int. J. Curr. Pharm. Res.* **2019**, *11*, 1–13.
149. Franco, E.D.S.; de Aquino, C.M.F.; Medeiros, P.L.D.; Evêncio, L.B.; Góes, A.J.D.S.; Maia, M.B.D.S. Effect of a semisolid formulation of *Linum usitatissimum* L. (Linseed) oil on the repair of skin wounds. *Evid.-Based Complement. Altern. Med.* **2012**, *2012*, 270752.
150. Soleimani, Z.; Hashemdokht, F.; Bahmani, F.; Taghizadeh, M.; Memarzadeh, M.R.; Asemi, Z. Clinical and metabolic response to flaxseed oil omega-3 fatty acids supplementation in patients with diabetic foot ulcer: A randomized, double-blind, placebo-controlled trial. *J. Diabetes Its Complicat.* **2017**, *31*, 1394–1400.
151. Ansari, P.; Hannon-Fletcher, M.P.; Flatt, P.R.; Abdel-Wahab, Y.H.A. Effects of 22 traditional anti-diabetic medicinal plants on DPP-IV enzyme activity and glucose homeostasis in high-fat fed obese diabetic rats. *Biosci. Rep.* **2021**, *41*, BSR20203824.
152. Manjunatha, B.; Krishna, V.; Vidya, S.; Mankani, K.; Manohara, Y. Wound healing activity of *Lycopodium serratum*. *Indian J. Pharm. Sci.* **2007**, *69*, 283.
153. Lodhi, S.; Singhai, A.K. Wound healing effect of flavonoid rich fraction and luteolin isolated from *Martynia annua* Linn. on streptozotocin induced diabetic rats. *Asian Pac. J. Trop. Med.* **2013**, *6*, 253–259.
154. Radhakrishna, K.; Vvs Nrk, B.M.; Kuppusamy, G. Potential use of herbal medicines in the treatment of diabetic foot ulcers. *History* **2014**, *14*, 34–42.
155. Chorepsima, S.; Tentolouris, K.; Dimitroulis, D.; Tentolouris, N. Melilotus: Contribution to wound healing in the diabetic foot. *J. Herb. Med.* **2013**, *3*, 81–86.
156. Al-Snafi, A.E. Chemical constituents and pharmacological effects of *Melilotus Officinalis*-A review. *IOSR J. Pharm.* **2020**, *10*, 26–36.

157. Zippel, J.; Deters, A.; Hensel, A. Arabinogalactans from *Mimosa tenuiflora* (Willd.) Poir bark as active principles for wound-healing properties: Specific enhancement of dermal fibroblast activity and minor influence on HaCaT keratinocytes. *J. Ethnopharmacol.* **2009**, *124*, 391–396.
158. Souza, R.S.O.D.; Albuquerque, U.P.D.; Monteiro, J.M.; Amorim, E.L.C.D. Jurema-Preta (*Mimosa tenuiflora* [Willd.] Poir.): A review of its traditional use, phytochemistry and pharmacology. *Braz. Arch. Biol. Technol.* **2008**, *51*, 937–947.
159. Singh, J.; Cumming, E.; Manoharan, G.; Kalasz, H.; Adeghate, E. Suppl 2: Medicinal chemistry of the anti-diabetic effects of *Momordica charantia*: Active constituents and modes of actions. *Open Med. Chem. J.* **2011**, *5*, 70.
160. Ly, H.T.; Pham Nguyen, M.T.; Nguyen, T.K.O.; Bui, T.P.Q.; Ke, X.; Le, V.M. Phytochemical analysis and wound-healing activity of noni (*Morinda citrifolia*) leaf extract. *J. Herbs Spices Med. Plants* **2020**, *26*, 379–393.
161. Deng, S.; West, B.J.; Jensen, C.J. Phytochemical, antioxidant and toxicological investigation of *Morinda citrifolia* L. blossoms. *Int. Sch. Res. Not.* **2012**, *2012*.
162. Raeiszadeh, M.; Esmaeili-Tarzi, M.; Bahrapour-Juybari, K.; Nematollahi-Mahani, S.; Pardakhty, A.; Nematollahi, M.; Mehrabani, M. Evaluation the effect of *Myrtus communis* L. extract on several underlying mechanisms involved in wound healing: An in vitro study. *S. Afr. J. Bot.* **2018**, *118*, 144–150.
163. Sisay, M.; Gashaw, T. Ethnobotanical, ethnopharmacological, and phytochemical studies of *Myrtus communis* Linn: A popular herb in Unani system of medicine. *J. Evid.-Based Complement. Altern. Med.* **2017**, *22*, 1035–1043.
164. Sallehuddin, N.; Nordin, A.; Bt Hj Idrus, R.; Fauzi, M.B. *Nigella sativa* and its active compound, thymoquinone, accelerate wound healing in an in vivo animal model: A comprehensive review. *Int. J. Environ. Res. Public Health* **2020**, *17*, 4160.
165. Kooti, W.; Hasanzadeh-Noohi, Z.; Sharafi-Ahvazi, N.; Asadi-Samani, M.; Ashtary-Larky, D. Phytochemistry, pharmacology, and therapeutic uses of black seed (*Nigella sativa*). *Chin. J. Nat. Med.* **2016**, *14*, 732–745.
166. Abdullah, F.I.; Chua, L.S.; Mohd Bohari, S.P.; Sari, E. Rationale of *Orthosiphon aristatus* for healing diabetic foot ulcer. *Nat. Prod. Commun.* **2020**, *15*. <https://doi.org/10.1177/1934578X209533>.
167. Asraf, M.H.; Sani, N.S.; Williams, C.D.; Jemon, K.; Malek, N.A.N.N. In situ biosynthesized silver nanoparticle-incorporated synthesized zeolite A using *Orthosiphon aristatus* extract for in vitro antibacterial wound healing. *Particuology* **2022**, *67*, 27–34.
168. Gupta, A.; Verma, S.; Gupta, A.; Jangra, M.; Pratap, R. Evaluation of *Prosopis cineraria* (Linn.) Druce leaves for wound healing activity in rats. *Ann. Pharm. Res* **2015**, *3*, 70–74.
169. Yadav, E.; Singh, D.; Yadav, P.; Verma, A. Antioxidant and anti-inflammatory properties of *Prosopis cineraria* based phenolic rich ointment in wound healing. *Biomed. Pharmacother.* **2018**, *108*, 1572–1583.
170. Manne, A.A.; Arigela, B.; Giduturi, A.K.; Komaravolu, R.K.; Mangamuri, U.; Poda, S. *Pterocarpus marsupium* Roxburgh heartwood extract/chitosan nanoparticles loaded hydrogel as an innovative wound healing agent in the diabetic rat model. *Mater. Today Commun.* **2021**, *26*, 101916.
171. Nayak, S.B.; Rodrigues, V.; Maharaj, S.; Bhogadi, V.S. Wound healing activity of the fruit skin of *Punica granatum*. *J. Med. Food* **2013**, *16*, 857–861.
172. Arun, N.; Singh, D. *Punica granatum*: A review on pharmacological and therapeutic properties. *Int. J. Pharm. Sci. Res.* **2012**, *3*, 1240.
173. Lau, T.; Lam, F.; Lau, K.; Chan, Y.; Lee, K.; Sahota, D.; Ho, Y.; Fung, K.; Leung, P.; Lau, C. Pharmacological investigation on the wound healing effects of *Radix Rehmanniae* in an animal model of diabetic foot ulcer. *J. Ethnopharmacol.* **2009**, *123*, 155–162.
174. Kim, S.-H.; Yook, T.-H.; Kim, J.-U. *Rehmanniae radix*, an effective treatment for patients with various inflammatory and metabolic diseases: Results from a review of Korean publications. *J. Pharmacopunct.* **2017**, *20*, 81.
175. Andrade, J.M.; Faustino, C.; Garcia, C.; Ladeiras, D.; Reis, C.P.; Rijo, P. *Rosmarinus officinalis* L.: An update review of its phytochemistry and biological activity. *Future Sci. OA* **2018**, *4*, FSO283.
176. de Macedo, L.M.; Santos, É.M.d.; Militão, L.; Tundisi, L.L.; Ataíde, J.A.; Souto, E.B.; Mazzola, P.G. Rosemary (*Rosmarinus officinalis* L., syn *Salvia rosmarinus* Spenn.) and its topical applications: A review. *Plants* **2020**, *9*, 651.
177. Shenoy, R.R.; Sudheendra, A.T.; Nayak, P.G.; Paul, P.; Kutty, N.G.; Rao, C.M. Normal and delayed wound healing is improved by sesamol, an active constituent of *Sesamum indicum* (L.) in albino rats. *J. Ethnopharmacol.* **2011**, *133*, 608–612.
178. Andargie, M.; Vinas, M.; Rathgeb, A.; Möller, E.; Karlovsky, P. Lignans of sesame (*Sesamum indicum* L.): A comprehensive review. *Molecules* **2021**, *26*, 883.
179. Singla, R.; Soni, S.; Patial, V.; Kulurkar, P.M.; Kumari, A.; Padwad, Y.S.; Yadav, S.K. Cytocompatible anti-microbial dressings of *Syzygium cumini* cellulose nanocrystals decorated with silver nanoparticles accelerate acute and diabetic wound healing. *Sci. Rep.* **2017**, *7*, 10457.
180. Ramya, S.; Neethirajan, K.; Jayakumararaj, R. Profile of bioactive compounds in *Syzygium cumini*—A review. *J. Pharm. Res.* **2012**, *5*, 4548–4553.
181. Singh, D.; Singh, D.; Choi, S.M.; Zo, S.M.; Painuli, R.M.; Kwon, S.W.; Han, S.S. Effect of extracts of *Terminalia chebula* on proliferation of keratinocytes and fibroblasts cells: An alternative approach for wound healing. *Evid.-Based Complement. Altern. Med.* **2014**, *2014*, 701656.
182. Sharma, P.; Dwivedee, B.P.; Bisht, D.; Dash, A.K.; Kumar, D. The chemical constituents and diverse pharmacological importance of *Tinospora cordifolia*. *Heliyon* **2019**, *5*, e02437.
183. Moghadam, F.H.; Vakili-Zarch, B.; Shafiee, M.; Mirjalili, A. Fenugreek seed extract treats peripheral neuropathy in pyridoxine induced neuropathic mice. *Excli J.* **2013**, *12*, 282.

184. Taranalli, A.; Kuppast, I. Study of wound healing activity of seeds of *Trigonella foenum graecum* in rats. *Indian J. Pharm. Sci.* **1996**, *58*, 117.
185. Mohamed, A.H.B.; Osman, A.A.F. Antibacterial and wound healing potential of ethanolic extract of *Zingiber Officinale* in albino rats. *J. Dis. Med. Plants* **2017**, *3*, 1–6.
186. Kumar Gupta, S.; Sharma, A. Medicinal properties of *Zingiber officinale* Roscoe-A review. *J. Pharm. Biol. Sci* **2014**, *9*, 124–129.
187. Kooti, W.; Farokhipour, M.; Asadzadeh, Z.; Ashtary-Larky, D.; Asadi-Samani, M. The role of medicinal plants in the treatment of diabetes: A systematic review. *Electron. Physician* **2016**, *8*, 1832.
188. Lim, J.Z.M.; Ng, N.S.L.; Thomas, C. Prevention and treatment of diabetic foot ulcers. *J. R. Soc. Med.* **2017**, *110*, 104–109.
189. Schaper, N.C.; van Netten, J.J.; Apelqvist, J.; Bus, S.A.; Hinchliffe, R.J.; Lipsky, B.A.; Board, I.E. Practical guidelines on the prevention and management of diabetic foot disease (IWGDF 2019 update). *Diabetes/Metab. Res. Rev.* **2020**, *36*, e3266.
190. Braun, L.; Kim, P.J.; Margolis, D.; Peters, E.J.; Lavery, L.A. What's new in the literature: An update of new research since the original WHS diabetic foot ulcer guidelines in 2006. *Wound Repair Regen.* **2014**, *22*, 594–604.
191. Everett, E.; Mathioudakis, N. Update on management of diabetic foot ulcers. *Ann. N. Y. Acad. Sci.* **2018**, *1411*, 153–165.
192. Mandal, M.D.; Mandal, S. Honey: Its medicinal property and antibacterial activity. *Asian Pac. J. Trop. Biomed.* **2011**, *1*, 154–160.
193. Van Netten, J.J.; Woodburn, J.; Bus, S.A. The future for diabetic foot ulcer prevention: A paradigm shift from stratified healthcare towards personalized medicine. *Diabetes/Metab. Res. Rev.* **2020**, *36*, e3234.
194. Eldor, R.; Raz, I.; Ben Yehuda, A.; Boulton, A. New and experimental approaches to treatment of diabetic foot ulcers: A comprehensive review of emerging treatment strategies. *Diabet. Med.* **2004**, *21*, 1161–1173.
195. Seyed, M.A.; Ayesha, S. Modern Phytomedicine in Treating Diabetic Foot Ulcer: Progress and Opportunities. In *Diabetic Foot Ulcer*; Zubair, M., Ahmad, J., Malik, A., Talluri, M.R., Eds.; Springer: Singapore, 2021; pp. 281–313.