University of Texas Rio Grande Valley

ScholarWorks @ UTRGV

School of Medicine Publications and Presentations

School of Medicine

2018

Non-congenital severe ocular complications of Zika virus infection

Mussaret B. Zaidi

c Gustavo De Moraes

Michele Petitto

Juan B. Yepez

Anavaj Sakuntabhai

See next page for additional authors

Follow this and additional works at: https://scholarworks.utrgv.edu/som_pub

Authors

Mussaret B. Zaidi, c Gustavo De Moraes, Michele Petitto, Juan B. Yepez, Anavaj Sakuntabhai, Etienne Simon-Loriere, Matthieu Prot, Claude Ruffie, Susan S. Kim, and Gladys E. Maestre

CASE QUIZ Zaidi et al., JMM Case Reports 2018;5 DOI 10.1099/jmmcr.0.005152



Non-congenital severe ocular complications of Zika virus infection

Mussaret B. Zaidi,^{1,2} C. Gustavo De Moraes,³ Michele Petitto,⁴ Juan B. Yepez,⁵ Anavaj Sakuntabhai,^{6,7} Etienne Simon-Loriere,^{6,7} Matthieu Prot,^{6,7} Claude Ruffie,^{5,7} Susan S. Kim,⁸ Rando Allikmets,³ Joseph D. Terwilliger,^{9,10,11,12} Joseph H. Lee^{10,13} and Gladys E. Maestre^{14,15,16,*}

CASE SUMMARY

In 2016, during a major Zika virus (ZIKV) outbreak in Maracaibo, Venezuela, a 49-year-old woman and an unrelated 4-year-old boy developed bilateral optic neuritis 2-3 weeks after presenting an acute febrile illness characterized by low-grade fever, rash and myalgia [1]. Both patients presented sudden, painless bilateral loss of vision with no corneal or uveal abnormalities. Fundoscopic examination revealed oedema of the optic nerve and optic disc pallor. Optical coherence tomography confirmed bilateral optic nerve head swelling in the case of the adult, but it was not carried out in the child. Automated perimetry performed in the adult revealed bilateral diffuse visual field loss. Magnetic resonance imaging of the brain in both cases was unremarkable. Both patients were diagnosed with bilateral optic neuritis of possible infectious or parainfectious origin. Differential diagnoses that were considered and subsequently discarded included arteritic and non-arteritic ischaemic optic neuropathy, and brain disorders such as multiple sclerosis and brain tumours. Both patients were seropositive for anti-ZIKV IgG and seronegative for anti-ZIKV IgM. In addition, both patients were positive for anti-dengue virus (DENV) IgG for all four DENV serotypes. Management included intravenous methylprednisolone for 3 days, followed by oral prednisolone for 11 days. Although the patients presented a modest improvement in their vision, they continued to have visual impairment after several months of follow-up [1].

QUESTION

Which of the following statements is accurate about noncongenital severe ocular complications of Zika virus (ZIKV) infection?

ANSWER OPTIONS

1. They are unique to ZIKV infection and readily distinguishable from complications caused by other flaviviruses.

2. Serious ocular complications are related to the severity of the acute exanthematous illness.

3. The diagnosis can be conclusively established by detecting anti-Zika IgM and/or IgG in the patient's serum.

4. These complications can lead to permanent visual impairment.

5. There is specific treatment for ocular manifestations caused by ZIKV infection.

DISCUSSION

Correct Answer: 4. These complications can lead to permanent visual impairment.

ZIKV is a mosquito-borne RNA virus belonging to the genus *Flavivirus* of the family *Flaviviridae* [2]. The classical

Received 1 April 2018; Accepted 23 April 2018

*Correspondence: Gladys E. Maestre, gladys.maestre@utrgv.edu or gladysmaestre@gmail.com

Keywords: Zika virus; dengue; Arbovirus; *Flavivirus*; uveitis; optic neuritis; vision loss; steroid treatment.

Abbreviations: DENV, dengue virus; ZIKV, Zika virus.

Author affiliations: ¹Infectious Diseases Research Laboratory, Hospital General O'Horan, Merida, Mexico; ²Department of Epidemiology and Biostatistics, Michigan State University, East Lansing, MI, USA; ³Department of Ophthalmology, Columbia University Medical Center, New York, NY, USA; ⁴Clinica de Ojos de Maracaibo, Maracaibo, Venezuela; ⁵Department of Ophthalmology, King Khaled Eye Specialist Hospital, Riyadh, Saudi Arabia; ⁶Functional Genetics of Infectious Diseases Unit, Institut Pasteur, Paris, France, CNRS, UMR2000, Paris, France; ⁷Viral Genomics and Vaccination Unit, Institut Pasteur, Paris, France, CNRS, UMR 3965, Paris, France; ⁸In-patient Diabetes Unit, St Peter's Hospital, Albany, NY, USA; ⁹Departments of Psychiatry and Genetics and Development, Columbia University Medical Center, New York, NY, USA; ¹⁰Sergievsky Center, Columbia University Medical Center, New York, NY, USA; ¹¹Division of Medical Genetics, New York State Psychiatric Institute, New York, NY, USA; ¹²Public Health Genomics Unit, National Institute for Health and Welfare, Helsinki, Finland; ¹³Taub Institute and Department of Epidemiology, Columbia University Medical Center, New York, NY, USA; ¹⁴Laboratory of Neuroscience, University of Zulia, Maracaibo, Venezuela; ¹⁶Department of Biomedical Sciences, Division of Neurosciences, University of Texas Rio Grande Valley School of Medicine, Brownsville, TX, USA; ¹⁶Department of Human Genetics, University of Texas Rio Grande Valley School of Medicine, Brownsville, TX, USA; ¹⁶Department of Human Genetics, University of Texas Rio Grande Valley School of Medicine, Brownsville, TX, USA; ¹⁶Department of Human Genetics, University of Texas

Type of virus	Complication
Zika	Uveitis [1, 41, 42]; macular atrophy [46]; chorioretinal scar [47]; chorioretinal macular atrophy [43]
Dengue	Maculopathy or neuropathy [3-8]; macular oedema, retinal detachment, retinal vascular occlusion, choroidal changes, optic disk swelling, optic neuritis and neuroretinitis [3-7, 9-20]
Chikungunya	Optic neuropathy [21]; conjunctivitis, episcleritis, keratitis, panuveitis, multifocal choroiditis, optic neuritis, neuroretinitis, central retinal artery occlusion, panophthalmitis, lagophthalmos and sixth nerve palsy [22–24]
West Nile	Foveal chorioretinal scar, choroidal neovascularization [25]; vitreous haemorrhage secondary to retinal neovascularization, severe ischaemic maculopathy [26]; macular oedema, optic atrophy or retrogeniculate damage; occlusive vasculitis [27–29]; uveitis [30]; vitritis [31]; chorioretinitis [31–35]; multifocal choroiditis [36]; chorioretinal lesions [37]; and optic neuritis [38, 39]
Rift Valley fever	Macular or paramacular scarring, retinal vascular occlusion or optic atrophy [9-11, 40]

Table 1. Non-congenital ocular complications of common Arbovirus infections

clinical picture of ZIKV infection includes fever, exanthema, headache and conjunctivitis. The most common noncongenital, ocular manifestation of ZIKV infection is a selflimiting conjunctivitis. Serious ocular complications have been reported for other arboviruses, such as DENV [3–20], chikungunya virus [21–24], West Nile virus [25–39] and Rift Valley fever virus [9–11, 40] (Table 1). To date, there is no specific ocular lesion that is pathognomonic for ZIKV infection [41–45].

Non-congenital ocular complications are infrequent, but serious, consequences of ZIKV and other arboviral infections. The complications may appear at the end of the acute febrile illness, but more commonly occur within 2 weeks to 1 month after the onset of symptoms. There is no evidence to suggest that serious ocular complications correlate with the severity of the acute febrile illness. One study, however, found that the white cell count and serum albumin are significant predictors of ocular complications of DENV [46].

Serological testing for arboviral diseases should be performed in all patients with ocular complications and a recent history of acute febrile exanthematous infection, who live, or have travelled to, endemic regions. The presence of IgM to ZIKV strongly suggests that the ocular manifestation is associated with this virus. A causative aetiology, however, can only be established by documenting the presence of the virus in body fluids, either by cell culture or by PCR. It should be noted that other viruses, such as herpes simplex virus and human immunodeficiency virus, can also cause retinal damage and optic neuritis. Furthermore, as in the cases presented here, the diagnosis is complicated by crossreactivity among flaviviruses, and by the co-circulation of arboviruses.

Most patients with ocular complications of arboviral infections recover completely. Nevertheless, physicians should be aware that a small percentage of patients have permanent damage with long-life visual impairment.

There is no specific or established treatment for optic neuritis caused by any arboviral infection. Systemic steroids may be used to reduce inflammation and resulting ischaemia. Corticosteroids have been used in combination with acyclovir to treat chikungunya-associated optic neuritis, but efficacy has not been proven [47].

Funding information

The authors received no specific grant from any funding agency.

Acknowledgements

We thank Elvira Alvarez and Javier Aracena for their help with literature searches. We also acknowledge the South Texas Diabetes and Obesity Institute for support in preparation of this Case Quiz.

Conflicts of interest

The authors declare that there are no conflicts of interest.

Ethical statement

Our study consists of observations only and no intervention/change to patient care was made.

References

- de Moraes CG, Pettito M, Yepez JB, Sakuntabhai A, Simon-Loriere E et al. Optic neuropathy and congenital glaucoma associated with probable Zika virus infection in Venezuelan patients. JMM Case Rep 2018;5:jmmcr.0.005145.
- Musso D, Gubler DJ. Zika virus. Clin Microbiol Rev 2016;29:487– 524.
- Cruz-Villegas V, Berrocal AM, Davis JL. Bilateral choroidal effusions associated with dengue fever. *Retina* 2003;23:576–578.
- Lim WK, Mathur R, Koh A, Yeoh R, Chee SP. Ocular manifestations of dengue fever. Ophthalmology 2004;111:2057–2064.
- Bacsal KE, Chee SP, Cheng CL, Flores JV. Dengue-associated maculopathy. Arch Ophthalmol 2007;125:501–510.
- Beral L, Laurence B, Merle H, Harold M, David T et al. Ocular complications of dengue fever. Ophthalmology 2008;115:1100– 1101.
- Teoh SC, Chee CK, Laude A, Goh KY, Barkham T et al. Optical coherence tomography patterns as predictors of visual outcome in dengue-related maculopathy. *Retina* 2010;30:390–398.
- 8. Chia A, Luu CD, Mathur R, Cheng B, Chee SP. Electrophysiological findings in patients with dengue-related maculopathy. *Arch Ophthalmol* 2006;124:1421–1426.
- Khairallah M, Jelliti B, Jenzeri S. Emergent infectious uveitis. Middle East Afr J Ophthalmol 2009;16:225–238.
- Khairallah M, Chee SP, Rathinam SR, Attia S, Nadella V. Novel infectious agents causing uveitis. Int Ophthalmol 2010;30:465–483.
- 11. Khairallah M, Yahia SB, Attia S. Arthropod vector-borne uveitis in the developing world. *Int Ophthalmol Clin* 2010;50:125–144.
- 12. Chan DP, Teoh SC, Tan CS, Nah GK, Rajagopalan R *et al.* Ophthalmic complications of dengue. *Emerg Infect Dis* 2006;12:285–289.
- Siqueira RC, Vitral NP, Campos WR, Oréfice F, de Moraes Figueiredo LT. Ocular manifestations in dengue fever. Ocul Immunol Inflamm 2004;12:323–327.

- Chlebicki MP, Ang B, Barkham T, Laude A. Retinal hemorrhages in 4 patients with dengue fever. *Emerg Infect Dis* 2005;11:770– 772.
- Loh BK, Bacsal K, Chee SP, Cheng BC, Wong D. Foveolitis associated with dengue fever: a case series. *Ophthalmologica* 2008;222: 317–320.
- Kanungo S, Shukla D, Kim R. Branch retinal artery occlusion secondary to dengue fever. *Indian J Ophthalmol* 2008;56:73–74.
- de Amorim Garcia CA, Gomes AH, de Oliveira AG. Bilateral stellar neuroretinitis in a patient with dengue fever. *Eye* 2006;20:1382– 1383.
- Venkatramani J, Lim WK. Bilateral vitreous haemorrhage associated with dengue fever. *Eye* 2006;20:1405–1406.
- Sanjay S, Wagle AM, Au Eong KG. Dengue optic neuropathy. Ophthalmology 2009;116:170.
- Tabbara K. Dengue retinochoroiditis. Ann Saudi Med 2012;32:530– 533.
- Mahendradas P, Avadhani K, Shetty R. Chikungunya and the eye: a review. J Ophthalmic Inflamm Infect 2013;3:35.
- Mahendradas P, Ranganna SK, Shetty R, Balu R, Narayana KM et al. Ocular manifestations associated with chikungunya. Ophthalmology 2008;115:287–291.
- Lalitha P, Rathinam S, Banushree K, Maheshkumar S, Vijayakumar R et al. Ocular involvement associated with an epidemic outbreak of chikungunya virus infection. Am J Ophthalmol 2007;144:552–556.
- Mittal A, Mittal S, Bharati MJ, Ramakrishnan R, Saravanan S et al. Optic neuritis associated with chikungunya virus infection in South India. Arch Ophthalmol 2007;125:1381–1386.
- Seth RK, Stoessel KM, Adelman RA. Choroidal neovascularization associated with West Nile virus chorioretinitis. *Semin Ophthalmol* 2007;22:81–84.
- Khairallah M, Ben Yahia S, Attia S, Zaouali S, Jelliti B et al. Indocyanine green angiographic features in multifocal chorioretinitis associated with West Nile virus infection. *Retina* 2006;26:358–359.
- Kaiser PK, Lee MS, Martin DA. Occlusive vasculitis in a patient with concomitant West Nile virus infection. Am J Ophthalmol 2003; 136:928–930.
- Teitelbaum BA, Newman TL, Tresley DJ. Occlusive retinal vasculitis in a patient with West Nile virus. *Clin Exp Optom* 2007; 90:463–467.
- Gohari AR, Willson RL, Gitter KA. West Nile virus occlusive retinal vasculitis. *Retin Cases Brief Rep* 2011;5:209–212.
- Kuchtey RW, Kosmorsky GS, Martin D, Lee MS. Uveitis associated with West Nile virus infection. Arch Ophthalmol 2003;121:1648– 1649.

- Bains HS, Jampol LM, Caughron MC, Parnell JR. Vitritis and chorioretinitis in a patient with West Nile virus infection. Arch Ophthalmol 2003;121:205–207.
- Adelman RA, Membreno JH, Afshari NA, Stoessel KM. West Nile virus chorioretinitis. *Retina* 2003;23:100–101.
- Shaikh S, Trese MT. West Nile virus chorioretinitis. Br J Ophthalmol 2004;88:1599–1560.
- Eidsness RB, Stockl F, Colleaux KM. West Nile chorioretinitis. Can J Ophthalmol 2005;40:721–724.
- Myers JP, Leveque TK, Johnson MW. Extensive chorioretinitis and severe vision loss associated with West Nile virus meningoencephalitis. Arch Ophthalmol 2005;123:1754–1756.
- Vandenbelt S, Shaikh S, Capone A, Williams GA. Multifocal choroiditis associated with West Nile virus encephalitis. *Retina* 2003;23: 97–99.
- Khairallah M, Ben Yahia S, Ladjimi A, Zeghidi H, Ben Romdhane F et al. Chorioretinal involvement in patients with West Nile virus infection. Ophthalmology 2004;111:2065–2070.
- Vaispapir V, Blum A, Soboh S, Ashkenazi H. West Nile virus meningoencephalitis with optic neuritis. Arch Intern Med 2002;162: 606-a-6607.
- Gilad R, Lampl Y, Sadeh M, Paul M, Dan M. Optic neuritis complicating West Nile virus meningitis in a young adult. *Infection* 2003; 31:55–56.
- Al-Hazmi A, Al-Rajhi AA, Abboud EB, Ayoola EA, Al-Hazmi M et al. Ocular complications of Rift Valley fever outbreak in Saudi Arabia. Ophthalmology 2005;112:313–318.
- Furtado JM, Espósito DL, Klein TM, Teixeira-Pinto T, da Fonseca BA. Uveitis associated with Zika virus infection. N Engl J Med 2016;375:394–396.
- Kodati S, Palmore TN, Spellman FA, Cunningham D, Weistrop B et al. Bilateral posterior uveitis associated with Zika virus infection. The Lancet 2017;389:125–126.
- Parke DW, Almeida DR, Albini TA, Ventura CV, Berrocal AM et al. Serologically confirmed Zika-related unilateral acute maculopathy in an adult. Ophthalmology 2016;123:2432–2433.
- Wong CW, Ng SR, Cheung CM, Wong TY, Mathur R. Zika-related maculopathy. *Retin Cases Brief Rep* 2017:doi: 10.1097/ ICB.000000000000552
- Marquezan MC, Ventura CV, Sheffield JS, Golden WC, Omiadze R et al. Ocular effects of Zika virus-a review. Surv Ophthalmol 2018; 63:166–173.
- Seet RC, Quek AM, Lim EC. Symptoms and risk factors of ocular complications following dengue infection. J Clin Virol 2007;38:101– 105.
- Mahendradas P, Ranganna SK, Shetty R, Balu R, Narayana KM et al. Ocular manifestations associated with chikungunya. Ophthalmology 2008;115:287–291.

Five reasons to publish your next article with a Microbiology Society journal

- 1. The Microbiology Society is a not-for-profit organization.
- 2. We offer fast and rigorous peer review average time to first decision is 4–6 weeks.
- 3. Our journals have a global readership with subscriptions held in research institutions around the world.
- 4. 80% of our authors rate our submission process as 'excellent' or 'very good'.
- 5. Your article will be published on an interactive journal platform with advanced metrics.

Find out more and submit your article at microbiologyresearch.org.