



Simulating the effects of mobility restrictions in the spread of SARS-CoV-2 in metropolitan areas



Sandra Oliveira*

Jorge Rocha

*Institute of Geography and Spatial Planning,
University of Lisbon, Portugal*



ISPUP

INSTITUTO DE SAÚDE PÚBLICA
DA UNIVERSIDADE DO PORTO

Ana Isabel Ribeiro

*Institute of Public Health,
University of Porto, Portugal*

In Portugal, strict lockdown periods have been established since March 2020:

- limitations on inter-municipal flows
- suspension of all non-essential travel
- compulsory remote work



Simulations of daily new infections in metropolitan areas for different mobility scenarios, to test the potential effect of travel restrictions on virus dissemination.

Metropolitan areas:

- the most densely populated areas in the country
- intense commuting flows
- virus dissemination across municipalities



Lisbon (LMA)


- 18 municipalities
- 3015 km²
- 2 840 005 inhabitants (28%)
- 81% of commuters
- 16% public transport



Porto (PMA)

- 17 municipalities
- 2041 km²
- 1 721 038 inhabitants (17%)
- 79% of commuters
- 11% public transport





Daily new cases. Active cases per day. Virus spread over time. Relation with mobility flows and restrictions.

Based on the regulations implemented by the Portuguese government, essential sectors are:

- ✓ Agriculture, fisheries, and forestry; extraction and transformation industry;
- ✓ electricity and gas supply; water supply, sanitation, and residues;
- ✓ construction; trade and auto repair;
- ✓ transport and storage;
- ✓ human health.



All population is initially susceptible

First infections based on nr. cases PT 23/03

First exposed calculated with model

Update of SEIR matrix
based on initial cases

```
# set up SEIR matrix
SEIR <- matrix(nrow = locs_len, ncol = 4) # initiate an empty SEIR matrix
colnames(SEIR) <- c("S", "E", "I", "R") # rename the vectors
SEIR[, "S"] <- N_k # assign the number of susceptible people in each cell
SEIR[, "E"] <- 0 # assign the number of exposed people in each cell
SEIR[, "I"] <- 0 # assign the number of infected people in each cell
SEIR[, "R"] <- 0 # assign the number of recovered people in each cell

# first infection
# assume no all are infectious and no one in incubation period at the beginning
first_infections <- (pop_cases$origin == zone0_id) * zone0_infected
str(first_infections)
str(zone0_infected)

#calculate first exposed based on sigma (rate exposed to infected=0.2) and first infections
first_exposed <- first_infections / sigma_vec
first_exposed

#Update the SEIR matrix with first infections and first exposed
#remove first infections and exposed from susceptible population
SEIR[, "S"] <- SEIR[, "S"] - (first_infections + first_exposed)
#add first infections to infected population
SEIR[, "I"] <- SEIR[, "I"] + first_infections
#add first exposed to exposed population
SEIR[, "E"] <- SEIR[, "E"] + first_exposed - first_infections
```

```
#Define OD to use
OD <- aml_all
```

Mobility scenario to use (origin/destination matrix)

```
# New E
infected_mat <- replicate(locs_len, SEIR_nsim[, "I"])
OD_all_infected <- round(aml_all*infected_mat)
flow_all_infected <- colsums(OD_all_infected)
total_flow_all_infected <- sum(flow_all_infected)
print(paste0("Total infected full flow: ", total_flow_all_infected))
new_exposed <-
  beta_vec * SEIR_sim[, "S"] * flow_all_infected / (N_k + colsums(aml_all)) + #exposed by contacting with imported infectious cases
  beta_vec * SEIR_sim[, "S"] * SEIR_sim[, "I"] / N_k #exposed by contacting with local infected cases
new_exposed[is.na(new_exposed)] <- 0
total_new_exposed <- round(sum(new_exposed))
print(paste0("New exposed: ", total_new_exposed))
new_exposed <- ifelse(new_exposed > SEIR_sim[, "S"], SEIR_sim[, "S"], new_exposed) #make sure the N exposed is not bigger than total susceptible
```

Calculation of new exposed based on OD matrix and
nr. infected (within and outside municipality)

```
# New I
new_infected <- sigma_vec * SEIR_sim[, "E"]
total_new_infected <- round(sum(new_infected, na.rm = T))
print(paste0("New infected: ", total_new_infected))
```

Calculation of new infected based on new exposed


```
# New R
new_recovered <- gamma_vec * SEIR_sim[, "I"]
total_new_recovered <- round(sum(new_recovered, na.rm = T))
print(paste0("New recovered: ", total_new_recovered))
```

Calculation of new recovered based on new infected


```
SEIR_sim[, "S"] <- SEIR_sim[, "S"] - new_exposed
SEIR_sim[, "E"] <- SEIR_sim[, "E"] + new_exposed - new_infected
SEIR_sim[, "I"] <- SEIR_sim[, "I"] + new_infected - new_recovered
SEIR_sim[, "R"] <- SEIR_sim[, "R"] + new_recovered
SEIR_sim <- ifelse(SEIR_sim < 0, 0, SEIR_sim)
```

Update of SEIR matrix, every day

New infections in LMA





New infections in PMA



Stricter lockdown periods (vertical dashed lines) cover the periods between **23rd March and 4th May 2020**, **28th November to 8th December 2020**, and from **27th December 2020 until 31st March 2021**

- ✓ Daily new cases - Estimations vs official records (23 March 2020 to 26 October 2020)
- ✓ 2 scenarios: Unrestricted mobility & Restricted to Essential activities




Estimated values considering the upper and lower thresholds of the confidence interval (95%) of R_t (dashed lines).


Official recorded cases of new infections are represented as points, daily values until the 6th July, and cumulative weekly values afterwards.

Lisbon (LMA)





Porto (PMA)



A) Unrestricted mobility LMA





A) Unrestricted mobility PMA



Lisbon (LMA)

Porto (PMA)



- ✓ The spread of the virus has responded to the mobility restrictions implemented. The number of new infections decreases after lockdown periods, with a certain delay
- ✓ Timeline of virus spread differs among metropolitan areas. PMA, closer to the scenario of mobility restricted to essential activities, in LMA closer to the unrestricted mobility scenario (lower threshold). Linked to secondary outbreaks in July 2020, especially in LMA, likely due to the ease of restrictions
- ✓ After Christmas, there was a substantial increase in the number of infected cases and fatalities, and the capacity of the health system was challenged. New infections started to decrease in mid-January 2021, following the implementation of mobility and social restrictions. Trends are captured by the simulations.
- ✓ In LMA, Lisbon absorbs most of the flows regardless of the scenario considered, higher number of cases. In PMA, Porto and VNGaia share the highest number of cases, depending on the scenario.
- ✓ **Further work:** How these restrictions can be adjusted to define suitable reopening strategies;
how other social distancing and hygiene measures can be of complement;
how the mobility conditions in metropolitan areas can be improved beyond the pandemic.

6-8 JULY 2021 | VIRTUAL



**17th International
Conference on
Urban Health**

Transforming our Collective Urban
Future: Learning from Covid-19



ISUH

**International Society
for Urban Health**



16:30 - 17:30 EDT

[O10] Pandemic lock-
downs in the city:
capturing the effects
of mobility restrictions

Thank you!



UNIVERSIDADE
DE LISBOA



Instituto de Geografia
e Ordenamento do Território
UNIVERSIDADE DE LISBOA



CEG
Centro de Estudos Geográficos



TERRA
Laboratory for sustainable
land use and ecosystem services



FCT

Fundação para a Ciência e a Tecnologia
Instituto de Saúde Pública e Ambiente



REPÚBLICA
PORTUGUESA

