

Levonorgestrel intrauterine device as a non-invasive approach of abnormal uterine bleeding caused by cesarean scar defect

Roxana-Elena Bohiltea¹, Bianca-Margareta Mihai², Valentin Varlas^{1,2}, Vlad Dima²,
Tiberiu Augustin Georgescu^{3,4}, Emilia-Maria Vladareanu⁵, Nicolae Bacalbasa¹, Corina Grigoriu^{1,6}

¹Department of Obstetrics and Gynecology, "Carol Davila" University of Medicine and Pharmacy, Bucharest, Romania

²Filantropia Clinical Hospital, Bucharest, Romania

³Department of Pathology, "Carol Davila" University of Medicine and Pharmacy, Bucharest, Romania

⁴"Alessandrescu-Rusescu" National Institute for Mother and Child Health, Bucharest, Bucharest

⁵Faculty of Medicine, "Carol Davila" University of Medicine and Pharmacy, Bucharest, Romania

⁶Department of Obstetrics and Gynecology, University Emergency Hospital, Bucharest, Romania

ABSTRACT

Cesarean scar defect, niche or isthmocele represents a poor healing in the anterior uterine wall after performing a cesarean section. The cesarean scar defect can be asymptomatic, or the patient could present abnormal uterine bleeding, chronic pelvic pain, dysmenorrhea, dyspareunia, cesarean scar pregnancy or abnormal placenta. Abnormal uterine bleeding caused by cesarean scar defect presents as a postmenstrual spotting and has become more and more common among women with a history of minimum one cesarean section delivery. The most studied risk factors are: multiple cesarean section deliveries, single layer suture, locked suture, retroflexed uterus and cesarean section delivery performed during active labor with a cervical dilatation of 5 cm. There have been described several surgical approaches: hysteroscopic, laparoscopic or vaginal. From our experience, we have treated successfully symptomatic patients with cesarean scar defect with the levonorgestrel-releasing intrauterine system. Although the therapeutic indications do not include this specific use, we have obtained significant improvement of abnormal uterine bleeding due to cesarean scar defect in our patients. Our results sustain the necessity of extensive interventional studies.

Keywords: cesarean scar defect, isthmocele, levonorgestrel, postmenstrual spotting, niche, abnormal uterine bleeding

INTRODUCTION

The number of newborns delivered by cesarean section (CS) has risen in the last years, currently being performed approximately 1.5 million CS every year around the world (1). Thus, the morbidity caused by the poor healing of the cesarean scar affects more and more women and presents in a wide range of symptoms: from asymptomatic to chronic pelvic pain, dyspareunia, dysmenorrhea, postmenstrual spotting, postmenstrual bleeding or even infertility (2-4). Regarding the obstetric complications,

there have been documented cesarean scar ectopic pregnancy, abnormal placenta (accrete and praevia) and scar dehiscence (5,6). The cesarean scar defect is described as a dome located in the anterior uterine wall after a cesarean section, which occurs as a result of an impaired healing of the uterine scar after CS (Fig. 1). The cesarean scar defect is also known as isthmocele, pouch, sacculation, uterine scar defect, cesarean scar dehiscence or uterine diverticulum niche (7-10). The isthmocele presents an incidence between 24-84% in women with previous cesarean section; 30% of women presenting a cesar-

Corresponding author:
Bianca-Margareta Mihai
E-mail: bmmihai@gmail.com

Article History:
Received: 7 November 2021
Accepted: 15 November 2021



FIGURE 1. Cesarean scar defect during pregnancy: amniotic diverticulum protruding into the uterine wall

ean scar defect suffer from abnormal uterine bleeding 6 months succeeding the cesarean section (11,12).

The diagnosis of a cesarean scar defect is frequently realized using transvaginal sonography or sonohysterography in nonpregnant patients and is described as 1 mm depth wedge-shaped anechogenic area with a minimum 2 mm indentation of the myometrium in the uterine isthmus located at the cesarean section scar site. Transvaginal ultrasonography is very valuable in determine the depth and size measurement of the dehiscent scar (Fig. 2) as well as the residual myometrium thickness (13-15). Other useful tools in diagnosing the uterine niche are represented by magnetic resonance imaging, hysterosalpingography and hysteroscopy (16,17). Concerning the classification of the cesarean scar defect size, it has been characterized by the ratio of the myometrial thickness at the scar (residual myometrium thickness) to the adjacent myometrium on transvaginal ultrasonography; the ratio $\leq 50\%$ has been used to describe severe deficiency (14). Another study has described a large defect if the residual myometrium thickness is 2.2 mm (18). Dehiscence has been described by Regnard et al. (19) on saline infusion sonohysterography as at least 80% of myometrial thinning.




FIGURE 2. Evaluation of residual and adjacent myometrium thickness (AMT = adjacent myometrium thickness; RMT = residual myometrium thickness)

In regard to risk factors for cesarean scar defect, many studies have concluded that multiple cesarean deliveries represent the essential risk factor for the uterine scar defect development (14,20). In addition, cesarean section performed during active labor with over 5 cm cervical dilatation has been correlated with a larger niche, that could be caused by the thin myometrium and defective healing capacity (11,21). Single layer myometrial closure compared to double layer has a tendency to increase the prevalence of isthmocele, but it was not associated to a larger defect (21). Roberge et al. (22) compared three techniques of uterine closure after a cesarean section: locked single layer, double layer with unlocked first layer and double layer with locked first layer; the conclusion was that double layer with unlocked first layer technique was correlated with enhanced uterine scar healing and firmer residual myometrium thickness. Locked sutures are linked to thinner myometrium and extended scar defects compared to unlocked sutures possibly due to elevated tissue hypoxia damaging the healing mechanism. Certain studies (14,20,21) have associated the retroflexed uterus with a defective uterine scar that can be explained by the mechanical tension intruding the uterine scar that diminishes blood perfusion and interferes with tissues regeneration.

ABNORMAL UTERINE BLEEDING

The correspondence between abnormal uterine bleeding and a patient's history of cesarean section has been reported and well investigated; an isthmocele is the most likely cause of typically postmenstrual spotting (10,23). Postmenstrual spotting presents as a persistent light vaginal bleeding after the menses has finished and has duration from 2 to 12 days (3). Fabres et al. (16) have conducted a retrospective study regarding sonohysterographic diagnosis of cesarean scar defect: 76% of women presented postmenstrual bleeding, 16% presented mid-cycle abnormal uterine bleeding, and 8% presented both. Uppal et al. (23) have concluded that the larger the uterine scar defect was, it was more likely for women to present prolonged menses or postmenstrual bleeding. The suggested mechanism of the abnormal uterine bleeding associated with cesarean scar defects is the delayed menstrual bleeding due to accumulation of blood in the lower uterine segment, in the isthmocele, which acts as a pouch (10,16,23). Due to defective drainage of normal menstrual blood, decreased contractility and in situ production, it has been implied the fact that blood accumulates in this pouch and slowly drains in the next days after the menses (3,16).

On the other hand, in some patients with asymptomatic cesarean scar defect, the accumulation of




FIGURE 3. Levonorgestrel-releasing intrauterine system insert in uterine cavity with niche

blood within the isthmocoele may not cause abnormal uterine bleeding, but the presence of isthmocoeles may support other determinants of abnormal uterine bleeding like chronic inflammation (16,23).

CURRENT APPROACHES

The management of abnormal uterine bleeding due to cesarean scar defect varies between hormonal treatment and hysterectomy. Some studies reported deficiency in the results of hormonal therapy regarding abnormal uterine bleeding caused by the niche and the patients required surgical therapy (3,24,25). Hysteroscopic repair is the most used approach for the cesarean scar defect revision and the results are promising, improving abnormal uterine bleeding and fertility (24,26,27). The hysteroscopic treatment consists in the scar resection surrounding the defect (27). The hysteroscopic approach imports the risk of uterine perforation with consecutive bladder injury; thus, it has been recommended to perform a hysteroscopic resection if the overlying myometrium is thicker than 2 mm (24). Another surgical approach is by using laparoscopy, either conventional or robotic-assisted, which consists in

fibrotic tissue resection, followed by the laparoscopically closing of the defect with the help of a Hegar dilator placed in the cervix to assure continuity maintenance between the uterine cavity and cervical canal (28). Finally, the cesarean scar defect may be repaired using a transvaginal approach. The technique requires the dissecting of the bladder off the cervix and the uterus in order to open the vesicovaginal space until the cesarean scar defect could be recognized. Further, the scar tissue is excised and the defect is closed using interrupted sutures (29).

Abnormal uterine bleeding is becoming more and more frequent in the current practice, as the number of cesarean sections increases worldwide. There are various methods of surgical treatment of the cesarean scar defect, which present benefits, along with the risks of any invasive procedure, for example the risk of uterine perforation that comes along with the hysteroscopic repair.

We propose the medical treatment of the cesarean scar defect symptomatology, the use of levonorgestrel-releasing intrauterine system, as we have had very good results in managing the abnormal uterine bleeding due to cesarean scar defect (Fig. 3).

CONCLUSIONS

After excluding other causes of abnormal uterine bleeding, our patients have presented a significant decrease and even disappearance of postmenstrual bleeding as well as intermenstrual bleeding caused by the uterine niche, starting with the second month after correctly placing the levonorgestrel intrauterine device. Although in the device's therapeutic indications the producer does not specify this intent to treat use, our patients have had only benefits after this hormonal therapy. The primary benefit for the patient is the avoidance of an invasive surgery such as laparoscopic or vaginal isthmoplasty or invasive procedure including hysteroscopic repair of the defect.

Conflict of interest: none declared
Financial support: none declared

REFERENCES

1. Guise JM, Eden K, Emeis C, Denman MA, Marshall N, Fu RR, Janik R, Nygren P, Walker M, McDonagh M. Vaginal birth after cesarean: new insights. *Evid Rep Technol Assess* (Full Rep). 2010 Mar;(191):1-397.
2. Bij de Vaate AJ, Brölmann HA, van der Voet LF, van der Slikke JW, Veersema S, Huirne JA. Ultrasound evaluation of the Cesarean scar: relation between a niche and postmenstrual spotting. *Ultrasound Obstet Gynecol*. 2011 Jan;37(1):93-9.
3. Thurmond AS, Harvey WJ, Smith SA. Cesarean section scar as a cause of abnormal vaginal bleeding: diagnosis by sonohysterography. *J Ultrasound Med*. 1999 Jan;18(1):13-8.
4. Toader O, Fetecau AC, Suci N, Voichițoiu A, Cîrstoiu M, Bohlîța R, Firta AR. Cesarean Scar Defects - an Underdiagnosed Cause for Abnormal Uterine Bleeding and Other Gynecologic Complications. *Filodiritto Editore-Proceedings. 5th Romanian Congress Of The Romanian Society Of Ultrasound In Obstetrics And Gynecology 2017*:660-663.
5. Francois K, Foley, M. Antepartum and postpartum hemorrhage. *Obstetrics: Normal and Problem Pregnancies*. PA: Churchill Livingstone Elsevier, 2007:456-485.
6. Landon M. Cesarean delivery. *Obstetrics: Normal and Problem Pregnancies*. PA: Churchill Livingstone Elsevier, 2008:486-520.

7. Chen Y, Chang Y, Yao S. Transvaginal management of cesarean scar section diverticulum: a novel surgical treatment. *Med Sci Monit.* 2014 Aug;20:1395-9.
8. Feng YL, Li MX, Liang XQ, Li XM. Hysteroscopic treatment of postcesarean scar defect. *J Minim Invasive Gynecol.* 2012 Jul-Aug; 19(4):498-502.
9. Cheng XY, Cheng L, Li WJ, Qian LH, Zhang YQ. The effect of surgery on subsequent pregnancy outcomes among patients with cesarean scar diverticulum. *Int J Gynaecol Obstet.* 2018 May;141(2):212-216.
10. Borges LM, Scapinelli A, de Baptista Depes D, Lippi UG, Coelho Lopes RG. Findings in patients with postmenstrual spotting with prior cesarean section. *J Minim Invasive Gynecol.* 2010 May-Jun;17(3):361-4.
11. Bij de Vaate AJ, van der Voet LF, Naji O, Witmer M, Veersema S, Brölmann HA, Bourne T, Huirne JA. Prevalence, potential risk factors for development and symptoms related to the presence of uterine niches following Cesarean section: systematic review. *Ultrasound Obstet Gynecol.* 2014 Apr;43(4):372-82.
12. Vervoort AJ, Uittenbogaard LB, Hehenkamp WJ, Brölmann HA, Mol BW, Huirne JA. Why do niches develop in Caesarean uterine scars? Hypotheses on the aetiology of niche development. *Hum Reprod.* 2015 Dec;30(12):2695-702.
13. Tulandi T, Cohen A. Emerging Manifestations of Cesarean Scar Defect in Reproductive-aged Women. *J Minim Invasive Gynecol.* 2016 Sep-Oct; 23(6):893-902.
14. Ofili-Yebovi D, Ben-Nagi J, Sawyer E, Yazbek J, Lee C, Gonzalez J, Jurkovic D. Deficient lower-segment Cesarean section scars: prevalence and risk factors. *Ultrasound Obstet Gynecol.* 2008 Jan;31(1):72-7.
15. Bohîlțea RE, Turcan G, Cîrstoiu MM, Ionescu C, Nemescu D, Turcan N, Vladareanu R. Clinical Implementation of Ultrasound Gynecological Examination Report (software REGU) Based on International Consensus of Tumor Study Groups. Filodiritto Editore-Proceedings. 5th Romanian Congress Of The Romanian Society Of Ultrasound In Obstetrics And Gynecology, 2017:99-104.
16. Fabres C, Aviles G, De La Jara C, Escalona J, Muñoz JF, Mackenna A, Fernández C, Zegers-Hochschild F, Fernández E. The cesarean delivery scar pouch: clinical implications and diagnostic correlation between transvaginal sonography and hysteroscopy. *J Ultrasound Med.* 2003 Jul;22(7):695-702.
17. Tower AM, Frishman GN. Cesarean scar defects: an underrecognized cause of abnormal uterine bleeding and other gynecologic complications. *J Minim Invasive Gynecol.* 2013 Sep-Oct;20(5):562-72.
18. Osser OV, Jokubkiene L, Valentin L. High prevalence of defects in Cesarean section scars at transvaginal ultrasound examination. *Ultrasound Obstet Gynecol.* 2009 Jul;34(1):90-7.
19. Regnard C, Nosbusch M, Fellemans C, Benali N, van Rysselberghe M, Barlow P, Rozenberg S. Cesarean section scar evaluation by saline contrast sonohysterography. *Ultrasound Obstet Gynecol.* 2004 Mar;23(3):289-92.
20. Wang CB, Chiu WW, Lee CY, Sun YL, Lin YH, Tseng CJ. Cesarean scar defect: correlation between Cesarean section number, defect size, clinical symptoms and uterine position. *Ultrasound Obstet Gynecol.* 2009 Jul;34(1):85-9.
21. Vikhareva Osser O, Valentin L. Risk factors for incomplete healing of the uterine incision after caesarean section. *BJOG.* 2010 Aug; 117(9):1119-26.
22. Roberge S, Demers S, Berghella V, Chaillet N, Moore L, Bujold E. Impact of single- vs double-layer closure on adverse outcomes and uterine scar defect: a systematic review and metaanalysis. *Am J Obstet Gynecol.* 2014 Nov;211(5):453-60.
23. Uppal T, Lanzarone V, Mongelli M. Sonographically detected caesarean section scar defects and menstrual irregularity. *J Obstet Gynaecol.* 2011 Jul;31(5):413-6.
24. Chang Y, Tsai EM, Long CY, Lee CL, Kay N. Resectoscopic treatment combined with sonohysterographic evaluation of women with postmenstrual bleeding as a result of previous cesarean delivery scar defects. *Am J Obstet Gynecol.* 2009 Apr;200(4):370.e1-4.
25. Morris H. Surgical pathology of the lower uterine segment caesarean section scar: is the scar a source of clinical symptoms? *Int J Gynecol Pathol.* 1995 Jan;14(1):16-20.
26. Fabres C, Arriagada P, Fernández C, Mackenna A, Zegers F, Fernández E. Surgical treatment and follow-up of women with intermenstrual bleeding due to cesarean section scar defect. *J Minim Invasive Gynecol.* 2005 Jan-Feb;12(1):25-8.
27. Shih CL, Chang YY, Ho M, Lin WC, Wang AM, Lin WC. Hysteroscopic transcervical resection. A straightforward method corrects bleeding related to cesarean section scar defects. *Am J Obstet Gynecol.* 2011 Mar;204(3):278.e1-2.
28. Donnez O, Jadoul P, Squifflet J, Donnez J. Laparoscopic repair of wide and deep uterine scar dehiscence after cesarean section. *Fertil Steril.* 2008 Apr;89(4):974-80.
29. Khoshnow Q, Pardey J, Uppal T. Transvaginal repair of caesarean scar dehiscence. *Aust N Z J Obstet Gynaecol.* 2010 Feb;50(1):94-5.