



Title	Relationship between adrenal steroid hormones in cord blood and birth weight : The Sapporo Cohort, Hokkaido Study on Environment and Children's Health
Author(s)	Mitsui, Takahiko; Araki, Atsuko; Goudarzi, Houman; Miyashita, Chihiro; Ito, Sachiko; Sasaki, Seiko; Kitta, Takeya; Moriya, Kimihiko; Cho, Kazutoshi; Morioka, Keita; Kishi, Reiko; Shinohara, Nobuo; Takeda, Masayuki; Nonomura, Katsuya
Citation	American Journal of Human Biology, 30(4), e23127 https://doi.org/10.1002/ajhb.23127
Issue Date	2018-07
Doc URL	http://hdl.handle.net/2115/87338
Rights	This is the peer reviewed version of the following article: Mitsui, T, Araki, A, Goudarzi, H, et al. Relationship between adrenal steroid hormones in cord blood and birth weight: The Sapporo Cohort, Hokkaido Study on Environment and Children's Health. Am J Hum Biol. 2018; 30:e23127, which has been published in final form at https://doi.org/10.1002/ajhb.23127 . This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Use of Self-Archived Versions.
Type	article (author version)
File Information	31_A J Human Biology, 2018.pdf



[Instructions for use](#)

Relationship between adrenal steroid hormones in cord blood and birth weight: The Sapporo Cohort, Hokkaido Study on Environment and Children's Health

Takahiko Mitsui^{1),2)}, Atsuko Araki³⁾, Houman Goudarzi^{3),4)}, Chihiro Miyashita³⁾,
Sachiko Ito³⁾, Seiko Sasaki⁵⁾, Takeya Kitta²⁾, Kimihiko Moriya²⁾, Kazutoshi Cho⁶⁾,
Keita Morioka⁶⁾, Reiko Kishi³⁾, Nobuo Shinohara²⁾, Masayuki Takeda¹⁾,
Katsuya Nonomura⁷⁾

- 1) Department of Urology, University of Yamanashi Graduate School of Medical Sciences
- 2) Department of Urology, Hokkaido University Graduate School of Medicine
- 3) Division of Respiratory Medicine, Graduate School of Medicine, Hokkaido University
- 4) Hokkaido University Center for Environmental and Health Sciences
- 5) Department of Public Health, Hokkaido University Graduate School of Medicine
- 6) Department of Obstetrics and Gynecology, Hokkaido University Graduate School of Medicine
- 7) Department of Urology, Kushiro Rosai Hospital

Acknowledgments

We thank the mothers and their children who participated in this study and the staff at Sapporo Toho Hospital. This work was financially supported, in part, by the Ministry of Health, Labour and Welfare, Health and Labour Sciences Research Grants, Grants-in-Aid for Scientific Research from the Japan Society for the Promotion of Science, and the Environment Research and Technology Development Fund (5C-1252) from the Ministry of the Environment, Japan.

Running title: prenatal hormones and birth weight

Word Count: 1495 words, 12 pages of manuscript , Tables: 2

Keywords: Steroid; Birth weight; Prenatal; Development; Sexual dimorphism;

Corresponding Author:

Takahiko Mitsui, MD

Address: 1110 Shimokato, Chuo-city, 409-3898, Japan

Phone: +81-55-273-9643 Fax: +81-55-273-9659

E-mail: tmitsui@yamanashi.ac.jp

Abstract

Objectives: We investigated the relationship between steroid hormone levels in cord blood and birth weight.

Methods: Among 514 participants in a prospective birth cohort study in Sapporo, the following hormone levels were measured in 294 stored cord blood samples from 135 males and 159 females: androstenedione, dehydroepiandrosterone (DHEA), cortisol, and cortisone. Birth weight information was obtained from medical records.

Results: androstenedione/DHEA was significantly higher in males than in females, while DHEA was significantly higher in females. Birth weight was significantly higher in males than in females. Regarding cortisone, androstenedione/DHEA, and cortisone/cortisol, a correlation was observed with birth weight in males but not in females.

Conclusions: Prenatal adrenal steroids as well as converting enzymes such as 11 β -hydrosteroid dehydrogenase type 2 and 3 β -hydrosteroid dehydrogenase may have an impact on prenatal physical development.

Introduction

The prenatal environment affects infant development. Adaptive changes to an adverse environment *in utero*, which results in low birth weight, have the potential to result in permanent changes in physiology, structure, and metabolism, which are collectively termed fetal 'programming' (Moisiadis and Matthews, 2014a; Moisiadis and Matthews, 2014b; O'Connor and Barrett, 2014) and are considered independent factors contributing to adult outcomes.

Glucocorticoids are an important factor in physiological functions, and the hypothalamic-pituitary-adrenal axis is a major system involved in the stress response and its regulation. Maternal cortisol passes through the placenta and affects both fetal cortisol levels and hypothalamic-pituitary-adrenal development, and is known as a glucocorticoid signal between the mother, placenta, and fetus (Moisiadis and Matthews, 2014a; Moisiadis and Matthews, 2014b; O'Connor and Barrett, 2014). Previous studies have reported that glucocorticoid levels in maternal blood and amniotic fluid correlate with birth weight (Baibazarova, et al., 2013; Field, et al., 2006). However, few studies have investigated the relationship between birth weight and glucocorticoid levels in cord blood which directly reflect the infant's hormonal environment at birth. In this study, we examined whether adrenal steroid hormone levels in cord blood were associated with infant birth weight.

Participants and Methods

Participants

This prospective birth cohort study was based on the Sapporo Cohort, Hokkaido Study on Environment and Children's Health (Kishi, et al., 2013). Details regarding the population, data collection, sampling of biological specimens, and contents of the questionnaire have been described previously (Kishi, et al., 2013). Among 514 pregnant women who were enrolled in this cohort study, we excluded women who had miscarriages, stillbirths, relocated, voluntarily withdrew ($n = 11$), and those who delivered twins ($n = 6$). Finally, 294 participants, who had data of hormones in cord blood and birth weight, were enrolled in the present study.

The current study was approved by the Institutional Ethical Board for Epidemiological Studies at Hokkaido University Graduate School of Medicine and Hokkaido University Center for Environmental and Health Sciences. All participants signed informed consent, and informed consent was provided by parents on behalf of the children enrolled.

Hormone measurements in cord blood samples

At the time of delivery, a 10-30 mL blood sample was collected from the umbilical cord and stored at -80°C for subsequent analysis. On 294 stored cord blood samples (135 males and 159 females), the following steroid hormones were measured using highly specific liquid chromatography-tandem mass spectrometry: cortisol, cortisone, androstenedione, and dehydroepiandrosterone (DHEA) (Yamashita, et al., 2007). Ratio of androstenedione/DHEA and cortisol/cortisone were also calculated for parameters of

11 β -hydrosteroid dehydrogenase type 2 (11 β -HSD2) and 3 β -hydrosteroid dehydrogenase (3 β -HSD) activities in the placenta and fetal adrenal gland, respectively.

All hormone measurements were performed by Aska Pharma Medical Co., Ltd. (Kanagawa, Japan). The detection limits for each hormone were as follows: cortisol, 0.250 ng/mL; cortisone, 0.100 ng/mL; androstenedione, 0.010 ng/mL; and DHEA, 0.010 ng/mL.

Statistical analyses

Participant characteristics and hormone values were analyzed using one-way ANOVA or Chi-square. Hormone values were converted to a log₁₀ scale because the data were not normally distributed. Half the detection limit listed above was used when levels were below the detection limit for individual hormones. The relationships between birth weight and steroid hormone levels in cord blood samples were calculated using multiple linear regression analyses. Inclusion of covariates was based on biological considerations, and adjustments were made for gestational age (continuous), maternal smoking and alcohol consumption during pregnancy (yes or no), annual household income (<5 or \geq 5 million yen/year), educational level (\leq 12 or \geq 13 years), and parents' body weights (continuous). Statistical analyses were performed using JMP pro 10 (SAS institute Inc., NC, USA). Significance levels were set to 0.05 for all comparisons.

Results

1) Characteristics of participants and their children

The characteristics of parents and their children were shown in Table 1. Birth weights of children ranged from 1784 g to 4162 g (median: 3096 g). Only 4 infants were born with low birth weight (<2500 g). Birth weights were significantly higher in males (mean \pm standard deviation: 3139.3 \pm 28.1 g) than in females (3072.0 \pm 25.9 g) ($p=0.002$). Maternal pre-pregnancy weight was significantly associated with female birth weight ($p=0.2685$, $p<0.001$) but not that of males ($p= 0.0977$, $p= 0.128$). Paternal weight was not associated with the birth weights of males ($p= 0.0310$, $p= 0.631$) or females ($p= 0.0400$, $p= 0.525$).

2) Sex hormones in cord blood samples

Median concentration of androstenedione/DHEA was significantly higher in males than in females, whereas that of DHEA was significantly higher in females. Other hormones were not different between males and females.

3) Relationship between birth weight and sex hormones

A multivariate regression model showed that birth weight was positively correlated with cortisone, cortisone/cortisol, and androstenedione/DHEA in males only. No significant correlations were observed between birth weight and any other hormones, in males or females (Table 2).

Discussion

The results of our investigation suggest that prenatal adrenal steroids are associated with birth weight in males but not in females. Furthermore, androstenedione/DHEA and cortisol/cortisone, which are indicative of converting enzymes in the placenta and fetal adrenal gland, also correlated with birth weight in males only.

Supraphysiologic levels of glucocorticoids may induce fetal growth retardation. High glucocorticoid concentrations in both maternal and fetal human studies have been associated with intrauterine growth retardation (Field, et al., 2006; McTernan, et al., 2001). Elevated concentrations of glucocorticoids have been found to be induced by stress, maternal malnutrition, and placental insufficiency due to restriction of placental blood flow. On the other hand, the fetus is generally protected from the effects of maternal glucocorticoids by 11β -HSD2 in the placenta. This enzyme rapidly catalyzes cortisol (in the form of active glucocorticoids) to cortisone, a physiologically inactive glucocorticoid. In an animal study using rats, maternal stress during gestation induced decreased fetal weight and placental 11β -HSD2 activity (Mairesse, et al., 2007). Thus, 11β -HSD2 functions to prevent the transport of maternal glucocorticoids to the fetus (Benediktsson and Seckl, 1998). Regarding 3β -HSD, there is no evidence for the relationship of 3β -HSD and fetal growth during the prenatal period, from either human epidemiological or animal studies. Although we speculate that some pathways in the steroidogenesis of progesterone and 3β -HSD in the placenta and fetus may be involved in fetal growth, this has not yet been demonstrated.

The sexually dimorphic effects of glucocorticoids on human fetal growth have been reported in previous studies. Ellman et al. (2008) reported that an increase in cortisol

levels in mothers during early pregnancy was correlated with physical maturity in newborn males only (Ellman, et al., 2008). Further, difficulty in adapting placental function to an adverse maternal environment may also induce sexual dimorphism in fetal growth (Clifton, 2010). Therefore, the effects of glucocorticoids during gestation may induce sexual dimorphism in fetal development, and glucocorticoid metabolic activity is less adaptive to high maternal glucocorticoid concentrations in the male placenta. However, the mechanisms underlying the sexually dimorphic effects of glucocorticoids on fetal growth remain unclear, and thus further studies are necessary.

There are some limitations to this study. First, the correlation between birth weight and prenatal hormone levels was relatively small, indicating the possibility of marked individual differences in human biological variation at birth. Second, the cord blood samples were collected throughout the day and night due to unpredictable deliveries. Therefore, the diurnal rhythms of circulating glucocorticoids may have affected these data. Third, stress from the delivery itself could have affected levels of steroid hormones in the cord blood (Keelan, et al., 2012). However, we did not investigate this factor, since there are no data establishing the impact of stress on steroid hormone levels during delivery. Fourth, the effects of hormone exposure during critical gestational periods on fetal growth are still unknown because we analyzed hormonal exposure using cord blood at delivery. Therefore, further studies are required to reveal the effects of steroid hormone levels *in utero* on physical changes in children.

Conclusions

Prenatal adrenal steroid exposure during the prenatal period has the potential to affect fetal growth. Converting enzymes, such as 11 β -HSD2 and 3 β -HSD in the placenta and fetal adrenal gland, respectively, may also impact birth weight. Since these relationships were observed in males only, sensitivity to steroid hormones and the activities of enzymes in steroidogenesis may induce sexually dimorphic effects of glucocorticoids on fetal growth.

Acknowledgments

We thank the mothers and their children who participated in this study and the staff at Sapporo Toho Hospital. This work was financially supported, in part, by the Ministry of Health, Labour and Welfare, Health and Labour Sciences Research Grants, Grants-in-Aid for Scientific Research from the Japan Society for the Promotion of Science, and the Environment Research and Technology Development Fund (5C-1252) from the Ministry of the Environment, Japan.

Abbreviations

3 β -HSD: 3 β -hydrosteroid dehydrogenase

11 β -HSD2: 11 β -hydrosteroid dehydrogenase type 2

DHEA: dehydroepiandrosterone

HPA: hypothalamic-pituitary-adrenal

References

- Baibazarova E, van de Beek C, Cohen-Kettenis PT, Buitelaar J, Shelton KH, van Goozen SH. 2013. Influence of prenatal maternal stress, maternal plasma cortisol and cortisol in the amniotic fluid on birth outcomes and child temperament at 3 months. *Psychoneuroendocrinology* 38(6):907-915.
- Benediktsson R, Seckl JR. 1998. Understanding human parturition. *Lancet* 351(9106):913-914.
- Clifton VL. 2010. Review: Sex and the human placenta: mediating differential strategies of fetal growth and survival. *Placenta* 31 Suppl:S33-39.
- Ellman LM, Schetter CD, Hobel CJ, Chicz-Demet A, Glynn LM, Sandman CA. 2008. Timing of fetal exposure to stress hormones: effects on newborn physical and neuromuscular maturation. *Dev Psychobiol* 50(3):232-241.
- Field T, Hernandez-Reif M, Diego M, Figueiredo B, Schanberg S, Kuhn C. 2006. Prenatal cortisol, prematurity and low birthweight. *Infant Behav Dev* 29(2):268-275.
- Keelan JA, Mattes E, Tan H, Dinan A, Newnham JP, Whitehouse AJ, Jacoby P, Hickey M. 2012. Androgen concentrations in umbilical cord blood and their association with maternal, fetal and obstetric factors. *PLoS One* 7(8):e42827.
- Kishi R, Kobayashi S, Ikeno T, Araki A, Miyashita C, Itoh S, Sasaki S, Okada E, Kashino I, Itoh K and others. 2013. Ten years of progress in the Hokkaido birth cohort study on environment and children's health: cohort profile--updated 2013. *Environ Health Prev Med* 18(6):429-450.
- Mairesse J, Lesage J, Breton C, Breant B, Hahn T, Darnaudery M, Dickson SL, Seckl J, Blondeau B, Vieau D and others. 2007. Maternal stress alters endocrine function of the fetoplacental unit in rats. *Am J Physiol Endocrinol Metab* 292(6):E1526-1533.
- McTernan CL, Draper N, Nicholson H, Chalder SM, Driver P, Hewison M, Kilby MD, Stewart PM. 2001. Reduced placental 11beta-hydroxysteroid dehydrogenase type 2 mRNA levels in human pregnancies complicated by intrauterine growth restriction: an analysis of possible mechanisms. *J Clin Endocrinol Metab* 86(10):4979-4983.
- Moisiadis VG, Matthews SG. 2014a. Glucocorticoids and fetal programming part 1: Outcomes. *Nat Rev Endocrinol* 10(7):391-402.
- Moisiadis VG, Matthews SG. 2014b. Glucocorticoids and fetal programming part 2: Mechanisms. *Nat Rev Endocrinol* 10(7):403-411.
- O'Connor TG, Barrett ES. 2014. Mechanisms of prenatal programming: identifying and distinguishing the impact of steroid hormones. *Front Endocrinol (Lausanne)* 5:52.

Yamashita K, Takahashi M, Tsukamoto S, Numazawa M, Okuyama M, Honma S. 2007. Use of novel picolinoyl derivatization for simultaneous quantification of six corticosteroids by liquid chromatography-electrospray ionization tandem mass spectrometry. *J Chromatogr A* 1173(1-2):120-128.

Table 1 Characteristics of participants

The values in brackets represent percentages.

		Participants with data of hormones and birth weight
Characteristics of parents		
	n	Mean \pm SD
Age at delivery (years old)	294	30.4 \pm 5.0
Body weights of mothers at pre-pregnancy (kg)	294	52.6 \pm 7.8
Body weights of fathers (kg)	289	68.5 \pm 11.0
		n (%)
Parity	Primiparous	149 (50.7)
	Multiparous	145 (49.3)
Annual household income (million yen per year)	<5	206 (70.6)
	\geq 5	86 (29.5)
Educational level (years)	\leq 12	133 (45.2)
	\geq 13	161 (54.8)
Smoking during pregnancy	Non-smoker	238 (81.0)
	Smoker	56 (19.0)
Alcohol consumption during pregnancy	Non-drinker	193 (65.7)
	Drinker	101 (34.3)
Infant characteristics		
		n (%)
Gender	Males	135 (45.9)
	Females	159 (54.1)
	n	Mean \pm SD
Birth weight (g)	294	3126.8 \pm 331.6
Gestational age (weeks)	294	39.3 \pm 1.0

SD: standard deviation

Table 2 Relationship between birth weight and sex hormones in cord blood

*: $p < 0.05$

Hormone levels	Males		Females	
	B	R ²	B	R ²
	95% CI		95% CI	
Cortisol	0.1203	0.1014	0.0060	0.2372
	-57.534 / 163.72		-78.199 / 84.852	
Cortisone	0.1741*	0.1432	0.0211	0.2375
	4.240 / 183.06		-51.437 / 68.884	
Androstenedione	0.1266	0.1089	-0.1115	0.2415
	-76.522 / 403.87		-357.66 / 43.907	
DHEA	-0.099	0.1232	-0.0148	0.2373
	-345.17 / 91.470		-162.87 / 132.84	
Cortisone / Cortisol	0.2125*	0.1553	0.0407	0.2386
	43.255 / 383.83		-103.24 / 179.88	
Androstenedione / DHEA	0.1746*	0.1436	--0.0593	0.2406
	10.269 / 385.74		-176.53 / 74.033	

CI: confidence interval, DHEA: dehydroepiandrosterone.

Co-variates: gestational age, maternal smoking and alcohol consumption during pregnancy, annual household income, educational level, and body weights of parents.