

## Federation University ResearchOnline

<https://researchonline.federation.edu.au>

Copyright Notice

This is the post-peer-review, pre-copyedit version of an article published in Jonathan Borwein Commemorative Conference. The final authenticated version is available online at:

[https://doi.org/10.1007/978-3-030-36568-4\\_4](https://doi.org/10.1007/978-3-030-36568-4_4)

Copyright © Springer Nature Switzerland AG 2020

See this record in Federation ResearchOnline at:

<http://researchonline.federation.edu.au/vital/access/HandleResolver/1959.17/179024>

# Characterizations of robust and stable duality for linearly perturbed uncertain optimization problems

N. Dinh\*, M.A. Goberna<sup>†</sup>, M.A. Lopez<sup>‡</sup>, M. Volle<sup>§</sup>

October 11, 2018

**Abstract.** We introduce a robust optimization model consisting in a family of perturbation functions giving rise to certain pairs of dual optimization problems in which the dual variable depends on the uncertainty parameter. The interest of our approach is illustrated by some examples, including uncertain conic optimization and infinite optimization via discretization. The main results characterize desirable robust duality relations (as robust zero-duality gap) by formulas involving the epsilon-minima or the epsilon-subdifferentials of the objective function. The two extreme cases, namely, the usual perturbational duality (without uncertainty), and the duality for the supremum of functions (duality parameter vanishing) are analyzed in detail.

**Key words.** Robust duality, strong robust duality, reverse strong robust duality, min-max robust duality.

**2000 Mathematics Subject Classification.** 90C26, 90C46, 90C31

## 1 Introduction

In this paper we consider a *family of perturbation functions*

$$F_u : X \times Y_u \rightarrow \mathbb{R}_\infty := \mathbb{R} \cup \{+\infty\}, \text{ with } u \in U,$$

and where  $X$  and  $Y_u$ ,  $u \in U$ , are given locally convex Hausdorff topological vector spaces (briefly, lcHtvs), the index set  $U$  is called the *uncertainty set* of the family,  $X$  is its *decision space*, and each  $Y_u$  is a *parameter space*. Note that our model includes a parameter space  $Y_u$ , depending on  $u \in U$ , which is a novelty with respect to

---

\*International University, Vietnam National University - HCMC, Linh Trung ward, Thu Duc district, Ho Chi Minh city, Vietnam (ndinh@hcmiu.edu.vn).

<sup>†</sup>Department of Mathematics, University of Alicante, Spain (mgoberna@ua.es)

<sup>‡</sup>Department of Mathematics, University of Alicante, Spain, and CIAO, Federation University, Ballarat, Australia (marco.antonio@ua.es)

<sup>§</sup>Avignon University, LMA EA 2151, Avignon, France (michel.volle@univ-avignon.fr)

the ‘‘classical’’ robust duality scheme (see [17] and references therein, where a unique parameter space  $Y$  is considered), allowing us to cover a wider range of applications including uncertain optimization problems under linear perturbations of the objective function. The significance of our approach is illustrated along the paper by relevant cases extracted from deterministic optimization with linear perturbations, uncertain optimization without linear perturbations, uncertain conic optimization and infinite optimization. The antecedents of the paper are described in the paragraphs devoted to the first two cases in Section 2.

We associate with each family  $\{F_u : u \in U\}$  of perturbation functions corresponding optimization problems whose definitions involve continuous linear functionals on the decision and the parameter spaces. We denote by  $0_X$ ,  $0_X^*$ ,  $0_u$ , and  $0_u^*$ , the null vectors of  $X$ , its topological dual  $X^*$ ,  $Y_u$ , and its topological dual  $Y_u^*$ , respectively. The optimal value of a minimization (maximization, respectively) problem (P) is denoted by  $\inf$  (P) ( $\sup$  (P)); in particular, we write  $\min$  (P) ( $\max$  (P)) whenever the optimal value of (P) is attained. We adopt the usual convention that  $\inf$  (P) =  $+\infty$  ( $\sup$  (P) =  $-\infty$ ) when the problem (P) has no feasible solution. The associated optimization problems are the following:

- *Linearly perturbed uncertain problems:* for each  $(u, x^*) \in U \times X^*$ ,

$$(P_u)_{x^*} : \quad \inf_{x \in X} \{F_u(x, 0_u) - \langle x^*, x \rangle\}.$$

- *Robust counterpart of  $\{(P_u)_{x^*}\}_{u \in U}$ :*

$$(RP)_{x^*} : \quad \inf_{x \in X} \left\{ \sup_{u \in U} F_u(x, 0_u) - \langle x^*, x \rangle \right\}.$$

Denoting by  $F_u^* : X^* \times Y_u^* \rightarrow \overline{\mathbb{R}}$ , where  $\overline{\mathbb{R}} := \mathbb{R} \cup \{\pm\infty\}$ , the *Fenchel conjugate* of  $F_u$ , namely,

$$F_u^*(x^*, y_u^*) := \sup_{(x, y_u) \in X \times Y_u} \left\{ \langle x^*, x \rangle + \langle y_u^*, y_u \rangle - F_u(x, y_u) \right\}, \quad (x^*, y_u^*) \in X^* \times Y_u^*,$$

we now introduce the corresponding dual problems:

- *Perturbational dual of  $(P_u)_{x^*}$ :*

$$(D_u)_{x^*} : \quad \sup_{y_u^* \in Y_u^*} -F_u^*(x^*, y_u^*).$$

Obviously,

$$\sup (D_u)_{x^*} \leq \inf (P_u)_{x^*} \leq \inf (RP)_{x^*}, \forall u \in U.$$

- *Optimistic dual of  $(RP)_{x^*}$ :*

$$(ODP)_{x^*} : \quad \sup_{(u, y_u^*) \in \Delta} -F_u^*(x^*, y_u^*),$$

where  $\Delta := \{(u, y_u^*) : u \in U, y_u^* \in Y_u^*\}$  is the disjoint union of the spaces  $Y_u^*$ . We have

$$\sup (ODP)_{x^*} = \sup_{u \in U} (D_u)_{x^*} \leq \inf (RP)_{x^*}.$$

We are interested in the following desirable robust duality properties:

- *Robust duality* is said to hold at  $x^*$  if  $\inf (\text{RP})_{x^*} = \sup (\text{ODP})_{x^*}$ ,
- *Strong robust duality* at  $x^*$  means  $\inf (\text{RP})_{x^*} = \max (\text{ODP})_{x^*}$ ,
- *Reverse strong robust duality* at  $x^*$  means  $\min (\text{RP})_{x^*} = \sup (\text{ODP})_{x^*}$ ,
- *Min-max robust duality* at  $x^*$  means  $\min (\text{RP})_{x^*} = \max (\text{ODP})_{x^*}$ .

Each of the above desirable properties is said to be *stable* when it holds for any  $x^* \in X^*$ . The main results of this paper characterize these properties in terms of formulas involving the  $\varepsilon$ -minimizers and  $\varepsilon$ -subdifferentials of the objective function of the robust counterpart problem  $(\text{RP})_{0_X^*}$ , namely, the function

$$p := \sup_{u \in U} F_u(\cdot, 0_u).$$

Theorem 3.1 characterizes robust duality at a given point  $x^* \in X^*$  as a formula for the inverse mapping of the  $\varepsilon$ -subdifferential at  $x^*$  without any convexity assumption. The same is done in Theorem 4.1 to characterize strong robust duality. In the case when a primal optimal solution does exist we give a formula for the exact minimizers of  $p - x^*$  to characterize dual strong (resp. min-max) robust duality at  $x^*$ , see Theorem 5.1 (resp. Theorem 5.2). We show that stable robust duality gives rise to a formula for the  $\varepsilon$ -subdifferential of  $p$  (Theorem 6.1, see also Theorem 3.1). The same is done for stable strong robust duality (Theorem 7.1). A formula for the exact subdifferential of  $p$  is provided in relation with robust duality at appropriate points (Theorem 8.1). The most simple possible formula for the exact subdifferential of  $p$  (the so-called *Basic Robust Qualification condition*) is studied in detail in Theorem 8.2. All the results from Sections 1-8 are specified for the two extreme cases (the case with no uncertainty and the one in absence of linear perturbations), namely, Cases 1 and 2 in Section 2 (for the sake of brevity, we do not give the specifications for Cases 3 and 4). It is worth noticing the generality of the mentioned results (as they do not require any assumption on the involved functions) and the absolute self containment of their proofs. The use of convexity in the data will be addressed in a forthcoming paper.

## 2 Special Cases and Applications

In this section we make explicit the meaning of the robust duality of the general model introduced in Section 1, composed by a family of perturbation functions together with its corresponding optimization problems. We are doing this by exploring the extreme case with no uncertainty, the extreme case in absence of perturbations, and two other significant situations. In all these cases, we propose *ad hoc* families of perturbation functions allowing to apply the duality results to given optimization problems, either turning back to variants of well-known formulas for conjugate functions or proposing new ones.

Let us recall the robust duality formula,  $\inf (\text{RP})_{x^*} = \sup (\text{ODP})_{x^*}$ , i.e.,

$$\inf_{x \in X} \sup_{u \in U} \{F_u(x, 0_u) - \langle x^*, x \rangle\} = \sup_{(u, y_u^*) \in \Delta} -F_u^*(x^*, y_u^*). \quad (2.1)$$

We firstly study the two extreme cases: the case with no uncertainty and the one with no perturbations.

**Case 1. The case with no uncertainty:** Deterministic optimization with linear perturbations deals with parametric problems of the form:

$$(P)_{x^*} : \quad \inf_{x \in X} \{f(x) - \langle x^*, x \rangle\},$$

where  $f : X \rightarrow \mathbb{R}_\infty$  (i.e.,  $f \in (\mathbb{R}_\infty)^X$ ) is the *nominal objective function* and the *parameter* is  $x^* \in X^*$ . Taking a singleton uncertainty set  $U = \{u_0\}$ ,  $Y_{u_0} = Y$  and  $F_{u_0} = F$  such that  $F(x, 0_Y) = f(x)$  for all  $x \in X$ , (2.1) reads

$$\inf_{x \in X} \{F(x, 0_Y) - \langle x^*, x \rangle\} = \sup_{y^* \in Y^*} -F^*(x^*, y^*), \quad (2.2)$$

which is *the fundamental perturbational duality formula* [3], [20], [24]. Stable and strong robust duality theorems are given in [5] (see also [7] and [16] for infinite optimization problems).

**Case 2. The case with no linear perturbations:** Uncertain optimization without linear perturbations deals with families of problems of the form

$$(P) : \quad \left\{ \inf_{x \in X} f_u(x) \right\}_{u \in U},$$

where  $f_u \in (\mathbb{R}_\infty)^X$ ,  $u \in U$ . Taking  $F$  such that  $F_u(x, 0_u) = f_u(x)$  for all  $u \in U$ , the problem  $(P_u)_{0_x^*}$  represents here the scenario of (P) corresponding to  $u \in U$ , while  $(RP)_{0_x^*}$  is the *robust counterpart* (also called *pessimistic* or *minmax counterpart* in the robust optimization literature) of (P), namely,

$$(RP) \quad \inf_{x \in X} \sup_{u \in U} f_u(x).$$

For instance, if  $F_u(x, y_u) = f_u(x)$ , for all  $y_u \in Y_u$ , and  $\text{dom } f_u \neq \emptyset$ , we have

$$F_u^*(x, y_u^*) = \begin{cases} f_u^*(x^*), & \text{if } y_u^* = 0_u^*, \\ +\infty, & \text{if } y_u^* \neq 0_u^*. \end{cases} \quad (2.3)$$

Then (2.1) writes

$$\left( \sup_{u \in U} f_u \right)^*(x^*) = \inf_{u \in U} f_u^*(x^*), \quad (2.4)$$

which amounts, for  $x^* = 0_x^*$ , to the *inf – sup duality in robust optimization*, also called *robust infimum* (recall that any constrained optimization problem can be reduced to an unconstrained one by summing up the indicator function of the feasible set to the objective function):

$$\inf_{x \in X} \sup_{u \in U} f_u(x) = \sup_{u \in U} \inf_{x \in X} f_u(x).$$

Robust duality theorems without linear perturbations are given in [23] for a special class of uncertain non-convex optimization problems while [7] provides robust strong

duality theorems for uncertain convex optimization problems which are expressed in terms of the closedness of suitable sets regarding the vertical axis of  $X^* \times \mathbb{R}$ .

**Case 3. Conic optimization problem with uncertain constraints:** *Consider the uncertain problem*

$$(P) : \quad \left\{ \inf_{x \in X} f(x) \text{ s.t. } H_u(x) \in -S_u \right\}_{u \in U},$$

where, for each  $u \in U$ ,  $S_u$  is an ordering convex cone in  $Y_u$ ,  $H_u: X \rightarrow Y_u$ , and  $f \in (\mathbb{R}_\infty)^X$ . Denote by  $S_u^+ := \{y_u^* \in Y_u^* : \langle y_u^*, y_u \rangle \geq 0, \forall y_u \in S_u\}$  the dual cone of  $S_u$ .

Problems of this type arise, for instance, in the production planning of firms producing  $n$  commodities from uncertain amounts of resources by means of technologies which depend on the available resources (e.g., the technology differs when the energy is supplied by either fuel gas or a liquid fuel). The problem associated with each parameter  $u \in U$  consists of maximizing the cash-flow  $c(x_1, \dots, x_n)$  of the total production, with  $x_i$  denoting the production level of the  $i$ -th commodity,  $i = 1, \dots, n$ . The decision vector  $x = (x_1, \dots, x_n)$  must satisfy a linear inequality system  $A_u x \leq b_u$ , where the matrix of technical coefficients  $A_u$  is  $m_u \times n$  and  $b_u \in \mathbb{R}^{m_u}$ , for some  $m_u \in \mathbb{N}$ . Denoting by  $i_{\mathbb{R}_+^n}$  the indicator function of  $\mathbb{R}_+^n$  (i.e.,  $i_{\mathbb{R}_+^n}(x) = 0$ , when  $x \in \mathbb{R}_+^n$ , and  $i_{\mathbb{R}_+^n}(x) = +\infty$ , otherwise), the uncertain production planning problem can be formulated as

$$(P) : \quad \left\{ \inf_{x \in \mathbb{R}^n} f(x) = -c(x) + i_{\mathbb{R}_+^n}(x) \text{ s.t. } A_u x - b_u \in -\mathbb{R}_+^{m_u} \right\}_{u \in U},$$

with the space  $Y_u = \mathbb{R}^{m_u}$  depending on the uncertain parameter  $u$ .

For each  $u \in U$ , define the perturbation function

$$F_u(x, y_u) = \begin{cases} f(x), & \text{if } H_u(x) + y_u \in -S_u, \\ +\infty, & \text{else.} \end{cases}$$

On the one hand,  $(RP)_{0_X^*}$  collapses to the robust counterpart of (P) in the sense of robust conic optimization with uncertain constraints:

$$(RP) : \quad \inf_{x \in X} f(x) \text{ s.t. } H_u(x) \in -S_u, \quad \forall u \in U.$$

On the other hand, it is easy to check that

$$F_u^*(x^*, y_u^*) = \begin{cases} (f + y_u^* \circ H_u)^*(x^*), & \text{if } y_u^* \in S_u^+, \\ +\infty, & \text{else,} \end{cases}$$

$(ODP)_{0_X^*}$  is nothing else than the optimistic dual in the sense of uncertain conic optimization:

$$(ODP) : \quad \sup_{u \in U, y_u^* \in S_u^+} \inf_{x \in X} \{f(x) + \langle y_u^*, H_u(x) \rangle\}$$

(a special case when  $Y_u = Y$ ,  $S_u = S$  for all  $u \in U$  is studied in [8, page 1097] and [17]). Thus,

- *Robust duality holds at  $0_X^*$*  means that  $\inf(\text{RP}) = \sup(\text{ODP})$ ,
- *Strong robust duality holds at  $0_X^*$*  means that

$$\inf \{f(x) : H_u(x) \in -S_u, \forall u \in U\} = \max_{\substack{u \in U \\ y_u^* \in S_u^+}} \inf_{x \in X} \{f(x) + \langle y_u^*, H_u(x) \rangle\}.$$

Conditions for having such an equality are provided in [8, Theorem 6.3], [9, Corollaries 5, 6], for the particular case  $Y_u = Y$  for all  $u \in U$ .

*Strong robust duality and uncertain Farkas lemma:* We focus again on the case where  $Y_u = Y$  and  $S_u = S$  for all  $u \in U$ . For a given  $r \in \mathbb{R}$ , let us consider the following statements:

- (i)  $H_u(x) \in -S, \forall u \in U \implies f(x) \geq r$ ,
- (ii)  $\exists u \in U, \exists y_u^* \in S^+$  such that  $f(x) + \langle y_u^*, H_u(x) \rangle \geq r, \forall x \in X$ .

Then, it is true that the Strong robust duality holds at  $0_X^*$  if and only if [(i)  $\iff$  (ii)] for each  $r \in \mathbb{R}$ , which can be seen as an uncertain Farkas lemma. For details see [8, Theorem 3.2] (also [9, Corollary 5 and Theorem 1]).

It is worth noticing that when return to problem (P), a given robust feasible solution  $\bar{x}$  is a minimizer if and only if  $f(\bar{x}) \leq f(x)$  for any robust feasible solution  $x$ . So, a robust (uncertain) Farkas lemma (with  $r = f(\bar{x})$ ) will lead automatically to an optimality test for (P). Robust conic optimization problems are studied in [2] and [21].

**Case 4. Discretizing infinite optimization problems:** Let  $f \in (\mathbb{R}_\infty)^X$  and  $g_t \in \mathbb{R}^X$  for all  $t \in T$  (a possibly infinite index set). Consider the set  $U$  of nonempty finite subsets of  $T$ , interpreted as admissible perturbations of  $T$ , and the parametric optimization problem

$$(P) : \left\{ \inf_{x \in X} f(x) \text{ s.t. } g_t(x) \leq 0, \forall t \in S \right\}_{S \in U}.$$

Consider the parameter space  $Y_S := \mathbb{R}^S$  (depending on  $S$ ) and the perturbation function  $F_S : X \times \mathbb{R}^S \rightarrow \mathbb{R}_\infty$  such that, for any  $x \in X$  and  $\mu := (\mu_s)_{s \in S} \in \mathbb{R}^S$ ,

$$F_S(x, \mu) = \begin{cases} f(x), & \text{if } g_s(x) \leq -\mu_s, \forall s \in S, \\ +\infty, & \text{else.} \end{cases}$$

We now interpret the problems associated with the family of function perturbations  $\{F_S : S \in U\}$ . One has  $Y_s^* = \mathbb{R}^S$  and

$$F_S^*(x^*, \lambda) = \begin{cases} \left( f + \sum_{s \in S} \lambda_s g_s \right)^*(x^*), & \text{if } \lambda \in \mathbb{R}_+^S, \\ +\infty, & \text{else.} \end{cases}$$

The robust counterpart at  $0_X^*$ ,

$$(\text{RP})_{0_X^*} : \inf f(x) \text{ s.t. } g_t(x) \leq 0 \text{ for all } t \in T,$$

is a general infinite optimization problem while the optimistic dual at  $0_X^*$  is

$$(\text{ODP})_{0_X^*} : \sup_{S \in U, \lambda \in \mathbb{R}_+^S} \left\{ \inf_{x \in X} \left( f(x) + \sum_{s \in S} \lambda_s g_s(x) \right) \right\},$$

or, equivalently, the Lagrange dual of  $(\text{RP})_{0_X^*}$ , i.e.,

$$(\text{ODP})_{0_{X^*}} : \sup_{\lambda \in \mathbb{R}_+^{(T)}} \left\{ \inf_{x \in X} \left( f(x) + \sum_{t \in T} \lambda_t g_t(x) \right) \right\},$$

where, for each  $\lambda = (\lambda_t)_{t \in T} \in \mathbb{R}_+^{(T)}$  (the subspace of  $\mathbb{R}^T$  formed by the functions  $\lambda$  whose support,  $\text{supp} \lambda := \{t \in T : \lambda_t \neq 0\}$ , is finite),

$$\sum_{t \in T} \lambda_t g_t(x) := \begin{cases} \sum_{t \in \text{supp} \lambda} \lambda_t g_t(x), & \text{if } \lambda \neq 0, \\ 0, & \text{if } \lambda = 0. \end{cases}$$

Following [10, Section 8.3], we say that  $(\text{RP})_{0_X^*}$  is *discretizable* if there exists a sequence  $(S_r)_{r \in \mathbb{N}} \subset U$  such that

$$\inf (\text{RP})_{0_X^*} = \liminf_r \{f(x) : g_t(x) \leq 0, \forall t \in S_r\}, \quad (2.5)$$

and it is *reducible* if there exists  $S \in U$  such that

$$\inf (\text{RP})_{0_X^*} = \inf \{f(x) : g_t(x) \leq 0, \forall t \in S\}.$$

Obviously,  $\inf (\text{RP})_{0_X^*} = -\infty$  entails that  $(\text{RP})_{0_X^*}$  is reducible which, in turn, implies that  $(\text{RP})_{0_X^*}$  is discretizable.

Discretizable and reducible problems are important in practice. Indeed, on the one hand, discretization methods generate sequences  $(S_r)_{r \in \mathbb{N}} \subset U$  satisfying 2.5 when  $(\text{RP})_{0_X^*}$  is discretizable; discretization methods for linear and nonlinear semi-infinite programs have been reviewed in [11, Subsection 2.3] and [19], while a hard infinite optimization problem has been recently solved via discretization in [18]. On the other hand, replacing the robust counterpart (a hard semi-infinite program when the uncertainty set is infinite) of a given uncertainty optimization problem, when it is reducible, by a finite subproblem allows many times to get the desired tractable reformulation (see e.g., [1] and [4]).

**Example 2.1 (Discretizing linear infinite optimization problems)** *Consider the problems introduced in Case 4 above, with  $f(\cdot) := \langle c^*, \cdot \rangle$  and  $g_t(x) := \langle a_t^*, \cdot \rangle - b_t$ , where  $c^*, a_t^* \in X^*$  and  $b_t \in \mathbb{R}$ , for all  $t \in T$ . Then,  $(\text{RP})_{0_X^*}$  collapses to the linear infinite programming problem*

$$(\text{RP})_{0_X^*} : \inf \langle c^*, x \rangle \quad \text{s.t.} \quad \langle a_t^*, x \rangle \leq b_t, \quad \forall t \in T,$$



whose feasible set we denote by  $A$ . So,  $\inf (\text{RP})_{0_X^*} = \inf_{x \in X} \{ \langle c^*, x \rangle + i_A(x) \}$ . We assume that  $A \neq \emptyset$ .

Given  $S \in U$  and  $\mu, \lambda \in \mathbb{R}^S$ ,

$$F_S(x, \mu) = \begin{cases} \langle c^*, x \rangle, & \text{if } \langle a_s^*, x \rangle \leq b_s - \mu_s, \forall s \in S, \\ +\infty, & \text{else,} \end{cases} \quad (2.6)$$

and

$$F_S^*(x^*, \lambda) = \begin{cases} \sum_{s \in S} \lambda_s b_s, & \text{if } \sum_{s \in S} \lambda_s a_s^* = x^* - c^* \text{ and } \lambda_s \geq 0, \forall s \in S, \\ +\infty, & \text{else.} \end{cases} \quad (2.7)$$

Hence,  $(\text{ODP})_{0_X^*}$  collapses to the so-called Haar dual problem [12] of  $(\text{RP})_{0_X^*}$ ,

$$(\text{ODP})_{0_X^*} : \sup \left\{ - \sum_{t \in \text{supp } \lambda} \lambda_t b_t : - \sum_{t \in \text{supp } \lambda} \lambda_t a_t^* = c^*, \lambda \in \mathbb{R}_+^{(T)} \right\},$$

i.e.,

$$\sup (\text{ODP})_{0_X^*} = - \inf_{S \in U, \lambda \in \mathbb{R}_+^S} \left\{ \sum_{s \in S} \lambda_s b_s : \sum_{s \in S} \lambda_s a_s^* = -c^* \right\}. \quad (2.8)$$

From (2.8), if  $\inf (\text{RP})_{0_X^*} = \max (\text{ODP})_{0_X^*} \in \mathbb{R}$ , then there exist  $S \in U$  and  $\lambda \in \mathbb{R}_+^S$  such that

$$\sum_{s \in S} \lambda_s (a_s^*, b_s) = - \left( c^*, \inf (\text{RP})_{0_X^*} \right). \quad (2.9)$$

Let  $A_S := \{x \in X : \langle a_s^*, x \rangle \leq b_s, \forall s \in S\}$ . Given  $x \in A_S$ , from (2.9),

$$0 \geq \sum_{s \in S} \lambda_s (\langle a_s^*, x \rangle - b_s) = - \langle c^*, x \rangle + \inf (\text{RP})_{0_X^*}.$$

Since

$$\begin{aligned} \inf (\text{RP})_{0_X^*} &\leq \langle c^*, x \rangle, \forall x \in A_S, \\ \inf (\text{RP})_{0_X^*} &= \inf \{ \langle c^*, x \rangle : \langle a_s^*, x \rangle \leq b_s, \forall s \in S \}, \end{aligned} \quad (2.10)$$

so that  $(\text{RP})_{0_X^*}$  is reducible. Conversely, if (2.10) holds with  $\inf (\text{RP})_{0_X^*} \in \mathbb{R}$  and cone  $\{(a_t^*, b_t) : t \in T\} + \mathbb{R}_+(0_X^*, 1)$  is weak\*-closed, since  $\inf (\text{RP})_{0_X^*} \leq \langle c^*, x \rangle$  is consequence of  $\{\langle a_s^*, x \rangle \leq b_s, \forall s \in S\}$ , by the non-homogeneous Farkas lemma in lcHtvs [6] and the closedness assumption, there exist  $\lambda \in \mathbb{R}_+^S$  and  $\mu \in \mathbb{R}_+$  such that

$$- \left( c^*, \inf (\text{RP})_{0_X^*} \right) = \sum_{s \in S} \lambda_s (a_s^*, b_s) + \mu (0_X^*, 1),$$

which implies that  $\mu = 0$  and  $\inf (\text{RP})_{0_X^*} = \max (\text{ODP})_{0_X^*}$ . The closedness assumption holds when  $X$  is finite dimensional (guaranteeing that any finitely generated convex cone in  $X^* \times \mathbb{R}$  is closed). So, as proved in [10, Theorem 8.3], a linear semi-infinite program  $(\text{RP})_{0_X^*}$  is reducible if and only if (2.10) holds if and only if  $\inf (\text{RP})_{0_X^*} = \max (\text{ODP})_{0_X^*}$ .

We now assume that  $\inf (\text{RP})_{0_X^*} = \sup (\text{ODP})_{0_X^*} \in \mathbb{R}$ . By (2.8), there exist sequences  $(S_r)_{r \in \mathbb{N}} \subset U$  and  $(\lambda_r)_{r \in \mathbb{N}}$ , with  $\lambda^r \in \mathbb{R}_+^{S_r}$  for all  $r \in \mathbb{N}$ , such that

$$\lim_r \inf_{\lambda^r \in \mathbb{R}_+^{S_r}} \left\{ \sum_{s \in S_r} \lambda_s^r b_s : \sum_{s \in S_r} \lambda_s^r a_s^* = -c^* \right\} = -\sup (\text{ODP})_{0_X^*}.$$

Denote  $v_r := -\sum_{s \in S_r} \lambda_s^r b_s$ . Then,

$$\sum_{s \in S_r} \lambda_s (a_s^*, b_s) = -(c^*, v_r), \quad (2.11)$$

with  $\lim_r v_r = \inf (\text{RP})_{0_X^*}$ . Let  $A_r := \{x \in X : \langle a_s^*, x \rangle \leq b_s, \forall s \in S_r\}$ ,  $r \in \mathbb{N}$ . Given  $x \in A_r$ , from (2.11),

$$0 \geq \sum_{s \in S_r} \lambda_s^r (\langle a_s^*, x \rangle - b_s) = -\langle c^*, x \rangle + v_r.$$

Since  $v_r \leq \langle c^*, x \rangle$  for all  $x \in A_r$ ,

$$v_r \leq \inf \{ \langle c^*, x \rangle : \langle a_s^*, x \rangle \leq b_s, \forall s \in S_r \} \leq \inf (\text{RP})_{0_X^*}.$$

Thus,

$$\lim_r \inf \{ \langle c^*, x \rangle : \langle a_s^*, x \rangle \leq b_s, \forall s \in S_r \} = \inf (\text{RP})_{0_X^*},$$

i.e.,  $(\text{RP})_{0_X^*}$  is discretizable. Once again, the converse is true in linear semi-infinite programming [10, Corollary 8.2.1], but not in linear infinite programming.

### 3 Robust Conjugate Duality

We now turn back to the general perturbation function  $F_u : X \times Y_u \rightarrow \mathbb{R}_\infty$ ,  $u \in U$  and let  $\Delta := \{(u, y_u^*) : u \in U, y_u^* \in Y_u^*\}$  be the disjoint union of the spaces  $Y_u^*$ . Recall that

$$(\text{RP})_{x^*} : \inf_{x \in X} \left\{ \sup_{u \in U} F_u(x, 0_u) - \langle x^*, x \rangle \right\}, \quad (3.1)$$

$$(\text{ODP})_{x^*} : \sup_{(u, y_u^*) \in \Delta} -F_u^*(x^*, y_u^*). \quad (3.2)$$

Define  $p \in \overline{\mathbb{R}}^X$  and  $q \in \overline{\mathbb{R}}^{X^*}$  such that

$$p := \sup_{u \in U} F_u(\cdot, 0_u) \quad \text{and} \quad q := \inf_{(u, y_u^*) \in \Delta} F_u^*(\cdot, y_u^*). \quad (3.3)$$

One then has

$$\begin{cases} p^*(x^*) = -\inf (\text{RP})_{x^*}, & q(x^*) = -\sup (\text{ODP})_{x^*} \\ q^* = \sup_{(u, y_u^*) \in \Delta} (F_u^*(\cdot, y_u^*))^* = \sup_{u \in U} F_u^{**}(\cdot, 0_u) \leq p, \end{cases} \quad (3.4)$$

and hence,

- *Weak robust duality* always holds:

$$p^*(x^*) \leq q^{**}(x^*) \leq q(x^*), \text{ for all } x^* \in X^*. \quad (3.5)$$

- *Robust duality at  $x^*$*  means:

$$p^*(x^*) = q(x^*). \quad (3.6)$$

Robust duality at  $x^*$  also holds when either  $p^*(x^*) = +\infty$  or  $q(x^*) = -\infty$ .

As an illustration, consider Case 4 with linear data, as in Example 2.1. Then,  $p(x) = \langle c^*, x \rangle + i_A(x)$ ,  $\text{dom } p = A$ , and so

$$p^*(0_X^*) = \sup_{x \in \mathbb{R}^n} (-p(x)) = - \inf_{x \in \mathbb{R}^n} \{ \langle c^*, x \rangle + i_A(x) \} = - \inf (\text{RP})_{0_X^*}.$$

Similarly, from (2.7),

$$q(x^*) = \inf_{S \in U, \lambda \in \mathbb{R}^S} \left\{ \sum_{s \in S} \lambda_s b_s : \sum_{s \in S} \lambda_s a_s^* = x^* - c^* \right\},$$

$\text{dom } q = c^* + \text{cone} \{a_t^* : t \in T\}$  and

$$q(0_X^*) = \inf_{S \in U, \lambda \in \mathbb{R}_+^S} \left\{ \sum_{s \in S} \lambda_s b_s : \sum_{s \in S} \lambda_s a_s^* = -c^* \right\} = - \sup (\text{ODP})_{0_X^*}. \quad (3.7)$$

### 3.1 Basic lemmas

Let us introduce the necessary notations. Given a lcHtvs  $Z$ , an extended real-valued function  $h \in \overline{\mathbb{R}}^Z$ , and  $\varepsilon \in \mathbb{R}_+$ , the set of  $\varepsilon$ -minimizers of  $h$  is defined by

$$\varepsilon - \text{argmin } h := \begin{cases} \{z \in Z : h(z) \leq \inf_Z h + \varepsilon\}, & \text{if } \inf_Z h \in \mathbb{R}, \\ \emptyset, & \text{if } \inf_Z h \notin \mathbb{R}, \end{cases}$$

or, equivalently,

$$\varepsilon - \text{argmin } h = \{z \in h^{-1}(\mathbb{R}) : h(z) \leq \inf_Z h + \varepsilon\}.$$

Note that  $\varepsilon - \text{argmin } h \neq \emptyset$  when  $\inf_Z h \in \mathbb{R}$  and  $\varepsilon > 0$ . Various calculus rules involving  $\varepsilon - \text{argmin}$  have been given in [22].

The  $\varepsilon$ -subdifferential of  $h$  at a point  $a \in Z$  is the set (see, for instance, [15])

$$\begin{aligned} \partial^\varepsilon h(a) &:= \begin{cases} \{z^* \in Z^* : h(z) \geq h(a) + \langle z^*, z - a \rangle - \varepsilon, \forall z \in Z\}, & \text{if } h(a) \in \mathbb{R}, \\ \emptyset, & \text{if } h(a) \notin \mathbb{R} \end{cases} \\ &= \left\{ z^* \in (h^*)^{-1}(\mathbb{R}) : h^*(z^*) + h(a) \leq \langle z^*, a \rangle + \varepsilon \right\}. \end{aligned}$$

It can be checked that if  $h \in \overline{\mathbb{R}}^X$  is convex and  $h(a) \in \mathbb{R}$ , then  $\partial^\varepsilon h(a) \neq \emptyset$  for all  $\varepsilon > 0$  if and only if  $h$  is lower semi-continuous at  $a$ .

The inverse of the set-valued mapping  $\partial^\varepsilon h : Z \rightrightarrows Z^*$  is denoted by  $M^\varepsilon h : Z^* \rightrightarrows Z$ . For each  $(\varepsilon, z^*) \in \mathbb{R}_+ \times Z^*$ , we have

$$\left(\partial^\varepsilon h\right)^{-1}(z^*) = \left(M^\varepsilon h\right)(z^*) = \varepsilon - \operatorname{argmin}(h - z^*).$$

Denoting by  $\partial^\varepsilon h^*(z^*)$  the  $\varepsilon$ -subdifferential of  $h^*$  at  $z^* \in Z^*$ , namely,

$$\partial^\varepsilon h^*(z^*) = \left\{ z \in (h^{**})^{-1}(\mathbb{R}) : h^{**}(z) + h^*(z^*) \leq \langle z^*, z \rangle + \varepsilon \right\},$$

where  $h^{**}(z) := \sup_{z^* \in Z^*} \{\langle z^*, z \rangle - h^*(z^*)\}$  is the biconjugate of  $h$ , we have

$$(M^\varepsilon h)(z^*) \subset (\partial^\varepsilon h^*)(z^*), \quad \forall (\varepsilon, z^*) \in \mathbb{R}_+ \times Z^*,$$

with equality if and only if  $h = h^{**}$ .

For each  $\varepsilon \in \mathbb{R}_+$ , we consider the set-valued mapping  $S^\varepsilon : X^* \rightrightarrows X$  as follows:

$$S^\varepsilon(x^*) := \{x \in p^{-1}(\mathbb{R}) : p(x) - \langle x^*, x \rangle \leq -q(x^*) + \varepsilon\}.$$

If  $q(x^*) = -\infty$ , then  $S^\varepsilon(x^*) = p^{-1}(\mathbb{R})$ . If  $q(x^*) = +\infty$ , then  $S^\varepsilon(x^*) = \emptyset$ .

Since  $p^* \leq q$ , it is clear that

$$S^\varepsilon(x^*) \subset (M^\varepsilon p)(x^*), \quad \forall \varepsilon \geq 0, \quad \forall x^* \in X^*. \quad (3.8)$$

**Lemma 3.1** *Assume that  $\operatorname{dom} p \neq \emptyset$ . Then, for each  $x^* \in X^*$ , the next statements are equivalent:*

- (i) *Robust duality holds at  $x^*$ , i.e.,  $p^*(x^*) = q(x^*)$ ,*
- (ii)  $(M^\varepsilon p)(x^*) = S^\varepsilon(x^*), \quad \forall \varepsilon \geq 0,$
- (iii)  $\exists \bar{\varepsilon} > 0 : (M^\varepsilon p)(x^*) = S^\varepsilon(x^*), \quad \forall \varepsilon \in ]0, \bar{\varepsilon}[.$

*Proof.* [(i)  $\Rightarrow$  (ii)] By definition

$$\begin{aligned} (M^\varepsilon p)(x^*) &= \varepsilon - \operatorname{argmin}(p - x^*) \\ &= \{x \in p^{-1}(\mathbb{R}) : p(x) - \langle x^*, x \rangle \leq -p^*(x^*) + \varepsilon\}. \end{aligned}$$

By (i) we thus have  $(M^\varepsilon p)(x^*) = S^\varepsilon(x^*)$ .

[(ii)  $\Rightarrow$  (iii)] It is obviously true.

[(iii)  $\Rightarrow$  (i)] Since  $p^*(x^*) \leq q(x^*)$ , (i) holds if  $p^*(x^*) = +\infty$ . Moreover, since  $\operatorname{dom} p \neq \emptyset$ , one has  $p^*(x^*) \neq -\infty$ . Let now  $p^*(x^*) \in \mathbb{R}$ . In order to get a contradiction, assume that  $p^*(x^*) \neq q(x^*)$ . Then  $p^*(x^*) < q(x^*)$  and there exists  $\varepsilon \in ]0, \bar{\varepsilon}[$  such that  $p^*(x^*) + \varepsilon < q(x^*)$ . Since  $\inf_{x \in X} \{p(x) - \langle x^*, x \rangle\} = -p^*(x^*) \in \mathbb{R}$  and  $\varepsilon > 0$ , we have

$\varepsilon - \operatorname{argmin}(p - x^*) \neq \emptyset$ . Let us pick  $x \in (M^\varepsilon p)(x^*) = \varepsilon - \operatorname{argmin}(p - x^*)$ . By (iii), we have  $x \in S^\varepsilon(x^*)$  and

$$-p^*(x^*) \leq p(x) - \langle x^*, x \rangle \leq -q(x^*) + \varepsilon,$$

which contradicts  $p^*(x^*) + \varepsilon < q(x^*)$ .  $\square$

For each  $\varepsilon \in \mathbb{R}_+$ , let us introduce now the the following set-valued mapping  $J^\varepsilon : U \rightrightarrows X$ :

$$J^\varepsilon(u) := \{x \in p^{-1}(\mathbb{R}) : p(x) \leq F_u(x, 0_u) + \varepsilon\}, \quad (3.9)$$

with the aim of making explicit the set  $S^\varepsilon(x^*)$ . To this purpose, given  $\varepsilon_1, \varepsilon_2 \in \mathbb{R}_+$ ,  $u \in U$ , and  $y_u^* \in Y_u^*$ , let us introduce the set-valued mapping  $A_{(u, y_u^*)}^{(\varepsilon_1, \varepsilon_2)} : X^* \rightrightarrows X$  such that

$$A_{(u, y_u^*)}^{(\varepsilon_1, \varepsilon_2)}(x^*) := \left\{x \in J^{\varepsilon_1}(u) : (x, 0_u) \in (M^{\varepsilon_2} F_u)(x^*, y_u^*)\right\}.$$

**Lemma 3.2** *For each  $x^* \in X^*$ ,  $\varepsilon_1, \varepsilon_2 \in \mathbb{R}_+$ ,  $u \in U$ , and  $y_u^* \in Y_u^*$ , one has*

$$A_{(u, y_u^*)}^{(\varepsilon_1, \varepsilon_2)}(x^*) \subset S^{\varepsilon_1 + \varepsilon_2}(x^*).$$

*Proof.* Let  $x \in J^{\varepsilon_1}(u)$  be such that  $(x, 0_u) \in (M^{\varepsilon_2} F_u)(x^*, y_u^*)$ . Then we have  $F_u^*(x^*, y_u^*) \in \mathbb{R}$  and  $F_u(x, 0_u) \in \mathbb{R}$ . Moreover

$$F_u(x, 0_u) + \varepsilon_1 \geq p(x) \geq F_u(x, 0_u) \in \mathbb{R},$$

implying  $p(x) \in \mathbb{R}$  and, by (3.4),

$$\begin{aligned} p(x) - \langle x^*, x \rangle &\leq F_u(x, 0_u) - \langle x^*, x \rangle + \varepsilon_1 \leq -F_u^*(x^*, y_u^*) + \varepsilon_1 + \varepsilon_2 \\ &\leq -q(x^*) + \varepsilon_1 + \varepsilon_2, \end{aligned}$$

that means  $x \in S^{\varepsilon_1 + \varepsilon_2}(x^*)$ .  $\square$

**Lemma 3.3** *Assume that*

$$\operatorname{dom} F_u \neq \emptyset, \quad \forall u \in U. \quad (3.10)$$

*Then, for each  $x^* \in X^*$ ,  $\varepsilon \in \mathbb{R}_+$ ,  $\eta > 0$ , one has*

$$S^\varepsilon(x^*) \subset \bigcup_{\substack{u \in U \\ y_u^* \in Y_u^*}} \bigcup_{\substack{\varepsilon_1 + \varepsilon_2 = \varepsilon + \eta \\ \varepsilon_1 \geq 0, \varepsilon_2 \geq 0}} A_{(u, y_u^*)}^{(\varepsilon_1, \varepsilon_2)}(x^*).$$

*Proof.* Let  $x \in p^{-1}(\mathbb{R})$  be such that  $x \in S^\varepsilon(x^*)$ , i.e.,

$$p(x) - \langle x^*, x \rangle \leq -q(x^*) + \varepsilon.$$

We then have, for any  $\eta > 0$ ,

$$q(x^*) < \langle x^*, x \rangle - p(x) + \varepsilon + \eta$$

and, by definition of  $q$  and  $p$ , there exist  $u \in U$ ,  $y_u^* \in Y_u^*$  such that

$$F_u^*(x^*, y_u^*) \leq \langle x^*, x \rangle - p(x) + \varepsilon + \eta \leq \langle x^*, x \rangle - F_u(x, 0_u) + \varepsilon + \eta. \quad (3.11)$$

Since  $p(x) \in \mathbb{R}$ ,  $F_u^*(x^*, y_u^*) \neq +\infty$ . In fact, by (3.10),  $F_u^*(x^*, y_u^*) \in \mathbb{R}$ . Similarly,  $F_u(x, 0_u) \in \mathbb{R}$ . Setting

$$\alpha_1 := p(x) - F_u(x, 0_u), \quad \alpha_2 := F_u^*(x^*, y_u^*) + F_u(x, 0_u) - \langle x^*, x \rangle,$$

we get  $\alpha_1 \in \mathbb{R}_+$ ,  $\alpha_2 \in \mathbb{R}$ . Actually  $\alpha_2 \geq 0$  since, by definition of conjugate,

$$F_u^*(x^*, y_u^*) = \sup_{z \in X, y_u \in Y_u} \{ \langle x^*, z \rangle + \langle y_u^*, y_u \rangle - F_u(z, y_u) \},$$

i.e., if  $z = x$  and  $y_u = 0_u$ ,

$$F_u^*(x^*, y_u^*) \geq \langle x^*, x \rangle - F_u(x, 0_u),$$

so that

$$F_u^*(x^*, y_u^*) + F_u(x, 0_u) - \langle x^*, x \rangle \geq 0.$$

Then, by (3.11),  $0 \leq \alpha_1 + \alpha_2 \leq \varepsilon + \eta$ . Consequently, there exist  $\varepsilon_1, \varepsilon_2 \in \mathbb{R}_+$  such that  $\alpha_1 \leq \varepsilon_1$ ,  $\alpha_2 \leq \varepsilon_2$ ,  $\varepsilon_1 + \varepsilon_2 = \varepsilon + \eta$ . Now  $\alpha_1 \leq \varepsilon_1$  means that  $x \in J^{\varepsilon_1}(u)$  and  $\alpha_2 \leq \varepsilon_2$  means that  $(x, 0_u) \in (M^{\varepsilon_2} F_u)(x^*, y_u^*)$ , and we have  $x \in A_{(u, y_u^*)}^{(\varepsilon_1, \varepsilon_2)}(x^*)$ .  $\square$

For each  $x^* \in X^*$ ,  $\varepsilon \in \mathbb{R}_+$ , let us define

$$\begin{aligned} \mathcal{A}^\varepsilon(x^*) &:= \bigcap_{\eta > 0} \bigcup_{\substack{u \in U \\ y_u^* \in Y_u^*}} \bigcup_{\substack{\varepsilon_1 + \varepsilon_2 = \varepsilon + \eta \\ \varepsilon_1 \geq 0, \varepsilon_2 \geq 0}} A_{(u, y_u^*)}^{(\varepsilon_1, \varepsilon_2)}(x^*) \\ &= \bigcap_{\eta > 0} \bigcup_{\substack{u \in U \\ y_u^* \in Y_u^*}} \bigcup_{\substack{\varepsilon_1 + \varepsilon_2 = \varepsilon + \eta \\ \varepsilon_1 \geq 0, \varepsilon_2 \geq 0}} \left\{ x \in J^{\varepsilon_1}(u) : (x, 0_u) \in (M^{\varepsilon_2} F_u)(x^*, y_u^*) \right\} \end{aligned}$$

### 3.2 Robust duality

We now can state the main result on characterizations of the robust conjugate duality.

**Theorem 3.1 (Robust duality)** *Assume that  $\text{dom } p \neq \emptyset$ . Then for each  $x^* \in X^*$ , the next statements are equivalent:*

- (i)  $\inf (\text{RP})_{x^*} = \sup (\text{ODP})_{x^*}$ ,
- (ii)  $(M^\varepsilon p)(x^*) = \mathcal{A}^\varepsilon(x^*)$ ,  $\forall \varepsilon \geq 0$ ,
- (iii)  $\exists \bar{\varepsilon} > 0 : (M^\varepsilon p)(x^*) = \mathcal{A}^\varepsilon(x^*)$ ,  $\forall \varepsilon \in ]0, \bar{\varepsilon}[$ .

*Proof.* We firstly claim that if  $\text{dom } p \neq \emptyset$  then for each  $x^* \in X^*$ ,  $\varepsilon \in \mathbb{R}_+$ , it holds:

$$S^\varepsilon(x^*) = \mathcal{A}^\varepsilon(x^*). \quad (3.12)$$

In deed, as  $\text{dom } p \neq \emptyset$ , (3.10) holds. It then follows from Lemma 3.3,  $S^\varepsilon(x^*) \subset \mathcal{A}^\varepsilon(x^*)$ . On the other hand, for each  $\eta > 0$ , one has, by Lemma 3.2,

$$\bigcup_{\substack{u \in U \\ y_u^* \in Y_u^*}} \bigcup_{\substack{\varepsilon_1 + \varepsilon_2 = \varepsilon + \eta \\ \varepsilon_1 \geq 0, \varepsilon_2 \geq 0}} A_{(u, y_u^*)}^{(\varepsilon_1, \varepsilon_2)}(x^*) \subset S^{\varepsilon + \eta}(x^*).$$

Taking the intersection over all  $\eta > 0$  we get

$$\mathcal{A}^\varepsilon(x^*) \subset \bigcap_{\eta > 0} S^{\varepsilon + \eta}(x^*) = S^\varepsilon(x^*),$$

and (3.12) follows. Taking into account the fact that (i) means  $p^*(x^*) = q(x^*)$ , the conclusions now follows from (3.12) and Lemma 3.1.  $\square$

For the deterministic optimization problem with linear perturbations (i.e., non-uncertain case where  $U$  is a singleton), the next result is a direct consequence of Theorem 3.1.

**Corollary 3.1 (Robust duality for Case 1)** *Let  $F: X \times Y \rightarrow \mathbb{R}_\infty$  be such that  $\text{dom } F(\cdot, 0_Y) \neq \emptyset$ . Then, for each  $x^* \in X^*$ , the fundamental duality formula (2.2) holds, i.e.,*

$$\inf_{x \in X} \{F(x, 0_Y) - \langle x^*, x \rangle\} = \sup_{y \in Y^*} -F^*(x^*, y^*),$$

if and only any of the (equivalent) conditions (ii) or (iii) in Theorem 3.1 holds, where

$$\mathcal{A}^\varepsilon(x^*) = \bigcap_{\eta > 0} \bigcup_{y^* \in Y^*} \left\{ x \in X : (x, 0_Y) \in (M^{\varepsilon + \eta} F)(x^*, y^*) \right\}. \quad (3.13)$$

*Proof.* Let  $F_u = F: X \times Y \rightarrow \mathbb{R}_\infty$ ,  $p = F(\cdot, 0_Y)$ . In this case, one has,

$$J^\varepsilon(u) = \{x \in X : F(x, 0_Y) \in \mathbb{R}\}, \quad \forall \varepsilon \geq 0,$$

and  $\mathcal{A}^\varepsilon(x^*)$  will take the form (3.13). The conclusion follows from Theorem 3.1.  $\square$

For uncertain optimization problem without linear perturbations, the following result is a consequence of Theorem 3.1.

**Corollary 3.2 (Robust duality for Case 2)** *Let  $(f_u)_{u \in U} \subset \mathbb{R}_\infty^X$  be a family of extended real-valued functions,  $p = \sup_{u \in U} f_u$  be such that  $\text{dom } p \neq \emptyset$ . Then, for each  $x^* \in X^*$ , the inf – sup duality in robust optimization (2.4) holds, i.e.,*

$$\left( \sup_{u \in U} f_u \right)^*(x^*) = \inf_{u \in U} f_u^*(x^*),$$

if and only any of the (equivalent) conditions (ii) or (iii) in Theorem 3.1 holds, where

$$\mathcal{A}^\varepsilon(x^*) = \bigcap_{\eta > 0} \bigcup_{u \in U} \bigcup_{\substack{\varepsilon_1 + \varepsilon_2 = \varepsilon + \eta \\ \varepsilon_1 \geq 0, \varepsilon_2 \geq 0}} \{J^{\varepsilon_1}(u) \cap (M^{\varepsilon_2} f_u)(x^*)\}, \quad (3.14)$$

with

$$J^{\varepsilon_1}(u) = \{x \in p^{-1}(\mathbb{R}) : f_u(x) \geq p(x) - \varepsilon_1\}.$$

*Proof.* Let  $F_u(x, y_u) = f_u(x)$ , for all  $u \in U$  and let  $p = \sup_{u \in U} f_u$ . Then, by (3.9),

$$J^\varepsilon(u) = \{x \in p^{-1}(\mathbb{R}) : f_u(x) \geq p(x) - \varepsilon\}, \quad \forall \varepsilon \geq 0. \quad (3.15)$$

Moreover, recalling (2.3), for each  $u \in U$  such that  $\text{dom } f_u \neq \emptyset$ ,  $(x^*, y_u^*) \in X^* \times Y_u^*$ , and  $\varepsilon \geq 0$ ,

$$(M^\varepsilon F_u)(x^*, y_u^*) = \begin{cases} (M^\varepsilon f_u)(x^*), & \text{if } y_u^* = 0_u^*, \\ \emptyset, & \text{else.} \end{cases} \quad (3.16)$$

Finally, for each  $(x^*, \varepsilon) \in X^* \times \mathbb{R}_+$ ,  $\mathcal{A}^\varepsilon(x^*)$  takes the form as in (3.14). The conclusion now follows from Theorem 3.1.  $\square$

## 4 Strong Robust Duality

We retain the notations in Section 3 and consider the robust problem  $(\text{RP})_{x^*}$  and its robust dual problem  $(\text{ODP})_{x^*}$  given in (3.1) and (3.2), respectively. Let  $p$  and  $q$  be the functions defined by (3.3) and recall the relations in (3.4), that is,

$$\begin{cases} p^*(x^*) = -\inf (\text{RP})_{x^*}, & q(x^*) = -\sup (\text{ODP})_{x^*} \\ q^* = \sup_{(u, y_u^*) \in \Delta} (F_u^*(\cdot, y_u^*))^* = \sup_{u \in U} F_u^{**}(\cdot, 0_u) \leq p. \end{cases}$$

In this section we will establish characterizations of *strong robust duality at  $x^*$* . Recall that the strong robust duality holds at  $x^*$  means that  $\inf (\text{RP})_{x^*} = \max (\text{ODP})_{x^*}$ , which is the same as:

$$\exists (u, y_u^*) \in \Delta : p^*(x^*) = F_u^*(x^*, y_u^*).$$

For this, we need a technical lemma, but firstly, given  $x^* \in X^*$ ,  $u \in U$ ,  $y_u^* \in Y_u^*$ , and  $\varepsilon \geq 0$ , let us introduce the set

$$B_{(u, y_u^*)}^\varepsilon(x^*) = \bigcup_{\substack{\varepsilon_1 + \varepsilon_2 = \varepsilon \\ \varepsilon_1 \geq 0, \varepsilon_2 \geq 0}} A_{(u, y_u^*)}^{(\varepsilon_1, \varepsilon_2)}(x^*) = \bigcup_{\substack{\varepsilon_1 + \varepsilon_2 = \varepsilon \\ \varepsilon_1 \geq 0, \varepsilon_2 \geq 0}} \left\{ x \in J^{\varepsilon_1}(u) : (x, 0_u) \in (M^{\varepsilon_2} F_u)(x^*, y_u^*) \right\}.$$

**Lemma 4.1** *Assume that  $\text{dom } F_u \neq \emptyset$ , for all  $u \in U$ , holds and let  $x^* \in X^*$  be such that*

$$q(x^*) = \min_{\substack{u \in U \\ y_u^* \in Y_u^*}} F_u^*(x^*, y_u^*).$$

*Then there exist  $u \in U$ ,  $y_u^* \in Y_u^*$  such that*

$$S^\varepsilon(x^*) = B_{(u, y_u^*)}^\varepsilon(x^*), \quad \forall \varepsilon \geq 0.$$

*Proof.* By Lemma 3.2 we have  $B_{(u, y_u^*)}^\varepsilon(x^*) \subset S^\varepsilon(x^*)$ . Conversely, let  $x \in S^\varepsilon(x^*)$ . By the exactness of  $q$  at  $x^*$ , there exist  $u \in U$  and  $y_u^* \in Y_u^*$  such that

$$p(x) - \langle x^*, x \rangle \leq -F_u^*(x^*, y_u^*) + \varepsilon.$$



Since  $p(x) \in \mathbb{R}$  and  $\text{dom } F_u \neq \emptyset$ , for all  $u \in U$ , we have  $F_u^*(x^*, y_u^*) \in \mathbb{R}$ ,  $F_u(x, 0_u) \in \mathbb{R}$ ,

$$\left( p(x) - F_u(x, 0_u) \right) + \left( F_u(x, 0_u) + F_u^*(x^*, y_u^*) - \langle x^*, x \rangle \right) \leq \varepsilon.$$

Consequently, there exist  $\varepsilon_1 \geq 0, \varepsilon_2 \geq 0$  such that  $\varepsilon_1 + \varepsilon_2 = \varepsilon$ ,

$$p(x) - F_u(x, 0_u) \leq \varepsilon_1 \text{ and } F_u(x, 0_u) + F_u^*(x^*, y_u^*) - \langle x^*, x \rangle \leq \varepsilon_2,$$

that is,  $x \in J^{\varepsilon_1}(u)$  and  $(x, 0_u) \in (M^{\varepsilon_2} F_u)(x^*, y_u^*)$ . Thus,  $x \in A_{(u, y_u^*)}^{(\varepsilon_1, \varepsilon_2)}(x^*) \subset B_{(u, y_u^*)}^\varepsilon(x^*)$ , since  $\varepsilon_1 + \varepsilon_2 = \varepsilon$ .  $\square$

**Theorem 4.1 (Strong robust duality)** *Assume that  $\text{dom } p \neq \emptyset$  and let  $x^* \in X^*$ . The next statements are equivalent:*

- (i)  $\inf (\text{RP})_{x^*} = \max (\text{ODP})_{x^*}$ ,
- (ii)  $\exists u \in U, \exists y_u^* \in Y_u^* : (M^\varepsilon p)(x^*) = B_{(u, y_u^*)}^\varepsilon(x^*), \forall \varepsilon \geq 0$ ,
- (iii)  $\exists \bar{\varepsilon} > 0, \exists u \in U, \exists y_u^* \in Y_u^* : (M^\varepsilon p)(x^*) = B_{(u, y_u^*)}^\varepsilon(x^*), \forall \varepsilon \in ]0, \bar{\varepsilon}[$ .

*Proof.* Observe firstly that (i) means that

$$p^*(x^*) = q(x^*) = \min_{\substack{u \in U \\ y_u^* \in Y_u^*}} F_u^*(x^*, y_u^*).$$

As  $\text{dom } p \neq \emptyset$ , (3.10) holds, and then by Lemmas 3.1 and 4.1, (i) implies the remaining conditions, which are equivalent to each other, and also that (iii) implies  $p^*(x^*) = q(x^*)$ .

We now prove that (iii) implies  $q(x^*) = F_u^*(x^*, y_u^*)$ . Assume by contradiction that there exists  $\varepsilon > 0$  such that  $q(x^*) + \varepsilon < F_u^*(x^*, y_u^*)$ , and without loss of generality one can take  $\varepsilon \in ]0, \bar{\varepsilon}[$ , where  $\bar{\varepsilon} > 0$  appeared in (iii). Then, by (iii),  $(M^\varepsilon p)(x^*) = B_{(u, y_u^*)}^\varepsilon(x^*)$ .

Pick  $x \in (M^\varepsilon p)(x^*) = B_{(u, y_u^*)}^\varepsilon(x^*)$ . Then, there are  $\varepsilon_1 \geq 0, \varepsilon_2 \geq 0, \varepsilon_1 + \varepsilon_2 = \varepsilon$  and  $x \in J^{\varepsilon_1}(u)$  and  $(x, 0_u) \in (M^{\varepsilon_2} F_u)(x^*, y_u^*)$ . In other words,

$$p(x) \leq F_u(x, 0_u) + \varepsilon_1, \tag{4.1}$$

$$F_u^*((x^*, y_u^*) + F_u(x, 0_u) \leq \langle x^*, x \rangle + \varepsilon_2. \tag{4.2}$$

It now follows from (4.1)-(4.2) that

$$\begin{aligned} p^*(x^*) &\geq \langle x^*, x \rangle - p(x) \geq \langle x^*, x \rangle - F_u(x, 0_u) - \varepsilon_1 \\ &\geq \langle x^*, x \rangle + F_u^*(x^*, y_u^*) - \langle x^*, x \rangle - \varepsilon_2 - \varepsilon_1 = F_u^*(x^*, y_u^*) - \varepsilon > q(x^*), \end{aligned}$$

which contradicts the fact that  $p^*(x^*) = q(x^*)$ .  $\square$

In deterministic optimization with linear perturbations we get the next consequence from Theorem 4.1.

**Corollary 4.1 (Strong robust duality for Case 1)** *Let  $F : X \times Y \rightarrow \mathbb{R}_\infty$ ,  $p = F(\cdot, 0_Y)$ , and assume that  $\text{dom } p \neq \emptyset$ . Then, for each  $x^* \in X^*$ , the strong duality for  $(P)_{x^*}$  in Case 1 holds at  $x^*$ , i.e.,*

$$\inf_{x \in X} \{F(x, 0_Y) - \langle x^*, x \rangle\} = \max_{y^* \in Y^*} -F^*(x^*, y^*),$$

*if and only if one of the (equivalent) conditions (ii) or (iii) in Theorem 4.1 holds with  $B_{(u, y_u^*)}^\varepsilon(x^*)$  being replaced by*

$$B_{y^*}^\varepsilon(x^*) := \{x \in X : (x, 0_Y) \in (M^\varepsilon F)(x^*, y^*)\}. \quad (4.3)$$

*Proof.* It is worth observing that we are in the non-uncertainty case (i.e.,  $U$  is a singleton), and the set  $B_{(u, y_u^*)}^\varepsilon(x^*)$  writes as in (4.3) for each  $(x^*, y^*) \in X^* \times Y^*$ ,  $\varepsilon \geq 0$ . The conclusion follows from Theorem 4.1.  $\square$

In the non-perturbation case, Theorem 4.1 gives rise to

**Corollary 4.2 (Strong robust duality for Case 2)** *Let  $(f_u)_{u \in U} \subset \mathbb{R}_\infty^X$ ,  $x^* \in X^*$ , and  $p = \sup_{u \in U} f_u$  such that  $\text{dom } p \neq \emptyset$ . Then, the robust duality formula*

$$\left(\sup_{u \in U} f_u\right)^*(x^*) = \min_{u \in U} f_u^*(x^*)$$

*holds if and only if one of the (equivalent) conditions (ii) or (iii) in Theorem 4.1 holds with  $B_{(u, y_u^*)}^\varepsilon(x^*)$  being replaced by*

$$B_u^\varepsilon(x^*) := \bigcup_{\substack{\varepsilon_1 + \varepsilon_2 = \varepsilon \\ \varepsilon_1 \geq 0, \varepsilon_2 \geq 0}} \left(J^{\varepsilon_1}(u) \cap (M^{\varepsilon_2} f_u)(x^*)\right). \quad (4.4)$$

*Proof.* Let  $F_u(x, y_u) = f_u(x)$ ,  $p = \sup_{u \in U} f_u$ , and, from (3.15) and (3.16) (see the proof of Corollary 3.2),

$$B_{(u, y_u^*)}^\varepsilon(x^*) = \begin{cases} \bigcup_{\substack{\varepsilon_1 + \varepsilon_2 = \varepsilon \\ \varepsilon_1 \geq 0, \varepsilon_2 \geq 0}} \left(J^{\varepsilon_1}(u) \cap (M^{\varepsilon_2} f_u)(x^*)\right), & \text{if } y_u^* = 0_u^*, \\ \emptyset, & \text{else,} \end{cases}$$

which in our situation, collapses to the set  $B_u^\varepsilon(x^*)$  defined by (4.4). The conclusion now follows from Theorem 4.1.  $\square$

## 5 Reverse strong and min-max robust duality

Given  $F_u : X \times Y_u \rightarrow (\mathbb{R}_\infty)^X$  for each  $u \in U$ ,  $p = \sup_{u \in U} F_u(\cdot, 0_u)$ , and  $x^* \in X^*$ , we assume in this section that the problem  $(\text{RP})_{x^*}$  is finite-valued and admits an optimal solution or, in other words, that  $\text{argmin}(p - x^*) = (M^0 p)(x^*) \neq \emptyset$ . For convenience, we set

$$(Mp)(x^*) := (M^0 p)(x^*), \quad S(x^*) := S^0(x^*), \text{ and}$$

$$\mathcal{A}(x^*) := \mathcal{A}^0(x^*) = \bigcap_{\eta > 0} \bigcup_{\substack{u \in U \\ y_u^* \in Y_u^*}} \bigcup_{\substack{\varepsilon_1 + \varepsilon_2 = \eta \\ \varepsilon_1 \geq 0, \varepsilon_2 \geq 0}} \left\{ x \in J^{\varepsilon_1}(u) : (x, 0_u) \in (M^{\varepsilon_2} F_u)(x^*, y_u^*) \right\}. \quad (5.1)$$

**Theorem 5.1 (Reverse strong robust duality)** *Let  $x^* \in X^*$  be such that  $(Mp)(x^*) \neq \emptyset$  and let  $\mathcal{A}(x^*)$  be as in (5.1). The next statements are equivalent:*

- (i)  $\min(\text{RP})_{x^*} = \sup(\text{ODP})_{x^*}$ ,
- (ii)  $(Mp)(x^*) = \mathcal{A}(x^*)$ .

*Proof.* Since  $(Mp)(x^*) \neq \emptyset$ ,  $\text{dom } p \neq \emptyset$ . It follows from Theorem 3.1 that [(i)  $\implies$  (ii)]. For the converse, let us pick  $x \in (Mp)(x^*)$ . Then by (ii), for each  $\eta > 0$  there exist  $u \in U$ ,  $y_u^* \in Y_u^*$ ,  $\varepsilon_1 \geq 0, \varepsilon_2 \geq 0$  such that  $\varepsilon_1 + \varepsilon_2 = \eta$ ,  $x \in J^{\varepsilon_1}(u)$ ,  $(x, 0_u) \in (M^{\varepsilon_2} F_u)(x^*, y_u^*)$  and we have

$$\begin{aligned} q(x^*) &\leq F_u^*(x^*, y_u^*) \leq \langle x^*, x \rangle - F_u(x, 0_u) + \varepsilon_2 \\ &\leq \langle x^*, x \rangle - p(x) + \varepsilon_1 + \varepsilon_2 \leq p^*(x^*) + \eta. \end{aligned}$$

Since  $\eta > 0$  is arbitrary we get  $q(x^*) \leq p^*(x^*)$ , which, together with the weak duality (see (3.5)), yields  $q(x^*) = \langle x^*, x \rangle - p(x) = p^*(x^*)$ , i.e., (i) holds and we are done.  $\square$

In the deterministic case we obtain from Theorem 5.1:

**Corollary 5.1 (Reverse strong robust duality for Case 1)** *Let  $F : X \times Y \rightarrow \mathbb{R}_\infty$ ,  $x^* \in X^*$ ,  $p = F(\cdot, 0_Y)$ , and*

$$\mathcal{A}(x^*) = \bigcap_{\eta > 0} \bigcup_{y^* \in Y^*} \left\{ x \in X : (x, 0_Y) \in (M^\eta F)(x^*, y^*) \right\}.$$

*Assume that  $(Mp)(x^*) \neq \emptyset$ . Then the next statements are equivalent:*

- (i)  $\min_{x \in X} \{ F(x, 0_Y) - \langle x^*, x \rangle \} = \sup_{y^* \in Y^*} -F^*(x^*, y^*)$ ,
- (ii)  $(Mp)(x^*) = \mathcal{A}(x^*)$ .

**Corollary 5.2 (Reverse strong robust duality for Case 2)** *Let  $(f_u)_{u \in U} \subset \mathbb{R}_\infty^X$ ,  $p = \sup_{u \in U} f_u$ ,  $x^* \in X^*$ , and*

$$\mathcal{A}(x^*) := \bigcap_{\eta > 0} \bigcup_{u \in U} \bigcup_{\substack{\varepsilon_1 + \varepsilon_2 = \eta \\ \varepsilon_1 \geq 0, \varepsilon_2 \geq 0}} \left( J^{\varepsilon_1}(u) \cap (M^{\varepsilon_2} f_u)(x^*) \right),$$

*where*

$$J^{\varepsilon_1}(u) = \left\{ x \in p^{-1}(\mathbb{R}) : f_u(x) \geq p(x) - \varepsilon_1 \right\}.$$

*Assume that  $(Mp)(x^*) \neq \emptyset$ . Then the next statements are equivalent:*

- (i)  $\left( \sup_{u \in U} f_u \right)^*(x^*) = \inf_{u \in U} f_u^*(x^*)$ , with attainment at the first member,
- (ii)  $(Mp)(x^*) = \mathcal{A}(x^*)$ .

Now, for each  $u \in U$ ,  $y_u^* \in Y_u^*$ ,  $x^* \in X^*$ , we set

$$J(u) := J^0(u) = \left\{ x \in p^{-1}(\mathbb{R}) : F_u(x, 0_u) = p(x) \right\},$$

$$(MF_u)(x^*, y_u^*) := (M^0 F_u)(x^*, y_u^*) = \operatorname{argmin} \left( F_u - \langle x^*, \cdot \rangle - \langle y_u^*, \cdot \rangle \right),$$

and

$$B_{(u, y_u^*)}(x^*) := B_{(u, y_u^*)}^0(x^*) = \left\{ x \in J(u) : (x, 0_u) \in (MF_u)(x^*, y_u^*) \right\}. \quad (5.2)$$

**Theorem 5.2 (Min-max robust duality)** *Let  $x^* \in X^*$  be such that  $(Mp)(x^*) \neq \emptyset$ . The next statements are equivalent:*

- (i)  $\min (\text{RP})_{x^*} = \max (\text{ODP})_{x^*}$ ,
  - (ii)  $\exists u \in U, \exists y_u^* \in Y_u^* : (Mp)(x^*) = B_{(u, y_u^*)}(x^*)$ ,
- where  $B_{(u, y_u^*)}(x^*)$  is the set defined in (5.2).

*Proof.* By Theorem 4.1 we know that [(i)  $\implies$  (ii)]. We now prove that [(ii)  $\implies$  (i)]. Pick  $x \in (Mp)(x^*)$  which is non-empty by assumption. Then by (ii),  $x \in B_{(u, y_u^*)}(x^*)$ , which yields  $x \in J(u)$  and  $(x, 0_u) \in (MF_u)(x^*, y_u^*)$ . Hence,

$$\begin{aligned} q(x^*) &\leq F_u^*(x^*, y_u^*) \leq \langle x^*, x \rangle - F_u(x, 0_u) \\ &\leq \langle x^*, x \rangle - p(x) \leq p^*(x^*) \leq q(x^*), \end{aligned}$$

which means that  $q(x^*) = F_u^*(x^*, y_u^*) = \langle x^*, x \rangle - p(x) = p^*(x^*)$  and (i) follows.  $\square$

**Corollary 5.3 (Min-max robust duality for Case 1)** *Let  $F : X \times Y \rightarrow \mathbb{R}_\infty$ ,  $x^* \in X^*$ ,  $p = F(\cdot, 0_Y)$ , and for each  $y^* \in Y^*$ ,*

$$B_{y^*}(x^*) := \left\{ x \in X : (x, 0_Y) \in (MF)(x^*, y^*) \right\}.$$

*Assume that  $(Mp)(x^*) \neq \emptyset$ . The next statements are equivalent:*

- (i)  $\min_{x \in X} \{ F(x, 0_Y) - \langle x^*, x \rangle \} = \max_{y^* \in Y^*} -F^*(x^*, y^*)$ ,
- (ii)  $\exists y^* \in Y^* : (Mp)(x^*) = B_{y^*}(x^*)$ .

**Corollary 5.4 (Min-max robust duality for Case 2)** *Let  $(f_u)_{u \in U} \subset \mathbb{R}_\infty^X$ ,  $p = \sup_{u \in U} f_u$ ,  $x^* \in X^*$ , and for each  $u \in U$ ,*

$$B_u(x^*) := J(u) \cap (Mf_u)(x^*),$$

*where  $J(u) = \{ x \in p^{-1}(\mathbb{R}) : f_u(x) = p(x) \}$ . Assume that  $(Mp)(x^*) \neq \emptyset$ . Then the next statements are equivalent:*

- (i)  $\left( \sup_{u \in U} f_u \right)^*(x^*) = \min_{u \in U} f_u^*(x^*)$ , with attainment at the first member,
- (ii)  $\exists u \in U : (Mp)(x^*) = B_u(x^*)$ .

## 6 Stable robust duality

Let us first recall some notations. Given  $F_u : X \times Y_u \rightarrow \mathbb{R}_\infty$ ,  $u \in U$ ,  $p = \sup_{u \in U} F_u(\cdot, 0_u)$  and  $q = \inf_{\substack{u \in U \\ y_u^* \in Y_u^*}} F_u^*(\cdot, y_u^*)$ . Remember that  $p^*(x^*) \leq q(x^*)$  for each  $x^* \in X^*$ . *Stable robust duality* means that  $\inf (\text{RP})_{x^*} = \sup (\text{ODP})_{x^*}$  for all  $x^* \in X^*$ , or equivalently,

$$p^*(x^*) = q(x^*), \quad \forall x^* \in X^*.$$

Theorem 3.1 says that, if  $\text{dom } p \neq \emptyset$ , then stable robust duality holds if and only if for each  $\varepsilon \geq 0$  the set-valued mappings  $M^\varepsilon p$ ,  $\mathcal{A}^\varepsilon : X^* \rightrightarrows X$  coincide, where, for each  $x^* \in X^*$ ,

$$\begin{aligned} (M^\varepsilon p)(x^*) &:= \varepsilon - \text{argmin}(p - x^*), \\ \mathcal{A}^\varepsilon(x^*) &:= \bigcap_{\eta > 0} \bigcup_{\substack{u \in U \\ y_u^* \in Y_u^*}} \bigcup_{\substack{\varepsilon_1 + \varepsilon_2 = \varepsilon + \eta \\ \varepsilon_1 \geq 0, \varepsilon_2 \geq 0}} \left\{ x \in J^{\varepsilon_1}(u) : (x, 0_u) \in (M^{\varepsilon_2} F_u)(x^*, y_u^*) \right\}. \end{aligned}$$

Consequently, stable robust duality holds if and only if for each  $\varepsilon \geq 0$ , the inverse set-valued mappings

$$(M^\varepsilon p)^{-1}, (\mathcal{A}^\varepsilon)^{-1} : X \rightrightarrows X^*,$$

coincide. Recall that  $(M^\varepsilon p)^{-1}$  is nothing but the  $\varepsilon$ -subdifferential of  $p$  at  $x$ .

Let us now make explicit  $(\mathcal{A}^\varepsilon)^{-1}$ . To this end we need to introduce for each  $\varepsilon \geq 0$  the ( $\varepsilon$ -active indexes) set-valued mapping  $I^\varepsilon : X \rightrightarrows U$  with

$$I^\varepsilon(x) = \begin{cases} \left\{ u \in U : F_u(x, 0_u) \geq p(x) - \varepsilon \right\}, & \text{if } p(x) \in \mathbb{R}, \\ \emptyset, & \text{if } p(x) \notin \mathbb{R}. \end{cases} \quad (6.1)$$

We observe that  $I^\varepsilon$  is nothing but the inverse of the set-valued mapping  $J^\varepsilon : U \rightrightarrows X$  defined in (3.9).

**Lemma 6.1** *For each  $(\varepsilon, x) \in \mathbb{R}_+ \times X$  one has*

$$(\mathcal{A}^\varepsilon)^{-1}(x) = \bigcap_{\eta > 0} \bigcup_{\substack{\varepsilon_1 + \varepsilon_2 = \varepsilon + \eta \\ \varepsilon_1 \geq 0, \varepsilon_2 \geq 0}} \bigcup_{u \in I^{\varepsilon_1}(x)} \text{proj}_{X^*}^u \partial^{\varepsilon_2} F_u(x, 0_u),$$

where  $\text{proj}_{X^*}^u : X^* \times Y_u^* \rightarrow X^*$  is the projection mapping  $\text{proj}_{X^*}^u(x^*, y_u^*) = x^*$ .

*Proof.* Let  $(\varepsilon, x, x^*) \in \mathbb{R}_+ \times X \times X^*$ . One has

$$\begin{aligned}
x^* \in (\mathcal{A}^\varepsilon)^{-1}(x) &\Leftrightarrow x \in \mathcal{A}^\varepsilon(x^*) \\
&\Leftrightarrow \begin{cases} \forall \eta > 0, \exists u \in U, \exists y_u^* \in Y_u^*, \exists (\varepsilon_1, \varepsilon_2) \in \mathbb{R}_+^2 \text{ such that} \\ \varepsilon_1 + \varepsilon_2 = \varepsilon + \eta, x \in J^{\varepsilon_1}(u) \text{ and } (x, 0_u) \in (M^{\varepsilon_2} F_u)(x^*, y_u^*) \end{cases} \\
&\Leftrightarrow \begin{cases} \forall \eta > 0, \exists u \in U, \exists y_u^* \in Y_u^*, \exists (\varepsilon_1, \varepsilon_2) \in \mathbb{R}_+^2 \text{ such that} \\ \varepsilon_1 + \varepsilon_2 = \varepsilon + \eta, u \in I^{\varepsilon_1}(x), \text{ and } (x^*, y_u^*) \in (\partial^{\varepsilon_2} F_u)(x, 0_u) \end{cases} \\
&\Leftrightarrow \begin{cases} \forall \eta > 0, \exists u \in U, \exists (\varepsilon_1, \varepsilon_2) \in \mathbb{R}_+^2 \text{ such that} \\ \varepsilon_1 + \varepsilon_2 = \varepsilon + \eta, u \in I^{\varepsilon_1}(x), \text{ and } x^* \in \text{proj}_{X^*}^u(\partial^{\varepsilon_2} F_u)(x, 0_u) \end{cases} \\
&\Leftrightarrow x^* \in \bigcap_{\eta > 0} \bigcup_{\substack{\varepsilon_1 + \varepsilon_2 = \varepsilon + \eta \\ \varepsilon_1 \geq 0, \varepsilon_2 \geq 0}} \bigcup_{u \in I^{\varepsilon_1}(x)} \text{proj}_{X^*}^u(\partial^{\varepsilon_2} F_u)(x, 0_u).
\end{aligned}$$

□

Now, for each  $(\varepsilon, x) \in \mathbb{R}_+ \times X$ , let us set

$$C^\varepsilon(x) := \bigcap_{\eta > 0} \bigcup_{\substack{\varepsilon_1 + \varepsilon_2 = \varepsilon + \eta \\ \varepsilon_1 \geq 0, \varepsilon_2 \geq 0}} \bigcup_{u \in I^{\varepsilon_1}(x)} \text{proj}_{X^*}^u(\partial^{\varepsilon_2} F_u)(x, 0_u). \quad (6.2)$$

Applying Theorem 3.1 and Lemma 6.1 we obtain:

**Theorem 6.1 (Stable robust duality)** *Assume that  $\text{dom } p \neq \emptyset$ . The next statements are equivalent:*

- (i)  $\inf (\text{RP})_{x^*} = \sup (\text{ODP})_{x^*}$  for all  $x^* \in X^*$ ,
- (ii)  $\partial^\varepsilon p(x) = C^\varepsilon(x)$ ,  $\forall (\varepsilon, x) \in \mathbb{R}_+ \times X$ ,
- (iii)  $\exists \bar{\varepsilon} > 0$ :  $\partial^\varepsilon p(x) = C^\varepsilon(x)$ ,  $\forall (\varepsilon, x) \in ]0, \bar{\varepsilon}[ \times X$ .

**Corollary 6.1 (Stable robust duality for Case 1)** *Let  $F : X \times Y \rightarrow \mathbb{R}_\infty$  be such that  $\text{dom } F(\cdot, 0_Y) \neq \emptyset$ . Let  $\text{proj}_{X^*} : X^* \times Y^* \rightarrow X^*$  be the projection mapping  $\text{proj}_{X^*}(x^*, y^*) = x^*$ . Then, the next statements are equivalent:*

- (i)  $\inf_{x \in X} \left\{ F(x, 0_Y) - \langle x^*, x \rangle \right\} = \sup_{y^* \in Y^*} -F^*(x^*, y^*)$ ,  $\forall x^* \in X^*$ ,
- (ii)  $(\partial^\varepsilon p)(x) = \bigcap_{\eta > 0} \text{proj}_{X^*}(\partial^\varepsilon F)(x, 0_Y)$ ,  $\forall (\varepsilon, x) \in \mathbb{R}_+ \times X$ ,
- (iii)  $\exists \bar{\varepsilon} > 0$ :  $(\partial^\varepsilon p)(x) = \bigcap_{\eta > 0} \text{proj}_{X^*}(\partial^\varepsilon F)(x, 0_Y)$ ,  $\forall (\varepsilon, x) \in ]0, \bar{\varepsilon}[ \times X$ .

*Proof.* Let  $U = \{u_0\}$  and  $F = F_{u_0} : X \times Y \rightarrow \mathbb{R}_\infty$ ,  $Y = Y_{u_0}$ ,  $p = F(\cdot, 0_Y)$ . Then for each  $(\varepsilon, x) \in \mathbb{R}_+ \times X$ ,

$$I^\varepsilon(x) = \begin{cases} \{u_0\}, & \text{if } p(x) \in \mathbb{R}, \\ \emptyset, & \text{if } p(x) \notin \mathbb{R}, \end{cases}$$

and

$$\bigcup_{\substack{\varepsilon_1 + \varepsilon_2 = \varepsilon \\ \varepsilon_1 \geq 0, \varepsilon_2 \geq 0}} \bigcup_{u \in I^{\varepsilon_1}(x)} \text{proj}_{X^*}^u(\partial^{\varepsilon_2} F_u)(x, 0_u) = \text{proj}_{X^*}(\partial^\varepsilon F)(x, 0_Y). \quad (6.3)$$

The conclusion now follows from (6.2)-(6.3) and Theorem 6.1. □

**Remark 6.1** Condition (ii) in Corollary 6.1 was quoted in [13, Theorem 4.3] for all  $(\varepsilon, x) \in ]0, +\infty[ \times \mathbb{R}$ , which is equivalent.

**Corollary 6.2 (Stable robust duality for Case 2)** Let  $(f_u)_{u \in U} \subset \mathbb{R}_\infty^X$ ,  $p = \sup_{u \in U} f_u$  and assume that  $\text{dom } p \neq \emptyset$ . The next statements are equivalent:

- (i)  $\left( \sup_{u \in U} f_u \right)^*(x^*) = \inf_{u \in U} f_u^*(x^*)$ ,  $\forall x^* \in X^*$ ,
  - (ii)  $(\partial^\varepsilon p)(x) = C^\varepsilon(x)$ ,  $\forall (\varepsilon, x) \in \mathbb{R}_+ \times X$ ,
  - (iii)  $\exists \bar{\varepsilon} > 0$ :  $(\partial^\varepsilon p)(x) = C^\varepsilon(x)$ ,  $\forall (\varepsilon, x) \in ]0, \bar{\varepsilon}[ \times X$ ,
- where  $C^\varepsilon(x)$  is the set

$$C^\varepsilon(x) = \bigcap_{\eta > 0} \bigcup_{\substack{\varepsilon_1 + \varepsilon_2 = \varepsilon + \eta \\ \varepsilon_1 \geq 0, \varepsilon_2 \geq 0}} \bigcup_{u \in I^{\varepsilon_1}(x)} (\partial^{\varepsilon_2} f_u)(x), \quad \forall (\varepsilon, x) \in \mathbb{R}_+ \times X. \quad (6.4)$$

*Proof.* Let  $F_u : X \times Y_u \rightarrow \mathbb{R}_\infty$  be such that  $F_u(x, y_u) = f_u(x)$  for all  $u \in U$ . Then for any  $(\varepsilon, x) \in \mathbb{R}_+ \times X$ ,

$$I^\varepsilon(x) = \begin{cases} \left\{ u \in U : f_u(x) \geq p(x) - \varepsilon \right\}, & \text{if } p(x) \in \mathbb{R}, \\ \emptyset, & \text{if } p(x) \notin \mathbb{R}, \end{cases}$$

$$(\partial^\varepsilon F_u)(x, 0_u) = (\partial^\varepsilon f_u)(x) \times \{0_u^*\}, \quad \forall (u, \varepsilon, x) \in U \times \mathbb{R}_+ \times X,$$

and  $C^\varepsilon(x)$  reads as in (6.4). The conclusion now follows from Theorem 6.1.  $\square$

## 7 Stable strong robust duality

We retain all the notations used in the Sections 3-6. Given  $(\varepsilon, u) \in \mathbb{R}_+ \times U$  and  $y_u^* \in Y_U^*$  we have introduced in Section 4 the set-valued mapping  $B_{(u, y_u^*)}^\varepsilon : X^* \rightrightarrows X$  defined by

$$B_{(u, y_u^*)}^\varepsilon(x^*) = \bigcup_{\substack{\varepsilon_1 + \varepsilon_2 = \varepsilon \\ \varepsilon_1 \geq 0, \varepsilon_2 \geq 0}} \left\{ x \in J^{\varepsilon_1}(u) : (x, 0_u) \in (M^{\varepsilon_2} F_u)(x^*, y_u^*) \right\}.$$

Let us now define  $B^\varepsilon : X^* \rightrightarrows X$  by setting

$$B^\varepsilon(x^*) := \bigcup_{\substack{u \in U \\ y_u^* \in Y_u^*}} B_{(u, y_u^*)}^\varepsilon(x^*), \quad \forall x^* \in X^*.$$

**Lemma 7.1** For each  $(\varepsilon, x) \in \mathbb{R}_+ \times X$  we have

$$(B^\varepsilon)^{-1}(x) = \bigcup_{\substack{\varepsilon_1 + \varepsilon_2 = \varepsilon \\ \varepsilon_1 \geq 0, \varepsilon_2 \geq 0}} \bigcup_{u \in I^{\varepsilon_1}(x)} \text{proj}_{X^*}^u(\partial^{\varepsilon_2} F_u)(x, 0_u).$$

*Proof.*  $x^* \in (B^\varepsilon)^{-1}(x)$  means that there exist  $u \in U$ ,  $y_u^* \in Y_u^*$   $\varepsilon_1 \geq 0$ ,  $\varepsilon_2 \geq 0$ , such that  $\varepsilon_1 + \varepsilon_2 = \varepsilon$ ,  $x \in J^{\varepsilon_1}(u)$ , and  $(x, 0_u) \in (M^{\varepsilon_2}F_u)(x^*, y_u^*)$ , or, equivalently,  $u \in I^{\varepsilon_1}(x)$ , and  $(x^*, y_u^*) \in (\partial^{\varepsilon_2}F_u)(x, 0_u)$ . In other words, there exist  $u \in U$ ,  $y_u^* \in Y_u^*$  such that  $x \in B_{(u, y_u^*)}^\varepsilon(x^*)$ , that is  $x \in B^\varepsilon(x^*)$ .  $\square$

For each  $\varepsilon \geq 0$  let us introduce the set-valued mapping  $D^\varepsilon := (B^\varepsilon)^{-1}$ . Now Lemma 7.1 writes:

$$D^\varepsilon(x) = \bigcup_{\substack{\varepsilon_1 + \varepsilon_2 = \varepsilon \\ \varepsilon_1 \geq 0, \varepsilon_2 \geq 0}} \bigcup_{u \in I^{\varepsilon_1}(x)} \text{proj}_{X^*}^u(\partial^{\varepsilon_2}F_u)(x, 0_u), \quad \forall (\varepsilon, x) \in \mathbb{R}_+ \times X. \quad (7.1)$$

Note that

$$C^\varepsilon(x) = \bigcap_{\eta > 0} D^{\varepsilon + \eta}(x), \quad \forall (\varepsilon, x) \in \mathbb{R}_+ \times X, \quad (7.2)$$

and that  $D^\varepsilon(x) = \emptyset$  whenever  $p(x) \notin \mathbb{R}$ .

We now provide a characterization of stable strong robust duality in terms of  $\varepsilon$ -subdifferential formulas.

**Theorem 7.1 (Stable strong robust duality)** *Assume that  $\text{dom } p \neq \emptyset$ , and let  $D^\varepsilon$  as in (7.1). The next statements are equivalent:*

- (i)  $\inf(\text{RP})_{x^*} = \max(\text{ODP})_{x^*} = \max_{\substack{u \in U \\ y_u^* \in Y_u^*}} -F_u^*(x^*, y_u^*), \quad \forall x^* \in X^*$ ,
- (ii)  $\partial^\varepsilon p(x) = D^\varepsilon(x), \quad \forall (\varepsilon, x) \in \mathbb{R}_+ \times X$ .

*Proof.* [(i)  $\implies$  (ii)] Let  $x^* \in \partial^\varepsilon p(x)$ . Then  $x \in (M^\varepsilon p)(x^*)$ . Since strong robust duality holds at  $x^*$ , Theorem 4.1 says that there exist  $u \in U$ ,  $y_u^* \in Y_u^*$  such that  $x \in B_{(u, y_u^*)}^\varepsilon(x^*) \subset B^\varepsilon(x^*)$ , and finally  $x^* \in D^\varepsilon(x)$  by Lemma 7.1. Thus  $\partial^\varepsilon p(x) \subset D^\varepsilon(x)$ .

Now, let  $x^* \in D^\varepsilon(x)$ . By Lemma 7.1 we have  $x \in B^\varepsilon(x^*)$  and there exist  $u \in U$ ,  $y_u^* \in Y_u^*$  such that  $x \in B_{(u, y_u^*)}^\varepsilon(x^*)$ . By Lemma 3.2 and the definition of  $B_{(u, y_u^*)}^\varepsilon(x^*)$  we have  $x \in S^\varepsilon(x^*)$ , and, by (3.8),  $x \in (M^\varepsilon p)(x^*)$  which means that  $x^* \in \partial^\varepsilon p(x)$ , and hence,  $D^\varepsilon(x) \subset \partial^\varepsilon p(x)$ . Thus (ii) follows.

[(ii)  $\implies$  (i)] If  $p^*(x^*) = +\infty$  then  $q(x^*) = +\infty$  and one has  $p^*(x^*) = F_u^*(x^*, y_u^*) = +\infty$  for all  $u \in U$ ,  $y_u^* \in Y_u^*$ , and (i) holds. Assume that  $p^*(x^*) \in \mathbb{R}$  and pick  $x \in p^{-1}(\mathbb{R})$  which is non-empty as  $\text{dom } p \neq \emptyset$  and  $p^*(x^*) \in \mathbb{R}$ . Let  $\varepsilon := p(x) + p^*(x^*) - \langle x^*, x \rangle$ . Then  $\varepsilon \geq 0$  and we have  $x^* \in \partial^\varepsilon p(x)$ . By (ii)  $x \in D^\varepsilon(x)$  and hence, there exist  $\varepsilon_1 \geq 0$ ,  $\varepsilon_2 \geq 0$ ,  $u \in U$ , and  $y_u^* \in Y_u^*$  such that  $\varepsilon_1 + \varepsilon_2 = \varepsilon$ ,  $u \in I^{\varepsilon_1}(x)$ ,  $(x^*, y_u^*) \in (\partial^{\varepsilon_2}F_u)(x, 0_u)$ . We have

$$\begin{aligned} q(x^*) &\leq F_u^*(x^*, y_u^*) \leq \langle x^*, x \rangle - F_u(x, 0_u) + \varepsilon_2 \\ &\leq \langle x^*, x \rangle - p(x) + \varepsilon_1 + \varepsilon_2 = p^*(x^*) \quad (\text{by definition of } \varepsilon) \\ &\leq q(x^*), \end{aligned}$$

and finally,  $q(x^*) = F_u^*(x^*, y_u^*) = p^*(x^*)$ , which is (i).  $\square$

Next, as usual, we give two consequences of Theorem 7.1 for the non-uncertainty and non-parametric cases.



**Corollary 7.1 (Stable strong duality for Case 1)** *Let  $F : X \times Y \rightarrow \mathbb{R}_\infty$ ,  $p = F(\cdot, 0_Y)$ ,  $\text{dom } p \neq \emptyset$ . The next statements are equivalent:*

- (i)  $\inf_{x \in X} \left\{ F(x, 0_Y) - \langle x^*, x \rangle \right\} = \max_{y^* \in Y^*} -F^*(x^*, y^*)$ ,  $\forall x^* \in X^*$ ,
- (ii)  $\partial^\varepsilon p(x) = \text{proj}_{X^*}(\partial^\varepsilon F)(x, 0_y)$ ,  $\forall (\varepsilon, x) \in \mathbb{R}_+ \times X$ .

*Proof.* This is the non-uncertainty case (i.e., the uncertainty set is a singleton) of the general problem  $(\text{RP})_{x^*}$ , with  $U = \{u_0\}$  and  $F_{u_0} = F : X \times Y \rightarrow \mathbb{R}_\infty$ . We have from (6.3),

$$D^\varepsilon(x) = \text{proj}_{X^*}(\partial^\varepsilon F)(x, 0_Y), \quad \forall (\varepsilon, x) \in \mathbb{R}_+ \times X. \quad (7.3)$$

The conclusion now follows from Theorem 7.1.  $\square$

**Corollary 7.2 (Stable strong duality for Case 2)** *Let  $(f_u)_{u \in U} \subset \mathbb{R}_\infty^X$ ,  $p = \sup_{u \in U} f_u$ , and  $\text{dom } p \neq \emptyset$ . The next statements are equivalent:*

- (i)  $(\sup_{u \in U} f_u)^*(x^*) = \min_{u \in U} f_u^*(x^*)$ ,  $\forall x^* \in X^*$ ,
- (ii)  $\partial^\varepsilon p(x) = D^\varepsilon(x)$ ,  $\forall (\varepsilon, x) \in \mathbb{R}_+ \times X$ , where

$$D^\varepsilon(x) = \bigcup_{\substack{\varepsilon_1 + \varepsilon_2 = \varepsilon \\ \varepsilon_1 \geq 0, \varepsilon_2 \geq 0}} \bigcup_{u \in I^{\varepsilon_1}(x)} (\partial^{\varepsilon_2} f_u)(x), \quad \forall (\varepsilon, x) \in \mathbb{R}_+ \times X, \quad (7.4)$$

and

$$I^\varepsilon(x) = \begin{cases} \left\{ u \in U : f_u(x) \geq p(x) - \varepsilon \right\} & \text{if } p(x) \in \mathbb{R}, \\ \emptyset & \text{if } p(x) \notin \mathbb{R}. \end{cases}$$

*Proof.* In this non-parametric situation, let  $F_u(x, y_u) = f_u(x)$ . It is easy to see that in this case, the set  $D^\varepsilon(x)$  can be expressed as in (7.4), and the conclusion follows from Theorem 7.1.  $\square$

## 8 Exact subdifferential formulas: Robust Basic Qualification condition

Given  $F_u : X \times Y_u \rightarrow \mathbb{R}_\infty$ ,  $u \in U$ , as usual, we let  $p = \sup_{u \in U} F_u(\cdot, 0_u)$ ,  $q := \inf_{(u, y_u^*) \in \Delta} F_u^*(\cdot, y_u^*)$ .

Again, we consider the robust problem  $(\text{RP})_{x^*}$  and its robust dual problem  $(\text{ODP})_{x^*}$  given in (3.1) and (3.2), respectively. Note that the reverse strong robust duality holds at  $x^*$  means that, for some  $\bar{x} \in X$ , it holds:

$$-p^*(x^*) = \min(\text{RP})_{x^*} = \sup_{u \in U} F_u(\bar{x}, 0_u) - \langle x^*, \bar{x} \rangle = p(\bar{x}) - \langle x^*, \bar{x} \rangle = \sup(\text{ODP})_{x^*} = -q(x^*). \quad (8.1)$$

Now, let us set, for each  $x \in X$ ,

$$D(x) := D^0(x) = \bigcup_{u \in I(x)} \text{proj}_{X^*}^u(\partial F_u)(x, 0_u), \quad (8.2)$$

$$C(x) := C^0(x) = \bigcap_{\eta > 0} \bigcup_{\substack{\varepsilon_1 + \varepsilon_2 = \eta \\ \varepsilon_1 \geq 0, \varepsilon_2 \geq 0}} \bigcup_{u \in I^{\varepsilon_1}(x)} \text{proj}_{X^*}^u(\partial^{\varepsilon_2} F_u)(x, 0_u), \quad (8.3)$$

where  $I^{\varepsilon_1}(x)$  is defined as in (6.1) and

$$I(x) := \begin{cases} \{u \in U : F_u(x, 0_u) = p(x)\}, & \text{if } p(x) \in \mathbb{R}, \\ \emptyset, & \text{if } p(x) \notin \mathbb{R}. \end{cases} \quad (8.4)$$

**Lemma 8.1** *For each  $x \in X$ , it holds*

$$D(x) \subset C(x) \subset \partial p(x).$$

*Proof.* The first inclusion is easy to check. Now let  $x^* \in C(x)$ . For each  $\eta > 0$  there exist  $(\varepsilon_1, \varepsilon_2) \in \mathbb{R}_+^2$ ,  $u \in I^{\varepsilon_1}(x)$ , and  $y_u^* \in Y_u^*$  such that  $\varepsilon_1 + \varepsilon_2 = \eta$  and  $(x^*, y_u^*) \in (\partial^{\varepsilon_2} F_u)(x, 0_u)$ . We then have  $F_u^*(x^*, y_u^*) + F_u(x, 0_u) - \langle x^*, x \rangle \leq \varepsilon_2$ ,  $p(x) \leq F_u(x, 0_u) + \varepsilon_1$  (as  $u \in I^{\varepsilon_1}(x)$ ), and  $p^*(x^*) \leq q(x^*) \leq F_u^*(x^*, y_u^*)$ . Consequently,

$$p^*(x^*) + p(x) - \langle x^*, x \rangle \leq F_u^*(x^*, y_u^*) + F_u(x, 0_u) + \varepsilon_1 - \langle x^*, x \rangle \leq \varepsilon_1 + \varepsilon_2 = \eta.$$

Since  $\eta > 0$  is arbitrary we get  $p^*(x^*) + p(x) - \langle x^*, x \rangle \leq 0$ , which means that  $x^* \in \partial p(x)$ . The proof is complete.  $\square$

**Theorem 8.1** *Let  $x \in p^{-1}(\mathbb{R})$  and  $C(x)$  be as in (8.3). The next statements are equivalent:*

- (i)  $\partial p(x) = C(x)$ ,
- (ii) *Reverse strong robust duality holds at each  $x^* \in \partial p(x)$ ,*
- (iii) *Robust duality holds at each  $x^* \in \partial p(x)$ .*

*Proof.* [(i)  $\implies$  (ii)] Let  $x^* \in \partial p(x)$ . We have  $x^* \in C(x) = (\mathcal{A})^{-1}(x)$  (see Lemma 6.1 with  $\varepsilon = 0$ ). Then  $x \in \mathcal{A}(x^*) = S(x^*)$  (see (3.12) with  $\varepsilon = 0$ ), and therefore,

$$\begin{aligned} -p^*(x^*) &\leq p(x) - \langle x^*, x \rangle \leq -q(x^*) \leq -p^*(x^*), \\ -p^*(x^*) &= \min_{z \in X} \{p(z) - \langle x^*, z \rangle\} = p(x) - \langle x^*, x \rangle = -q(x^*), \end{aligned}$$

that means that reverse strong robust duality holds at  $x^*$  (see (8.1)).

[(ii)  $\implies$  (iii)] is obvious.

[(iii)  $\implies$  (i)] By Lemma 8.1 it suffices to check that the inclusion “ $\subset$ ” holds. Let  $x^* \in \partial p(x)$ . We have  $x \in (Mp)(x^*)$ . Since robust duality holds at  $x^*$ , Theorem 3.1 (with  $\varepsilon = 0$ ) says that  $x \in \mathcal{A}(x^*)$ . Thus,  $x^* \in \mathcal{A}^{-1}(x)$ , and, by Lemma 6.1,  $x^* \in C(x)$ .  $\square$

In the deterministic and the non-parametric cases, we get the next results from Theorem 8.1.

**Corollary 8.1** Let  $F : X \times Y \rightarrow \mathbb{R}_\infty$ ,  $p = F(\cdot, 0_Y)$ , and  $x \in p^{-1}(\mathbb{R})$ . The next statements are equivalent:

- (i)  $\partial p(x) = \bigcap_{\eta > 0} \text{proj}_{X^*}(\partial^\eta F)(x, 0_Y)$ ,
- (ii)  $\min_{z \in X} \left\{ F(z, 0_Y) - \langle x^*, z \rangle \right\} = \sup_{y^* \in Y^*} -F^*(x^*, y^*)$ ,  $\forall x^* \in \partial p(x)$ ,
- (iii)  $\inf_{z \in X} \left\{ F(z, 0_Y) - \langle x^*, z \rangle \right\} = \sup_{y^* \in Y^*} -F^*(x^*, y^*)$ ,  $\forall x^* \in \partial p(x)$ .

*Proof.* Let  $F_u = F : X \times Y \rightarrow \mathbb{R}_\infty$  and  $p = F(\cdot, 0_Y)$ . We then have

$$C(x) = \bigcap_{\eta > 0} \text{proj}_{X^*}(\partial^\eta F)(x, 0_Y), \quad \forall x \in X,$$

(see Corollary 6.1) and the conclusion follows directly from Theorem 8.1.  $\square$

**Corollary 8.2** Let  $(f_u)_{u \in U} \subset \mathbb{R}_\infty^X$ ,  $p = \sup_{u \in U} f_u$ ,  $x \in p^{-1}(\mathbb{R})$ . The next statements are equivalent:

- (i)  $\partial \left( \sup_{u \in U} f_u \right) (x) = C(x)$ ,
- (ii)  $\max_{z \in X} \left\{ \langle x^*, z \rangle - p(z) \right\} = \inf_{u \in U} f_u^*(x^*)$ ,  $\forall x^* \in \partial p(x)$ ,
- (iii)  $\left( \sup_{u \in U} f_u \right)^*(x^*) = \inf_{u \in U} f_u^*(x^*)$ ,  $\forall x^* \in \partial p(x)$ ,

where

$$C(x) = \bigcap_{\eta > 0} \bigcup_{\substack{\varepsilon_1 + \varepsilon_2 = \eta \\ \varepsilon_1 \geq 0, \varepsilon_2 \geq 0}} \bigcup_{u \in I^{\varepsilon_1}(x)} \text{proj}_{X^*}^u(\partial^{\varepsilon_2} f_u)(x), \quad \forall x \in X. \quad (8.5)$$

*Proof.* Let  $F_u(x, y_u) = f_u(x)$ . Then it is easy to see that in this case,  $C(x)$  can be expressed as in (8.5). The conclusion now follows from Theorem 8.1.  $\square$

Let us come back to the general case and consider the most simple subdifferential formula one can expect for the robust objective function  $p = \sup_{u \in U} F_u(\cdot, 0_u)$ :

$$\partial p(x) = \bigcup_{u \in I(x)} \text{proj}_{X^*}^u(\partial F_u)(x, 0_u), \quad (8.6)$$

where the set of active indexes at  $x$ ,  $I(x)$ , is defined by (8.4).

In Case 3 we have

$$p(x) = \begin{cases} f(x), & \text{if } H_u(x) \in -S_u, \forall u \in U, \\ +\infty, & \text{else,} \end{cases}$$

$I(x) = U$  for each  $x \in p^{-1}(\mathbb{R})$ , and (8.6) writes

$$\partial p(x) = \bigcup_{\substack{u \in U, z_u^* \in S_u^+ \\ \langle z_u^*, H_u(x) \rangle = 0}} \partial(f + z_u^* \circ H_u)(x),$$

which has been called *Basic Robust Subdifferential Condition* (BRSC) in [4] (see [14, page 307] for the deterministic case). More generally, let us introduce the following terminology:

**Definition 8.1** *Given  $F_u : X \times Y_u \rightarrow \mathbb{R}_\infty$  for each  $u \in U$ , and  $p = \sup_{u \in U} F_u(\cdot, 0_u)$ , we will say that **Basic Robust Subdifferential Condition holds at a point**  $x \in p^{-1}(\mathbb{R})$  if (8.6) is satisfied, that is  $\partial p(x) = D(x)$ .*

Recall that, in Example 2.1,  $p(x) = \langle c^*, x \rangle + i_A(x)$ , where  $A = p^{-1}(\mathbb{R})$  is the feasible set of the linear system. So, given  $x \in A$ ,  $\partial p(x)$  is the sum of  $c^*$  with the normal cone of  $A$  at  $x$ , i.e., Basic Robust Subdifferential Condition (at  $x$ ) asserts that such a cone can be expressed in some prescribed way.

**Theorem 8.2** *Let  $x \in p^{-1}(\mathbb{R})$ . The next statements are equivalent:*

- (i) *Basic Robust Subdifferential Condition holds at  $x$ ,*
- (ii) *Min-max robust duality holds at each  $x^* \in \partial p(x)$ ,*
- (iii) *Strong robust duality holds at each  $x^* \in \partial p(x)$ .*

*Proof.* [(i)  $\implies$  (ii)] Let  $x^* \in \partial p(x)$ . We have  $x^* \in D(x)$  and by (8.2) there exist  $u \in I(x)$  (i.e.,  $p(x) = F_u(x, 0_u)$ ),  $y_u^* \in Y_u^*$ , such that  $(x^*, y_u^*) \in \partial F_u(x, 0_u)$ . Then,

$$\begin{aligned} p^*(x^*) &\geq \langle x^*, x \rangle - p(x) = \langle x^*, x \rangle - F_u(x, 0_u) = F_u^*(x^*, y_u^*) \\ &\geq q(x^*) \geq p^*(x^*). \end{aligned}$$

It follows that

$$\max_{z \in X} \{ \langle x^*, z \rangle - p(z) \} = \langle x^*, x \rangle - p(x) = F_u^*(x^*, y_u^*) = q(x^*),$$

and min-max robust duality holds at  $x^*$ .

[(ii)  $\implies$  (iii)] is obvious.

[(iii)  $\implies$  (i)] By Lemma 8.1, it suffices to check that  $\partial p(x) \subset D(x)$ . Let  $x^* \in \partial p(x)$ . We have  $x \in (Mp)(x^*)$ . Since strong robust duality holds at  $x^*$ , Theorem 4.1 says that there exist  $u \in U$ ,  $y_u^* \in Y_u^*$  such that  $x \in B_{(u, y_u^*)}^0(x^*)$ , that means (see (5.2))

$$(x, 0_u) \in (MF_u)((x^*, y_u^*), (x^*, y_u^*) \in \partial F_u(x, 0_u),$$

and by (8.2),  $x^* \in D(x)$ . □

As usual, Theorem 8.2 gives us corresponding results for the two extreme cases: non-uncertainty and non-perturbation cases.

**Corollary 8.3** Let  $F : X \times Y \rightarrow \mathbb{R}_\infty$ ,  $p = F(\cdot, 0_Y)$ , and  $x \in p^{-1}(\mathbb{R})$ . The next statements are equivalent:

- (i)  $\partial p(x) = \text{proj}_{X^*}(\partial F)(x, 0_Y)$ ,
- (ii)  $\max_{z \in X} \left\{ \langle x^*, z \rangle - F(z, 0_Y) \right\} = \min_{y^* \in Y^*} F^*(x^*, y^*)$ ,  $\forall x^* \in \partial p(x)$ ,
- (iii)  $p^*(x^*) = \min_{y^* \in Y^*} F^*(x^*, y^*)$ ,  $\forall x^* \in \partial p(x)$ .

*Proof.* In this case we have, by (7.3),  $D(x) = \text{proj}_{X^*}(\partial F)(x, 0_Y)$  and the conclusion is a direct consequence of Theorem 8.2.  $\square$

**Corollary 8.4** Let  $(f_u)_{u \in U} \subset \mathbb{R}_\infty^X$ ,  $p = \sup_{u \in U} f_u$ ,  $x \in p^{-1}(\mathbb{R})$ . The next statements are equivalent:

- (i)  $\partial p(x) = \bigcup_{u \in I(x)} \partial f_u(x)$ ,
- (ii)  $\max_{z \in X} \left\{ \langle x^*, z \rangle - p(z) \right\} = \min_{u \in U} f_u^*(x^*)$ ,  $\forall x^* \in \partial p(x)$ ,
- (iii)  $(\sup_{u \in U} f_u)^*(x^*) = \min_{y^* \in Y^*} f_u^*(x^*)$ ,  $\forall x^* \in \partial p(x)$ .

*Proof.* In this non-parametric case, let  $F_u(x, y_u) = f_u(x)$ ,  $p = \sup_{u \in U} f_u$ . We have

$$D(x) = \bigcup_{u \in I(x)} \partial f_u(x), \quad I(x) = \{u \in U : f_u(x) = p(x) \in \mathbb{R}\}$$

and the Theorem 8.2 applies.  $\square$

**Acknowledgements** This research was supported by the National Foundation for Science & Technology Development (NAFOSTED) from Vietnam, Project 101.01-2015.27 *Generalizations of Farkas lemma with applications to optimization*, by the Ministry of Economy and Competitiveness of Spain and the European Regional Development Fund (ERDF) of the European Commission, Project MTM2014-59179-C2-1-P, and by the Australian Research Council, Project DP160100854.

## References

- [1] Ben-Tal, A., El Ghaoui, L., Nemirovski, A.: Robust Optimization, Princeton U.P., Princeton (2009)
- [2] Bertsimas, D., Sim, M.: Tractable approximations to robust conic optimization problems. *Math. Programming* **107B**, 5-36 (2006)
- [3] Bot, R.I.: Conjugate duality in convex optimization. Springer, Berlin (2010)
- [4] Bot, R.I., Jeyakumar, V., Li, G.Y.: Robust duality in parametric convex optimization. *Set-Valued Var. Anal.* **21**, 177-189 (2013)

- [5] Burachick, R.S., Jeyakumar, V., Wu, Z.-Y.: Necessary and sufficient condition for stable conjugate duality. *Nonlinear Anal.* **64**, 1998-2006 (2006)
- [6] Chu, Y.C.: Generalization of some fundamental theorems on linear inequalities. *Acta Math Sinica* **16**, 25-40 (1966)
- [7] Dinh, N., Goberna, M.A., López, M.A., Volle, M.: A unifying approach to robust convex infinite optimization duality. *J. Optim. Theory Appl.* **174**, 650-685 (2017)
- [8] Dinh, N., Mo, T.H., Vallet, G., Volle, M.: A unified approach to robust Farkas-type results with applications to robust optimization problems. *SIAM J. Optim.* **27**, 1075-1101 (2017)
- [9] Dinh, N., Long, D.H.: Complete characterizations of robust strong duality for robust optimization problems. *Vietnam J. Math.*, to appear. DOI: 10.1007/s10013-018-0283-1
- [10] Goberna, M.A., López, M.A.: *Linear Semi-Infinite Optimization*. J. Wiley, Chichester (1998)
- [11] Goberna, M.A., López, M.A.: Recent contributions to linear semi-infinite optimization. *4OR-Q. J. Oper. Res.* **15**, 221-264 (2017)
- [12] Goberna, M.A., López, M.A., Volle, M.: Primal attainment in convex infinite optimization duality. *J. Convex Anal.* **21**, 1043-1064 (2014)
- [13] Grad, S.-M.: Closedness type regularity conditions in convex optimization and beyond. *Front. Appl. Math. Stat.* **16**, September (2016) <https://doi.org/10.3389/fams.2016.00014>.
- [14] Hiriart-Urruty, J.-B., Lemarechal, C., *Convex Analysis and Minimization Algorithms I*, Springer, Berlin (1993).
- [15] Hiriart-Urruty, J.-B., Moussaoui, M., Seeger, A., Volle, M.: Subdifferential calculus without qualification conditions, using approximate subdifferentials: a survey. *Nonlinear Anal.* **24**, 1727-1754 (1995)
- [16] Jeyakumar, V., Li, G.Y.: Stable zero duality gaps in convex programming: complete dual characterisations with applications to semidefinite programs. *J. Math. Anal. Appl.* **360**, 156-167 (2009)
- [17] Li, G.Y., Jeyakuma, V., Lee, G.M.: Robust conjugate duality for convex optimization under uncertainty with application to data classification. *Nonlinear Anal.* **74**, 2327-2341 (2011)
- [18] Lindsey, M., Rubinstein, Y.A.: Optimal transport via a Monge-Ampère optimization problem. *SIAM J. Math. Anal.* **49**, 3073-3124 (2017)
- [19] López, M.A., Still, G.: Semi-infinite programming. *European J. Oper. Res.* **180**, 491-518 (2007)

- [20] Rockafellar, R.T.: Conjugate Duality and Optimization. CBMS Lecture Notes Series No. 162, SIAM (1974)
- [21] Vera, J.R.: Geometric measures of convex sets and bounds on problem sensitivity and robustness for conic linear optimization. Math. Programming **147A**, 47-79 (2014)
- [22] Volle, M.: Calculus rules for global approximate minima and applications to approximate subdifferential calculus. J. Global Optim. **5**, 131-157 (1994)
- [23] Wang, Y., Shi, R., Shi, J.: Duality and robust duality for special nonconvex homogeneous quadratic programming under certainty and uncertainty environment. J. Global Optim. **62**, 643-659 (2015)
- [24] C. Zălinescu, Convex Analysis in General Vector Spaces. World Scientific, River Edge, NJ (2002)