

REPRESENTATION THEORY

An Electronic Journal of the American Mathematical Society

Volume 20, Pages 39–93 (February 12, 2016)

<http://dx.doi.org/10.1090/ert/477>

PROJECTIVE NORMALITY OF MODEL VARIETIES AND RELATED RESULTS

PAOLO BRAVI, JACOPO GANDINI, AND ANDREA MAFFEI

ABSTRACT. We prove that the multiplication of sections of globally generated line bundles on a model wonderful variety M of simply connected type is always surjective. This follows by a general argument which works for every wonderful variety and reduces the study of the surjectivity for every couple of globally generated line bundles to a finite number of cases. As a consequence, the cone defined by a complete linear system over M or over a closed G -stable subvariety of M is normal. We apply these results to the study of the normality of the compactifications of model varieties in simple projective spaces and of the closures of the spherical nilpotent orbits. Then we focus on a particular case proving two specific conjectures of Adams, Huang and Vogan on an analogue of the model orbit of the group of type E_8 .

INTRODUCTION

Let G be a complex linear algebraic group, semisimple and simply connected. A G -variety M is called *wonderful* of rank n if it satisfies the following conditions:

- M is smooth and projective;
- M possesses an open orbit whose complement is a union of n smooth prime divisors (the *boundary divisors*) with non-empty transversal intersections;
- any orbit closure in M equals the intersection of the prime divisors which contain it.

Examples of wonderful varieties are the *flag varieties*, which are the wonderful varieties of rank zero, and the *complete symmetric varieties* introduced by C. De Concini and C. Procesi [20], which we rather call *adjoint symmetric wonderful varieties*. Wonderful varieties were then considered in full generality by D. Luna, who started a program of classification in terms of combinatorial invariants [35].

Consider the following.

Question. Let M be a wonderful variety and, for $\mathcal{L}, \mathcal{L}' \in \text{Pic}(M)$, consider the multiplication map

$$m_{\mathcal{L}, \mathcal{L}'} : \Gamma(M, \mathcal{L}) \otimes \Gamma(M, \mathcal{L}') \longrightarrow \Gamma(M, \mathcal{L} \otimes \mathcal{L}').$$

Is $m_{\mathcal{L}, \mathcal{L}'}$ surjective for all globally generated $\mathcal{L}, \mathcal{L}'$?

In the case of a flag variety, the answer to the previous question is affirmative, indeed by the Borel-Weil theorem $\Gamma(M, \mathcal{L})$ is a simple G -module for all $\mathcal{L} \in \text{Pic}(M)$. In the case of the wonderful compactification of an adjoint group (regarded as a symmetric $G \times G$ -variety) a still affirmative answer was obtained by S.S. Kannan [27] with a very explicit analysis. In the case of any adjoint symmetric wonderful

Received by the editors February 19, 2015 and, in revised form, December 29, 2015.

2010 *Mathematics Subject Classification.* Primary 14M27; Secondary 20G05.

variety the same was obtained by R. Chirivì and the third named author in [15] with an inductive argument. In general, when all the above-mentioned multiplication maps are surjective, it follows that the image of M in the dual projective space of the complete linear system associated to any globally generated line bundle is projectively normal.

Another remarkable class of wonderful varieties is that of the *model wonderful varieties*, introduced by Luna [36]. Given a central subgroup $\Gamma \subset G$, a *model homogeneous space* for the algebraic group $G_\Gamma := G/\Gamma$ is a quasi-affine homogeneous space G/H such that $\Gamma \subset H$ and the coordinate ring $\mathbb{C}[G/H]$ is a model of the representations of G_Γ in the sense of I.M. Gel'fand (see [4], [24], [25]), that is, $\mathbb{C}[G/H]$ is isomorphic as a G -module to the direct sum of all the irreducible representations of G_Γ . Main examples of model homogeneous spaces arise as nilpotent orbits for the adjoint action of G ; see also [16] where a standard monomial theory for some classical model homogeneous spaces was developed. The model homogeneous spaces for G_Γ were classified in [36], where Luna constructed a variety $M_{G_\Gamma}^{\text{mod}}$, which is wonderful for the action of G , whose orbits parametrize the model homogeneous spaces for G_Γ . Varieties of the shape $M_{G_\Gamma}^{\text{mod}}$ for some central subgroup $\Gamma \subset G$ are called model wonderful varieties. Given a model wonderful variety $M_{G_\Gamma}^{\text{mod}}$, we say that it is of *simply connected type* if $M_{G_\Gamma}^{\text{mod}}$ and M_G^{mod} are G -equivariantly isomorphic. For G almost simple (i.e., a non-commutative group having no proper closed connected normal subgroups), it follows by Luna's description that $M_{G_\Gamma}^{\text{mod}}$ is not of simply connected type if and only if $G_\Gamma = \text{SO}(2r+1)$ (in which case $G = \text{Spin}(2r+1)$ and $\Gamma \simeq \mathbb{Z}/2\mathbb{Z}$ is the center of G).

The first result of the paper is the following.

Theorem 1 (see Theorem 4.1). *Let M be a model wonderful variety of simply connected type. The multiplication of global sections*

$$m_{\mathcal{L}, \mathcal{L}'} : \Gamma(M, \mathcal{L}) \otimes \Gamma(M, \mathcal{L}') \longrightarrow \Gamma(M, \mathcal{L} \otimes \mathcal{L}')$$

is surjective for all globally generated line bundles $\mathcal{L}, \mathcal{L}'$ on M .

This is false if M is a model wonderful variety not of simply connected type. Indeed, in case $M = M_{\text{SO}(2r+1)}^{\text{mod}}$ the multiplication of global sections $m_{\mathcal{L}, \mathcal{L}'}$ is surjective for all globally generated line bundles $\mathcal{L}, \mathcal{L}'$ on M if and only if $r < 4$ (see Section 9.1). This is essentially a consequence of the fact that the tensor product of an almost simple group of type B_r does not satisfy the saturation property in the sense of A. Klyachko (see [31] and [33]).

Theorem 1 follows from a general argument which works for every wonderful variety M and reduces the surjectivity of the maps $m_{\mathcal{L}, \mathcal{L}'}$ for all globally generated line bundles $\mathcal{L}, \mathcal{L}'$ on M to the surjectivity for a finite number of couples. Indeed, we also prove similar theorems for other classes of wonderful varieties arising in some specific applications.

We now explain in more detail our reduction in the study of the surjectivity of the multiplication maps on a wonderful variety, and how we apply Theorem 1 and its analogues in different directions.

A general reduction. Fix a Borel subgroup $B \subset G$ and a maximal torus $T \subset B$. Denote by Σ the set of G -stable prime divisors of M and by Δ the set of B -stable prime divisors of M which are not G -stable: since M possesses an open B -orbit, Δ is a finite set. Then the Picard group $\text{Pic}(M)$ is freely generated by the line

bundles \mathcal{L}_D with $D \in \Delta$ and the free group $\mathbb{Z}\Sigma$ embeds in $\text{Pic}(M)$, so that we may regard $\mathbb{Z}\Sigma$ as a sublattice of $\mathbb{Z}\Delta$. Moreover, via the isomorphism $\text{Pic}(M) \simeq \mathbb{Z}\Delta$, the semigroup of globally generated line bundles is identified with the semigroup $\mathbb{N}\Delta$. We denote by \mathcal{L}_E the globally generated line bundle associated to an element $E \in \mathbb{N}\Delta$. Define the following partial order relation on $\mathbb{N}\Delta$:

$$E \leq_{\Sigma} F \quad \text{if} \quad E - F \in \mathbb{N}\Sigma.$$

This partial order is tightly related to the isotypic decomposition of the spaces of global sections of the globally generated line bundles on M , which we may always assume linearized (see Proposition 1.1).

In case M is the wonderful compactification of the adjoint group $G_{\text{ad}} = G/Z(G)$ regarded as a $G \times G$ -variety, then Σ is naturally identified with the basis of the root system of G , while Δ is naturally identified with the set of the fundamental weights of G . More generally, this is true whenever M is the wonderful compactification of a non-Hermitian symmetric space, in which case there always exists a root system Φ_{Σ} (the *reduced root system*) such that Σ is a basis of Φ_{Σ} and Δ is the corresponding set of fundamental weights. This is no longer true in the case of a general wonderful variety: while there always exists a root system $\Phi_{\Sigma} \subset \mathbb{Z}\Delta$ with Σ as set of simple roots, the fundamental weights of Φ_{Σ} associated to Σ may differ from Δ . Therefore we may think the couple (Σ, Δ) as a generalization of a root datum.

Suppose that $E, F \in \mathbb{N}\Delta$ are such that $E <_{\Sigma} F$ and there is no D with $E <_{\Sigma} D <_{\Sigma} F$: then we say that $F - E \in \mathbb{N}\Sigma$ is a *covering difference*. The set of the covering differences is finite and in the case of an usual root system it was studied by J.R. Stembridge in [41].

Given $E = \sum_{D \in \Delta} n_D D \in \mathbb{Z}\Delta$, define the *positive part* $E^+ = \sum_{n_D > 0} n_D D$ and the *height* $\text{ht}(E) = \sum_{D \in \Delta} n_D$. We prove the following.

Lemma 2 (see Lemma 2.3). *Let M be a wonderful variety and let n be such that $\text{ht}(\gamma^+) \leq n$ for every covering difference γ . If the multiplication map*

$$m_{\mathcal{L}_E, \mathcal{L}_F} : \Gamma(M, \mathcal{L}_E) \otimes \Gamma(M, \mathcal{L}_F) \longrightarrow \Gamma(M, \mathcal{L}_{E+F})$$

is surjective for all $E, F \in \mathbb{N}\Delta$ with $\text{ht}(E + F) \leq n$, then it is surjective for all $E, F \in \mathbb{N}\Delta$.

We use Lemma 2 to prove Theorem 1. We first study the covering relation in the case of a model wonderful variety proving that $\text{ht}(\gamma^+) \leq 2$ for all covering differences γ , then we study the multiplication maps $m_{\mathcal{L}_E, \mathcal{L}_F}$ in the fundamental cases $E, F \in \Delta$. To check the inclusions arising in the exceptional group cases we use the computer.

The fact that $\text{ht}(\gamma^+) \leq 2$ for all covering differences γ is an easy exercise in case the couple (Σ, Δ) corresponds to a root system, and as far as we know it could be a general fact which holds for all wonderful varieties.

Proceeding inductively on the partial order \leq_{Σ} , it is easy to reduce the surjectivity of the multiplication map $m_{\mathcal{L}_E, \mathcal{L}_F}$ for every $E, F \in \mathbb{N}\Delta$ to the fact that some special submodules of $\Gamma(M, \mathcal{L}_{E+F})$ occur in the image of $m_{\mathcal{L}_E, \mathcal{L}_F}$. This leads to the definition of *low triple* (see Definition 2.2), which was already introduced in [15] to treat the case of an adjoint symmetric wonderful variety. To prove Lemma 2 we show that it is possible to treat inductively (w.r.t. the height) the low triples of M .

Spherical orbits in simple projective spaces. Our first application of the surjectivity of the multiplication maps regards the study of the normality of the closure of a spherical orbit in a simple projective space, i.e., the projective space of a simple G -module. Recall that a G -variety X is called *spherical* if it is normal and possesses an open B -orbit, wonderful varieties form a very relevant class of spherical varieties and play a prominent role in their classification.

Let V be a simple G -module, let $G \cdot [v] \subset \mathbb{P}(V)$ be a spherical orbit and consider its closure $X = \overline{G \cdot [v]}$, which is not necessarily normal. Let H be the stabilizer of $[v]$, then the spherical homogeneous space G/H can always be embedded as the open orbit of a wonderful variety M , called the *wonderful compactification* of G/H , which dominates any other projective compactification of G/H with a unique closed orbit.

If the multiplication of global sections of globally generated line bundles on M is surjective, the normality of every orbit closure $X \subset \mathbb{P}(V)$ dominated by M can be reduced to the normality of some “fundamental” orbit closures. On the other hand, sometimes such fundamental orbit closures are easily shown to be normal. In particular, we have the following.

Theorem 3 (see Corollary 6.9). *Let V be a simple G -module, let $X \subset \mathbb{P}(V)$ be the closure of a spherical orbit and consider a wonderful variety M which dominates X . If M is symmetric with reduced root system of type A or model for a connected semisimple group of type AD , then X is normal.*

Spherical nilpotent orbits. Our second application regards the study of the normality of the closure of a spherical nilpotent orbit $\mathcal{O} \subset \mathfrak{g}$, where \mathfrak{g} denotes the Lie algebra of G . In particular, if we consider the projectivization of \mathcal{O} , we get again a spherical orbit $\mathcal{U} \subset \mathbb{P}(\mathfrak{g})$, and the closure $\overline{\mathcal{O}}$ coincides with the affine cone over $\overline{\mathcal{U}}$.

Following [13] and a suggestion of D. Luna, to study the normality of $\overline{\mathcal{O}}$ we may consider a wonderful variety $M_{\mathcal{O}}$, namely the wonderful compactification of \mathcal{U} . It turns out that the wonderful variety $M_{\mathcal{O}}$ is either symmetric or model, or closely related to one of these. If the surjectivity of the multiplication maps of $M_{\mathcal{O}}$ is known, the description of the G -module structure of the coordinate ring of $\overline{\mathcal{O}}$ is an easy task.

Therefore, we prove the following (with computer-aided computations in the exceptional group cases).

Theorem 4 (see Theorem 7.1). *Let $\mathcal{O} \subset \mathfrak{g}$ be a spherical nilpotent orbit and let $M_{\mathcal{O}}$ be the associated wonderful variety. Then the multiplication map*

$$m_{\mathcal{L}, \mathcal{L}'} : \Gamma(M_{\mathcal{O}}, \mathcal{L}) \otimes \Gamma(M_{\mathcal{O}}, \mathcal{L}') \longrightarrow \Gamma(M_{\mathcal{O}}, \mathcal{L} \otimes \mathcal{L}')$$

is surjective for all globally generated line bundles $\mathcal{L}, \mathcal{L}'$ on $M_{\mathcal{O}}$.

As a corollary (see Corollary 7.2) we obtain the classification of the spherical nilpotent adjoint orbits \mathcal{O} whose closure is normal (these results are already known; see for instance [18, Theorem 5.1] or [38, Table 2]). As a particular case, among these orbits we also find the model orbit of E_8 studied by J. Adams, J-S. Huang and D.A. Vogan Jr. in [1].

The real model orbit of type E_8 . Following [1], we also consider an analogue of the model nilpotent adjoint orbit \mathcal{O} of E_8 . More explicitly, we consider a K -orbit, where K (an algebraic group) is the complexification of a maximal compact

subgroup $K_{\mathbb{R}}$ of the split real form of E_8 . Then K is the fixed point subgroup of an involution of E_8 , this involution passes to the Lie algebra and K acts on the eigenspace \mathfrak{p} of eigenvalue -1 . The analogue of the model orbit that we consider is the intersection $\mathcal{O}_{\mathfrak{p}} = \mathcal{O} \cap \mathfrak{p}$, which is a K -orbit.

For the closure of $\mathcal{O}_{\mathfrak{p}}$ we prove its normality and we describe its coordinate ring (Theorem 8.6). Furthermore, we describe the space of $K_{\mathbb{R}}$ -finite vectors of the unitary representation of the split real form of E_8 that should be associated to this K -orbit via the so-called orbit method (Theorem 8.10). Both descriptions were already present in [1] as consequences of some conjectures, which as far as we know are still open.

In order to prove these theorems, we are led to consider another class of wonderful varieties, which we call *comodel wonderful varieties* since they are somewhat dual to the model wonderful varieties (see Theorem 5.1 for a precise definition). For the comodel wonderful varieties we also show that the multiplication of global sections of globally generated line bundles is surjective (see Theorem 5.2).

It turns out indeed that $\mathcal{O}_{\mathfrak{p}}$ is the cone over a homogeneous space whose wonderful compactification is closely related to a comodel wonderful variety.

Multiplication of functions on a spherical homogeneous space. Suppose that G/H is a spherical homogeneous space, in the last section of the paper we also consider the problem of multiplying G -modules of functions in the coordinate ring $\mathbb{C}[G/H]$. Indeed, $\mathbb{C}[G/H]$ is a multiplicity-free G -module (that is, every isotypical component is irreducible), and given two irreducible components V_{λ} and $V_{\lambda'}$ of highest weight λ and λ' it is well defined their product $V_{\lambda}V_{\lambda'} \subset \mathbb{C}[G/H]$, that is, the G -module generated by the products ff' with $f \in V_{\lambda}$ and $f' \in V_{\lambda'}$.

More generally, given $\xi \in \mathcal{X}(H)$, one may consider the corresponding G -module of the H -semi-invariant function

$$\mathbb{C}[G]_{\xi}^{(H)} = \{f \in \mathbb{C}[G] : f(gh) = \xi(h)f(g) \quad \forall h \in H\},$$

and given $\xi, \xi' \in \mathcal{X}(H)$ there is a natural multiplication map

$$\mathbb{C}[G]_{\xi}^{(H)} \otimes \mathbb{C}[G]_{\xi'}^{(H)} \longrightarrow \mathbb{C}[G]_{\xi+\xi'}^{(H)}.$$

As in the case $\xi = 0$, the G -module $\mathbb{C}[G]_{\xi}^{(H)}$ is multiplicity free, and we denote by $V_{\lambda, \xi}$ its irreducible component of highest weight λ . Therefore, given $\xi, \xi' \in \mathcal{X}(H)$, it is well defined as the product $V_{\lambda, \xi}V_{\lambda', \xi'} \subset \mathbb{C}[G]_{\xi+\xi'}^{(H)}$.

After discussing some non-degeneracy property of such multiplication and giving some examples, we describe the semigroup generated by the set of differences

$$\{\lambda + \lambda' - \mu : V_{\mu, \xi+\xi'} \subset V_{\lambda, \xi}V_{\lambda', \xi'} \quad \exists \xi, \xi' \in \mathcal{X}(H)\}$$

in terms of the spherical roots of a suitable wonderful variety as follows. There is a canonical spherical subgroup of G containing H , called the *spherical closure* of H and denoted \overline{H} , such that G/\overline{H} admits a wonderful compactification M . Moreover, $\mathcal{X}(H) = \mathcal{X}(\overline{H})$, and for all $\xi \in \mathcal{X}(H)$ it holds that $\mathbb{C}[G]_{\xi}^{(H)} = \mathbb{C}[G]_{\xi}^{(\overline{H})}$. This allows us to reformulate the multiplication of H -semi-invariant functions in terms of sections of line bundles on M , and we prove the following fact.

Proposition 5 (see Proposition 9.2). *Let \mathcal{M} be the semigroup generated by the set of differences*

$$\{\lambda + \lambda' - \mu : V_{\mu, \xi + \xi'} \subset V_{\lambda, \xi} V_{\lambda', \xi'} \quad \exists \xi, \xi' \in \mathcal{X}(H)\},$$

and denote by Σ the set of spherical roots of M . Then $\mathcal{M} = \mathbb{N}\Sigma$.

Structure of the paper. The first part of the paper is entirely devoted to the proof of Theorem 1 and Lemma 2, whereas in the second part we have collected the various applications.

In Section 1 we fix the notation and recall some results about the wonderful varieties and their line bundles. In Section 2 we define the low triples and prove Lemma 2. In Section 3 and in Section 4 we focus on the case of a model wonderful variety, first classifying the covering differences and then classifying the low fundamental triples and studying the associated inclusions.

In Section 5 we introduce the comodel wonderful varieties and prove the surjectivity of the multiplication for this class of varieties.

In Section 6 we explain in general how the surjectivity of the multiplication map can give information on the normality of the closure of a spherical orbit in the projective space of a simple G -module (proving, by the way, Theorem 3). In Section 7 we prove Theorem 4 and we use our results to deduce the normality and the non-normality of the spherical nilpotent orbit closures. In Section 8 we concentrate on the real model orbit of type E_8 .

In Section 9 we give the above-mentioned counterexample to the surjectivity of the multiplication in the case of a model wonderful variety of not simply connected type. This leads us to discuss some general properties of the multiplication map and to prove Proposition 5.

1. GENERALITIES

Let G be a simply connected semisimple complex algebraic group. Fix a maximal torus T of G and a Borel subgroup $B \supset T$. For any group K we denote by $\mathcal{X}(K)$ the multiplicative characters of K . We denote also by $\mathcal{X}(T)^+$ the set of dominant characters w.r.t. B and if $\lambda \in \mathcal{X}(T)^+$ we denote by $V(\lambda)$ an irreducible representation of highest weight λ and by $-\lambda^*$ the lowest weight of $V(\lambda)$. We denote by S the set of simple roots.

Let M be a wonderful G -variety with (unique) closed G -orbit Y . We assume that the center $Z(G)$ acts trivially on M , and we will keep this assumption throughout the paper: all the wonderful varieties we will deal with will be G_{ad} -varieties. By [34], M is *spherical*, i.e., it possesses an open B -orbit, say $B \cdot x_0 \subset G \cdot x_0 \subset M$. We denote by H the stabilizer of x_0 in G .

1.1. Colors and spherical roots. Since $B \cdot x_0$ is affine, $G \cdot x_0 \setminus B \cdot x_0$ is a union of finitely many B -stable prime divisors and we denote by Δ the set of their closures in M :

$$\Delta = \{D \subset M : D \text{ is a } B\text{-stable prime divisor, } D \cap G \cdot x_0 \neq \emptyset\}.$$

The elements of Δ are called the *colors* of M .

Denote by B^- the opposite Borel subgroup of B and let $y_0 \in Y$ be the unique B^- -fixed point of M . The normal space of Y in M at y_0 , $T_{y_0}M/T_{y_0}Y$, is a multiplicity-free T -module. Set

$$\Sigma = \{T\text{-weights of } T_{y_0}M/T_{y_0}Y\} :$$

the elements of Σ are called the *spherical roots* of M and they naturally correspond to local equations of the boundary divisors of M , which are G -stable. If $\sigma \in \Sigma$, we denote by M^σ the associated boundary divisor of M such that $T_{y_0}M/T_{y_0}M^\sigma$ is the 1-dimensional T -module of weight σ .

The rank one wonderful varieties were classified by D.N. Akhiezer [2], and we denote by $\Sigma(G) \subset \mathcal{X}(T)$ the finite set of all possible weights occurring as the spherical root of a rank one wonderful variety. These are called the *spherical roots* of G . Every spherical root is either a positive root, or the sum of two orthogonal positive roots, we refer to [8, Table 1] for a list of the spherical roots of G .

1.2. Picard group and Cartan pairing. Recall that every line bundle on M or on Y has a unique G -linearization.

We may identify $\text{Pic}(Y)$ with a sublattice of $\mathcal{X}(T)$ and $\text{Pic}(G \cdot x_0)$ with $\mathcal{X}(H)$ (see [30]): we identify $\mathcal{L} \in \text{Pic}(Y)$ with the character of T acting on the fiber of \mathcal{L} over y_0 , and we identify $\mathcal{L} \in \text{Pic}(G \cdot x_0)$ with the character of H acting on the fiber over x_0 .

Consider now the maps $\omega: \text{Pic}(M) \rightarrow \mathcal{X}(T)$ and $\xi: \text{Pic}(M) \rightarrow \mathcal{X}(H)$ defined by the restriction to the closed and to the open orbit. We may regard $\text{Pic}(M)$ as a sublattice of $\mathcal{X}(T) \times \mathcal{X}(H)$ by identifying $\mathcal{L} \in \text{Pic}(M)$ with the couple $(\omega(\mathcal{L}), \xi(\mathcal{L}))$ (see [11]). Moreover, we have the following exact sequence

$$0 \rightarrow \mathbb{Z}\Sigma \rightarrow \text{Pic}(M) \rightarrow \mathcal{X}(H) \rightarrow 0.$$

Given $E \in \mathbb{Z}\Delta$, denote by $\mathcal{L}_E := \mathcal{O}(E)$ the associated line bundle. As a group, $\text{Pic}(M)$ is freely generated by the line bundles \mathcal{L}_D with $D \in \Delta$ (see [10, Proposition 2.2]). For all $E \in \mathbb{Z}\Delta$, the associated line bundle \mathcal{L}_E is globally generated (resp. ample) if and only if E is a non-negative (resp. positive) combination of colors. We set $\omega_E = \omega(\mathcal{L}_E)$, $\xi_E = \xi(\mathcal{L}_E)$.

There exists a natural \mathbb{Z} -bilinear pairing (called the *Cartan pairing* of M)

$$c: \mathbb{Z}\Delta \times \mathbb{Z}\Sigma \rightarrow \mathbb{Z}$$

which maps the couple (D, σ) to the coefficient of $[M^\sigma]$ along $[D]$. So, regarding $\mathbb{Z}\Sigma$ as a sublattice of $\mathbb{Z}\Delta$, for $\sigma \in \mathbb{Z}\Sigma$ we have

$$\sigma = \sum_{D \in \Delta} c(D, \sigma)D.$$

1.3. Global sections and multiplication. We now recall the description of the space of global sections $\Gamma(M, \mathcal{L})$ of a line bundle \mathcal{L} . Notice first that since M is spherical the decomposition of $\Gamma(M, \mathcal{L})$ into simple G -modules is multiplicity free for all $\mathcal{L} \in \text{Pic}(M)$. If $E \in \mathbb{N}\Delta$ then \mathcal{L}_E is generated by global sections. In particular $\Gamma(M, \mathcal{L}_E)$ must contain a copy of $V(\omega_E)$ (which is the space of sections of \mathcal{L}_E on Y), hence ω_E is dominant. We denote by V_E the unique simple G -submodule of $\Gamma(M, \mathcal{L}_E)$ of highest weight ω_E . Notice also that the image of x_0 in $\mathbb{P}(\Gamma(M, \mathcal{L}_E)^*)$ is a point fixed by H . In particular, since $BH \subset G$ is open, it follows that the *space of spherical vectors*

$$V(\omega_E^*)_{\xi_E}^{(H)} = \{v \in V(\omega_E^*) : hv = \xi_E(h)v \quad \forall h \in H\}$$

has dimension one and we denote by h_E a generator of this line.

If $\gamma = \sum a_\sigma \sigma \in \mathbb{N}\Sigma$, we denote by $s^\gamma \in \Gamma(M, \mathcal{L}_{M^\gamma})$ a section whose divisor is equal to $M^\gamma = \sum a_\sigma M^\sigma$. Notice that this section is G -invariant. Recall, as defined in the introduction, that we say that $F \leq_\Sigma E$ if $E - F \in \mathbb{N}\Sigma$. If $E \in \mathbb{N}\Delta$ and

$E \leq_{\Sigma} F$ the multiplication by s^{F-E} induces a G -equivariant map from the sections of \mathcal{L}_E to the sections of \mathcal{L}_F , in particular, we have $s^{F-E}V_E \subset \Gamma(M, \mathcal{L}_F)$.

Proposition 1.1 ([10, Proposition 2.4]). *Let $F \in \mathbb{Z}\Delta$. Then*

$$\Gamma(M, \mathcal{L}_F) = \bigoplus_{E \in \mathbb{N}\Delta : E \leq_{\Sigma} F} s^{F-E}V_E.$$

If $E, F \in \mathbb{N}\Delta$, consider the multiplication of sections

$$m_{E,F}: \Gamma(M, \mathcal{L}_E) \otimes \Gamma(M, \mathcal{L}_F) \longrightarrow \Gamma(M, \mathcal{L}_{E+F})$$

and denote by $V_E V_F$ the image of $V_E \otimes V_F$. A way to translate the description of this map into a problem on spherical vectors is the following.

Lemma 1.2 ([14, Lemma 19]). *Let $D, E, F \in \mathbb{N}\Delta$ be such that $D \leq_{\Sigma} E + F$. Then $s^{E+F-D}V_D \subset V_E V_F$ if and only if the projection of $h_E \otimes h_F \in V(\omega_E^*) \otimes V(\omega_F^*)$ onto the isotypic component of highest weight ω_D^* is non-zero.*

1.4. Distinguished sets of colors. Let M' be a wonderful G -variety together with a surjective equivariant morphism $\phi: M \rightarrow M'$ with connected fibers and denote $\Delta_{\phi} \subset \Delta$ the set of colors which map dominantly onto M' . Then the semi-group

$$(\mathbb{N}\Sigma)/\Delta_{\phi} = \{\gamma \in \mathbb{N}\Sigma : c(D, \gamma) = 0 \quad \forall D \in \Delta_{\phi}\}$$

is free and its basis, which we denote by Σ/Δ_{ϕ} , coincides with the set of spherical roots of M' , while the set of colors of M' is identified with $\Delta \setminus \Delta_{\phi}$ (see [35, Proposition 3.3.2]).

Conversely, if $\Delta_0 \subset \Delta$, then there exists a wonderful variety M' (unique up to isomorphism) together with a surjective equivariant morphism $\phi: M \rightarrow M'$ with connected fibers if and only if Δ_0 is *distinguished*, that is, there exists an element $D \in \mathbb{N}_{>0}\Delta_0$ such that $c(D, \sigma) \geq 0$ for every $\sigma \in \Sigma$ (see [35, Proposition 3.3.2] and [5, Theorem 3.1]).

If $\Delta_0 \subset \Delta$ is distinguished, then we say that the associated wonderful variety M' is the *quotient* of M by Δ_0 , denoted by M/Δ_0 .

Recall the following general fact.

Proposition 1.3. *Let X, Y be normal varieties and suppose that $\phi: X \rightarrow Y$ is a surjective proper morphism with connected fibers. If $\mathcal{L} \in \text{Pic}(Y)$, then $\Gamma(Y, \mathcal{L}) = \Gamma(X, \phi^*\mathcal{L})$.*

If $E = \sum_{D \in \Delta} a_D D \in \mathbb{Z}\Delta$ we define $\text{supp}(E)$, the *support* of E , as the set of colors D such that $a_D \neq 0$.

Corollary 1.4. *Let $E \in \mathbb{N}\Delta$ and suppose that Δ_0 is a distinguished subset of Δ such that $\Delta_0 \cap \text{supp}(E) = \emptyset$. Then $\Gamma(M, \mathcal{L}_E) = \Gamma(M/\Delta_0, \mathcal{L}_E)$.*

1.5. Parabolic induction: spherical roots and colors. We describe now a standard way to construct a wonderful variety for the group G from a wonderful variety for a Levi subgroup of a parabolic subgroup of G .

Let L be a proper Levi subgroup of G which contains T . Let Q be the parabolic subgroup associated to L and containing B , and let Q^- be the opposite parabolic subgroup. Denote by R_Q and R_{Q^-} the solvable radicals and by U_Q and U_{Q^-} the unipotent radicals of Q and Q^- , respectively. Finally let L^{ss} be the semisimple part of L and denote by $L_{\text{ad}} = L^{\text{ss}}/Z(L^{\text{ss}}) = L/Z(L)$ the adjoint quotient of L .

Let N be a wonderful variety for the group L_{ad} . Although wonderful varieties have been defined for semisimple groups, in this case it is convenient to look at N as an L -variety and to consider L -linearized line bundles on N . The definitions given for a wonderful variety can be extended to this situation.

Extend the action of L on N to Q^- , with U_Q^- acting trivially, and define the parabolic induction of N as

$$M = G \times_{Q^-} N.$$

We have a projection $\pi: M \rightarrow G/Q^-$, N is the fiber over Q^- and, moreover, it is the set of points of M fixed by U_Q^- . One can easily show the following properties.

- (1) M is a wonderful G -variety.
- (2) The intersection of a subset of M with N induces a bijection from the set of G -stable subsets of M to the set of L -stable subsets of N . For every spherical root σ of M we denote by N^σ the intersection $M^\sigma \cap N$.
- (3) The restriction of line bundles from M to N induces an isomorphism ρ between the class groups of linearized line bundles $\text{Pic}_G(M)$ and $\text{Pic}_L(N)$.
- (4) We have an injective map ε from the set $\Delta(N)$ of colors of N to the set $\Delta(M)$ of colors of M given by $\varepsilon(D) = \overline{B \cdot D}$ for all $D \in \Delta(N)$.
- (5) The exact sequence $0 \rightarrow \mathcal{X}(L/L^{\text{ss}}) \rightarrow \text{Pic}_L(N) \rightarrow \text{Pic}_{L^{\text{ss}}}(N) \rightarrow 0$ has a natural splitting given by $\mathcal{L}_D \mapsto \mathcal{L}_{\varepsilon(D)}|_N$ for every color $D \in \Delta(N)$.
- (6) If $D \in \Delta(M) \setminus \varepsilon(\Delta(N))$ then the restriction of $\mathcal{O}(D)$ to N is a trivial line bundle on N .
- (7) The set of colors $\varepsilon(\Delta(N))$ is distinguished and $M/\varepsilon(\Delta(N)) \simeq G/Q^-$.
- (8) Since L^{ss} is simply connected we have $\text{Pic}(N) \simeq \text{Pic}_{L^{\text{ss}}}(N)$.

Denote

$$\Sigma = \{T\text{-weights of } T_{y_0}N/T_{y_0}(L \cdot y_0)\},$$

where we recall that y_0 is the point of M fixed by B^- , hence $L \cdot y_0$ is the unique closed orbit of N . The set Σ is in correspondence with the set of L -stable divisors and, in particular, σ is the T -weight of $T_{y_0}N/T_{y_0}N^\sigma$. Moreover, we have that Σ equals the set of spherical roots of M and every element of Σ is a sum of simple roots of L .

By the above properties (3) and (4) for every color D in $\Delta(N)$ we have a canonical choice of a linearization of the associated line bundle and we have a natural decomposition

$$\text{Pic}_L(N) \simeq \mathcal{X}(L/L^{\text{ss}}) \oplus \bigoplus_{D \in \Delta(N)} \mathbb{Z}[D].$$

For all $\sigma \in \Sigma$, $[N^\sigma] = \sum_{D \in \Delta(M)} c(D, \sigma) \rho([D])$. If $\mathcal{L}, \mathcal{L}' \in \text{Pic}_L(N)$ we define $\mathcal{L} \geq_\Sigma \mathcal{L}'$ if, using the additive notation, $\mathcal{L} - \mathcal{L}' = \sum a_\sigma \mathcal{O}(N^\sigma)$ with $a_\sigma \geq 0$ for all σ , similarly to what we have done in Section 1.3. In this way Proposition 1.1 holds for N without any change.

It is easy to see when a G -wonderful variety M can be obtained by parabolic induction. Let P^- be the stabilizer of y_0 and let S^p the set of simple roots that are in the set of roots of P^- . Assume that $S' = S^p \cup \text{supp}_\Sigma \neq S$ and let L be the standard Levi subgroup associated to S' . Let $Q, Q^-, R_Q, R_Q^-, U_Q, U_Q^-, L^{\text{ss}}, L_{\text{ad}}$ be defined as above. Set $\tilde{\Delta} = \{D \in \Delta : \text{if } P_\alpha \cdot D \neq D \text{ then } \alpha \in S'\}$, then $\tilde{\Delta}$ is a distinguished set and $M/\tilde{\Delta} = G/Q^-$. Moreover, $N = M^{U_Q^-}$ is a wonderful variety for L_{ad} and M is obtained by parabolic induction from N as above.

1.6. Parabolic induction: global sections. Now let M be obtained by parabolic induction from N as above. Let \mathcal{M} be a line bundle on M and denote by \mathcal{N} its restriction to N . We want to compare the sections of \mathcal{M} and \mathcal{N} . The restriction of sections induces a map $r_{\mathcal{M}}: \Gamma(M, \mathcal{M}) \rightarrow \Gamma(N, \mathcal{N})$. Notice that $U \cdot N$ is a dense subset of M , hence the restriction of $r_{\mathcal{M}}$ to $\Gamma(M, \mathcal{M})^U$ is injective. We first describe the kernel of $r_{\mathcal{M}}$.

Let $\mathcal{M} = \mathcal{L}_F$, then $\Gamma(M, \mathcal{M}) = \bigoplus s^{F-E} V_E$ where the sum is over all $E \in \mathbb{N}\Delta$ such that $E \leq_{\Sigma} F$. For each $E \in \mathbb{N}\Delta$ set $W_E = V_E^{U_Q}$. This is an irreducible L -submodule with the same highest weight as V_E . Let I_E be the L -stable complement of W_E in V_E .

Lemma 1.5. *We have*

$$\ker r_{\mathcal{M}} = \bigoplus_{E \in \mathbb{N}\Delta : E \leq_{\Sigma} F} s^{F-E} I_E.$$

Proof. By the above discussion it is enough to prove that $r_{\mathcal{L}_E}(I_E) = 0$. Let v be a highest weight vector (for the action of L) in I_E of weight λ . Then $\omega_E - \lambda = \sum_{\alpha \in S} a_{\alpha} \alpha$ with $a_{\alpha} \in \mathbb{N}$. If S' is the set of simple roots for L notice that, since $v \in I_E$, there exists $\alpha \in S \setminus S'$ such that $a_{\alpha} \neq 0$. On the other hand, by the generalization of Proposition 1.1 for N discussed in the previous section, the weights of the highest weight vectors in $\Gamma(N, \mathcal{L}_E)$ are of the form $\omega_E - \beta$ with $\beta \in \mathbb{N}S'$. In particular, we must have $r_{\mathcal{L}_E}(v) = 0$. \square

Since the restriction commutes with the multiplication we recover the following.

Proposition 1.6 ([15, Proposition 2.9]). *For all $E, F \in \mathbb{N}\Delta$ and $\gamma \in \mathbb{N}\Sigma$ such that $E + F - \gamma \in \mathbb{N}\Delta$, $s^{\gamma} V_{E+F-\gamma} \subset V_E V_F$ if and only if $s^{\gamma} W_{E+F-\gamma} \subset W_E W_F$.*

In particular, using the property (6) of the previous section, we have the following.

Corollary 1.7. *For all $D \in \mathbb{N}(\Delta \setminus \tilde{\Delta})$ and $E \in \mathbb{N}\Delta$, $V_D V_E = V_{D+E}$.*

2. PROJECTIVE NORMALITY AND THE COVERING RELATION

2.1. The covering relation. Let $\{\omega_{\alpha} : \alpha \in S\}$ be the set of fundamental weights w.r.t. the simple roots S .

For all $\lambda = \sum k_{\alpha} \omega_{\alpha} \in \mathcal{X}(T)$, denote by $\text{supp}(\lambda)$ the set of $\alpha \in S$ such that $k_{\alpha} \neq 0$ and define its positive part λ^+ , respectively, its negative part λ^- , as the dominant weights

$$\lambda^+ = \sum_{k_{\alpha} > 0} k_{\alpha} \omega_{\alpha} \quad \text{and} \quad \lambda^- = \lambda^+ - \lambda.$$

If $\lambda \in \mathcal{X}(T)^+$, define also the height of λ as the number $\text{ht}(\lambda) = \sum_{\alpha \in S} k_{\alpha}$.

Suppose that λ and μ are dominant weights with $\lambda < \mu$ (w.r.t. the usual dominance order) and suppose that there is no dominant weight ν such that $\lambda < \nu < \mu$: then one says that μ covers λ and we call $\mu - \lambda$ a covering difference in $\mathcal{X}(T)^+$. Notice that an element $\gamma \in \mathbb{N}S$ is a covering difference in $\mathcal{X}(T)^+$ if and only if γ^+ covers γ^- . Although the following proposition is an immediate consequence of [41, Theorem 2.6], here we give an easy independent proof.

Proposition 2.1. *If $\gamma \in \mathbb{N}S$ is a covering difference in $\mathcal{X}(T)^+$, then $\text{ht}(\gamma^+) \leq 2$.*

Proof. Let $\gamma \in \mathbb{N}S$ be a covering difference and suppose that $\alpha \in S$ is such that $\langle \gamma^+, \alpha^\vee \rangle \geq 2$: then $\gamma^+ - \alpha \in \mathcal{X}(T)^+$, hence we must have $\gamma = \alpha$ and $\text{ht}(\gamma^+) = 2$. Hence we may assume that $\langle \gamma^+, \alpha^\vee \rangle \leq 1$ for every $\alpha \in S$. Suppose that $\text{ht}(\gamma^+) \geq 3$, (up to reindexing the simple roots) we can take $\alpha_i, \alpha_j \in \text{supp}_S(\gamma^+)$ with $i < j$ such that $\{\alpha_i, \dots, \alpha_j\}$ generates an irreducible subsystem of type A ; then $\gamma^+ - (\alpha_i + \dots + \alpha_j) \in \mathcal{X}(T)^+$ and it follows $\gamma = \alpha_i + \dots + \alpha_j$ and $\text{ht}(\gamma^+) = 2$ (a contradiction). \square

We now consider the covering relation in the more general context of wonderful varieties. Let M be a wonderful variety with set of spherical roots Σ and with set of colors Δ . For all $E = \sum_{D \in \Delta} k_D D \in \mathbb{Z}\Delta$, define its *positive part* E^+ and its *negative part* E^- as

$$E^+ = \sum_{k_D > 0} k_D D \quad \text{and} \quad E^- = E^+ - E.$$

If $E \in \mathbb{N}\Delta$, define the *height* of E as the number $\text{ht}(E) = \sum_{D \in \Delta} k_D$. On the other hand, for all $\gamma = \sum_{\sigma \in \Sigma} a_\sigma \sigma \in \mathbb{N}\Sigma$ define its Σ -*height* as $\text{ht}_\Sigma(\gamma) = \sum_{\sigma \in \Sigma} a_\sigma$.

Let E and F be in $\mathbb{N}\Delta$ with $E <_\Sigma F$ and suppose that there is no $D \in \mathbb{N}\Delta$ such that $E <_\Sigma D <_\Sigma F$; then we say that F covers E and we call $F - E$ a covering difference in $\mathbb{N}\Delta$. Again, $\gamma \in \mathbb{N}\Sigma$ is a covering difference in $\mathbb{N}\Delta$ if and only γ^+ covers γ^- .

Remark 1. Notice that there are only finitely many covering differences. Indeed, consider the lattice $X = \{(D, E) \in \mathbb{Z}\Delta \times \mathbb{Z}\Delta : E - D \in \mathbb{Z}\Sigma\}$ and the finitely generated monoid $X^+ = \{(D, E) \in \mathbb{N}\Delta \times \mathbb{N}\Delta : D - E \in \mathbb{N}\Sigma\}$. If γ is a covering difference then (γ^+, γ^-) is an indecomposable element of X^+ .

Finally, the number of indecomposable elements in X^+ is finite and can be controlled as follows: let ℓ_1, \dots, ℓ_t be the half-lines generating the cone X^+ , let $v_i = (D_i, E_i)$ be a generator of $\ell_i \cap X^+$ and let $n_i = \text{ht}(D_i) + \text{ht}(E_i)$. Then if (D, E) is indecomposable, $\text{ht}(D) + \text{ht}(E) \leq \sum_i n_i$.

As already said in the introduction, in the case of the wonderful compactification of a non-Hermitian symmetric space, there exists a root system Φ_Σ (the *restricted root system*) which is generated by the spherical roots and such that Δ is naturally identified with the set of fundamental weights of Φ_Σ and the pairing between Σ and Δ is the Cartan pairing of Φ_Σ . Although Σ is always the basis of a root system Φ_Σ , in the general case it is not possible to identify Δ with the fundamental weights of Φ_Σ ; in particular, it may happen that $\mathbb{Z}\Sigma \subset \mathbb{Z}\Delta$ is not a sublattice of finite index and that the semigroup of dominant weights of Φ_Σ contained in $\mathbb{Z}\Sigma$ is not even contained in $\mathbb{N}\Delta$. However, one can consider the property of Proposition 2.1 without modifications in this context:

(2-ht) If $\gamma \in \mathbb{N}\Sigma$ is a covering difference in $\mathbb{N}\Delta$, then $\text{ht}(\gamma^+) \leq 2$.

Notice that by the above discussion the property (2-ht) holds if M is the wonderful compactification of a non-Hermitian symmetric space. In the case of the wonderful compactifications of Hermitian symmetric spaces the same argument works without serious complications. In the subsequent section we show that it is true also in the case of a model wonderful variety (of simply connected type). We have also checked many other examples and, as far as we know, it is possible that it holds for all wonderful varieties.

In analogy with the case of a root system, we say that a non-zero element $D \in \mathbb{N}\Delta$ is *minuscule* if it is minimal in $\mathbb{N}\Delta$ w.r.t. the partial order \leq_Σ .

2.2. Projective normality. The notion of low triple has been introduced in [15] in the case of a symmetric wonderful variety. Here we use the same terminology with a slightly weaker definition.

Definition 2.2. Let M be a wonderful variety with set of spherical roots Σ and with set of colors Δ . Let $D, E, F \in \mathbb{N}\Delta$ be such that $F \leq_\Sigma D + E$. The triple (D, E, F) is called a *low triple* if:

for all $D' \leq_\Sigma D$ and $E' \leq_\Sigma E$ such that $F \leq_\Sigma D' + E'$, it holds that $D' = D$ and $E' = E$.

The triple (D, E, F) is called a *fundamental triple* if $D, E \in \Delta$.

Notice that if (D, E, F) is a low triple, then $s^{D+E-F}V_F \subset \Gamma(M, \mathcal{L}_D)\Gamma(M, \mathcal{L}_E)$ if and only if $s^{D+E-F}V_F \subset V_DV_E$.

Lemma 2.3. *Let M be a wonderful variety and let n be such that $\text{ht}(\gamma^+) \leq n$ for every covering difference γ . If*

$$(2.1) \quad s^{D+E-F}V_F \subset V_DV_E,$$

for all low triples (D, E, F) with $\text{ht}(D + E) \leq n$, then the multiplication map

$$m_{D,E}: \Gamma(M, \mathcal{L}_D) \otimes \Gamma(M, \mathcal{L}_E) \longrightarrow \Gamma(M, \mathcal{L}_{D+E})$$

is surjective for all $D, E \in \mathbb{N}\Delta$.

Proof. For any $D \in \mathbb{Z}\Delta$ let $\Gamma_D = \Gamma(M, \mathcal{L}_D)$. For any triple (D, E, F) with $F \leq_\Sigma D + E$ we want to show that $s^{D+E-F}V_F \subset \Gamma_D\Gamma_E$. We proceed by induction first on $\text{ht}_\Sigma(D + E - F)$ and then on $\text{ht}(D + E)$. If $\text{ht}_\Sigma(D + E - F) = 0$ the claim is trivial.

If (D, E, F) is not a low triple then there exist $D' \leq_\Sigma D$ and $E' \leq_\Sigma E$ such that $F \leq_\Sigma D' + E'$ and $\text{ht}_\Sigma(D' + E' - F) < \text{ht}_\Sigma(D + E - F)$. Hence the claim is true for (D', E', F) , and

$$s^{D+E-F}V_F = s^{D-D'+E-E'}s^{D'+E'-F}V_F \subset s^{D-D'+E-E'}\Gamma_{D'}\Gamma_{E'} \subset \Gamma_D\Gamma_E.$$

If (D, E, F) is a low triple and $\text{ht}(D + E) \leq n$ then the claim is true by assumption.

Assume now that $\text{ht}(D + E) > n$ and that (D, E, F) is a low triple. Let $F_1 \in \mathbb{N}\Delta$ be an element covered by $D + E$ such that $F_1 \geq_\Sigma F$. Since $\text{ht}(D + E) > \text{ht}(\gamma^+)$ for every covering difference $\gamma \in \mathbb{N}\Sigma$, it follows that $\text{supp}(F_1) \cap \text{supp}(D + E) \neq \emptyset$. Fix $D_0 \in \text{supp}(F_1) \cap \text{supp}(D + E)$ and set

$$(D_1, E_1) = \begin{cases} (D - D_0, E) & \text{if } D_0 \in \text{supp}(E), \\ (D, E - D_0) & \text{otherwise.} \end{cases}$$

Choose F_2 in $\mathbb{N}\Delta$ minimal w.r.t. \leq_Σ such that $F - D_0 \leq_\Sigma F_2 \leq_\Sigma F_1 - D_0$. Notice that $\text{ht}_\Sigma(D_0 + F_2 - F) \leq \text{ht}_\Sigma(F_1 - F) < \text{ht}_\Sigma(D + E - F)$. Moreover, (D_0, F_2, F) is a low triple; indeed if there exist $F'_2 \leq_\Sigma F_2$ and $D'_0 \leq_\Sigma D_0$ in $\mathbb{N}\Delta$ such that $F \leq_\Sigma D'_0 + F'_2$, then

$$F \leq_\Sigma D'_0 + F'_2 \leq_\Sigma D'_0 + D_1 + E_1 \leq_\Sigma D + E.$$

Since (D, E, F) is a low triple, we get $D'_0 = D_0$, while since F_2 is minimal we get $F'_2 = F_2$. Hence, by induction, it follows that $s^{D_0+F_2-F}V_F \subset V_{D_0}V_{F_2}$.

Notice that we have $\text{ht}_\Sigma(D_1 + E_1 - F_2) \leq \text{ht}_\Sigma(D + E - F)$ and that $\text{ht}(D_1 + E_1) = \text{ht}(D + E) - 1$, hence we can apply the inductive hypothesis to (D_1, E_1, F_2) . So $s^{D+E-F}V_F = s^{D_1+E_1-F_2}s^{D_0+F_2-F}V_F \subset s^{D_1+E_1-F_2}V_{F_2}V_{D_0} \subset \Gamma_{D_1}\Gamma_{E_1}V_{D_0} \subset \Gamma_D\Gamma_E$. \square

Together with Remark 1, the lemma implies that to prove the surjectivity of $m_{D,E}$ for all $D, E \in \mathbb{N}\Delta$ it is enough to check a finite number of cases. In particular if the property (2-ht) holds, it is enough to check the above inclusion (2.1) only for the low fundamental triples.

In Section 9.1 we show that there exist wonderful varieties possessing low fundamental triples (D, E, F) such that $s^{D+E-F}V_F \not\subset V_DV_E$; in particular, the multiplication of sections of line bundles on a wonderful variety is not necessarily surjective.

3. THE COVERING RELATION FOR MODEL WONDERFUL VARIETIES

As said in the introduction, a model homogeneous space for a reductive group G (not necessarily simply connected) is a quasi affine G -homogeneous space whose coordinate ring is a model representation of G : each irreducible representation of G appears exactly with multiplicity one. By [36], for every group G there exists a wonderful variety M such that for every model homogeneous space G/H_0 there exists a point in M whose stabilizer is equal to $N_G(H_0)$ and, vice versa, if H is the stabilizer of a point in M and we set H_0 to be the intersections of the kernels of the multiplicative characters of H , then G/H_0 is a model homogeneous space. The variety M is called the model wonderful variety of G .

Notice that H_0 has no characters, hence the set of spherical vectors (i.e., the eigenvectors of H) in a simple G -module V , which is a subspace of dimension at most one, can be also characterized as the set of vectors in V fixed by H_0 .

The description of the model wonderful varieties of an almost simple group G is given in [36]. For type **B** there are two different model wonderful varieties, according to whether G is isomorphic to the special orthogonal group or to the spin group. In all the other cases the model wonderful variety of G is the same as the model wonderful variety of the adjoint group of G .

In this section we describe the covering relation for a model wonderful variety M of simply connected type, and we prove that the property (2-ht) holds in this case. For model wonderful varieties of simply connected type the set of colors Δ is in bijection, via ω , with the set of fundamental weights, and the set of spherical roots Σ is the set of the sums $\alpha + \beta$ with α, β non-orthogonal simple roots.

Clearly, in order to study the multiplication of sections of line bundles and the partial order of dominant weights in the case of a model wonderful variety, we may reduce to the case where G is almost simple. Moreover, if $\gamma = \sum_{\alpha \in S} a_\alpha \alpha \in \mathbb{Z}\Sigma$ we denote by $\text{supp}_S(\gamma)$ the set of simple roots α such that $a_\alpha \neq 0$. We have the following lemma, whose proof is immediate.

Lemma 3.1. *Let γ be a covering difference in $\mathbb{N}\Delta$ w.r.t. Σ , then $\text{supp}_S(\gamma)$ is a connected subset of S and γ is a covering difference for the model variety of the simply connected group associated to the root subsystem generated by $\text{supp}_S(\gamma)$.*

Therefore, we only classify the covering differences γ such that $\text{supp}_S(\gamma) = S$. In the following two sections, r denotes the semisimple rank of the group and $\Delta = \{D_1, \dots, D_r\}$ is the set of colors, labelled in such a way that ω_{D_i} is the

fundamental weight ω_i . For simplicity we also set $D_i = 0$ for all $i \leq 0$ and for all $i > r$. Similarly, we denote $\Sigma = \{\sigma_1, \dots, \sigma_{r-1}\}$ (see the following subsections for the definition of the single spherical roots σ_i), and we set $\sigma_i = 0$ for all $i \leq 0$ and for all $i \geq r$.

For the rest of the section, γ will be a covering difference in $\mathbb{N}\Delta$ w.r.t. Σ , and we set

$$\gamma = \sum_{i=1}^{r-1} a_i \sigma_i = \sum_{i=1}^r c_i D_i.$$

For convenience we also set $a_i = c_j = 0$ for all $i, j \leq 0$, for all $i \geq r$ and for all $j > r$.

3.1. Type A_r ($r \geq 2$). Let $\Sigma = \{\sigma_1, \dots, \sigma_{r-1}\}$ be the set of spherical roots where $\sigma_i = \alpha_i + \alpha_{i+1}$.

Proposition 3.2. *Let $\gamma \in \mathbb{N}\Sigma$ be a covering difference in $\mathbb{N}\Delta$ with $\text{supp}_S(\gamma) = S$. Then either*

- (1) r is even and $\gamma = \sum_{i=1}^{r/2} \sigma_{2i-1} = D_1 + D_r$, or
- (2) r is odd and $\gamma = \sum_{i=1}^{r-1} \sigma_i = D_2 + D_{r-1}$.

Proof. Clearly, $a_1 > 0$. Assume first that $a_2 > 0$. Take the maximum integer k such that $a_i > 0$ for all $i \leq k$. Then $c_j \geq 0$ for all $3 \leq j \leq k+2$ otherwise $\gamma^- + \sum_{i=1}^{j-2} \sigma_i \in \mathbb{N}\Delta$.

If k were $< r-1$, then $c_{k+1} + c_{k+2} = -a_{k-1} - a_{k+3}$ would be < 0 , a contradiction. Therefore, $\text{supp}_\Sigma(\gamma) = \Sigma$.

Now, since $\gamma^- + \sum_{i=1}^{r-1} \sigma_i \in \mathbb{N}\Delta$, γ must necessarily be equal to $\sum_{i=1}^{r-1} \sigma_i$ which is a covering difference if and only if r is odd; indeed, if r is even, $\sum_{i=1}^{r/2} \sigma_{2i-1} \in \mathbb{N}\Delta$.

Assume now that $a_2 = 0$. Take the maximum odd integer k such that $a_i = 0$ for all even $i < k$. Then $c_k \geq 0$ otherwise $\gamma^- + \sum_{i=1}^{(k-1)/2} \sigma_{2i-1} \in \mathbb{N}\Delta$. Furthermore, $c_{k-1} \geq 0$ otherwise, reasoning as above, $a_j > 0$ for all $j \geq k$ and $\gamma^- + \sum_{i=k}^{r-1} \sigma_i \in \mathbb{N}\Delta$.

If k were $< r-1$, then $c_{k-1} + c_k = -a_{k+1}$ would be < 0 , a contradiction. Therefore, $a_i > 0$ iff i is odd, and r is even.

Now, since $\gamma^- + \sum_{i=1}^{r/2} \sigma_{2i-1} \in \mathbb{N}\Delta$, γ must necessarily be equal to $\sum_{i=1}^{r/2} \sigma_{2i-1}$ which is a covering difference. \square

3.2. Type B_r ($r \geq 2$). Let $\Sigma = \{\sigma_1, \dots, \sigma_{r-1}\}$ be the set of spherical roots where $\sigma_i = \alpha_i + \alpha_{i+1}$.

Proposition 3.3. *Let $\gamma \in \mathbb{N}\Sigma$ be a covering difference in $\mathbb{N}\Delta$ with $\text{supp}_S(\gamma) = S$. Then r is even and $\gamma = \sum_{i=1}^{r/2} \sigma_{2i-1} = D_1$.*

Proof. Clearly, $a_{r-1} > 0$. Suppose that $k < r$ is such that $r-k$ is odd and $c_i \leq 0$ for every $k < i < r$ with $r-i$ odd. Then we have the inequalities (where $a_i = 0$ if $i \leq 0$)

$$a_{r-1} \leq a_{r-3} - a_{r-2} \leq a_{r-5} - a_{r-4} \leq \dots \leq a_{k+2} - a_{k+3} \leq a_k - a_{k+1}$$

and it follows that $a_j \geq a_{r-1} > 0$ for every $j \geq k$ with $r-j$ odd. In particular, k must be > 0 .

Let k be maximal with $r-k$ odd and $c_k > 0$. Denote $\gamma_0 = \sum_{i=0}^{(r-k-1)/2} \sigma_{k+2i} = -D_{k-1} + D_k$; then $\gamma_0 \leq_\Sigma \gamma$ and $\gamma^+ - \gamma_0 \in \mathbb{N}\Delta$. Hence we must have $\gamma = \gamma_0$, $k = 1$ and r even. \square

3.3. Type C_r ($r \geq 3$). Let $\Sigma = \{\sigma_1, \dots, \sigma_{r-1}\}$ be the set of spherical roots where $\sigma_i = \alpha_i + \alpha_{i+1}$.

Proposition 3.4. *Let $\gamma \in \mathbb{N}\Sigma$ be a covering difference in $\mathbb{N}\Delta$ with $\text{supp}_S(\gamma) = S$. Then $\gamma = \sum_{i=1}^{r-1} \sigma_i = D_2$.*

Proof. Clearly, $a_{r-1} > 0$. Let k be maximal such that $c_k > 0$. Then we have the following inequalities (for $k < r$)

$$a_{k-1} \geq a_k \geq \dots \geq a_{r-1}.$$

It follows that $a_i > 0$ for every $k-1 \leq i < r$. Therefore, $\gamma^+ - \sum_{i=k-1}^{r-1} \sigma_i = \gamma^+ + c_{k-2} - c_k \in \mathbb{N}\Delta$ and we get $k = 2$ with $\gamma = \sum_{i=1}^{r-1} \sigma_i$. \square

3.4. Type D_r ($r \geq 4$). Let $\Sigma = \{\sigma_1, \dots, \sigma_{r-1}\}$ be the set of spherical roots where $\sigma_i = \alpha_i + \alpha_{i+1}$ if $i < r-1$ and $\sigma_{r-1} = \alpha_{r-2} + \alpha_r$.

Proposition 3.5. *Let $\gamma \in \mathbb{N}\Sigma$ be a covering difference in $\mathbb{N}\Delta$ with $\text{supp}_S(\gamma) = S$. Then r is odd and $\gamma = \sum_{i=1}^{(j-1)/2} \sigma_{2i-1} + \sum_{i=(j+1)/2}^{(r-3)/2} 2\sigma_{2i-1} + \sigma_{r-2} + \sigma_{r-1} = D_1 - D_{j-1} + D_j$ for j odd $\leq r-2$.*

Proof. Clearly, we have the inequalities $a_{r-2} > 0$ and $a_{r-1} > 0$. One has $c_{r-2} \leq 1$, otherwise $\gamma^+ - (-2D_{r-3} + 2D_{r-2}) \in \mathbb{N}\Delta$, thus $\gamma = -2D_{r-3} + 2D_{r-2} = \sigma_{r-2} + \sigma_{r-1}$, but $\text{supp}_S(\gamma) \neq S$. Since $c_{r-2} = -a_{r-4} + a_{r-3} + a_{r-2} + a_{r-1}$ we get $a_{r-4} > 0$ and, moreover, if $c_{r-2} \leq 0$, $a_{r-4} > 1$.

We go along this way step-by-step. Let k be $< (r-1)/2$. Assume $a_{r-2i} > 0$ for all $i \leq k$ and, moreover, if for some $j < k$, $c_{r-2i} \leq 0$ for all $i \leq j$, then $a_{r-2i-2} > 1$ for all $i \leq j$. One has $\sum_{i=1}^k c_{r-2i} \leq 1$, otherwise there would exist $1 \leq i_1 \leq i_2 \leq k$ such that $\gamma^+ - (-D_{r-2i_2-1} + D_{r-2i_2} - D_{r-2i_1-1} + D_{r-2i_1}) \in \mathbb{N}\Delta$, thus $\gamma = -D_{r-2i_2-1} + D_{r-2i_2} - D_{r-2i_1-1} + D_{r-2i_1} = \sum_{i=i_1+1}^{i_2} \sigma_{r-2i} + \sum_{i=2}^{i_1} 2\sigma_{r-2i} + \sigma_{r-2} + \sigma_{r-1}$, but $\text{supp}_S(\gamma) \neq S$. Since

$$\sum_{i=1}^k c_{r-2i} = -a_{r-2k-2} + a_{r-2k-1} + a_{r-2} + a_{r-1},$$

we get $a_{r-2k-2} > 0$ and, moreover, if $c_{r-2i} \leq 0$ for all $i \leq k$, $a_{r-2k-2} > 1$.

Therefore, we have that r is odd, $\sum_{i=1}^{(r-1)/2} c_{r-2i} > 1$ and $\gamma = -D_{r-2i_2-1} + D_{r-2i_2} - D_{r-2i_1-1} + D_{r-2i_1}$ with $1 \leq i_1 \leq i_2 \leq (r-1)/2$. Since $\text{supp}_S(\gamma) = S$, we must have $i_2 = (r-1)/2$. \square

3.5. Type E_r ($6 \leq r \leq 8$). Let $\Sigma = \{\sigma_1, \dots, \sigma_{r-1}\}$ be the set of spherical roots where $\sigma_1 = \alpha_1 + \alpha_3$, $\sigma_2 = \alpha_2 + \alpha_4$ and, for $i \geq 3$, $\sigma_i = \alpha_i + \alpha_{i+1}$.

Proposition 3.6. *Let $\gamma \in \mathbb{N}\Sigma$ be a covering difference in $\mathbb{N}\Delta$ with $\text{supp}_S(\gamma) = S$.*

If $r = 6$,

- (1) *either $\gamma = \sigma_1 + \sigma_2 + \sigma_3 + \sigma_4 + \sigma_5 = D_4 - D_2$*
- (2) *or $\gamma = 2\sigma_1 + 2\sigma_2 + \sigma_3 + \sigma_4 + 2\sigma_5 = D_1 + D_6$.*

If $r = 7$,

- (3) $\gamma = \sigma_1 + \sigma_2 + \sigma_4 + \sigma_5 + \sigma_6 = D_1 + D_6 - D_3$.

If $r = 8$,

- (4) *either $\gamma = \sigma_1 + \sigma_2 + \sigma_3 + \sigma_4 + 2\sigma_5 + \sigma_6 + \sigma_7 = D_6 - D_2$*
- (5) *or $\gamma = 2\sigma_1 + 2\sigma_2 + \sigma_3 + \sigma_4 + 2\sigma_5 + \sigma_7 = D_1 + D_8 - D_7$.*

Proof. Suppose $r = 6$. Clearly, a_1, a_2, a_5 are > 0 . Moreover, $c_1 + c_4 + c_6 = a_2$.

If c_2 were > 0 , then c_1, c_4, c_6 would all be ≤ 0 ; indeed, either $\gamma^+ - \sigma_2$ or $\gamma^+ - (\sigma_1 + \sigma_2)$ or $\gamma^+ - (\sigma_2 + \sigma_5)$ would be in $\mathbb{N}\Delta$, contradicting $\text{supp}_S(\gamma) = S$ (we will use such an argument repeatedly in the rest of the proof). Therefore, $c_2 \leq 0$ and $a_3 + a_4 > 0$. By symmetry, we can assume $a_3 > 0$.

Assume $c_4 > 0$. Then $c_3 \leq 0$. If a_4 were zero, $c_2 + c_3$ would be equal to a_1 (which is > 0). Therefore, $a_4 > 0$ and $\gamma = D_4 - D_2$.

Assume now $c_4 \leq 0$. We can also assume $c_6 > 0$ (by symmetry, if $a_4 > 0$). Then both c_3 and c_5 are ≤ 0 , hence a_4 is > 0 . Since $c_2 + c_3 + c_4 + c_5 = 0$, all $c_i = 0$, for $2 \leq i \leq 5$. This implies $a_1, a_2, a_5 \geq 2$, $c_6 \leq 1$ (since $(2\sigma_1 + \sigma_2 + \sigma_4)^+ = 2D_1$) and $c_1 > 0$. Therefore, $\gamma = D_1 + D_6$.

Suppose $r = 7$. Clearly, a_1, a_2, a_6 are > 0 . Moreover, $c_1 + c_4 + c_6 = a_2 + a_6$.

Assume $a_3 = 0$. We get $c_1 > 0$. Then both c_2 and c_3 are ≤ 0 , hence a_4 is > 0 . Then $c_7 \leq 0$, hence $a_5 > 0$. Moreover, $c_4 \leq 0$ (since $(\sigma_1 + \sigma_2 + \sigma_4)^+ = D_1 + D_4$), $c_1 \leq 1$ (since $(2\sigma_1 + \sigma_2 + \sigma_4)^+ = 2D_1$) and $c_6 > 0$. Therefore, $\gamma = D_1 + D_6 - D_3$.

Assume now $a_3 > 0$. If c_4 were > 0 , then both c_2 and c_3 would be ≤ 0 , hence $a_4 > 0$. This would imply $c_7 \leq 0$, hence $a_5 > 0$, which is impossible (since $(\sigma_1 + \sigma_2 + \sigma_3 + \sigma_4 + \sigma_5)^+ = D_4$). Then $c_4 \leq 0$.

If c_1 were > 0 , then both c_2 and c_3 would be ≤ 0 , hence $a_4 > 0$. This would imply $c_7 \leq 0$ hence $a_5 > 0$, and $c_1 \leq 1$ (since $(2\sigma_1 + \sigma_2 + \sigma_4)^+ = 2D_1$) hence $c_6 > 0$, which is impossible (since $(\sigma_1 + \sigma_3 + \sigma_4 + \sigma_5 + \sigma_6)^+ = D_1 + D_6$). Then $c_1 \leq 0$.

If c_6 were > 0 , we would get $c_7 \leq 0$, $a_5 > 0$ and $c_6 \leq 1$, which is impossible (since $c_1 + c_4 + c_6 = a_2 + a_6 \geq 2$).

Suppose $r = 8$. Clearly, a_1, a_2, a_7 are > 0 . Moreover, $c_1 + c_4 + c_6 + c_8 = a_2$.

If c_4 were > 0 , we would get $c_2 \leq 0$, $a_3 + a_4 > 0$, actually $a_3 > 0$ ($a_3 = 0$ would imply $c_1 > 0$, but also $a_4 > 0$ and $c_1 \leq 0$), hence $c_3 \leq 0$, $a_4 > 0$, $c_4 \leq 1$ and $c_1 \leq 0$, therefore $a_5 > 0$ (since $a_2 + a_4 - a_5 = c_1 + c_4 \leq 1$), which is impossible (since $(\sigma_1 + \sigma_2 + \sigma_3 + \sigma_4 + \sigma_5)^+ = D_4$). Then $c_4 \leq 0$.

Assume $c_6 > 0$. Then $a_5 > 0$ (since $a_6 > 0$ implies $c_7 \leq 0$), hence both c_2 and c_5 are ≤ 0 .

If a_3 were zero, we would get $c_1 > 0$, $a_4 > 0$, $a_6 > 0$ (since $a_5 + a_6 = c_2 + c_5 \leq 0$), which is impossible (since $(\sigma_1 + \sigma_3 + \sigma_4 + \sigma_5 + \sigma_6)^+ = D_1 + D_6$). Then $a_3 > 0$, hence $c_3 \leq 0$ and $a_4 > 0$.

If a_6 were zero, since $c_2 + c_3 + c_4 + c_5 = -a_6$, all c_i would be zero, for $2 \leq i \leq 5$. This would imply $a_5 \geq 2$ (since $-2a_2 + a_5 = c_2 + c_5 = 0$), on the other hand c_8 would be > 0 , which is impossible (since $(\sigma_2 + \sigma_3 + 2\sigma_5 + \sigma_7)^+ = D_6 + D_8$). Then $a_6 > 0$, hence $c_7 \leq 0$ and $a_5 \geq 2$. Therefore, $\gamma = D_6 - D_2$.

Assume now $c_6 \leq 0$ and $c_1 > 0$. Then both c_2 and c_3 are ≤ 0 , hence $a_4 > 0$ and $c_5 \leq 0$. If a_6 were > 0 , we would get $c_7 \leq 0$, $a_5 > a_6$, $a_3 > 0$ (since $-2a_3 + a_5 - a_6 = c_2 + c_5 \leq 0$), this would imply $a_1 \geq 2$ hence $c_1 \leq 1$ (since $(2\sigma_1 + \sigma_2 + \sigma_4)^+ = 2D_1$) and on the other hand $a_2 \geq 2$ hence $c_8 > 0$ (since $c_1 + c_4 + c_6 + c_8 = a_2$), but this is impossible (since $(2\sigma_1 + 2\sigma_2 + \sigma_3 + \sigma_4 + 2\sigma_5 + \sigma_7)^+ = D_1 + D_8$). Then $a_6 = 0$, hence $a_5 > 0$ and $c_8 > 0$. Furthermore, since $c_i = 0$ for all $2 \leq i \leq 5$, $a_3 > 0$ and $a_1, a_2, a_5 \geq 2$. Therefore, $\gamma = D_1 + D_8 - D_7$.

Finally, assume both c_1 and $c_6 \leq 0$. Recall that $c_4 \leq 0$, therefore $c_8 \geq a_2 > 0$. Furthermore, $a_3 > 0$, $c_7 \leq 0$, $a_5 > 0$, $c_2 \leq 0$, $c_3 \leq 0$ and $a_4 > 0$. Then $a_2 \geq 2$, hence $c_8 \geq 2$, $a_7 \geq 2$ and $a_5 \geq 2$, which is impossible (since $(\sigma_2 + \sigma_3 + 2\sigma_5 + 2\sigma_7)^+ = 2D_8$). \square

3.6. Type F_4 . Let $\Sigma = \{\sigma_1, \sigma_2, \sigma_3\}$ be the set of spherical roots where $\sigma_i = \alpha_i + \alpha_{i+1}$.

Proposition 3.7. *Let $\gamma \in \mathbb{N}\Sigma$ be a covering difference in $\mathbb{N}\Delta$ with $\text{supp}_S(\gamma) = S$. Then either*

- (1) $\gamma = \sigma_1 + \sigma_2 + 2\sigma_3 = D_4$ or
- (2) $\gamma = \sigma_1 + \sigma_3 = D_1 + D_4 - D_3$.

Proof. Clearly, $a_1, a_3 > 0$. Since $\sigma_2^+ = D_2$, it follows that $c_2 \leq 0$. Then $c_4 > 0$ (since $c_2 + c_4 = a_1$), hence $a_3 > a_2$. If $a_2 > 0$, then $\gamma = D_4$, whereas if $a_2 = 0$, then $c_1 > 0$ and $\gamma = D_1 + D_4 - D_3$. \square

3.7. Type G_2 . Let $\Sigma = \{\sigma\}$ be the set of spherical roots where $\sigma = \alpha_1 + \alpha_2$. The proof of the following proposition is trivial.

Proposition 3.8. *Let $\gamma \in \mathbb{N}\Sigma$ be a covering difference in $\mathbb{N}\Delta$ with $\text{supp}_S(\gamma) = S$. Then $\gamma = \sigma = D_2 - D_1$.*

4. PROJECTIVE NORMALITY OF MODEL WONDERFUL VARIETIES

In this section we prove the following.

Theorem 4.1. *Let M be a model wonderful variety of simply connected type. The multiplication of global sections*

$$m_{\mathcal{L}, \mathcal{L}'} : \Gamma(M, \mathcal{L}) \otimes \Gamma(M, \mathcal{L}') \longrightarrow \Gamma(M, \mathcal{L} \otimes \mathcal{L}')$$

is surjective for all globally generated line bundles $\mathcal{L}, \mathcal{L}' \in \text{Pic}(M)$.

Following Lemma 2.3, we first classify all the low fundamental triples for the model wonderful varieties of simply connected type, and prove that these triples all satisfy the condition (2.1). As in the previous section, we can restrict ourselves to the case of an almost simple group G . As shown in the proof of Theorem 4.1 at the end of the section, it is enough to consider only low fundamental triples (D, E, F) such that $\text{supp}_S(D + E - F) = S$.

We keep the notation of the previous section. We denote by H the stabilizer of a point x_0 in the open G -orbit of M and by H_0 the intersection of the kernels of the multiplicative characters of H .

4.1. Type A_r .

Lemma 4.2. *Let (D_p, D_q, F) be a low fundamental triple with $\text{supp}_S(D_p + D_q - F) = S$. Then $F = 0$ and $p + q = r + 1$. If, moreover, r is odd, then p and q are even.*

Proof. Notice that every fundamental triple is low; indeed, by Proposition 3.2, for every $\eta \in \mathbb{N}\Sigma$ covering the difference in $\mathbb{N}\Delta$ one has $\text{ht}(\eta^+) = 2$, hence every color is minimal in $\mathbb{N}\Delta$ w.r.t. \leq_Σ . Therefore, we only need to compute the fundamental triples (D_p, D_q, F) with $\text{supp}_S(D_p + D_q - F) = S$.

Take a sequence

$$F = F_n <_\Sigma F_{n-1} <_\Sigma \dots <_\Sigma F_0 = D_p + D_q$$

such that $F_i \in \mathbb{N}\Delta$ and $\gamma_i = F_{i-1} - F_i$ is a covering difference for every $i \leq n$. By Proposition 3.2, it follows that if $\gamma_i^+ = F_{i-1} = D_s + D_t$ (with $1 \leq s \leq t \leq r$)

then $\gamma_i^- = F_i = D_{s-j} + D_{t+j}$ (with j equal to 1 or 2). Therefore, $\text{supp}_S(\gamma_n) = \text{supp}_S(D_p + D_q - F) = S$, $F = \gamma_n^- = 0$ and $q = r + 1 - p$.

If r is odd, all the covering differences $\gamma_i = F_{i-1} - F_i$ are of the type (2) of Proposition 3.2, then p is even. \square

Proposition 4.3. *Let r be odd and let (D, E, F) be a low fundamental triple with $\text{supp}_S(D + E - F) = S$. Then $s^{D+E-F}V_F \subset V_D V_E$.*

Proof. By the previous lemma, $(D, E, F) = (D_p, D_{r+1-p}, 0)$ and p is even (as well as $r + 1 - p$).

Set $\Delta_{\text{odd}} = \{D_i \in \Delta : i \text{ is odd}\}$; the subset $\Delta_{\text{odd}} \subset \Delta$ is distinguished and the quotient $M' = M/\Delta_{\text{odd}}$ is a symmetric wonderful variety (with spherical roots $\alpha_{2k-1} + 2\alpha_{2k} + \alpha_{2k+1}$).

By Corollary 1.4 together with the surjectivity of the multiplication map in the symmetric case we have

$$\Gamma(M, \mathcal{L}_D)\Gamma(M, \mathcal{L}_E) = \Gamma(M', \mathcal{L}_D)\Gamma(M', \mathcal{L}_E) = \Gamma(M', \mathcal{L}_{D+E}) = \Gamma(M, \mathcal{L}_{D+E}).$$

\square

Proposition 4.4. *Let r be even and let (D, E, F) be a low fundamental triple with $\text{supp}_S(D + E - F) = S$. Then $s^{D+E-F}V_F \subset V_D V_E$.*

Proof. By Lemma 4.2, we have $F = 0$ and $V(\omega_E) \simeq V(\omega_D)^*$, hence $V(\omega_D)^* \otimes V(\omega_E)^* \simeq \text{End}(V(\omega_D))$.

If r is even, the stabilizer H of a point in the open G -orbit of M is the normalizer in G of $\text{Sp}(r)$ and in particular is reductive (see [36]). Therefore, the one-dimensional H -submodules of $V(\omega_D)^*$ and of $V(\omega_E)^*$ associated respectively with D and E are dual to each other, hence we may choose the H -eigenvectors $h_D \in V(\omega_D)^*$ and $h_E \in V(\omega_E)^* \simeq V(\omega_D)$ in such a way that $h_D(h_E) = 1$. If we complete h_E and h_D to dual bases $\{h_E, v_1, \dots, v_n\} \subset V(\omega_D)$ and $\{h_D, v_1^*, \dots, v_n^*\} \subset V(\omega_D)^*$, then the identity $\text{Id} \in \text{End}(V(\omega_D))$ is a G -invariant element which, as tensor, is described as follows:

$$\text{Id} = h_D \otimes h_E + \sum_{i=1}^n v_i^* \otimes v_i.$$

Therefore, $h_D \otimes h_E$ has a non-zero projection on the isotypic component of highest weight zero and, by Lemma 1.2, we get that $s^{D+E}V_0 \subset V_D V_E$. \square

4.2. Type B_r .

Proposition 4.5. *There are no low triples (D, E, F) with $\sigma_{r-1} \in \text{supp}_\Sigma(D + E - F)$.*

Proof. Set

$$\Delta^{\text{even}} = \{D_i \in \Delta : r - i \text{ is even}\},$$

$$\Delta^{\text{odd}} = \{D_i \in \Delta : r - i \text{ is odd}\}.$$

If $F <_\Sigma D + E$ with $\sigma_{r-1} \in \text{supp}_\Sigma(D + E - F)$, then $\text{supp}(D + E) \cap \Delta^{\text{odd}} \neq \emptyset$. Indeed, if $\text{supp}(D + E)$ were included in Δ^{even} , take a sequence

$$F = F_n <_\Sigma F_{n-1} <_\Sigma \dots <_\Sigma F_0 = D + E$$

of coverings in $\mathbb{N}\Delta$: every covering difference $\gamma_i = F_{i-1} - F_i$ would necessarily be of the type (2) of Proposition 3.2, hence $\sigma_{r-1} \notin \text{supp}_\Sigma(D + E - F)$.

Let $k < r$ be the maximum such that $D_k \in \text{supp}(E+F) \cap \Delta^{\text{odd}}$. Set $D+E-F = \gamma = \sum_{i=1}^{r-1} a_i \sigma_i$. Since $c(\gamma, D_i) \leq 0$ for every $k < i < r$ with $r-i$ odd, as in the proof of Proposition 3.3 it follows that $a_j \geq a_{r-1} > 0$ for every $j \geq k$ with $r-j$ odd.

Denote $\gamma_0 = \sum_{i=0}^{(r-k-1)/2} \sigma_{k+2i} = -D_{k-1} + D_k$ and $F' = D + E - \gamma_0$. Then $F' \in \mathbb{N}\Delta$ and $F \leq_{\Sigma} F' <_{\Sigma} D + E$. Set

$$(D', E') = \begin{cases} (D - \gamma_0, E) & \text{if } D_k \in \text{supp}(D), \\ (D, E - \gamma_0) & \text{otherwise.} \end{cases}$$

Then $D' \leq_{\Sigma} D$, $E' \leq_{\Sigma} E$ and $F \leq_{\Sigma} D' + E' <_{\Sigma} D + E$, hence (D, E, F) is not a low triple. \square

4.3. Type C_r .

Proposition 4.6. *There are no low triples (D, E, F) with $\sigma_{r-1} \in \text{supp}_{\Sigma}(D+E-F)$.*

Proof. Let $D, E, F \in \mathbb{N}\Delta$ with $F <_{\Sigma} D+E$ and $\sigma_{r-1} \in \text{supp}_{\Sigma}(D+E-F)$. Denote $k \leq r$ the maximum such that $D_k \in \text{supp}(D+E)$. Reasoning as in the proof of Proposition 3.4, it follows that $\sigma_i \in \text{supp}_{\Sigma}(D+E-F)$ for every $k-1 \leq i < r$. Therefore, if we set $\gamma_0 = \sum_{i=k-1}^{r-1} \sigma_i = -D_{k-2} + D_k$ and $F' = D + E - \gamma_0$, then $F' \in \mathbb{N}\Delta$ and $F \leq_{\Sigma} F' <_{\Sigma} D + E$. Set

$$(D', E') = \begin{cases} (D - \gamma_0, E) & \text{if } D_k \in \text{supp}(E), \\ (D, E - \gamma_0) & \text{otherwise.} \end{cases}$$

Then $D' \leq_{\Sigma} D$, $E' \leq_{\Sigma} E$ and $F \leq_{\Sigma} D' + E' <_{\Sigma} D + E$, hence (D, E, F) is not a low triple. \square

4.4. Type D_r .

Lemma 4.7. *Let (D_p, D_q, F) be a low fundamental triple with $\text{supp}_S(D_p + D_q - F) = S$. Then p, q, r are odd, $p, q \leq r-2$ and either*

- (1) $p+q \leq r-1$ with $F = D_{p+q-2}$ or
- (2) $p+q = r+1$ with $F = D_{r-1} + D_r$.

Proof. Notice that, as in type A, every fundamental triple is low. Therefore, we only need to compute the fundamental triples (D_p, D_q, F) with $\text{supp}_S(D_p + D_q - F) = S$. Assume $p \leq q$.

Take a sequence

$$F = F_n <_{\Sigma} F_{n-1} <_{\Sigma} \dots <_{\Sigma} F_0 = D_p + D_q$$

such that $F_i \in \mathbb{N}\Delta$ and $\gamma_i = F_{i-1} - F_i$ is a covering difference for every $i \leq n$. Recall the classification of covering differences of Propositions 3.2 and 3.5.

(1) $p, q \leq r-2$. If q were equal to $r-1$ then $(r-1)-p$ should be non-zero and odd, thus $\gamma_1 = -D_{p-1} + D_p + D_{r-1} - D_r$ and $F_1 = D_{p-1} + D_r$, which is impossible since the distance between the vertices r and $p-1$ is even. By symmetry, the same argument works if $q = r$.

(2) $q-p$ is even. If $q-p$ were odd, there would exist i such that F_i is $D_{p'} + 2D_{q'}$ with q' equal to $r-1$ or r and $(r-1)-p'$ even, which is impossible.

(3) $r-p$ is even (as well as $r-q$). Here again if $r-p$ and $r-q$ were odd, there would exist i such that F_i is $D_{p'} + 2D_{q'}$ with q' equal to $r-1$ or r and $(r-1)-p'$ even.

Therefore, there exists i such that γ_j is of the type (2) of Proposition 3.2 for every $j < i$ and γ_i is either of type (1) of Proposition 3.2 or of type D.

In the first case, $F_{i-2} = D_{p'} + D_{q'}$ with $q' = r - 2$ and $p' + q' = p + q$, $F_{i-1} = D_{p'-2} + 2D_{q''}$ where q'' is equal to $r - 1$ or r . Then necessarily $p' - 2 = 1$ and $F_i = F_n = D_{r-1} + D_r$.

In the second case, $F_{i-1} = D_{p'} + D_{q'}$ with $p' + q' = p + q$ and $F_i = D_{p'-1} + D_{q'-1}$. Then γ_j is of the type (2) of Proposition 3.2 for every $j > i$ until $F_{n-1} = D_{p'-1-2k} + D_{q'-1+2k}$ where $p' - 1 - 2k = 2$ ($k = n - 1 - i$) and F_n is equal to either $D_{q'-1+2k+2}$ if $q' - 1 + 2k \leq r - 5$ or $D_{r-1} + D_r$ if $q' - 1 + 2k = r - 3$. \square

Proposition 4.8. *Let (D, E, F) be a low fundamental triple with $\text{supp}_S(D + E - F) = S$, then $s^{D+E-F}V_F \subset V_D V_E$.*

Proof. We need an explicit computation.

Denote by $U = \mathbb{C}^{2r}$ the first fundamental representation of G (that is, the standard representation of $\text{SO}(2r)$) and fix $b \in S^2U$ a G -invariant non-degenerate symmetric 2-form. If $W \subset U$ is a maximal isotropic subspace, we then get a decomposition $U = W \oplus W^*$. Fix a non-zero vector $e_0 \in W$ and consider the corresponding decomposition $U = V \oplus \mathbb{C}e_0 \oplus V^* \oplus \mathbb{C}e_0^*$, where $V \subset W$ is a complement of the line $\mathbb{C}e_0$ and where $e_0^* \in V^*$ is defined by $e_0^*(e_0) = 1$, $e_0^*|_V = 0$.

Then (see [36]) the Lie algebra \mathfrak{h}_0 of H_0 can be described as

$$\mathfrak{h}_0 = \mathfrak{sp}(V) \oplus V^* \oplus \text{Skew}(V, V^*),$$

where $\text{Skew}(V, V^*) \subset \text{Hom}(V, V^*)$ denotes the subspace of skew-symmetric linear maps and where \mathfrak{h}_0 is embedded in $\mathfrak{so}(U)$ as follows (here we denote by $u = (v, \lambda e_0, \psi, \mu e_0^*)$ a generic element in $U = V \oplus \mathbb{C}e_0 \oplus V^* \oplus \mathbb{C}e_0^*$):

- if $f \in \mathfrak{sp}(V)$, then $f(u) = f(v) + f \cdot \psi$,
- if $\phi \in V^*$, then $\phi(u) = \phi(v)e_0 - (\lambda + \mu)\phi + \phi(v)e_0^*$,
- if $\Phi \in \text{Skew}(V, V^*)$, then $\Phi(u) = \Phi(v)$.

As already mentioned at the beginning of Section 3 every simple G -module possesses a unique \mathfrak{h}_0 -invariant element, up to scalars. In particular, if we denote $h_1 = e_0 - e_0^* \in U$ and $h_2 \in \Lambda^2 V^*$ a $\mathfrak{sp}(V)$ -invariant non-degenerate 2-form, then $h_1 \in U^{\mathfrak{h}_0}$ and $h_2 \in (\Lambda^2 U)^{\mathfrak{h}_0}$. In this way we may describe the \mathfrak{h}_0 -invariant vectors in every exterior power $\Lambda^i U$ with $i \leq r - 1$. Indeed, set

$$h_i = \begin{cases} h_2^{\wedge k} & \text{if } i = 2k \text{ is even,} \\ h_1 \wedge h_2^{\wedge k} & \text{if } i = 2k + 1 \text{ is odd,} \end{cases}$$

then $h_i \in (\Lambda^i U)^{\mathfrak{h}_0}$.

Set $\omega_0 = 0$ and recall that if $i \geq 0$ then

$$\Lambda^i U = \begin{cases} V(\omega_i) & \text{if } i \leq r - 2, \\ V(\omega_{r-1} + \omega_r) & \text{if } i = r - 1. \end{cases}$$

To conclude the proof, by Lemma 1.2, we only need to show that, if i, j are odd integers with $i + j \leq r + 1$, then there exists an equivariant projection $\pi: \Lambda^i U \otimes \Lambda^j U \rightarrow \Lambda^{i+j-2} U$ such that $\pi(h_i \otimes h_j) \neq 0$. Define π as follows:

$$\pi((u_1 \wedge \cdots \wedge u_i) \otimes (w_1 \wedge \cdots \wedge w_j)) = \sum_{h,k} (-1)^{h+k} b(u_h, w_k) u_1 \wedge \cdots \wedge \hat{u}_h \wedge \cdots \wedge \hat{w}_k \wedge \cdots \wedge w_j.$$

Suppose that i, j are odd with $i + j \leq r + 1$ and set $k = (i + j - 2)/2$. Notice that $\pi(h_i \otimes h_j) = b(h_1, h_1)h_2^{\wedge k} + q$, where $q \in \Lambda^{i+j-2}U$ is linearly independent with $h_2^{\wedge k}$; since $b(h_1, h_1) \neq 0$ and since $h_2^{\wedge k} \neq 0$, it follows then $\pi(h_i \otimes h_j) \neq 0$. \square

4.5. Type E_r .

Lemma 4.9. *The low fundamental triples (D, E, F) with $\text{supp}_S(D + E - F) = S$ are the following:*

- If $r = 6$: (D_1, D_3, D_2) (D_1, D_5, D_3) $(D_1, D_6, 0)$ (D_3, D_6, D_5) (D_5, D_6, D_2) ,
- If $r = 7$: (D_1, D_6, D_3) $(D_6, D_6, D_2 + D_7)$,
- If $r = 8$: (D_1, D_1, D_2) $(D_1, D_5, 2D_2)$ (D_1, D_7, D_3) (D_1, D_8, D_7) (D_3, D_8, D_5)
 $(D_5, D_8, D_2 + D_7)$ (D_7, D_8, D_2) .

By making use of Proposition 3.6, the proof of this lemma is a quite long but trivial case-by-case computation, which we do not report here.

Proposition 4.10. *Let (D, E, F) be a low fundamental triple with $\text{supp}_S(D + E - F) = S$, then $s^{D+E-F}V_F \subset V_D V_E$.*

Let us first deal with the triple $(D_1, D_6, 0)$ of E_6 . The set $\Delta_0 = \{D_2, D_3, D_4, D_5\}$ is distinguished and the quotient $M' = M/\Delta_0$ is a symmetric wonderful variety (with spherical roots $\{2\alpha_1 + \alpha_2 + 2\alpha_3 + 2\alpha_4 + \alpha_5, \alpha_2 + \alpha_3 + 2\alpha_4 + 2\alpha_5 + 2\alpha_6\}$). Therefore, we can conclude by Corollary 1.4 and the surjectivity of the multiplication map in the symmetric case.

For all the other triples of Lemma 4.9 we have to use Lemma 1.2. Since the dimension of the involved representations is quite high, we have used the computer and, more precisely, GAP [23], a software for computations which contains built-in functions to construct and deal with representations of simple Lie algebras (see also [21]).

Although the dimension of some of the involved representations is very high, we have succeeded to make the computation accessible with a currently available home computer. A quite convenient tool is the quadratic Casimir operator c , which acts as the scalar $(\lambda + 2\rho, \lambda)$ on every irreducible representation $V(\lambda)$. Let (D, E, F) be a low fundamental triple, once the irreducible representations V_D^* and V_E^* are constructed, and the vectors h_D and h_E are explicitly found, there is no need to construct the whole tensor product $V_D^* \otimes V_E^*$ in order to project $h_D \otimes h_E$ onto the eigenspace of c relative to $(\omega_F^* + 2\rho, \omega_F^*)$. Moreover, in all our cases there exists only one isotypical component in $V_D^* \otimes V_E^*$ of eigenvalue $(\omega_F^* + 2\rho, \omega_F^*)$ with respect to c .

Sometimes for E_8 the dimension is so high that it is even quite costly to construct the irreducible representation V_D^* itself (V_{D_5} has dimension 146,325,270). Here we use a further escamotage. The set $\Delta_0 = \{D_1, D_4, D_6, D_8\}$ is distinguished and the quotient $M' = M/\Delta_0$ is the parabolic induction of a symmetric wonderful $SL(8)$ -variety with spherical roots $\{\alpha_1 + 2\alpha_3 + \alpha_4, \alpha_4 + 2\alpha_5 + \alpha_6, \alpha_6 + 2\alpha_7 + \alpha_8\}$. This means that, if D equals D_3, D_5 or D_7 , then h_D can be chosen to be the $Sp(8)$ -invariant vector in the simple $SL(8)$ -submodule $W_D \subset V_D$ of highest weight ω_D . Furthermore, V_{D_1} and V_{D_8} are still accessible (the former has dimension 3875 and the latter is the adjoint representation), $V_{D_7} \subset \Lambda^2 V_{D_8}$, $V_{D_5} \subset \Lambda^4 V_{D_8}$, $V_{D_3} \subset \Lambda^2 V_{D_1}$, and respectively $W_{D_7} \subset \Lambda^2 W_{D_8}$, $W_{D_5} \subset \Lambda^4 W_{D_8}$, $W_{D_3} \subset \Lambda^2 W_{D_1}$. Therefore, the vectors h_D can be explicitly determined if D equals D_1, D_3, D_5, D_7 or D_8 , and notice that this is enough to treat all the above triples.

4.6. Type F_4 .

Lemma 4.11. *The only low fundamental triple (D, E, F) with $\text{supp}_S(D+E-F) = S$ is (D_1, D_4, D_3) .*

The proof is trivial, after Proposition 3.7.

Proposition 4.12. $s^{D_1+D_4-D_3}V_{D_3} \subset V_{D_1}V_{D_4}$.

This can easily be checked via computer (as explained above).

4.7. Type G_2 .

Proposition 4.13. *There are no low triples (D, E, F) with $F \neq D + E$.*

The proof is trivial.

4.8. Projective normality of model wonderful varieties. We are now ready to prove that the multiplication of sections on a model wonderful variety of simply connected type is surjective.

A *localization* of a wonderful variety M is a G -stable subvariety of M , which is a wonderful G -variety by itself. Notice that we have a bijective correspondence between localizations of M and subsets of Σ . More precisely, for all subsets Σ' of Σ , the intersection of the boundary divisors M^σ for $\sigma \in \Sigma'$ gives a wonderful variety with $\Sigma \setminus \Sigma'$ as a set of spherical roots.

Proof of Theorem 4.1. By the classification of the covering differences given in the previous section, it follows that the model wonderful varieties of simply connected type satisfy the property (2-ht). Hence by Lemma 2.3 it is enough to show the inclusion $s^{D+E-F}V_F \subset V_DV_E$ for all the low fundamental triples (D, E, F) . Here we only need to show that we can reduce to the low fundamental triples (D, E, F) such that $\text{supp}_S(D+E-F) = S$, whose case has been proved above in this section.

Let (D, E, F) be a low fundamental triple, denote $\gamma = D + E - F$ and $S' = \text{supp}_S(\gamma)$ and suppose that $S' \neq S$. Let Q be the parabolic subgroup associated to S' and set $G' = Q/R_Q$. Since (D, E, F) is a low triple, it follows that S' is connected, hence G' is an almost simple group. Consider the localization M' of M with $\Sigma' = \{\sigma \in \Sigma : \text{supp}_S(\sigma) \subset S'\}$ as a set of spherical roots. Then M' is the parabolic induction of the model wonderful G' -variety N of simply connected type.

The restrictions $\text{Pic}(M) \rightarrow \text{Pic}(M')$ and $\text{Pic}(M') \rightarrow \text{Pic}(N)$ identify $\text{Pic}(N)$ with a sublattice of $\text{Pic}(M)$. More precisely, if $\tilde{\Delta} \subset \Delta$ is the set of colors \tilde{D} such that $\omega_{\tilde{D}} = \omega_\alpha$ for some $\alpha \in S'$, then $\text{Pic}(N)$ is identified with $\mathbb{Z}\tilde{\Delta}$. Moreover, an element of $\mathbb{Z}\tilde{\Delta}$ induces a globally generated line bundle on N if and only if its coefficients are non-negative. If $\tilde{D} \in \mathbb{N}\tilde{\Delta}$, set $W_{\tilde{D}} = V_{\tilde{D}}^{U_Q}$.

Notice that $D, E \in \tilde{\Delta}$ and consider the difference $\tilde{F} = D + E - \gamma \in \mathbb{Z}\tilde{\Delta}$ (where γ is regarded as sum of spherical roots of N hence as an element of $\mathbb{Z}\tilde{\Delta}$). Since (D, E, F) is a low triple in $\mathbb{N}\Delta$, it follows that (D, E, \tilde{F}) is a low triple in $\mathbb{N}\tilde{\Delta}$. Since $\text{supp}_{S'}(\gamma) = S'$, we have that $s^\gamma W_{\tilde{F}} \subset W_D W_E$, therefore Proposition 1.6 implies that $s^\gamma V_F \subset V_D V_E$ with respect to the multiplication map $m'_{D,E}: \Gamma(M', \mathcal{L}_D) \otimes \Gamma(M', \mathcal{L}_E) \rightarrow \Gamma(M', \mathcal{L}_{D+E})$. This concludes the proof, since $m'_{D,E}$ is just the restriction of the multiplication map $m_{D,E}: \Gamma(M, \mathcal{L}_D) \otimes \Gamma(M, \mathcal{L}_E) \rightarrow \Gamma(M, \mathcal{L}_{D+E})$. \square

Let M be a wonderful variety and let N be a quotient of M . Then the pull-back of line bundles identifies $\text{Pic}(N)$ with a sublattice of $\text{Pic}(M)$ and a line bundle $\mathcal{L} \in \text{Pic}(N)$ is generated by global sections if and only if its pull-back (which we still denote by \mathcal{L}) is. Moreover, by Corollary 1.4 we have $\Gamma(N, \mathcal{L}) = \Gamma(M, \mathcal{L})$. It follows that if $\mathcal{L}, \mathcal{L}' \in \text{Pic}(N)$ are generated by global sections and if the multiplication $\Gamma(M, \mathcal{L}) \otimes \Gamma(M, \mathcal{L}') \rightarrow \Gamma(M, \mathcal{L} \otimes \mathcal{L}')$ is surjective, then the multiplication $\Gamma(N, \mathcal{L}) \otimes \Gamma(N, \mathcal{L}') \rightarrow \Gamma(N, \mathcal{L} \otimes \mathcal{L}')$ is also surjective.

Now let N be a localization of M and let $\mathcal{L}, \mathcal{L}' \in \text{Pic}(M)$ be generated by global sections. Then the restriction of sections to N is surjective, therefore if the multiplication $\Gamma(M, \mathcal{L}) \otimes \Gamma(M, \mathcal{L}') \rightarrow \Gamma(M, \mathcal{L} \otimes \mathcal{L}')$ is surjective, the multiplication $\Gamma(N, \mathcal{L}|_N) \otimes \Gamma(N, \mathcal{L}'|_N) \rightarrow \Gamma(N, \mathcal{L}|_N \otimes \mathcal{L}'|_N)$ is also surjective.

If, moreover, M is a model wonderful variety of simply connected type, then by the description of the restriction $\omega: \text{Pic}(M) \rightarrow \mathcal{X}(T)$ (see [42, Lemma 30.24]) we get an isomorphism $\text{Pic}(M) \simeq \mathcal{X}(T)$, which identifies the semigroup of globally generated (resp. ample) line bundles on M with $\mathcal{X}(T)^+$ (resp. with the semigroup of regular dominant weights). If N is any localization of M , then ω factors through N and we get an isomorphism $\text{Pic}(M) \simeq \text{Pic}(N)$, which identifies the globally generated (resp. the ample) line bundles on M and on N . Up to replacing $\mathcal{X}(T)$ with some sublattice generated by fundamental weights, the same is true whenever M' is the quotient of a localization of a model wonderful variety of simply connected type and N' is a localization of M' . Therefore, proceeding inductively, we get the following corollary of Theorem 4.1.

Corollary 4.14. *Let N be a wonderful variety obtained from a model wonderful variety of simply connected type via operations of localization and quotient by distinguished sets of colors. Then the multiplication map*

$$m_{\mathcal{L}, \mathcal{L}'}: \Gamma(N, \mathcal{L}) \otimes \Gamma(N, \mathcal{L}') \rightarrow \Gamma(N, \mathcal{L} \otimes \mathcal{L}')$$

is surjective for all globally generated line bundles $\mathcal{L}, \mathcal{L}' \in \text{Pic}(N)$.

5. PROJECTIVE NORMALITY OF COMODEL WONDERFUL VARIETIES

Motivated by an application which we illustrate below in Section 8, here we study the surjectivity of the multiplication of sections in another class of wonderful varieties, which we call *comodel*.

Let G be simply connected of simply-laced type, and fix T and B as usual. Let M be a model wonderful variety of G , with set of colors Δ (in bijection with the set of fundamental weights), set of spherical roots Σ and Cartan pairing $c: \Delta \times \Sigma \rightarrow \mathbb{Z}$. Let G^\vee be a simply connected group whose root system is isomorphic to Φ_Σ , the root system generated by Σ . Once fixed T^\vee and B^\vee , its set of simple roots S^\vee is thus in bijective correspondence with Σ . From [5] and [9] it follows that:

Theorem 5.1. *There exists a wonderful G^\vee -variety M^\vee whose set of spherical roots Σ^\vee is equal to the set of simple roots S^\vee of G^\vee , its set of colors Δ^\vee is in bijective correspondence with Δ and, under these correspondences, its Cartan pairing $c^\vee: \Delta^\vee \times \Sigma^\vee \rightarrow \mathbb{Z}$ equals the Cartan pairing c of the model wonderful G -variety M .*

We say that the wonderful variety M^\vee of the previous theorem is a *comodel wonderful variety* of G^\vee , and the type of G is called the *cotype* of M^\vee .

Forgetting the model wonderful variety, in the following M is a comodel wonderful variety of G , H denotes the stabilizer of a point x_0 in the open orbit of M , and H_0 the kernel of its multiplicative characters; \mathfrak{h}_0 denotes the Lie algebra of H_0 in the Lie algebra \mathfrak{g} of G .

The comodel wonderful varieties correspond to the following cases in [5], an explicit description of the corresponding subgroups H can be found in [9].

- Cotype A_{2m} : type $A_{m-1} \times A_m$, case S-5.
- Cotype A_{2m+1} : type $A_m \times A_m$, case S-4.
- Cotype D_{2m} : type $A_{m-1} \times D_m$, case S-10.
- Cotype D_{2m+1} : type $A_{m-1} \times D_{m+1}$, case S-11.
- Cotype E_6 : type A_5 , case S-50.
- Cotype E_7 : type A_6 , case S-49.
- Cotype E_8 : type D_7 , case S-58.

In this section we prove the following.

Theorem 5.2. *Let M be a comodel wonderful variety and let $\mathcal{L}, \mathcal{L}'$ be line bundles generated by global sections. Then the multiplication map $m_{\mathcal{L}, \mathcal{L}'}$ is surjective.*

Let M be a comodel wonderful variety, we enumerate the colors of M as in the corresponding model wonderful variety. Similarly, we denote by $h_i \in V_{D_i}^*$ the H -semi-invariant associated to D_i . This vector is invariant under H_0 .

More explicitly, set the map $\omega: \text{Pic}(M) \rightarrow \mathcal{X}(T)$ as follows (if the Dynkin diagram of G has two connected components, we distinguish with a superscript the fundamental weights coming from the different components).

- Cotype A_{2m} : $\omega(D_1) = \omega'_1$, $\omega(D_{2i}) = \omega_i + \omega'_i$ for $i < m$, $\omega(D_{2i-1}) = \omega_{i-1} + \omega'_i$ for $i > 1$, $\omega(D_{2m}) = \omega'_m$.
- Cotype A_{2m+1} : $\omega(D_1) = \omega_1$, $\omega(D_{2i}) = \omega_i + \omega'_i$, $\omega(D_{2i-1}) = \omega_i + \omega'_{i-1}$ for $1 < i < m+1$, $\omega(D_{2m+1}) = \omega'_m$.
- Cotype D_{2m} : $\omega(D_1) = \omega_1$, $\omega(D_{2i}) = \omega_i + \omega'_i$ for $i < m-1$, $\omega(D_{2i-1}) = \omega_i + \omega'_{i-1}$ for $1 < i < m$, $\omega(D_{2m-2}) = \omega_{m-1} + \omega'_{m-1} + \omega'_m$, $\omega(D_{2m-1}) = \omega'_{m-1}$, $\omega(D_{2m}) = \omega'_m$.
- Cotype D_{2m+1} : $\omega(D_1) = \omega'_1$, $\omega(D_{2i}) = \omega_i + \omega'_i$ for $i < m$, $\omega(D_{2i-1}) = \omega_{i-1} + \omega'_i$ for $1 < i < m$, $\omega(D_{2m-1}) = \omega_{m-1} + \omega'_m + \omega'_{m+1}$, $\omega(D_{2m}) = \omega'_m$, $\omega(D_{2m+1}) = \omega'_{m+1}$.
- Cotype E_6 : $\omega(D_1) = \omega_2$, $\omega(D_2) = \omega_3$, $\omega(D_3) = \omega_2 + \omega_5$, $\omega(D_4) = \omega_1 + \omega_3 + \omega_5$, $\omega(D_5) = \omega_1 + \omega_4$, $\omega(D_6) = \omega_4$.
- Cotype E_7 : $\omega(D_1) = \omega_3$, $\omega(D_2) = \omega_4$, $\omega(D_3) = \omega_3 + \omega_6$, $\omega(D_4) = \omega_2 + \omega_4 + \omega_6$, $\omega(D_5) = \omega_2 + \omega_5$, $\omega(D_6) = \omega_1 + \omega_5$, $\omega(D_7) = \omega_1$.
- Cotype E_8 : $\omega(D_1) = \omega_3$, $\omega(D_2) = \omega_4$, $\omega(D_3) = \omega_3 + \omega_6$, $\omega(D_4) = \omega_2 + \omega_4 + \omega_6$, $\omega(D_5) = \omega_2 + \omega_5$, $\omega(D_6) = \omega_1 + \omega_5$, $\omega(D_7) = \omega_1 + \omega_7$, $\omega(D_8) = \omega_7$.

Since the Cartan pairing of the comodel wonderful varieties is the same as that of the model varieties, the classification of the covering differences is also the same and the property (2-ht) holds. As shown in Section 5.6, in order to apply Lemma 2.3 it is enough to test the surjectivity on the same low fundamental triples arising in Section 4 in the model case.

In the computations below we use the following conventions. We denote by e_1, \dots, e_n the standard basis of $V = \mathbb{C}^n$ and by $\varphi_1, \dots, \varphi_n$ the dual basis. We also denote by $e_{i_1 i_2 \dots i_k}$ the vector $e_{i_1} \wedge e_{i_2} \wedge \dots \wedge e_{i_k} \in \Lambda^k V$ and similarly for $\varphi_{i_1 i_2 \dots i_k}$. On $V \oplus V^*$ the symmetric bilinear form $(u, \varphi) \cdot (v, \psi) = \varphi(v) + \psi(u)$ is

defined. A pairing between $\Lambda^k V$ and $\Lambda^{n-k} V$ is defined by $\langle x, y \rangle = \frac{x \wedge y}{e_{1, \dots, n}}$ and we denote by $\gamma_k: \Lambda^k V \rightarrow \Lambda^{n-k} V^*$ the associated map. We also identify $V_1 \otimes V_2$ with $\text{Hom}(V_1^*, V_2)$ in the usual way, and if $V_1 = V_2 = V$ we identify $S^2 V$ and $\Lambda^2 V$ with symmetric and antisymmetric linear maps in $\text{Hom}(V^*, V)$.

Starting with the vector space V we construct a new vector space $W = V \oplus V^* \oplus Z$ as the direct sum of V , V^* and another piece Z that is zero, one- or two-dimensional. In particular, the dual of W can be identified with $V^* \oplus V \oplus Z^*$. Once a basis of Z is fixed, say z_1, \dots, z_t , we denote the dual basis of $e_1, \dots, e_n, \varphi_1, \dots, \varphi_n, z_1, \dots, z_t$ by $e_1^*, \dots, e_n^*, \varphi_1^*, \dots, \varphi_n^*, z_1^*, \dots, z_t^*$.

For a vector space U we have contraction maps $\kappa_U^{ij}: \Lambda^i U \otimes \Lambda^j U^* \rightarrow \Lambda^{i-1} U \otimes \Lambda^{j-1} U^*$ given by

$$\begin{aligned} & \kappa_U^{ij}(u_1 \wedge \dots \wedge u_i \otimes \varphi_1 \wedge \dots \wedge \varphi_j) \\ &= \sum_{k,l} (-1)^{k+l} \varphi_l(u_k) u_1 \wedge \dots \wedge \hat{u}_k \wedge \dots \wedge u_i \otimes \varphi_1 \wedge \dots \wedge \hat{\varphi}_l \wedge \dots \wedge \varphi_j. \end{aligned}$$

In particular, we set $\kappa_U = \kappa_U^{21}$.

Similarly, for a vector space U with a symmetric bilinear form $(,)$ we have $\tilde{\kappa}_U^{ij}: \Lambda^i U \otimes \Lambda^j U \rightarrow \Lambda^{i+j-2} U$ such that

$$\begin{aligned} & \tilde{\kappa}_U^{ij}(u_1 \wedge \dots \wedge u_i \otimes v_1 \wedge \dots \wedge v_j) \\ &= \sum_{k,l} (-1)^{k+l} (u_k, v_l) u_1 \wedge \dots \wedge \hat{u}_k \wedge \dots \wedge u_i \wedge v_1 \wedge \dots \wedge \hat{v}_l \wedge \dots \wedge v_j. \end{aligned}$$

5.1. Cotype A_r . We test the triples of Lemma 4.2: $(D_p, D_{r+1-p}, 0)$ where, if r is odd, p is even.

If r is odd, the set Δ_{odd} of odd-indexed colors is distinguished and the quotient is a symmetric wonderful variety, so the surjectivity follows as in Proposition 4.3.

If r is even, H is reductive and the surjectivity follows as in Proposition 4.4.

5.2. Cotype D_r . We test the triples of Lemma 4.7: (D_p, D_q, F) where p, q, r are odd, $p, q \leq r-2$ and either

- $p+q \leq r-1$ with $F = D_{p+q-2}$ or
- $p+q = r+1$ with $F = D_{r-1} + D_r$.

Let $m > 1$ and set $r = 2m + 1$. Let $V = \mathbb{C}^m$, and set $W = V \oplus \mathbb{C}e \oplus V^* \oplus \mathbb{C}\varepsilon$. On W is defined a quadratic form such that V and V^* are orthogonal to e and ε , moreover, $(e, \varepsilon) = 1$ and $(e, e) = (\varepsilon, \varepsilon) = 0$. Let $G = \text{SL}(V) \times \text{Spin}(W)$. We have $\mathfrak{h}_0 = \mathfrak{sl}(V) \oplus \Lambda^2 V \oplus V$, which is embedded as a subalgebra of \mathfrak{g} as follows. First notice that we have a natural immersion of $\text{SL}(V)$ in $\text{Spin}(W)$ so that $\mathfrak{sl}(V)$ can be included diagonally in \mathfrak{g} . The action of $\Lambda^2 V$ on W is zero on V , e and ε , while its action on V^* is given by the identification of $\Lambda^2 V$ with the antisymmetric maps from V^* to V . The action of V on W is given as follows: if $v, u \in V$ and $\varphi \in V^*$ then

$$v \cdot u = 0 \quad v \cdot e = v \quad v \cdot \varphi = -\varphi(v)(e + \varepsilon) \quad v \cdot \varepsilon = v.$$

Now let $\tilde{D}_i = D_i$ for i odd or for $i \neq 2m$ while $\tilde{D}_{2m} = D_{2m} + D_{2m+1}$. We have

$$V_{\tilde{D}_{2i}}^* = \Lambda^i V^* \otimes \Lambda^i W \quad V_{\tilde{D}_{2i+1}}^* = \Lambda^i V^* \otimes \Lambda^{i+1} W.$$

The \mathfrak{h}_0 -invariants in $V_{\tilde{D}_{2i}}^*$ are generated by the vector h_{2i} corresponding to the identity element in $\Lambda^i V^* \otimes \Lambda^i V \subset \Lambda^i V^* \otimes \Lambda^i W$. Similarly, the \mathfrak{h}_0 -invariants in

$V_{D_{2i+1}}^*$ are generated by the vector $h_{2i+1} = h_{2i} \wedge (e - \varepsilon)$. So that we have

$$h_{2i} = \sum_{j_1 < \dots < j_i} \varphi_{j_1 \dots j_i} \otimes e_{j_1 \dots j_i} \quad \text{and} \quad h_{2i+1} = \sum_{j_1 < \dots < j_i} \varphi_{j_1 \dots j_i} \otimes (e_{j_1 \dots j_i} \wedge (e - \varepsilon)).$$

Finally, if $p = 2t + 1$ and $q = 2s + 1$ then the projection $\Phi: V_{D_p}^* \otimes V_{D_q}^* \longrightarrow V_{D_{p+q-2}}^*$ is given by

$$\Phi((x \otimes w) \otimes (y \otimes z)) = x \wedge y \otimes \tilde{\kappa}_W^{t+1, s+1}(w \otimes z)$$

for $x \in \Lambda^t V^*$, $y \in \Lambda^s V^*$, $w \in \Lambda^{t+1} W$ and $z \in \Lambda^{s+1} W$. A direct computation shows that

$$\Phi(h_p \otimes h_q) = (-1)^{t+s+1} 2 \binom{t+s}{t} h_{p+q-2}.$$

5.3. Cotype E_6 . Let $V = \mathbb{C}^3$ and $W = V \oplus V^*$. Let $\mathfrak{g} = \mathfrak{sl}(W)$ and $\mathfrak{h}_0 = \mathfrak{sl}(V) \oplus S^2 V$ where the action of $\mathfrak{sl}(V)$ is the natural one and the action of $S^2 V$ is given by $b \cdot (v, \varphi) = (b(\varphi), 0)$ while the action on $W^* = V^* \oplus V$ is given by $b \cdot (\varphi, v) = (0, -b(\varphi))$.

We analyze the triples of Lemma 4.9: (D_1, D_3, D_2) , (D_1, D_5, D_3) , $(D_1, D_6, 0)$, (D_3, D_6, D_5) , (D_5, D_6, D_2) .

The set of colors $\{D_2, D_3, D_4, D_5\}$ is distinguished and the associated quotient is a symmetric wonderful variety, so the third triple follows as in Proposition 4.3. Therefore, by symmetry it is enough to analyze the last two triples. We need to compute h_3, h_5, h_6 .

We have $V_{D_6}^* = \Lambda^2 W$. Looking at the action of $\mathfrak{sl}(V)$ we find only one invariant in $V \otimes V^* \subset \Lambda^2 W$ corresponding to the identity in $\text{End}(V^*)$. Hence we get $h_6 = e_1 \wedge \varphi_1 + e_2 \wedge \varphi_2 + e_3 \wedge \varphi_3$.

The representation $V_{D_5}^*$ is contained in $\Lambda^2 W \otimes W^*$ and it is the kernel of κ_W . It contains two invariants under the action of $\mathfrak{sl}(V)$: $x \in \Lambda^2 V \otimes V \simeq \text{End}(V)$ and $y \in \Lambda^2 V^* \otimes V^* \simeq \text{End}(V^*)$. Notice that $\Lambda^2 W \otimes W^* \simeq V_{D_5}^* \oplus W$ so both $x, y \in V_{D_5}^*$. Moreover, the action of $S^2 V$ on x is clearly trivial while it is not on y . So we have $h_5 = e_{12} \otimes \varphi_3^* - e_{13} \otimes \varphi_2^* + e_{23} \otimes \varphi_1^*$.

Similarly, we have $V_{D_3}^* = \ker \kappa_{W^*} \subset \Lambda^2 W^* \otimes W$ and $h_3 = \varphi_{12}^* \otimes e_3 - \varphi_{13}^* \otimes e_2 + \varphi_{23}^* \otimes e_1$.

5.3.1. Analysis of the triple (D_5, D_6, D_2) . We have $V_{D_2}^* = \Lambda^3 W$. Consider the map $\Phi: \Lambda^2 W \otimes \Lambda^2 W \otimes W^* \longrightarrow \Lambda^3 W$ given by

$$\Phi(x \otimes y \otimes \varphi) = \kappa_W(x \otimes \varphi) \wedge y$$

for $x, y \in \Lambda^2 W$ and $\varphi \in W^*$. A direct computation shows that $\Phi(h_6 \otimes h_5) = 3e_{123} \neq 0$.

5.3.2. Analysis of the triple (D_3, D_6, D_5) . Consider the map $\Phi: \Lambda^2 W \otimes \Lambda^2 W^* \otimes W \longrightarrow \Lambda^2 W \otimes W^*$ given by

$$\Phi((x \wedge y) \otimes (\varphi \wedge \psi) \otimes w) = (\kappa_W(x \wedge y \otimes \varphi) \wedge w) \otimes \psi - (\kappa_W(x \wedge y \otimes \psi) \wedge w) \otimes \varphi$$

for $x, y, w \in W$ and $\varphi, \psi \in W^*$. A direct computation shows that $\Phi(h_6 \otimes h_3) = 2h_5$.

5.4. **Cotype E₇.** Let $V = \mathbb{C}^3$ and set $W = V \oplus V^* \oplus \mathbb{C}e$, $\mathfrak{h}_0 = \mathfrak{sl}(V) \oplus \mathbb{S}^2V \oplus \Lambda^2V \oplus V$ and $\mathfrak{g} = \mathfrak{sl}(W)$. Recall that we identify $\mathbb{S}^2V \oplus \Lambda^2V$ with the decomposition of $\text{Hom}(V^*, V)$ into symmetric and antisymmetric matrices. We have an action of \mathfrak{h}_0 on W given as follows. Let $(a, b, \omega, u) \in \mathfrak{h}_0$ and $(v, \varphi, \lambda e) \in W$, then

$$\begin{array}{llll} a \cdot v = a(v) & b \cdot v = 0 & \omega \cdot v = 0 & u \cdot v = 0 \\ a \cdot \varphi = -a^t(\varphi) & b \cdot \varphi = b(\varphi) & \omega \cdot \varphi = \omega(\varphi) & u \cdot \varphi = 0 \\ a \cdot e = 0 & b \cdot e = 0 & \omega \cdot e = \gamma(\omega) & u \cdot e = u \end{array}$$

where $\gamma = \gamma_2: \Lambda^2V \rightarrow V^*$ is defined as above. This defines an immersion of \mathfrak{h}_0 into $\mathfrak{sl}(W)$ whose image is closed under the Lie bracket. Indeed, if $(a, b, \omega, u) \in \mathfrak{h}_0$ and $(a', b', \omega', u') \in \mathfrak{h}_0$, then

$$\begin{array}{llllll} [a, a'] = aa' - a'a & [a, b] = ab + ba^t & [a, \omega] = a\omega + \omega a^t & [a, u] = a(u) \\ [b, b'] = 0 & [b, \omega] = b(\gamma(\omega)) & [b, u] = 0 & [\omega, \omega'] = 0 & [\omega, u] = 0 & [u, u'] = 0. \end{array}$$

It is also useful to write the action on W^* which is given as follows. We have $W^* = V^* \oplus V \oplus \mathbb{C}e^*$, then

$$\begin{array}{llll} a \cdot \varphi = -a^t(\varphi) & b \cdot \varphi = -b(\varphi) & \omega \cdot \varphi = \omega(\varphi) & u \cdot \varphi = -\varphi(u)e^* \\ a \cdot v = a(v) & b \cdot v = 0 & \omega \cdot v = -\langle \omega, v \rangle e^* & u \cdot v = 0 \\ a \cdot e^* = 0 & b \cdot e^* = 0 & \omega \cdot e^* = \gamma(\omega) & u \cdot e^* = u \end{array}$$

The triples of Lemma 4.9 are (D_1, D_6, D_3) and $(D_6, D_6, D_2 + D_7)$.

5.4.1. *Computation of h_1 .* The representation associated to D_1 is Λ^4W . Looking at the action of $\mathfrak{sl}(V)$ we get three invariants: the vector $x \in \Lambda^2V \otimes \Lambda^2V^*$ corresponding to the identity in $\text{End}(\Lambda^2V)$, the vector $y = e_{123} \wedge e \in \Lambda^3V \otimes \mathbb{C}e$ and the vector $z = \varphi_{123} \wedge e \in \Lambda^3V^* \otimes \mathbb{C}e$. Hence the invariant is a linear combination of these vectors. A small computation shows that

$$h_1 = 2y - x = 2e_{123} \wedge e - e_{12} \wedge \varphi_{12} - e_{13} \wedge \varphi_{13} - e_{23} \wedge \varphi_{23}.$$

5.4.2. *Computation of h_6 .* The representation associated to D_6 is the kernel of the map κ_W in $\Lambda^2W \otimes W^*$. In $\Lambda^2W \otimes W^*$ there are five invariant vectors under $\mathfrak{sl}(V)$: $x_1 \in \Lambda^2V \otimes V$, $x_2 \in \Lambda^2V^* \otimes V^*$, $y_1 \in (V \wedge \mathbb{C}e) \otimes V^*$, $y_2 \in (V^* \wedge \mathbb{C}e) \otimes V$ and $z \in (V \wedge V^*) \otimes \mathbb{C}e^*$. A small computation shows that

$$\begin{aligned} h_6 = 2x_1 + z = 2e_{12} \otimes \varphi_3^* - 2e_{13} \otimes \varphi_2^* + 2e_{23} \otimes \varphi_1^* + (e_1 \wedge \varphi_1) \otimes e^* \\ + (e_2 \wedge \varphi_2) \otimes e^* + (e_3 \wedge \varphi_3) \otimes e^*. \end{aligned}$$

5.4.3. *Analysis of the triple (D_1, D_6, D_3) .* The representation associated to the color D_3 is the kernel of the wedge product in $W \otimes \Lambda^4W \rightarrow \Lambda^5W$. We consider the map $\Phi: \Lambda^4W \otimes \Lambda^2W \otimes W^* \rightarrow W \otimes \Lambda^4W$ given by

$$\Phi(u \otimes (v_1 \wedge v_2) \otimes \varphi) = v_1 \otimes (\kappa_W^{41}(u \otimes \varphi) \wedge v_2) - v_2 \otimes (\kappa_W^{41}(u \otimes \varphi) \wedge v_1).$$

A direct computation shows that

$$\Phi(h_1 \otimes h_6) = -4(e_1 \otimes (e_{123} \wedge \varphi_1) + e_2 \otimes (e_{123} \wedge \varphi_2) + e_3 \otimes (e_{123} \wedge \varphi_3)) \neq 0.$$

5.4.4. *Analysis of the triple* $(D_6, D_6, D_2 + D_7)$. The representation $V_{D_2+D_7}^*$ associated to the color $D_2 + D_7$ is the kernel of the map $\kappa_W^{31}: \Lambda^3 W \otimes W^* \longrightarrow \Lambda^2 W$. We consider the map $\Psi: (\Lambda^2 W \otimes W^*) \otimes (\Lambda^2 W \otimes W^*) \longrightarrow \Lambda^3 W \otimes W^*$ given by

$$\Psi((u \otimes \varphi) \otimes (v \otimes \psi)) = (u \wedge \kappa_W(v \otimes \varphi)) \otimes \psi.$$

A direct computation shows that

$$\Psi(h_6 \otimes h_6) = -6 e_{123} \otimes e^* \in V_{D_2+D_7}^*.$$

5.5. **Cotype E_8 .** Let $V = \mathbb{C}^4$ and set $W = V \oplus \Lambda^2 V \oplus V^* = X \oplus Y \oplus Z$. On W it is defined a non-degenerate symmetric bilinear form such that $V \oplus V^*$ is orthogonal to $\Lambda^2 V$ and such that restricted to $V \oplus V^*$ and to $\Lambda^2 V$ is the one introduced at the beginning of this section.

Let A be the kernel of the wedge product $V \otimes \Lambda^2 V \longrightarrow \Lambda^3 V$ and set $\mathfrak{h}_0 = \mathfrak{sl}(V) \oplus A \oplus \Lambda^2 V$. There is an action of \mathfrak{h}_0 on W given as follows. Let $(a, b, \alpha) \in \mathfrak{h}_0$ and $(v, \omega, \varphi) \in W$ then

$$\begin{array}{lll} a \cdot v = a(v) & b \cdot v = 0 & \alpha \cdot v = 0 \\ a \cdot \omega = a(\omega) & b \cdot \omega = \gamma(b \otimes \omega) & \alpha \cdot \omega = 0 \\ a \cdot \varphi = -a^t(\varphi) & b \cdot \varphi = \delta(b \otimes \varphi) & \alpha \cdot \varphi = \kappa_V(\alpha \otimes \varphi) \end{array}$$

where γ, δ are defined as follows:

$$\begin{array}{ll} \gamma: V \otimes \Lambda^2 V \otimes \Lambda^2 V \longrightarrow V & \gamma(v \otimes \alpha \otimes \alpha') = (\alpha, \alpha') v \\ \delta: V \otimes \Lambda^2 V \otimes V^* \longrightarrow \Lambda^2 V & \delta(v \otimes \alpha \otimes \varphi) = \varphi(v) \alpha \end{array}$$

The action of \mathfrak{h}_0 on W defines an immersion of \mathfrak{h}_0 into $\mathfrak{so}(W)$, whose image is closed under the Lie bracket and more explicitly for $a, a' \in \mathfrak{sl}(V)$, $b, b' \in A$ and $\alpha, \alpha' \in \Lambda^2 V$ we have

$$\begin{array}{lll} [a, a'] = aa' - a'a & [a, b] = a(b) & [a, \alpha] = a(\alpha) \\ [b, b'] = -\zeta(b \otimes b') & [b, \alpha] = 0 & [\alpha, \alpha'] = 0 \end{array}$$

where ζ is the map

$$\zeta: (V \otimes \Lambda^2 V) \otimes (V \otimes \Lambda^2 V) \longrightarrow \Lambda^2 V \quad \zeta((u \otimes \alpha) \otimes (u' \otimes \alpha')) = (\alpha, \alpha') u \wedge u'.$$

The triples of Lemma 4.9 follow: (D_1, D_1, D_2) , $(D_1, D_5, 2D_2)$, (D_1, D_7, D_3) , (D_1, D_8, D_7) , (D_3, D_8, D_5) , $(D_5, D_8, D_2 + D_7)$ and (D_7, D_8, D_2) .

5.5.1. *Computation of h_1 .* The representation associated to D_1 is $\Lambda^3 W$. Looking at the action of $\mathfrak{sl}(V)$ we get two invariants: $x \in \Lambda^2 X \otimes Y \simeq \Lambda^2 V \otimes \Lambda^2 V$ and $y \in Y \otimes \Lambda^2 Z \simeq \Lambda^2 V \otimes \Lambda^2 V^*$. We notice now that X is invariant also by the action of A and $\Lambda^2 V$. Indeed, if $b \in A$, then $b \cdot x \in \Lambda^3 X \simeq \Lambda^3 V$; in particular, we get a $\mathfrak{sl}(V)$ -equivariant map from A to $\Lambda^3 V$, which must be zero. A similar argument proves that $\alpha \cdot x = 0$ for $\alpha \in \Lambda^2 V$. Finally, it is easy to check that if $b = e_1 \wedge e_{12}$, which belongs to A , then $b \cdot y \neq 0$. Hence

$$h_1 = x = e_1 \wedge e_2 \wedge e_{34} - e_1 \wedge e_3 \wedge e_{24} + e_1 \wedge e_4 \wedge e_{23} + e_2 \wedge e_3 \wedge e_{14} - e_2 \wedge e_4 \wedge e_{13} + e_3 \wedge e_4 \wedge e_{12}.$$

5.5.2. *Computation of h_3, h_5, h_7 .* Let P be the parabolic of $\text{Spin}(W)$ defined by $g(V) \subset V$. Notice that $H \subset P$. Let U be the unipotent radical of P , L its Levi subgroup, and L^{ss} its semisimple part. Notice that $L^{\text{ss}} \simeq \text{SL}(4) \times \text{SL}(4)$. Let $T \subset G$ be the maximal torus of elements acting diagonally on W with respect to the basis $e_1, \dots, e_4, e_{12}, e_{13}, e_{14}, e_{23}, -e_{24}, e_{34}, \varphi_4, \dots, \varphi_1$ and $B \subset G$ the subgroup of elements whose action on W is upper triangular with respect to this basis. The natural action of $\text{SL}(V)$ on W induces an embedding $\text{SL}(V) \rightarrow L$, that on diagonal elements takes the form

$$(t_1, t_2, t_3, t_4) \mapsto (t_1, t_2, t_3, t_4, t_1 t_2, t_1 t_3, t_1 t_4, t_2 t_3, t_2 t_4, t_3 t_4, t_4^{-1}, t_3^{-1}, t_2^{-1}, t_1^{-1}).$$

The set of colors $\{D_1, D_4, D_6, D_8\}$ is distinguished, and $K_0 = H_0 U$ is the intersection of the kernels of the multiplicative characters of the stabilizer K of a point in the open orbit of the quotient. Therefore, h_3, h_5, h_7 must be K_0 -invariant vectors in $V_{D_i}^*$. Let $W_i = (V_{D_i}^*)^U$. This is an irreducible representation of L of the same highest weight as $V_{D_i}^* : \omega_3 + \omega_7$ for $i = 3$, $\omega_2 + \omega_5$ for $i = 5$, and $\omega_1 + \omega_6$ for $i = 7$. When we restrict these representations to $\text{SL}(V)$ we get

$$W_3 \simeq \Lambda^3 V \otimes V \quad W_5 \simeq \Lambda^2 V \otimes \Lambda^2 V \quad W_7 \simeq V \otimes \Lambda^3 V;$$

in particular, there is an invariant element under \mathfrak{h}_0 .

For notational convenience, here and below, set

$$e_5 = e_{12}, e_6 = e_{13}, e_7 = e_{14}, \varphi_7 = e_{23}, \varphi_6 = -e_{24}, \varphi_5 = e_{34}.$$

Let $U \subset W$ be the subspace spanned by e_1, \dots, e_7 , so that W becomes $U \oplus U^*$.

Now we need to describe the spin representations. Consider the whole exterior algebra ΛU^* . It decomposes into odd and even degree parts $\Lambda^{\text{odd}} U^* \oplus \Lambda^{\text{even}} U^*$. Since the G -action we are going to define is not the natural one, we stress the difference by using a different notation: set $\psi_{i_1 \dots i_k} = \varphi_{i_1 \dots i_k}$. Define the map $\sigma : W \otimes \Lambda U^* \rightarrow \Lambda U^*$ such that

$$\sigma(e_i \otimes \psi_{i_1 \dots i_k}) = \kappa_{U^*}^{k1}(\varphi_{i_1 \dots i_k} \otimes e_i),$$

$$\sigma(\varphi_i \otimes \psi_{i_1 \dots i_k}) = \varphi_i \wedge \varphi_{i_1 \dots i_k},$$

and more generally the map $\sigma^n : \otimes^n W \otimes \Lambda U^* \rightarrow \Lambda U^*$ such that

$$\sigma^n(w_1 \otimes \dots \otimes w_n \otimes y) = \sigma(w_1 \otimes \sigma(\dots \otimes \sigma(w_n \otimes y) \dots))$$

which we can restrict to $\Lambda^n W$ if we think of $w_1 \wedge \dots \wedge w_n$ as the corresponding antisymmetric tensor, with coefficient $\frac{1}{n!}$.

To get the spin representations we can just take the map σ^2 ; indeed, notice that $\Lambda^2 W$ identifies with $\mathfrak{so}(W)$ through $w_1 \wedge w_2 = (w_2,)w_1 - (w_1,)w_2$. We thus have that the vector $\psi_{i_1 \dots i_k}$ has weight

$$\frac{1}{2} \left(\sum_{i \notin \{i_1, \dots, i_k\}} \varepsilon_i - \sum_{i \in \{i_1, \dots, i_k\}} \varepsilon_i \right),$$

$V(\omega_6) \simeq \Lambda^{\text{odd}} U^*$ and $V(\omega_7) \simeq \Lambda^{\text{even}} U^*$.

We get the following expressions of the H_0 -invariants:

$$\begin{aligned} h_3 &= e_{123} \otimes \psi_{65} + e_{124} \otimes \psi_{75} + e_{134} \otimes \psi_{76} - e_{234} \otimes \psi_{\emptyset}, \\ h_5 &= e_{12} \otimes e_{1234} \wedge \varphi_5 + e_{13} \otimes e_{1234} \wedge \varphi_6 + e_{14} \otimes e_{1234} \wedge \varphi_7 + e_{23} \otimes e_{12347} \\ &\quad - e_{24} \otimes e_{12346} + e_{34} \otimes e_{12345}, \\ h_7 &= e_1 \otimes \psi_{765} - e_2 \otimes \psi_5 - e_3 \otimes \psi_6 - e_4 \otimes \psi_7. \end{aligned}$$

5.5.3. *Computation of h_8 .* The representation $V_{D_8}^*$ is the spin representation of highest weight ω_6 . By direct computation one can show that the only H_0 -invariant is given by

$$h_8 = \psi_1 + \psi_{762} - \psi_{753} + \psi_{654}.$$

5.5.4. *Analysis of the triple (D_1, D_1, D_2) .* The representation associated with D_2 is $\Lambda^4 W$. The H_0 -invariant h_2 is e_{1234} . Indeed, the set of colors $\Delta \setminus \{D_2\}$ is distinguished and the quotient is homogeneous, hence an H -semi-invariant in $V_{D_2}^*$ must be P -semiinvariant.

Here we get

$$\tilde{\kappa}_W^{33}(h_1 \otimes h_1) = 3h_2.$$

5.5.5. *Analysis of the triple $(D_1, D_5, 2D_2)$.* Consider the map $\Phi: \Lambda^3 W \otimes \Lambda^2 W \otimes \Lambda^5 W \rightarrow \Lambda^4 W \otimes \Lambda^4 W$ such that

$$\begin{aligned} \Phi((w_1 \wedge w_2 \wedge w_3) \otimes x \otimes y) &= (w_1 \wedge w_2 \wedge x) \otimes \kappa_W^{51}(y \otimes w_3) \\ &\quad - (w_1 \wedge w_3 \wedge x) \otimes \kappa_W^{51}(y \otimes w_2) + (w_2 \wedge w_3 \wedge x) \otimes \kappa_W^{51}(y \otimes w_1). \end{aligned}$$

We have

$$\Phi(h_1 \otimes h_5) = 6h_2 \otimes h_2.$$

5.5.6. *Analysis of the triple (D_1, D_7, D_3) .* Consider the map

$$\Phi: \Lambda^3 W \otimes W \otimes \Lambda^{\text{odd}} U^* \rightarrow \Lambda^3 W \otimes \Lambda^{\text{even}} U^*$$

such that

$$\begin{aligned} \Phi((w_1 \wedge w_2 \wedge w_3) \otimes w \otimes \psi) &= (w_1 \wedge w_2 \wedge w) \otimes \sigma(w_3 \otimes \psi) \\ &\quad - (w_1 \wedge w_3 \wedge w) \otimes \sigma(w_2 \otimes \psi) + (w_2 \wedge w_3 \wedge w) \otimes \sigma(w_1 \otimes \psi). \end{aligned}$$

We get

$$\Phi(h_1 \otimes h_7) = 3h_3.$$

5.5.7. *Analysis of the triple (D_1, D_8, D_7) .* Consider the map $\Phi: \Lambda^3 W \otimes \Lambda^{\text{odd}} U^* \rightarrow W \otimes \Lambda^{\text{odd}} U^*$ such that

$$\begin{aligned} \Phi((w_1 \wedge w_2 \wedge w_3) \otimes \psi) &= w_1 \otimes \sigma^2((w_2 \wedge w_3) \otimes \psi) \\ &\quad - w_2 \otimes \sigma^2((w_1 \wedge w_3) \otimes \psi) + w_3 \otimes \sigma^2((w_1 \wedge w_2) \otimes \psi). \end{aligned}$$

We get

$$\Phi(h_1 \otimes h_8) = -3h_7.$$

5.5.8. *Analysis of the triple (D_3, D_8, D_5) .* On $\Lambda^6 W$ there is a symmetric bilinear form (\cdot, \cdot) naturally induced by the given form on W . On the other hand, $V(\omega_6)$ and $V(\omega_7)$ are dual to each other, so there is a natural non-degenerate pairing $\langle \cdot, \cdot \rangle$. Consider the map $\Psi: \Lambda^{\text{odd}} U^* \otimes \Lambda^{\text{even}} U^* \rightarrow \Lambda^6 W$ such that

$$(u, \Psi(\psi \otimes \psi')) = \langle \sigma^6(u)\psi, \psi' \rangle,$$

and the map $\Phi: \Lambda^3 W \otimes \Lambda^{\text{odd}} U^* \otimes \Lambda^{\text{even}} U^* \rightarrow \Lambda^2 W \otimes \Lambda^5 W$ such that

$$\begin{aligned} \Phi((w_1 \wedge w_2 \wedge w_3) \otimes \psi \otimes \psi') \\ = (w_1 \wedge w_2) \otimes \tilde{\kappa}_W^{2,1}(\Psi(\psi \otimes \psi') \otimes w_3) \\ - (w_1 \wedge w_3) \otimes \tilde{\kappa}_W^{2,1}(\Psi(\psi \otimes \psi') \otimes w_2) \\ + (w_2 \wedge w_3) \otimes \tilde{\kappa}_W^{2,1}(\Psi(\psi \otimes \psi') \otimes w_1). \end{aligned}$$

We get

$$\Phi(h_3 \otimes h_8) = h_5.$$

5.5.9. *Analysis of the triple $(D_5, D_8, D_2 + D_7)$.* Consider the map $\Phi: \Lambda^2 W \otimes \Lambda^5 W \otimes \Lambda^{\text{odd}} U^* \rightarrow \Lambda^4 W \otimes W \otimes \Lambda^{\text{odd}} U^*$ such that

$$\begin{aligned} \Phi((w_1 \wedge w_2) \otimes (z_1 \wedge \cdots \wedge z_5) \otimes \psi) \\ = \sum_i (-1)^{i+1} (z_1 \wedge \cdots \wedge \hat{z}_i \wedge \cdots \wedge z_5) \otimes w_2 \otimes \sigma^2((z_i \wedge w_1) \otimes \psi) \\ - \sum_i (-1)^{i+1} (z_1 \wedge \cdots \wedge \hat{z}_i \wedge \cdots \wedge z_5) \otimes w_1 \otimes \sigma^2((z_i \wedge w_2) \otimes \psi). \end{aligned}$$

We get

$$\Phi(h_5 \otimes h_8) = -3h_2 \otimes h_7.$$

5.5.10. *Analysis of the triple (D_7, D_8, D_2) .* Consider the map $\Phi: W \otimes \Lambda^{\text{odd}} U^* \otimes \Lambda^{\text{odd}} U^* \rightarrow \Lambda^{\text{odd}} U^* \otimes \Lambda^{\text{even}} U^*$ such that

$$\Phi(w \otimes \psi \otimes \psi') = \sigma(w \otimes \psi') \otimes \psi.$$

Here we get that $\Phi(h_7 \otimes h_8)$ is a U -invariant vector of weight ω_4 .

5.6. Projective normality of comodel wonderful varieties.

Proof of Theorem 5.2. As in the case of the model wonderful varieties, by Lemma 2.3, we are reduced to study the low fundamental triples.

Recall that in the model case we have classified, for every G of connected Dynkin type, the low fundamental triples (D, E, F) of the model wonderful G -variety (of simply connected type) such that $\text{supp}_S(D + E - F) = S$.

In the comodel case, these correspond to the low fundamental triples (D, E, F) of the comodel wonderful varieties of connected Dynkin cotype such that $\text{cosupp}(D + E - F) = \Delta$, where

$$\text{cosupp } \gamma = \{D \in \Delta : \langle \alpha^\vee, \omega_D \rangle \neq 0 \text{ for some } \alpha \in \text{supp}_S \gamma\}.$$

Let us go back to the model case. Let S be the set of simple roots. Let M be a model wonderful variety with set of colors Δ . Recall that for every low fundamental triple (D, E, F) , $\text{supp}_S(D + E - F) = S'$ is connected. Hence (D, E, F) corresponds to a low fundamental triple (D, E, \tilde{F}) , of a model wonderful variety N (with set of colors $\tilde{\Delta} \subset \Delta$) of connected Dynkin type (with S' as set of simple roots) such

that M is parabolic induction of N , with $D, E \in \tilde{\Delta}$, $D + E - F = D + E - \tilde{F}$ and $\text{supp}_{S'}(D + E - \tilde{F}) = S'$.

This can be translated into the comodel case. Every low fundamental triple (D, E, F) of a comodel wonderful variety M (with set of colors Δ) corresponds to a low fundamental triple (D, E, \tilde{F}) of a comodel wonderful variety N of connected Dynkin cotype (with set of colors $\tilde{\Delta} \subset \Delta$) such that M is parabolic induction of N , with $D, E \in \tilde{\Delta}$, $D + E - F = D + E - \tilde{F}$ and $\text{cosupp}(D + E - \tilde{F}) = \tilde{\Delta}$.

Since we have already checked the surjectivity for all such low fundamental triples, we can conclude by applying Proposition 1.6 as in the proof of Theorem 4.1 (see Section 4.8). \square

6. ON THE NORMALITY OF SPHERICAL ORBIT CLOSURES IN SIMPLE PROJECTIVE SPACES

Let M be a wonderful G -variety with set of spherical roots Σ and set of colors Δ . Denote by H the stabilizer of a point x_0 in the open G -orbit.

If D is in $\mathbb{N}\Delta$ and $h_D \in (V_D^*)_{\xi_D}^{(H)}$ is the associated H -eigenvector, consider the orbit closure

$$X_D = \overline{G \cdot [h_D]} \subset \mathbb{P}(V_D^*),$$

which is a simple possibly non-normal spherical variety (recall that a spherical variety is called *simple* if it contains a unique closed orbit). We have a natural morphism $\phi_D: M \rightarrow X_D$ such that $\phi_D^* \mathcal{O}(1) = \mathcal{L}_D$.

By [29, Corollary 7.6] and [8, Corollary 2.4.2.2], every spherical orbit in a simple projective space always admits a wonderful compactification M , so it is of the shape $G \cdot [h_D]$ for some $D \in \mathbb{N}\Delta$. As a consequence of the results of the previous sections, here we show that under some special assumptions on M the variety X_D is always normal.

The variety X_D was studied by G. Pezzini in [39] when D is ample, that is, $D \in \mathbb{N}_{>0}\Delta$. Under this assumption, either X_D is isomorphic to M or it is not even normal. In case $X_D \simeq M$, then M is called *strict*; this is equivalent to the conditions $H = N_G(H)$ and $\Sigma \cap S = \emptyset$. There are essentially two main classes of examples of strict wonderful varieties: the adjoint symmetric wonderful varieties and the model wonderful varieties.

When D is not ample, the variety X_D was then studied in [37] in the symmetric case and in [22] in general. More precisely, the orbit structure of X_D and that of its normalization \tilde{X}_D were analyzed. In particular, it was proved that the normalization $\tilde{X}_D \rightarrow X_D$ is always bijective if M is adjoint symmetric or if G is of simply laced type and M is strict, while the main counterexamples where bijectivity fails, in the strict case, arise with the model wonderful varieties for groups of not simply laced type.

We now consider a special class of big divisors on M . We say that $D \in \mathbb{N}\Delta$ is a *faithful divisor* on M if ϕ_D restricts to an open embedding of G/H in $\mathbb{P}(V_D^*)$, i.e., if H equals the stabilizer of $[h_D]$. The formalism of distinguished sets of colors allows us to characterize combinatorially the faithful divisors. For simplicity, we restrict ourselves to the case of a strict wonderful variety.

Proposition 6.1 (see [8, Proposition 2.4.3]). *Let M be a strict wonderful variety and let $D \in \mathbb{N}\Delta$. Then D is faithful if and only if every distinguished subset of Δ intersects $\text{supp}(D)$.*

Let $D \in \mathbb{N}\Delta$, denote $\tilde{A}(D) = \bigoplus_{n \in \mathbb{N}} \Gamma(M, \mathcal{L}_{nD})$ and denote $A(D) \subset \tilde{A}(D)$ the subalgebra generated by $V_D \subset \Gamma(M, \mathcal{L}_D)$. Let \tilde{X}_D be the image of M in $\mathbb{P}(\Gamma(M, \mathcal{L}_D)^*)$ via the morphism $\tilde{\phi}_D$ associated to the complete linear system of D . Then $\tilde{A}(D)$ is identified with the projective coordinate ring of \tilde{X}_D , whereas $A(D)$ is identified with the projective coordinate ring of X_D . Notice that we have a natural projection $\beta_D: \tilde{X}_D \rightarrow X_D$ such that $\phi_D = \tilde{\phi}_D \circ \beta_D$. Since M is smooth, $\tilde{A}(D)$ is an integrally closed algebra, therefore \tilde{X}_D is normal. Moreover, we have the following.

Proposition 6.2 (see [13, Proposition 2.1]). *The algebra $\tilde{A}(D)$ is integral over $A(D)$.*

It follows that $\beta_D: \tilde{X}_D \rightarrow X_D$ is the normalization if and only if it is birational. Clearly this is the case if D is minuscule or faithful. On the other hand, β_D is not necessarily birational: if M is the model wonderful variety of C_4 , then $A(D_2) \subset \tilde{A}(D_2)$ do not have the same quotient field. However, we have the following.

Proposition 6.3. *Suppose that M is adjoint symmetric or that G is simply laced and M is strict. Then $\tilde{A}(D)$ and $A(D)$ have the same quotient field.*

Proof. Let $\Delta_D \subset \Delta$ be the maximal distinguished subset which does not intersect $\text{supp}(D)$, and denote by M' the quotient of M by Δ_D and by M'' the wonderful compactification of $G \cdot [h_D]$. By [22, Corollary 3.7] and [22, Remark 3.8], M' and M'' have the same dimension, and we have $M' \neq M''$ if and only if some spherical root of M'' is the double of a spherical root of M' .

On the other hand, by the quotient construction, the spherical roots of M' are contained in $\mathbb{N}\Sigma$. By the description of the possible spherical roots occurring in a symmetric variety or in a strict variety for a simply laced G , it follows that $\mathbb{N}\Sigma$ contains no elements of $\Sigma(G)$ whose double is still in $\Sigma(G)$ (see [8, Table 1]). Therefore, M' and M'' must have the same spherical roots and we get $M' = M''$, and by Corollary 1.4 it follows that $\Gamma(M, \mathcal{L}_{nD}) \simeq \Gamma(M', \mathcal{L}_{nD})$ for every $n \geq 0$. Therefore, we are reduced to the case of a faithful divisor, and the claim follows. \square

In the adjoint symmetric case, the above proposition was proved in [13, Theorem 2.6].

If $D_1, \dots, D_m \in \mathbb{N}\Delta$, consider the variety

$$X_{D_1, \dots, D_m} = \overline{G \cdot ([h_{D_1}] \times \dots \times [h_{D_m}])} \subset \mathbb{P}(V_{D_1}^*) \times \dots \times \mathbb{P}(V_{D_m}^*)$$

and denote by $\phi_{D_1, \dots, D_m}: M \rightarrow X_{D_1, \dots, D_m}$ the map such that $\phi_{D_1, \dots, D_m}(x) = (\phi_{D_1}(x), \dots, \phi_{D_m}(x))$.

Proposition 6.4 ([7, Proposition 1.2]). *Let M be a wonderful variety and let $D, E \in \mathbb{N}\Delta$.*

- i) *If $\text{supp}(\omega_D) \cap \text{supp}(\omega_E) = \emptyset$, then $X_{D+E} \simeq X_{D,E}$.*
- ii) *If M is strict and $\text{supp}(D) = \{D_1, \dots, D_m\}$, then $X_D \simeq X_{D_1, \dots, D_m}$. In particular, if M is strict, $X_D \simeq X_E$ if and only if $\text{supp}(D) = \text{supp}(E)$.*

Proof. i) Since $\text{supp}(\omega_D) \cap \text{supp}(\omega_E) = \emptyset$, by [7, Lemma 1.1] we have a closed equivariant embedding $\psi_{D,E}: \mathbb{P}(V_D^*) \times \mathbb{P}(V_E^*) \rightarrow \mathbb{P}(V_{D+E}^*)$, and since $\psi_{D,E}([h_D], [h_E]) = [h_{D+E}]$ the isomorphism $X_{D+E} \simeq X_{D,E}$ follows.

ii) Since M is strict, by the description of the restriction $\omega: \text{Pic}(M) \rightarrow \mathcal{X}(T)$ (see [42, Lemma 30.24]) it follows that $\text{supp}(D) \cap \text{supp}(E) = \emptyset$ if and only if

$\text{supp}(\omega_D) \cap \text{supp}(\omega_E) = \emptyset$. Therefore, by [7, Lemma 1.1] we have a closed equivariant embedding $\psi_D: \mathbb{P}(V_{D_1}^*) \times \cdots \times \mathbb{P}(V_{D_m}^*) \rightarrow \mathbb{P}(V_D^*)$ such that $\psi_D([h_{D_1}], \dots, [h_{D_m}]) = [h_D]$ and it follows that $X_D \simeq X_{D_1, \dots, D_m}$. \square

Assume now that the multiplication of sections is surjective for every couple of globally generated line bundles. In particular $\tilde{A}(D)$ is generated by its degree one component $\Gamma(M, \mathcal{L}_D)$ and it follows that $A(D) = \tilde{A}(D)$ if and only if $\Gamma(M, \mathcal{L}_D) = V_D$ if and only if D is minuscule or $D = 0$. Since $\tilde{A}(D)$ is integrally closed, we get the following proposition which we need for later use.

Proposition 6.5. *Let M be a wonderful variety and suppose that the multiplication of sections is surjective for every couple of globally generated line bundles. Let $D \in \mathbb{N}\Delta$, $D \neq 0$.*

- i) *If D is minuscule, then X_D is projectively normal.*
- ii) *If β_D is birational and X_D is projectively normal, then D is minuscule.*

If $D_1, \dots, D_m \in \mathbb{N}\Delta$ and if V_1, \dots, V_m are G -modules of sections such that $V_{D_i} \subset V_i \subset \Gamma(M, \mathcal{L}_{D_i})$, consider the associated morphisms $\phi_{V_i}: M \rightarrow \mathbb{P}(V_i^*)$ and denote $X_{V_i} = \phi_{V_i}(M)$. Also, we denote by

$$X_{V_1, \dots, V_m} \subset \mathbb{P}(V_1^*) \times \cdots \times \mathbb{P}(V_m^*)$$

the image of M via the map $\phi_{V_1, \dots, V_m}(x) = (\phi_{V_1}(x), \dots, \phi_{V_m}(x))$ and by

$$X_{V_1 \otimes \cdots \otimes V_m} \subset \mathbb{P}(V_1^* \otimes \cdots \otimes V_m^*)$$

the image of X_{V_1, \dots, V_m} via the Segre embedding.

Lemma 6.6. *Let $D_1, \dots, D_m \in \mathbb{N}\Delta$ and denote $D = \sum_{i=1}^m D_i$. If V_1, \dots, V_m are G -modules such that $V_{D_i} \subset V_i \subset \Gamma(M, \mathcal{L}_{D_i})$, then the projective coordinate ring of $X_{V_1 \otimes \cdots \otimes V_m}$ is the subalgebra $A(V_1, \dots, V_m) \subset \tilde{A}(D)$ generated by the product $V_1 \cdots V_m \subset \Gamma(M, \mathcal{L}_D)$.*

Proof. Consider the map $\phi_{V_1 \otimes \cdots \otimes V_m}: M \rightarrow X_{V_1 \otimes \cdots \otimes V_m}$. The lemma follows by noticing that $\phi_{V_1 \otimes \cdots \otimes V_m}^* \mathcal{O}(1) = \mathcal{L}_D$ and that $\phi_{V_1 \otimes \cdots \otimes V_m}^*: V_1 \otimes \cdots \otimes V_m \rightarrow \Gamma(M, \mathcal{L}_D)$ is the multiplication map. \square

Proposition 6.7. *Let M be a wonderful variety and suppose that the multiplication of sections is surjective for every couple of globally generated line bundles. Let $D_1, \dots, D_m \in \mathbb{N}\Delta$ and denote $\Gamma_i = \Gamma(M, \mathcal{L}_{D_i})$. Then the variety $X_{\Gamma_1 \otimes \cdots \otimes \Gamma_m} \subset \mathbb{P}(\Gamma_1^* \otimes \cdots \otimes \Gamma_m^*)$ is projectively normal.*

Proof. Denote $D = \sum_{i=1}^m D_i$. Since the multiplication of sections is surjective, $\Gamma_1 \cdots \Gamma_m = \Gamma(M, \mathcal{L}_D)$, hence by the previous lemma $A(\Gamma_1, \dots, \Gamma_m) = \tilde{A}(D)$ and $X_{\Gamma_1 \otimes \cdots \otimes \Gamma_m}$ is a projectively normal variety. \square

Corollary 6.8. *Let M be a wonderful variety and suppose that the multiplication of sections is surjective for every couple of globally generated line bundles.*

- i) *Let $D, E \in \mathbb{N}\Delta$ be such that $\text{supp}(\omega_D) \cap \text{supp}(\omega_E) = \emptyset$. If X_D, X_E are normal, then X_{D+E} is normal as well.*
- ii) *If M is strict, then X_D is normal for all $D \in \mathbb{N}\Delta$ if and only if it is normal for all $D \in \Delta$.*

Proof. i) Denote $\Gamma_D = \Gamma(M, \mathcal{L}_D)$ and $\Gamma_E = \Gamma(M, \mathcal{L}_E)$. By the previous proposition, we have that $X_{\Gamma_D, \Gamma_E} \simeq X_{\Gamma_D \otimes \Gamma_E}$ is a normal variety. On the other hand, since X_D and X_E are normal, we have that $X_{D, E} \simeq X_{\Gamma_D, \Gamma_E}$, while by Proposition 6.4.i we have that $X_{D+E} \simeq X_{D, E}$.

ii) Since M is strict, it follows by the description of $\omega: \text{Pic}(M) \rightarrow \mathcal{X}(T)$ that $\text{supp}(\omega_D) \cap \text{supp}(\omega_E) = \emptyset$ if and only if $\text{supp}(D) \cap \text{supp}(E) = \emptyset$. Therefore the claim follows straightforwardly from i). \square

Corollary 6.9. *Suppose that M is a symmetric variety with reduced root system of type A or that it is a model wonderful variety for a connected semisimple group of type AD. Then X_D is normal for all $D \in \mathbb{N}\Delta$.*

Proof. By the description of the covering relation, it follows that under the assumptions on M it holds $\text{ht}(\gamma^+) = 2$ for every covering difference γ in $\mathbb{N}\Delta$. Therefore every $D \in \Delta$ is minuscule and $\Gamma(M, \mathcal{L}_D) = V_D$. Therefore, X_D is projectively normal for all $D \in \Delta$ and it follows by the previous corollary that X_D is normal for all $D \in \mathbb{N}\Delta$. \square

7. ON THE NORMALITY OF CONES AND NILPOTENT ORBIT CLOSURES

Following the same approach as in [19] and [13], we can apply Theorem 4.1 to study the normality of cones over model varieties. In particular, as pointed out by Luna some years ago, we can apply our theory to study the normality of the closure of spherical nilpotent orbits in the Lie algebra of G .

Let M be a wonderful variety with set of colors Δ and set of spherical roots Σ and assume that the multiplication of sections is surjective for every couple of globally generated line bundles. Let $D \in \mathbb{N}\Delta$ and denote by $C_D \subset V_D^*$ the cone over the variety X_D introduced in the previous section. Analogously, denote by $\tilde{C}_D \subset \Gamma(M, \mathcal{L}_D)^*$ the cone over the variety \tilde{X}_D . Then the coordinate ring of C_D is identified with $A(D)$, whereas that of \tilde{C}_D is identified with $\tilde{A}(D)$, which is an integrally closed ring. This yields a map $\alpha_D: \tilde{C}_D \rightarrow C_D$ that is birational if and only if $\beta_D: \tilde{X}_D \rightarrow X_D$ is birational. As already recalled in the previous section, $\tilde{A}(D)$ is the integral closure of $A(D)$ if and only if α_D is birational, whereas $A(D) = \tilde{A}(D)$ if and only if D is minuscule or $D = 0$.

In the case of the model wonderful varieties of simply connected type, after the results of Section 3, we have the following classification of minuscule weights, where $a, b, c \in \mathbb{N}$.

Case A_r , r even: $D_1, D_2, \dots, D_r, aD_1, aD_r, aD_1 + D_d$ with d odd, $D_m + bD_r$ with m even;

Case A_r , r odd: $D_1, D_2, \dots, D_r, aD_1, aD_r, aD_1 + D_d$ with d odd, $D_m + bD_r$ with m odd, $aD_1 + bD_r$;

Case B_r , r even: $aD_r, D_m + aD_r$ with m even;

Case B_r , r odd: $aD_1 + bD_r, aD_1 + D_m + bD_r$ with m odd;

Case C_r : aD_1 ;

Case D_r , r even: $D_1, D_2, \dots, D_r, aD_1 + bD_{r-1} + cD_r, aD_1 + D_m + bD_{r-1} + cD_r$ with m odd;

Case D_r , r odd: $D_1, D_2, \dots, D_r, aD_{r-1} + bD_r, D_m + aD_{r-1} + bD_r$ with m even;

Case E_6 : $D_1, D_6, aD_2, aD_2 + D_3, aD_2 + D_5$;

Case E_7 : $D_1, D_6, aD_2 + bD_7, aD_2 + D_3 + bD_7, aD_2 + D_5 + bD_7$;

Case E_8 : $D_1, D_8, aD_2, aD_2 + D_3, aD_2 + D_5, aD_2 + D_7$;

Case F_4 : aD_1 ;

Case G_2 : aD_1 .

Let \mathfrak{g} be the Lie algebra of G , as a general reference about nilpotent orbits in \mathfrak{g} ; see [17]. If \mathfrak{g} is not simple, then its nilpotent orbits are products of the nilpotent orbits of its simple factors, and so are their closures. Therefore we may assume that \mathfrak{g} is simple.

Let $e \in \mathfrak{g}$ be a non-zero nilpotent element and let \mathcal{O} be its adjoint orbit. By the Jacobson-Morozov theorem there exists an $\mathfrak{sl}(2)$ -triple of the form (e, h, f) . Choose a maximal toral subalgebra \mathfrak{t} of \mathfrak{g} containing h and a Borel subalgebra \mathfrak{b} containing \mathfrak{t} and e and such that $\alpha(h) \geq 0$ for every $\alpha \in S$, where we denote by $S = \{\alpha_1, \dots, \alpha_r\}$ the set of simple roots defined by the choice of \mathfrak{t} and \mathfrak{b} . The string $(\alpha_1(h), \dots, \alpha_r(h))$ is called the Kostant–Dynkin diagram of \mathcal{O} and it uniquely determines the orbit \mathcal{O} . Moreover, every $\alpha_i(h)$ is 0, 1 or 2. Let θ be the highest root corresponding to the choice of S and define the *height* of \mathcal{O} as $\text{height}(\mathcal{O}) = \theta(h)$. The height does not depend on the various choices we have made (see [17, § 3.5]); furthermore, \mathcal{O} is spherical if and only if $\text{height}(\mathcal{O}) \leq 3$, and this last condition is equivalent to saying that it has height equal to 2 or to 3; see [38].

By making use of the projective normality of the symmetric wonderful varieties, in [13] it has been proved that the closure $\overline{\mathcal{O}}$ is normal if $\text{height}(\mathcal{O}) = 2$, which is originally due to W. Hesselink [26]. We now study the normality of $\overline{\mathcal{O}}$ in the case of $\text{height}(\mathcal{O}) = 3$ (see [38, Table 2]) by making use of the projective normality of the model wonderful varieties.

Denote by $\mathcal{U} \simeq G/H$ the orbit of the line $[e] \in \mathbb{P}(\mathfrak{g}) = \mathbb{P}(V(\theta))$, namely the image of \mathcal{O} via the natural projection. As every spherical orbit in the projective space of a simple G -module, \mathcal{U} possesses a wonderful compactification, which we denote by $M_{\mathcal{O}}$. In [6] we can find a description of the stabilizer of $[e]$ as well as the associated Luna diagram. In particular, $M_{\mathcal{O}}$ turns out to be a strict wonderful variety and, in particular, the restriction of line bundles to the closed orbit $\text{Pic}(M_{\mathcal{O}}) \rightarrow \mathcal{X}(B)$ is always injective. Therefore, we may regard θ as an element of $\mathbb{N}\Delta$ and we have $\overline{\mathcal{U}} = X_{\theta}$ and $\overline{\mathcal{O}} = C_{\theta}$.

In order to study the normality of $\overline{\mathcal{O}}$, we prove the following.

Theorem 7.1. *Let $\mathcal{O} \subset \mathfrak{g}$ be a spherical nilpotent orbit and let $M_{\mathcal{O}}$ be the associated wonderful variety. Then the multiplication map*

$$m_{\mathcal{L}, \mathcal{L}'} : \Gamma(M_{\mathcal{O}}, \mathcal{L}) \otimes \Gamma(M_{\mathcal{O}}, \mathcal{L}') \longrightarrow \Gamma(M_{\mathcal{O}}, \mathcal{L} \otimes \mathcal{L}')$$

is surjective for all globally generated line bundles $\mathcal{L}, \mathcal{L}'$ on $M_{\mathcal{O}}$.

Notice that by construction θ is identified with a faithful divisor on $M_{\mathcal{O}}$. Therefore the normality (resp. the non-normality) of $\overline{\mathcal{O}} = C_{\theta}$ follows by noticing case-by-case that θ is minuscule (resp. non-minuscule) in $\mathbb{N}\Delta$, and applying Proposition 6.5. Given a nilpotent orbit \mathcal{O} of height 3, we summarize the results on the normality of its closure in Table 1. Given a spherical nilpotent orbit \mathcal{O} , there we write in the second column its Kostant–Dynkin diagram, in the third column its Bala–Carter label (see [15, §8.4]), in the fourth column the corresponding case in [6] and in the fifth column the normality of $\overline{\mathcal{O}}$. In particular we have the following.

Corollary 7.2 (see [18, Theorem 5.1] or [38, Table 2]). *Let $\mathcal{O} \subset \mathfrak{g}$ be a spherical nilpotent orbit with $\text{height}(\mathcal{O}) = 3$. Then the closure $\overline{\mathcal{O}}$ is always normal except in cases I and XI of Table 1.*

TABLE 1. Nilpotent orbits of height 3

	type of G	Kostant-Dynkin diagram	partition/ Bala-Carter label	case in [6]	norm.
I	B_{2n+1}	(10...01)	$(3, 2^{2n})$	(13)	no
II	B_{2n+m+1} ($m > 0$)	(10...01 $\underbrace{0\dots 0}_m$)	$(3, 2^{2n}, 1^{2m})$	(18)	yes
III	D_{2n+2}	(10...011)	$(3, 2^{2n}, 1)$	(41)	yes
IV	D_{2n+m+2} ($m > 0$)	(10...01 $\underbrace{0\dots 0}_{m+1}$)	$(3, 2^{2n}, 1^{2m+1})$	(43)	yes
V	E_6	(000100)	A_1	(53)	yes
VI	E_7	(0010000)	$(3A_1)'$	(54)	yes
VII	E_7	(0100001)	$4A_1$	(51)	yes
VIII	E_8	(00000010)	$3A_1$	(52)	yes
IX	E_8	(01000000)	$4A_1$	(51)	yes
X	F_4	(0100)	$A_1 + \tilde{A}_1$	(60)	yes
XI	G_2	(10)	\tilde{A}_1	(66)	no

The rest of the section is devoted to the proof of Theorem 7.1 and Corollary 7.2. Since the case $\text{height}(\mathcal{O}) = 2$ was already considered in [13], we consider only the case $\text{height}(\mathcal{O}) = 3$.

7.1. Cases I, III, VII, IX, XI: model orbits. By the description in [6], in these cases we have that $M_{\mathcal{O}}$ is the model wonderful variety of the simply connected group G . Notice that θ is always minuscule in $\mathbb{N}\Delta$ but in the cases B_{2n+1} and G_2 : it follows that $\overline{\mathcal{O}}$ is normal in the cases III, VII, IX, whereas it is not normal in the cases I, XI.

7.2. Cases IV (m even), VI, VIII: localization of model wonderful varieties. In this case $M_{\mathcal{O}}$ is not a model wonderful variety, however, it is a quotient of a localization of a model wonderful variety and we still may proceed as in the case of a model orbit thanks to Corollary 1.4 and Corollary 4.14. In order to conclude the argument, we now describe which localizations and quotients we have to take into account in each of the considered cases. We denote by M the model wonderful variety of G .

7.2.1. Case IV (m even). Let N be the boundary divisor corresponding to the spherical root $\sigma_{2n+1} = \alpha_{2n+1} + \alpha_{2n+2}$. Then $M_{\mathcal{O}}$ is the quotient of N by the distinguished subset of colors $\Delta' = \{D_{2n+2}, D_{2n+3}, \dots, D_{2n+m+2}\}$. Since $\theta = D_2$ is minuscule in $\mathbb{N}\Delta$, it follows that $\overline{\mathcal{O}}$ is normal.

7.2.2. Case VI. Let N be the boundary divisor corresponding to the spherical root $\sigma_3 = \alpha_3 + \alpha_4$. Then $M_{\mathcal{O}}$ is the quotient of N by the distinguished subset $\Delta' = \{D_2, D_5, D_7\}$. Since $\theta = D_1$ is minuscule in $\mathbb{N}\Delta$, it follows that $\overline{\mathcal{O}}$ is normal.

7.2.3. *Case VIII.* Let N be the boundary divisor corresponding to the spherical root $\sigma_6 = \alpha_6 + \alpha_7$. Then $M_{\mathcal{O}}$ is the quotient of N by the distinguished subset $\Delta' = \{D_2, D_3, D_4, D_5\}$. Since $\theta = D_8$ is minuscule in $\mathbb{N}\Delta$, it follows that $\overline{\mathcal{O}}$ is normal.

7.3. **Cases II, IV.** In these cases we need to prove the surjectivity of the multiplication for other classes of wonderful varieties. Let $G = \text{Spin}(k)$, let r be the semisimple rank of G (i.e., $k = 2r + 1$ if k odd or $k = 2r$ if k even) and let $2 \leq s \leq (k - 3)/2$. Consider the wonderful variety M corresponding in [6] to the case (18) when k is odd and to the case (43) when k is even.

Its spherical roots and colors are given as follows: $\Sigma = \{\sigma_1, \dots, \sigma_s\}$ and $\Delta = \{D_1, \dots, D_{s+1}\}$, where $\sigma_i = \alpha_i + \alpha_{i+1} = D_i + D_{i+1} - D_{i-1} - D_{i+2}$ for $i = 1, \dots, s-1$ and

$$\sigma_s = \begin{cases} 2(\alpha_{s+1} + \dots + \alpha_r) = 2D_{s+1} - 2D_s & \text{if } k \text{ is odd,} \\ 2(\alpha_{s+1} + \dots + \alpha_{r-2}) + \alpha_{r-1} + \alpha_r = 2D_{s+1} - 2D_s & \text{if } k \text{ is even.} \end{cases}$$

Notice that the Cartan pairing of M does not depend on the parity of k . Also, notice that $\omega(D_i) = \omega_i$ for $i = 1, \dots, s+1$.

First we need to classify the covering differences for M . We omit the proof of the following proposition, which is essentially the same as that of Proposition 3.5.

Proposition 7.3. *Let $\gamma \in \mathbb{N}\Sigma$ be a covering difference in $\mathbb{N}\Delta$ with $\text{supp}_S(\gamma) = S$. Then s is even and $\gamma = \sum_{i=1}^{(j-1)/2} \sigma_{2i-1} + \sum_{i=(j+1)/2}^{s/2} 2\sigma_{2i-1} + \sigma_s = D_1 + D_j - D_{j-1}$ for j odd $\leq s+1$.*

In particular, it follows that every covering difference for M satisfies the property (2-ht). Therefore, by Lemma 2.3, in order to prove the surjectivity of the multiplication of sections of globally generated line bundles on M , we are reduced to the study of the low fundamental triples. Here, as in Lemma 4.7, we have the following.

Lemma 7.4. *Let (D_p, D_q, F) be a low fundamental triple with $\text{supp}_S(D_p + D_q - F) = S$. Then s is even, p, q are odd, $p + q \leq s + 2$ and $F = D_{p+q-2}$.*

Proposition 7.5. *Let (D, E, F) be a low fundamental triple. Then $s^{D+E-F}V_F \subset V_DV_E$.*

Proof. As in the proof of Theorem 4.1, if $\text{supp}(D + E - F) \neq S$ we can proceed by localization and parabolic induction. Therefore it is enough to consider the triples of Lemma 7.4.

Denote by $U = \mathbb{C}^k$ the standard representation of $\text{Spin}(k)$ with the invariant symmetric bilinear form b . Let $V \subset U$ be a totally isotropic subspace of dimension s , let $\omega_V \in \Lambda^2 V$ be a symplectic form on V and fix $e_0 \in U \setminus (V \oplus V^*)$ with $b(e_0, e_0) = 1$.

Let H be the generic stabilizer of M . Then H contains the center of $\text{Spin}(k)$ and its image \overline{H} in $\text{SO}(k)$ is described as follows:

$$\overline{H} = \{g \in \text{SO}(k) : g e_0 = e_0, \quad gV \subset V, \quad g\omega_V \in \mathbb{C}\omega_V\}.$$

If $j \leq s + 1$, then we have $V_{D_j} = V(\omega_j) = \Lambda^j U$. Let h_j be a non-zero H -semi-invariant vector in V_{D_j} . If $2j \leq s$, then up to a scalar factor we have $h_{2j} = \omega_V^{\wedge j}$

and $h_{2j+1} = e_0 \wedge h_{2j}$. The projection $\pi: \Lambda^p U \otimes \Lambda^q U \longrightarrow \Lambda^{p+q-2} U$ is given by contraction as follows:

$$\pi((v_1 \wedge \cdots \wedge v_p) \otimes (u_1 \wedge \cdots \wedge u_q)) = \sum_{i,j} (-1)^{i+j} b(v_i, u_j) v_1 \wedge \cdots \wedge \hat{v}_i \wedge \cdots \wedge \hat{u}_j \wedge \cdots \wedge u_q.$$

Therefore, if (D_p, D_q, D_{p+q-2}) is as in Lemma 7.4, we get $\pi(h_p \otimes h_q) = h_{p+q-2} \neq 0$, and by Lemma 1.2 it follows that $s^{D_p+D_q-D_{p+q-2}} V_{D_{p+q-2}} \subset V_{D_p} V_{D_q}$. \square

If $k = 4n + 2m + 3$ and $s = 2n + 1$, then $M = M_{\mathcal{O}}$ is the wonderful variety corresponding to the nilpotent orbit \mathcal{O} of the case II, while if $k = 4n + 2m + 4$ and $s = 2n + 1$, then $M = M_{\mathcal{O}}$ is the wonderful variety corresponding to the nilpotent orbit \mathcal{O} of the case IV. By the previous proposition, the multiplication of sections is surjective for every couple of globally generated line bundles on $M_{\mathcal{O}}$. Since $\theta = D_2$ is minuscule in $\mathbb{N}\Delta$, it follows that $\overline{\mathcal{O}}$ is normal.

7.4. Case V. The variety $M_{\mathcal{O}}$ corresponds to the case (53) in [6]. We have $\Sigma = \{\sigma_1, \sigma_2, \sigma_3\}$ where

$$\begin{aligned} \sigma_1 &= \alpha_1 + \alpha_6 = 2D_1 - D_3, \\ \sigma_2 &= \alpha_2 + \alpha_4 = D_2 - D_3 + D_4, \\ \sigma_3 &= \alpha_3 + \alpha_5 = -D_1 + 2D_3 - 2D_4, \end{aligned}$$

and $\omega(D_1) = \omega_1 + \omega_6$, $\omega(D_2) = \omega_2$, $\omega(D_3) = \omega_3 + \omega_5$, $\omega(D_4) = \omega_4$. The covering differences are

$$\sigma_1, \sigma_2, \sigma_3, \sigma_1 + \sigma_3, \sigma_2 + \sigma_3, 2\sigma_2 + \sigma_3, \sigma_1 + \sigma_2 + \sigma_3;$$

therefore, we have $\text{ht}(\sigma^+) = 2$ for every covering difference. Correspondingly, we get the following low fundamental triples:

$$\begin{aligned} &(D_1, D_1, D_3), (D_2, D_4, D_3), (D_3, D_3, D_1 + 2D_4), (D_1, D_3, 2D_4), \\ &(D_2, D_3, D_1 + D_4), (D_2, D_2, D_1), (D_1, D_2, D_4). \end{aligned}$$

We need to show that $s^{E+D-F} V_F \subset V_D V_E$ for all low fundamental triples (D, E, F) . As in the proof of Theorem 4.1, if $\text{supp}(D + E - F) \neq S$ we can proceed by localization and parabolic induction. In this way, the triples (D_1, D_1, D_3) , $(D_3, D_3, D_1 + 2D_4)$, $(D_1, D_3, 2D_4)$ reduce to the case of a symmetric wonderful variety, the triple (D_2, D_4, D_3) reduces to a model wonderful variety and the triples $(D_2, D_3, D_1 + D_4)$, (D_2, D_2, D_1) reduce to a wonderful variety studied in the case IV. So we are left to study the cases $D = D_1$, $E = D_2$ and $F = D_4$, which can easily be checked via computer (as explained in Section 4.5). Since $\theta = D_2$ is minuscule in $\mathbb{N}\Delta$, it follows that $\overline{\mathcal{O}}$ is normal.

7.5. Case X. The variety $M_{\mathcal{O}}$ corresponds to the case (60) in [6]. We have $\Sigma = \{\sigma_1, \sigma_2, \sigma_3\}$ where

$$\begin{aligned} \sigma_1 &= 2\alpha_4 = 2D_1 - D_3, \\ \sigma_2 &= \alpha_1 + \alpha_2 = D_2 - D_3 + D_4, \\ \sigma_3 &= 2\alpha_3 = -D_1 + 2D_3 - 2D_4, \end{aligned}$$

and $\omega(D_1) = 2\omega_4$, $\omega(D_2) = \omega_2$, $\omega(D_3) = 2\omega_3$, $\omega(D_4) = \omega_1$. Since the Cartan pairing of $M_{\mathcal{O}}$ is the same as that of the previous case, it follows that the covering differences and the low fundamental triples are also the same.

In order to prove that $s^{E+D-F}V_F \subset V_DV_E$ for all low fundamental triples (D, E, F) , if $\text{supp}(D + E - F) \neq S$ we can proceed by localization and parabolic induction. In this way, the triples (D_1, D_1, D_3) , $(D_3, D_3, D_1 + 2D_4)$, $(D_1, D_3, 2D_4)$ reduce to the case of a symmetric wonderful variety, the triple (D_2, D_4, D_3) reduces to a model wonderful variety and the triples $(D_2, D_3, D_1 + D_4)$, (D_2, D_2, D_1) reduce to a wonderful variety studied in the case III. The only remaining cases, $D = D_1$, $E = D_2$ and $F = D_4$, can easily be checked via computer (as explained in Section 4.5). Since $\theta = D_4$ is minuscule in $\mathbb{N}\Delta$, it follows that $\overline{\mathcal{O}}$ is normal.

8. APPLICATION TO THE REAL MODEL ORBIT OF TYPE E_8

In [1], Adams, Huang and Vogan study the model orbit in the Lie algebra of type E_8 . Their study is motivated by the so-called orbit method to construct representations of reductive Lie groups. In their case the group is the complex algebraic group of type E_8 considered as a real Lie group. In particular they describe the decomposition into irreducible modules of the coordinate ring of the nilpotent orbit of type IX (see Section 7) and prove it is indeed a model orbit. In the same paper they also make some conjectures about another orbit which is the analogue of the model orbit for the split real form of E_8 .

We start with some general preliminaries, we refer to [1] and to [43] for the motivation of these constructions coming from the representation theory of Lie groups. Let $\tilde{G}_{\mathbb{R}}$ be a real form of a connected and complex algebraic semisimple group \tilde{G} and let σ be the associated Galois involution of \tilde{G} . There exists a complex algebraic involution θ of \tilde{G} which commutes with σ such that the subgroup $K_{\mathbb{R}}$ of points of $\tilde{G}_{\mathbb{R}}$ fixed by θ is a maximal compact subgroup of $\tilde{G}_{\mathbb{R}}$. Then the subgroup K of points of \tilde{G} fixed by θ is a complexification of $K_{\mathbb{R}}$. The Lie algebra $\tilde{\mathfrak{g}}$ of \tilde{G} decomposes as $\mathfrak{k} \oplus \mathfrak{p}$ where \mathfrak{k} is the Lie algebra of K and \mathfrak{p} is the eigenspace of eigenvalue -1 of θ . An analogue of the nilpotent cone \mathcal{N} is defined as

$$\mathcal{N}_{\theta} = \mathcal{N} \cap \mathfrak{p}.$$

Fix a point $e \in \mathcal{N}_{\theta}$, let \mathcal{O} be its K -orbit and $K(e)$ its stabilizer. Consider the multiplicative character γ_e of $K(e)$ given by $\gamma_e(g) = \det(\text{Ad}_g|_{\mathfrak{k}(e)}) \det(\text{Ad}_g|_{\mathfrak{k}})^{-1}$. If $\chi: K(e) \rightarrow \mathbb{C}^*$ is any multiplicative character we can consider the algebraic line bundle on \mathcal{O} given by $\mathcal{V}_{\chi} = K \times_{K(e)} \mathbb{C}_{\chi}$. As in [1], the pair (e, χ) is said to be *admissible* if $\overline{\mathcal{O}} \setminus \mathcal{O}$ has codimension at least two in $\overline{\mathcal{O}}$ and $\chi^2(g) = \gamma_e(g)$ for all g in the identity component of $K(e)$. In this paper we need a slightly more general definition of admissible pair.

Let G be a double covering of K and $G(e)$ be the inverse image of $K(e)$ in G so that $\mathcal{O} \simeq G/G(e)$. Let γ'_e denote the character of $G(e)$ induced by γ_e . Given a character χ of $G(e)$ we can construct the line bundle \mathcal{V}_{χ} as above. We say that the pair (e, χ) is admissible if $\overline{\mathcal{O}} \setminus \mathcal{O}$ has codimension at least two in $\overline{\mathcal{O}}$ and $\chi^2(g) = \gamma'_e(g)$ for all g in the identity component of $G(e)$. Let $G_{\mathbb{R}}$ also be the inverse image of $K_{\mathbb{R}}$ in G . Notice that G is the complexification of $G_{\mathbb{R}}$. The coverings of $\tilde{G}_{\mathbb{R}}$ are in correspondence with the coverings of $K_{\mathbb{R}}$ hence there exists a Lie group $\hat{G}_{\mathbb{R}}$ which is a double covering of $\tilde{G}_{\mathbb{R}}$ whose maximal compact subgroup is $G_{\mathbb{R}}$.

Assume that $R(e, \chi)$ is the irreducible unitary representation of $\hat{G}_{\mathbb{R}}$, attached to an admissible pair (e, χ) according to the orbit method. Then the decomposition of the $G_{\mathbb{R}}$ -finite vectors of $R(e, \chi)$ into irreducible submodules should be equal to

the decomposition into irreducible G -submodules of the space of algebraic sections $\Gamma(\mathcal{O}, \mathcal{V}_\chi)$ (see [1, Conjecture 2.9]).

In [1], Adams, Huang and Vogan analyze the geometric side of two particular cases of this construction. They obtain a complete description of the decomposition of $\Gamma(\mathcal{O}, \mathcal{V}_\chi)$ into G -modules in one of these cases and a conjectural description in the second case. In Section 8.1 we recall the result obtained by Adams, Huang and Vogan in the first case and we analyze it using our techniques. In Sections 8.3, 8.4 and 8.6 we analyze the second case considered by Adams, Huang and Vogan and we prove that their conjectural description of $\Gamma(\mathcal{O}, \mathcal{V}_\chi)$ is correct.

8.1. The case of the complex model orbit. Let $\tilde{G}_\mathbb{R}$ be the complex algebraic group of type E_8 . Hence $\tilde{G} = \tilde{G}_\mathbb{R} \times \tilde{G}_\mathbb{R}$ and $G = K$ is the diagonal subgroup. So \mathfrak{p} is isomorphic to the Lie algebra of G and one can consider the nilpotent orbit \mathcal{O} with Kostant-Dynkin diagram equal to $(0, 1, 0, 0, 0, 0, 0, 0)$ (case IX of Section 7). In this case γ_e is trivial and the stabilizer $K(e)$ is connected, so χ also has to be trivial and $\Gamma(\mathcal{O}, \mathcal{V}_\chi) = \mathbb{C}[\mathcal{O}]$. In [1] it is proved that all simple modules of G appear in $\mathbb{C}[\mathcal{O}]$ with multiplicity one.

We fix a maximal torus T , and a Borel subgroup of G containing T . We denote by Φ the set of roots and by S the set of simple roots determined by these choices. We denote also by $\varepsilon_1, \dots, \varepsilon_8$ an orthonormal basis of $\mathcal{X}(T)$ such that Φ and S have the following description (with respect to the choice given in [1] we have changed the sign of ε_1):

$$\begin{aligned} \Phi &= A \cup B \text{ where } A = \{\pm\varepsilon_i \pm \varepsilon_j : i \neq j\} \text{ and} \\ B &= \left\{ \frac{1}{2} \left(\sum_{i=1}^8 a_i \varepsilon_i \right) : a_i = \pm 1 \text{ and } \prod_{i=1}^8 a_i = 1 \right\}; \\ S &= \{\alpha_1, \dots, \alpha_8\} \text{ where } \alpha_1 = -\varepsilon_1 - \varepsilon_2, \\ \alpha_2 &= \frac{1}{2}(\varepsilon_1 - \varepsilon_2 - \varepsilon_3 + \varepsilon_4 + \varepsilon_5 + \varepsilon_6 + \varepsilon_7 + \varepsilon_8) \text{ and} \\ \alpha_i &= \varepsilon_{i-1} - \varepsilon_i \text{ for } i = 3, \dots, 8. \end{aligned}$$

For each root α choose also an $\mathfrak{sl}(2)$ -triple $x_\alpha, \alpha^\vee, y_\alpha$ where x_α has weight α and y_α has weight $-\alpha$. Let $\beta_1 = -\varepsilon_1 + \varepsilon_2$, $\beta_2 = \varepsilon_3 + \varepsilon_4$, $\beta_3 = \varepsilon_5 + \varepsilon_6$, $\beta_4 = \varepsilon_7 + \varepsilon_8$ and define

$$\begin{aligned} e_0 &= x_{\beta_1} + x_{\beta_2} + x_{\beta_3} + x_{\beta_4}, \quad f_0 = y_{\beta_1} + y_{\beta_2} + y_{\beta_3} + y_{\beta_4} \\ h_0 &= \beta_1^\vee + \beta_2^\vee + \beta_3^\vee + \beta_4^\vee = -\varepsilon_1^* + \varepsilon_2^* + \dots + \varepsilon_8^*. \end{aligned}$$

These elements are an $\mathfrak{sl}(2)$ -triple and it is clear that e_0 is an element with associated Kostant-Dynkin diagram equal to $(0, 1, 0, 0, 0, 0, 0, 0)$. Let \mathcal{O} be its orbit. As we have already recalled in Section 7, the orbit of the line spanned by e_0 in $\mathbb{P}(\mathfrak{g})$ is the open orbit of the model wonderful variety of E_8 .

The colors of the model wonderful variety of E_8 are D_1, \dots, D_8 with $\omega(D_i) = \omega_i^\alpha$, where $\omega_1^\alpha, \dots, \omega_8^\alpha$ are the fundamental weights w.r.t. the simple system $\alpha_1, \dots, \alpha_8$. Notice also that, in the notation of the previous section, we have $\overline{\mathcal{O}} = C_{D_8}$. Finally, the spherical roots of M are

$$\begin{aligned} \sigma_1 &= \alpha_6 + \alpha_7, \quad \sigma_2 = \alpha_4 + \alpha_5, \quad \sigma_3 = \alpha_1 + \alpha_3, \quad \sigma_4 = \alpha_2 + \alpha_4, \\ \sigma_5 &= \alpha_5 + \alpha_6, \quad \sigma_6 = \alpha_3 + \alpha_4, \quad \sigma_7 = \alpha_7 + \alpha_8. \end{aligned}$$

Here we have ordered them so that it is clear that they are a basis of a root system of type D_7 . Notice also that, since in this case ω is injective, the list above also determines the Cartan pairing c of the wonderful variety M .

Theorem 8.1 (see [1, Theorem 1.1]). *The variety $\overline{\mathcal{O}}$ is normal and*

$$\mathbb{C}[\overline{\mathcal{O}}] = \mathbb{C}[\mathcal{O}] \simeq \bigoplus_{\lambda \in \mathcal{X}(T)^+} V(\lambda).$$

Proof. We know that $\overline{\mathcal{O}}$ is normal by the discussion in Section 7, in particular, we have $\overline{\mathcal{O}} = C_{D_8} = \tilde{C}_{D_8}$. Moreover, each adjoint orbit has even dimension so the first equality follows by normality. Since $\overline{\mathcal{O}} = \tilde{C}_{D_8}$ we also have

$$\mathbb{C}[\overline{\mathcal{O}}] \simeq \bigoplus_{n \geq 0} \Gamma(M, \mathcal{L}_{nD_8}) \simeq \bigoplus_{\substack{n \geq 0, \gamma \in \mathbb{N}\Sigma: \\ nD_8 - \gamma \in \mathbb{N}\Delta}} s^\gamma V_{nD_8 - \gamma}.$$

Now notice that D_8 and the spherical roots are linearly independent and that ω is injective in this case, so all irreducible G -representations occur with multiplicity at most one. Moreover, since the variety is irreducible, if $V(\lambda)$ and $V(\mu)$ occur in this decomposition then $V(\lambda + \mu)$ also occurs. Finally, we have

$$\begin{aligned} D_1 &= 2D_8 - (\sigma_4 + 2\sigma_5 + \sigma_6 + 2\sigma_7), \\ D_2 &= 4D_8 - (\sigma_1 + 2\sigma_2 + 3\sigma_3 + 4\sigma_4 + 6\sigma_5 + 3\sigma_6 + 5\sigma_7), \\ D_3 &= 5D_8 - (\sigma_1 + 2\sigma_2 + 3\sigma_3 + 5\sigma_4 + 7\sigma_5 + 3\sigma_6 + 6\sigma_7), \\ D_4 &= 7D_8 - (\sigma_1 + 2\sigma_2 + 4\sigma_3 + 6\sigma_4 + 9\sigma_5 + 4\sigma_6 + 8\sigma_7), \\ D_5 &= 6D_8 - (\sigma_1 + 2\sigma_2 + 4\sigma_3 + 6\sigma_4 + 8\sigma_5 + 4\sigma_6 + 7\sigma_7), \\ D_6 &= 4D_8 - (\sigma_2 + 2\sigma_3 + 3\sigma_4 + 4\sigma_5 + 2\sigma_6 + 4\sigma_7), \\ D_7 &= 3D_8 - (\sigma_2 + 2\sigma_3 + 3\sigma_4 + 4\sigma_5 + 2\sigma_6 + 3\sigma_7), \\ D_8 &= D_8 - 0, \end{aligned}$$

hence $V(\omega_i)$ occurs in the decomposition of $\mathbb{C}[\overline{\mathcal{O}}]$, for all i . \square

As we have already recalled above, the second isomorphism in the statement of the theorem was already proved in [1]. Notice that our proof of this isomorphism does not use the projective normality proved in Section 4 but only general considerations on wonderful varieties.

The normality of $\overline{\mathcal{O}}$ was not studied in [1], however, Panyushev in the last section of [38] sketches how to deduce it by their result. Notice that our proof of the normality of $\overline{\mathcal{O}}$, relies on Theorem 4.1 for which, in the case of E_8 , we used a computer. The normality of $\overline{\mathcal{O}}$ is due to A. Broer (see [12, Section 7.8, Remark iii])), however, we could not find a proof in the literature.

8.2. From complex to real orbits: general considerations. Let $\tilde{G}, K, \tilde{\mathfrak{g}}, \mathfrak{p}$ be as at the beginning of this section. Let $\mathcal{O} \subset \tilde{\mathfrak{g}}$ be a nilpotent adjoint orbit of the complex algebraic group \tilde{G} . We want to make some general standard remarks on the intersection $\mathcal{O} \cap \mathfrak{p}$ (see for example [40, Section 9]). Fix $e \in \mathcal{O} \cap \mathfrak{p}$ and let $\tilde{G}(e)$ and $K(e)$ be the stabilizers of e in \tilde{G} and K , respectively. The subgroup $\tilde{G}(e)$ is stable under θ and we define $Z = \{z \in \tilde{G}(e) : \theta(z) = z^{-1}\}$. We have an action of $\tilde{G}(e)$ on Z by $g \cdot z = gz\theta(g)^{-1}$ and we define \mathbb{H}^1 as Z modulo the action of $\tilde{G}(e)$.

- Lemma 8.2.** i) *Every connected component of $\mathcal{O} \cap \mathfrak{p}$ is a single K -orbit.*
 ii) *The map $g \cdot x \mapsto g^{-1}\theta(g)$ induces a bijection from the set of K -orbits in $\mathcal{O} \cap \mathfrak{p}$ to \mathbb{H}^1 .*
 iii) *Every connected component of Z is a single $\tilde{G}(e)^0$ -orbit (where $\tilde{G}(e)^0$ is the identity component of $\tilde{G}(e)$).*

Proof. Set $\eta = -\theta$. First notice that $\mathcal{O} \cap \mathfrak{p} = \mathcal{O}^\eta$. Being \mathcal{O} smooth and η an involution we deduce that \mathcal{O}^η is smooth. Take $e \in \mathcal{O}^\eta$. We prove that

$$(\tilde{\mathfrak{g}} \cdot e) \cap \mathfrak{p} = \mathfrak{k} \cdot e.$$

Let $\tilde{\mathfrak{g}}(e)$ be the annihilator of e in $\tilde{\mathfrak{g}}$ and let $y \in \tilde{\mathfrak{g}}$ be such that $y \cdot e \in \mathfrak{p}$. Then $\theta(y \cdot e) = -y \cdot e$, hence $z = \theta(y) - y \in \tilde{\mathfrak{g}}(e)$. Take $u = \frac{1}{2}z$, then $u \in \tilde{\mathfrak{g}}(e)$ is such that $v = y + u \in \mathfrak{k}$ and $v \cdot e = y \cdot e$.

This proves that any K -orbit in $\tilde{G} \cdot e \cap \mathfrak{p}$ is open. This implies i).

Point iii) can be proved similarly and ii) is trivial. \square

In particular, notice that if we prove that Z is connected then it follows that $\mathcal{O} \cap \mathfrak{p}$ is a single K -orbit.

We refine this lemma, using the Jordan decomposition. Choose an $\mathfrak{sl}(2)$ -triple, e, h, f such that $\theta(h) = h$ and $\theta(f) = -f$, this is always possible; see [32, Proposition 4]. Let U be the unipotent radical of $\tilde{G}(e)$ and $L = \{g \in \tilde{G}(e) : g \cdot h = h\}$. Then $\tilde{G}(e) = L \cdot U$ is a Levi decomposition of $\tilde{G}(e)$ (see [3, Proposition 2.4]). Notice also that L and U are stable under the action of θ . Define $Z_L = Z \cap L$ and \mathbb{H}_L^1 as Z_L modulo the action of L given by $g \cdot z = gz\theta(g)^{-1}$.

Lemma 8.3. i) *The inclusion $Z_L \subset Z$ induces a bijection from \mathbb{H}_L^1 to \mathbb{H}^1 .*

ii) *If Z_L is connected, then $\mathcal{O} \cap \mathfrak{p}$ is a single K -orbit.*

Proof. Let $x, y \in Z_L$ and assume that $x = hy\theta(h)^{-1}$ with $h = \ell u$, $\ell \in L$ and $u \in U$. It follows immediately that $x = \ell y\theta(\ell)^{-1}$. This prove the injectivity of the map $\mathbb{H}_L^1 \rightarrow \mathbb{H}^1$ induced by the inclusion $Z_L \subset Z$.

Now let $z = \ell u \in Z$ with $\ell \in L$ and $u \in U$. From $\theta(z) = z^{-1}$ we deduce that $\theta(\ell) = \ell^{-1}$ and $\theta(u) = \ell u^{-1}\ell^{-1}$. Now U being unipotent there exists a unique $v \in U$ such that $v^{-2} = u$. Using again that U is unipotent we deduce that $\theta(v) = \ell v^{-1}\ell^{-1}$. Set $h' = \ell v\ell^{-1}$, then

$$h'(\ell u)\theta(h')^{-1} = \ell v\ell^{-1}\ell u\ell^{-1}\ell v\ell^{-1}\ell = \ell.$$

Hence the map $\mathbb{H}_L^1 \rightarrow \mathbb{H}^1$ is surjective. This proves i). Now ii) follows from Lemma 8.2. \square

8.3. The case of the real model orbit of E_8 . In the last two sections of [1] a real version of the model orbit is considered. We now recall from [1] some of the structural results about this orbit. We also prove Proposition 8.5 which is probably well known and is somehow implicit (even if not necessary) in the discussion in [1].

From now on \tilde{G} is the complex algebraic group of type E_8 and $\tilde{G}_{\mathbb{R}}$ is its split real form. Then K is isogenous to $\text{Spin}(16)$. For the complex group of type E_8 we have already introduced some notation in Section 8.1 (notice that this group had the role of the group G and not of \tilde{G} in that section). We keep that notation, so T is a maximal torus of \tilde{G} , Φ , the associated root system, and $\varepsilon_1, \dots, \varepsilon_8$, A , B , $S = \{\alpha_1, \dots, \alpha_8\}$ are as in Section 8.1.

Then we can choose $\tilde{G}_{\mathbb{R}}$ so that \mathfrak{k} is the Lie algebra spanned by \mathfrak{t} , the Lie algebra of T , and by the vectors x_{α} with $\alpha \in A$. With this choice \mathfrak{p} is spanned by the vectors x_{α} with $\alpha \in B$.

As a simple system for the root system of $\mathfrak{k} \simeq \mathfrak{so}(16)$ we choose the usual basis, but we enumerate it starting from zero, that is,

$$\tau_i = \varepsilon_{i+1} - \varepsilon_{i+2} \text{ for } i = 0, \dots, 6 \text{ and } \tau_7 = \varepsilon_7 + \varepsilon_8.$$

We denote by $\omega_i^{\mathbb{D}}$ the associated fundamental weights. In particular we obtain that \mathfrak{p} is the spin representation associated to the weight $\omega_7^{\mathbb{D}}$. Moreover, since \tilde{G} is simply connected the subgroup K is connected. Finally, notice that $\omega_8^{\mathbb{D}} \notin \mathcal{X}(T)$ while $\omega_7^{\mathbb{D}} \in \mathcal{X}(T)$. Hence $\text{Spin}(16)$ is a double covering of K . We set $G = \text{Spin}(16)$. Notice that in [1] it is claimed that K is isomorphic to $\text{Spin}(16)$, but this does not affect any of their arguments.

In order to prove that the roots τ_1, \dots, τ_7 are conjugated to the roots $\sigma_1, \dots, \sigma_7$ introduced in Section 8.1, we introduce a new simple system of the root system of type E_8 . The vectors

$$\begin{aligned} \zeta_1 &= \frac{1}{2}(-\varepsilon_2 - \varepsilon_3 - \varepsilon_4 + \varepsilon_5), & \zeta_2 &= \frac{1}{2}(\varepsilon_1 + \varepsilon_6 + \varepsilon_7 - \varepsilon_8), \\ \zeta_3 &= \frac{1}{2}(\varepsilon_2 + \varepsilon_3 - \varepsilon_4 + \varepsilon_5), & \zeta_4 &= \frac{1}{2}(\varepsilon_1 + \varepsilon_6 - \varepsilon_7 + \varepsilon_8), \\ \zeta_5 &= \frac{1}{2}(\varepsilon_2 - \varepsilon_3 + \varepsilon_4 + \varepsilon_5), & \zeta_6 &= \frac{1}{2}(\varepsilon_1 - \varepsilon_6 + \varepsilon_7 + \varepsilon_8), \\ \zeta_7 &= \frac{1}{2}(-\varepsilon_2 + \varepsilon_3 + \varepsilon_4 + \varepsilon_5), & \zeta_8 &= \frac{1}{2}(\varepsilon_1 - \varepsilon_6 - \varepsilon_7 - \varepsilon_8), \end{aligned}$$

form an orthonormal basis of \mathfrak{t}^* , and $\pm\zeta_i \pm \zeta_j \in \Phi$ for i odd and j even. Notice also that

$$\begin{aligned} \gamma_2 &= \frac{1}{2}(\zeta_1 - \zeta_2 - \zeta_3 + \zeta_4 + \zeta_5 + \zeta_6 + \zeta_7 + \zeta_8) \\ &= \frac{1}{2}(\varepsilon_1 - \varepsilon_2 - \varepsilon_3 + \varepsilon_4 + \varepsilon_5 - \varepsilon_6 - \varepsilon_7 + \varepsilon_8) \in \Phi, \\ \gamma_1 &= -\zeta_1 - \zeta_2, \text{ and } \gamma_i = \zeta_{i-1} - \zeta_i, \text{ for } i = 3, \dots, 8, \end{aligned}$$

form a simple system of Φ (since they are elements of Φ with the right scalar products). Hence there exists an element w in the Weyl group such that $w(\varepsilon_i) = \zeta_i$, for all i . Finally, notice that we have

$$\begin{aligned} \tau_1 &= \gamma_6 + \gamma_7, & \tau_2 &= \gamma_4 + \gamma_5, & \tau_3 &= \gamma_1 + \gamma_3, & \tau_4 &= \gamma_2 + \gamma_4, \\ \tau_5 &= \gamma_5 + \gamma_6, & \tau_6 &= \gamma_3 + \gamma_4, & \tau_7 &= \gamma_7 + \gamma_8, \end{aligned}$$

so that $w(\sigma_i) = \tau_i$. Notice also that

$$\tau_0 = \frac{1}{2}(\zeta_1 - \zeta_3 - \zeta_5 + \zeta_7 + \zeta_2 + \zeta_4 + \zeta_6 + \zeta_8) = \gamma_2 + \gamma_3 + \gamma_4 + \gamma_5.$$

We denote by Λ the weight lattice of $\text{Spin}(16)$ and by $\Lambda_E = \mathcal{X}(T)$ the sublattice given by the weights of E_8 . We denote also by Λ^+ the dominant weights of Λ w.r.t. the simple system τ_0, \dots, τ_7 and by Λ_E^+ the dominant weights of Λ_E w.r.t. the simple system $\gamma_1, \dots, \gamma_8$. We have $\Lambda_E^+ \subset \Lambda^+$. We denote by ω_i^{γ} the fundamental weights of Λ_E w.r.t. the simple system $\gamma_1, \dots, \gamma_8$. Notice that $\omega_7^{\mathbb{D}} = \omega_8^{\gamma}$.

Now let \mathcal{O} be the adjoint orbit of $\tilde{\mathfrak{g}}$ considered in Section 8.1. We want to study the intersection $\mathcal{O} \cap \mathfrak{p}$. Let us first choose an element $e \in \mathcal{O} \cap \mathfrak{p}$.

Consider the semisimple part K_0 of the standard Levi factor of the maximal parabolic subgroup P_0 of K associated with τ_0 . This is a group isogeneous to $\text{Spin}(14)$, we denote its Lie algebra by \mathfrak{k}_0 . Let \mathfrak{p}_0 be the subspace of \mathfrak{p} spanned

by the root vectors of weight of the form $\frac{1}{2}(\sum a_i \varepsilon_i)$ with $a_1 = -1$. Its highest weight is $\frac{1}{2}(-\varepsilon_1 + \varepsilon_2 + \varepsilon_3 + \varepsilon_4 + \varepsilon_5 + \varepsilon_6 + \varepsilon_7 - \varepsilon_8)$. Hence it is an irreducible Spin(14)-submodule of \mathfrak{p} isomorphic to the module $V_{D_8}^*$ of Section 5.5.3.

Lemma 8.4. *The vector h_8 of Section 5.5.3 belongs to \mathcal{O} .*

Proof. The vectors

$$(8.1a) \quad \eta_1 = \frac{1}{2}(\varepsilon_1 - \varepsilon_6 - \varepsilon_7 - \varepsilon_8), \quad \eta_2 = \frac{1}{2}(-\varepsilon_2 + \varepsilon_3 + \varepsilon_4 + \varepsilon_5),$$

$$(8.1b) \quad \eta_3 = \frac{1}{2}(-\varepsilon_1 + \varepsilon_6 - \varepsilon_7 - \varepsilon_8), \quad \eta_4 = \frac{1}{2}(\varepsilon_2 - \varepsilon_3 + \varepsilon_4 + \varepsilon_5),$$

$$(8.1c) \quad \eta_5 = \frac{1}{2}(-\varepsilon_1 - \varepsilon_6 + \varepsilon_7 - \varepsilon_8), \quad \eta_6 = \frac{1}{2}(\varepsilon_2 + \varepsilon_3 - \varepsilon_4 + \varepsilon_5),$$

$$(8.1d) \quad \eta_7 = \frac{1}{2}(-\varepsilon_1 - \varepsilon_6 - \varepsilon_7 + \varepsilon_8), \quad \eta_8 = \frac{1}{2}(\varepsilon_2 + \varepsilon_3 + \varepsilon_4 - \varepsilon_5)$$

form an orthonormal basis of \mathfrak{t}^* , and $\pm\eta_i \pm \eta_j \in \Phi$ if i is odd and j is even. Notice also that

$$\hat{\gamma}_2 = \frac{1}{2}(\eta_1 - \eta_2 - \eta_3 + \eta_4 + \eta_5 + \eta_6 + \eta_7 + \eta_8) = \varepsilon_2 - \varepsilon_6,$$

$$\hat{\gamma}_1 = -\eta_1 - \eta_2, \text{ and } \hat{\gamma}_i = \eta_{i-1} - \eta_i, \text{ for } i = 3, \dots, 8,$$

is a simple system of Φ . Hence there exists an element w of the Weyl group such that $w(\varepsilon_i) = \eta_i$, for all i . Choose a representative \dot{w} of w in \tilde{G} . Define $e = \dot{w}(e_0)$, $h = w(h_0) = -\eta_1^* + \eta_2^* + \dots + \eta_8^*$ and $f = \dot{w}(f_0)$. Then $\theta(h) = h$ and $\theta(f) = -f$. More explicitly, we have $h = -2\varepsilon_1^* + \varepsilon_2^* + \varepsilon_3^* + \varepsilon_4^* + \varepsilon_5^*$, $e = x_{\delta_1} + x_{\delta_2} + x_{\delta_3} + x_{\delta_4}$, where $w(\beta_1) = \delta_1 = -\eta_1 + \eta_2$, $w(\beta_2) = \delta_2 = \eta_3 + \eta_4$, $w(\beta_3) = \delta_3 = \eta_5 + \eta_6$, $w(\beta_4) = \delta_4 = \eta_7 + \eta_8$ and similarly for f .

Notice that $e \in \mathfrak{p}_0$. Moreover, the two vectors e and h_8 are linear combinations with non-zero coefficients of vectors of the same weights $\delta_1, \delta_2, \delta_3, \delta_4$ and these weights are linearly independent, so they are conjugated under the action of the maximal torus. \square

Let $w \in W$ be the element defined in the previous proof, as shown there we can choose a representative \dot{w} of w such that $\dot{w}(e_0) = h_8$. Set $e = h_8$ and $h = w(h_0)$. In particular, the stabilizer of e is the parabolic induction of the stabilizer of h_8 in Spin(14). More explicitly, we have

$$(8.2) \quad \mathfrak{k}(e) = \mathbb{C}\omega_0^\vee \oplus \mathfrak{u}_0^- \oplus \mathfrak{h}_0,$$

where \mathfrak{u}_0^- is the Lie algebra of the unipotent radical of the parabolic subgroup opposite to P_0 , \mathfrak{h}_0 is the annihilator of h_8 in \mathfrak{k}_0 and ω_0^\vee is orthogonal to τ_1, \dots, τ_7 . In particular, the Levi factor of $\mathfrak{k}(e)$ is isomorphic to $\mathfrak{gl}(4)$.

We now want to describe in some detail the stabilizer $K(e)$ and apply Lemma 8.3 to prove that $\mathcal{O} \cap \mathfrak{p}$ is a single K -orbit. As recalled in Section 8.1, the Levi factor \tilde{L} of $\tilde{G}(e)$ is Sp(8) and $L = \tilde{L}^\theta$. Furthermore, notice that there is only one involution of Sp(8) such that the Lie algebra of the fixed point subgroup is isomorphic to $\mathfrak{gl}(4)$. This involution can be described as follows. Let $I_4 \in \text{SL}(4)$ be the identity matrix, and define the 8×8 matrices

$$J = \begin{pmatrix} 0 & I_4 \\ -I_4 & 0 \end{pmatrix}, \quad \rho = \begin{pmatrix} I_4 & 0 \\ 0 & -I_4 \end{pmatrix},$$

and Sp(8) as the matrices preserving the form J . Then Sp(8) is stable under the conjugation by the matrix ρ , which is an involution. Moreover, L is isomorphic to GL(4).

Proposition 8.5. *If \mathcal{O} is the model orbit of E_8 and $\tilde{G}_{\mathbb{R}}$ is the split real form of E_8 , then $\mathcal{O} \cap \mathfrak{p}$ is a single K -orbit.*

Proof. By Lemma 8.3 and the above discussion, it is enough to prove that

$$Z_L = \{z \in \mathrm{Sp}(8) : \theta(z) = z^{-1}\}$$

is connected. Let $z = \begin{pmatrix} A & B \\ C & D \end{pmatrix}$ where A, B, C, D are 4×4 matrices. The condition $z \in Z_L$ is equivalent to

(8.3)

$$A^2 = I_4 + BC, \quad D = A^t, \quad B = B^t, \quad C = C^t, \quad AB = BA^t \text{ and } A^t C = CA.$$

Acting by $g \in \mathrm{Sp}(8)$, via $g \cdot z = gz\theta(g)^{-1}$, we remain in the same connected component. Using g of the form $\begin{pmatrix} \alpha & & & \\ & \iota\alpha^{-1} & & \\ & & & \\ & & & \end{pmatrix}$ with $\alpha \in \mathrm{GL}(4)$ we see that we can assume B is of the form $\begin{pmatrix} I_r & 0 \\ & 0 \end{pmatrix}$ and $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$, and using equations (8.3) we see that $c = 0$, $a = a^t$ and $d^2 = 1$. Then using g of the forms $\begin{pmatrix} \beta & & & \\ & \iota\beta^{-1} & & \\ & & & \\ & & & \end{pmatrix}$ and $\beta = \begin{pmatrix} I_r & 0 \\ & \gamma \end{pmatrix}$ we can also assume that d is diagonal. Now we choose g of the form $\begin{pmatrix} I & sI \\ & I \end{pmatrix}$. We get

$$g \cdot z = \begin{pmatrix} A + sC & B + s(A + A^t) + s^2C \\ C & A^t + sC \end{pmatrix}.$$

If we compute the determinant of $B + s(A + A^t) + s^2C$ we see that it is a polynomial in s and its lowest degree term is $4 \det(d)$. Hence there exists s such that $B + s(A + A^t) + s^2C$ is invertible. So we can assume B invertible and arguing as before we can assume $B = I_4$. Now, for $B = I_4$, the equations in (8.3) take the form $A = A^t = D$, and $C = -I + A^2$. Such equations define an algebraic subset which is isomorphic to an affine space and, in particular, connected. Therefore Z_L is connected. \square

8.4. The coordinate ring of the real model orbit. Here we describe the coordinate ring of $\mathcal{O} \cap \mathfrak{p}$. In [1] it is shown that this description follows from the vanishing of certain cohomology groups which in the case of the real model orbit is conjectural (see Conjecture 3.13 and Theorem 7.13 in [1]).

Let M_0 be the wonderful comodel variety of cotype E_8 . This is a wonderful variety for the group $\mathrm{Spin}(14)$. Consider the parabolic induction M of M_0 to $\mathrm{Spin}(16)$. This is a wonderful variety with spherical roots equal to τ_1, \dots, τ_7 and colors D'_0, \dots, D'_8 , where $\omega(D'_0) = \omega_0^D$ and, for $i > 0$, D'_i is induced by the respective color of M_0 (see Section 5).

Let $\mathcal{O}_{\mathfrak{p}} = \mathcal{O} \cap \mathfrak{p}$. Then $X_{\mathfrak{p}} = \mathbb{P}(\mathcal{O}_{\mathfrak{p}}) \subset \mathbb{P}(\mathfrak{p})$ is the open orbit of M , and $\overline{\mathcal{O}_{\mathfrak{p}}}$ is the cone C_D for $D = D'_8$. Indeed, $e = h_8 \in V_{\omega^*(D'_8)} = V_{\omega_7^*} = V_{\omega_7^D} \simeq \mathfrak{p}$ and D'_8 is faithful.

Theorem 8.6. *The cone $\overline{\mathcal{O}_{\mathfrak{p}}}$ is normal and we have the following isomorphism of K -modules:*

$$\mathbb{C}[\overline{\mathcal{O}_{\mathfrak{p}}}] \simeq \bigoplus_{\lambda \in \Lambda_{\mathbb{E}}^+} V(\lambda).$$

Proof. Notice that the combinatorics of colors and spherical roots is essentially the same as that of the model wonderful variety of type E_8 . In particular, since D'_8 is minuscule, from Theorem 5.2 it follows that $\overline{\mathcal{O}_{\mathfrak{p}}}$ is normal. Hence, as a K -module we have that its coordinate ring is the sum of all the modules $V_{nD'_8 - \tau}$ with $n \geq 0$, $\tau \in \mathbb{N}[\tau_1, \dots, \tau_7]$ and $nD'_8 - \tau \in \mathbb{N}[D'_0, \dots, D'_8]$. Moreover, with τ as above, the latter condition is equivalent to $nD'_8 - \tau \in \mathbb{N}[D'_1, \dots, D'_8]$. Now notice that $\omega(D'_8)$ and τ_1, \dots, τ_7 are linearly independent (this is not true for M_0 but it is true for M

because of the presence of the extra color D'_0). So we obtain that the coordinate ring of $\overline{\mathcal{O}_p}$ is the sum of all modules $V(n\omega_s^D - \tau)$ where n and τ are as above. Finally, the computation is exactly the same as that given in the proof of Theorem 8.1 for the model orbit of type E_8 , since the two situations are conjugated by an element of the Weyl group. \square

Notice that our proof of the normality of $\overline{\mathcal{O}_p}$ via Theorem 5.2 did not require any computer calculation. Moreover, the description of the coordinate ring of the normalization of $\overline{\mathcal{O}_p}$ is independent of Theorem 5.2.

The combinatorics of distinguished subsets of colors allows us to describe completely the K -orbits in the closure of \mathcal{O}_p (see for example [22]), and in particular to prove that $\overline{\mathcal{O}_p} \setminus \mathcal{O}_p$ has codimension at least two in $\overline{\mathcal{O}_p}$. Indeed, one sees that this property depends only on the combinatorics of colors and spherical roots, and in this case the combinatorics is the same as that of the complex model orbit, whose boundary has codimension at least two. More precisely, here $\text{codim}_{\overline{\mathcal{O}}}(\overline{\mathcal{O}} \setminus \mathcal{O}) = 16$ and $\text{codim}_{\overline{\mathcal{O}_p}}(\overline{\mathcal{O}_p} \setminus \mathcal{O}_p) = 8$.

Here we can avoid such an argument. Below we prove that $\mathbb{C}[\mathcal{O}_p] = \mathbb{C}[\overline{\mathcal{O}_p}]$ and this also implies that $\overline{\mathcal{O}_p} \setminus \mathcal{O}_p$ has codimension at least two.

8.5. Computation of the space of sections of a line bundle associated to an admissible pair: general considerations.

For the next lemma we put ourselves in a general setting. Let G be simply connected and let M be a wonderful compactification of G/H . Let $E \in \mathbb{N}\Delta$. Let $\mathbb{C}h$ be a line in $V(\omega_E^*)$ where H acts with character ξ_E . Assume that the stabilizer of $\mathbb{C}h$ is equal to H and let H_0 be the stabilizer of h . Furthermore, assume that ξ_E induces an isomorphism of H/H_0 with \mathbb{C}^* : we identify H/H_0 with \mathbb{C}^* choosing such an isomorphism. Finally, notice that H/H_0 acts on the right on G/H_0 .

Lemma 8.7. *Let $D \in \mathbb{Z}\Delta$ and let χ denote the restriction of ξ_D to H_0 . We have the following isomorphism of G -modules*

$$\Gamma(G/H_0, \mathcal{V}_\chi) \simeq \bigoplus_{\substack{n \in \mathbb{Z}, \sigma \in \mathbb{Z}\Sigma: \\ D+nE-\sigma \in \mathbb{N}\Delta}} V_{D+nE-\sigma}.$$

Proof. We have

$$\Gamma(G/H_0, \mathcal{V}_\chi) \simeq (\mathbb{C}[G] \otimes \mathbb{C}_{-\chi})^{H_0} \simeq \bigoplus_{\lambda \in \mathcal{X}(T)^+} V(\lambda^*) \otimes (V(\lambda) \otimes \mathbb{C}_{-\chi})^{H_0}.$$

Now notice that H acts on $(V(\lambda) \otimes \mathbb{C}_{-\chi})^{H_0}$ and so the latter decomposes into H -stable subspaces where H acts by a character of the form $\xi_D + n\xi_E$. Moreover, all these eigenspaces are of dimension one, since H is spherical. Hence we have

$$\Gamma(G/H_0, \mathcal{V}_\chi) \simeq \bigoplus_{\lambda \in \mathcal{X}(T)^+, n \in \mathbb{Z}} V(\lambda^*) \otimes (V(\lambda) \otimes \mathbb{C}_{-\xi_D - n\xi_E})^H.$$

Now recall that the space of spherical vectors $V(\omega_F^*)_{\xi_F}^{(H)}$ is non-zero if and only if $F \in \mathbb{N}\Delta$, and $\xi_F = \xi_{D+nE}$ if and only if $F = D + nE - \sigma$ for $\sigma \in \mathbb{Z}\Sigma$. \square

Let us specialize this identity to the case $D = 0$ and compare the coordinate ring of G/H_0 with the coordinate ring of the normalization of its closure in $V(\omega_E^*)$.

Since

$$\mathbb{C}[\tilde{\mathcal{C}}_E] = \bigoplus_{n \geq 0} \Gamma(M, \mathcal{L}_{nE}) \simeq \bigoplus_{\substack{n \geq 0, \sigma \in \mathbb{N}\Sigma: \\ nE - \sigma \in \mathbb{N}\Delta}} V_{nE - \sigma},$$

we have the following.

Lemma 8.8. *The equality $\mathbb{C}[G/H_0] = \mathbb{C}[\tilde{\mathcal{C}}_E]$ is equivalent to the fact that, for all $n \in \mathbb{Z}$ and $\sigma \in \mathbb{Z}\Sigma$, if $nE - \sigma \in \mathbb{N}\Delta$ then $n \geq 0$ and $\sigma \in \mathbb{N}\Sigma$.*

We now apply this lemma to our special case in which G is $\text{Spin}(16)$, $H_0 = G(e)$, M is the wonderful compactification of $X_{\mathfrak{p}}$ and $E = D'_8$.

Lemma 8.9. *For all $n \in \mathbb{Z}$ and $\sigma \in \mathbb{Z}\Sigma$, if $nD'_8 - \sigma \in \mathbb{N}\Delta$ then $n \geq 0$ and $\sigma \in \mathbb{N}\Sigma$.*

Proof. Write $nD'_8 - \sigma = \sum_{i=0}^8 a_i D'_i$ and notice that the conditions $a_1, \dots, a_8 \geq 0$ are equivalent to the conditions obtained from Theorem 8.1 by applying Lemma 8.8 to the complex model orbit of type E_8 . \square

From this and Theorem 8.6, by applying Lemma 8.8 to the real model orbit, we get that

$$\mathbb{C}[\mathcal{O}_{\mathfrak{p}}] = \mathbb{C}[\overline{\mathcal{O}_{\mathfrak{p}}}]$$

8.6. Sections of the line bundle associated to the admissible pair for the real model orbit.

We now want to compute the characters γ_e and γ'_e . For this we further analyze the stabilizer $T(e)$ of e in T . Let Λ_E^\vee be the lattice dual to Λ_E . Recall the definition of the vectors η_i given in the equations (8.1) and denote by η_i^* the vectors of the dual basis. Let $x_1^* = \eta_1^* + \eta_2^*$ and $x_i^* = \eta_{2i-1}^* - \eta_{2i}^*$ for $i = 2, 3, 4$ and define R^\vee as the sublattice of Λ_E^\vee generated by x_1^*, \dots, x_4^* and R_0^\vee as the sublattice generated by $x_1^* + x_2^*, x_1^* + x_3^*, x_1^* + x_4^*$. Then R^\vee and R_0^\vee are direct factors of Λ_E^\vee . Finally, let T_0 be the maximal torus of K_0 (the subgroup of K isogenous to $\text{Spin}(14)$ introduced above) contained in T such that $T_0(e) = T(e) \cap T_0$. We have

$$T = \Lambda_E^\vee \otimes_{\mathbb{Z}} \mathbb{C}^*, \quad T(e) = R^\vee \otimes_{\mathbb{Z}} \mathbb{C}^* \quad \text{and} \quad T_0(e) = R_0^\vee \otimes_{\mathbb{Z}} \mathbb{C}^*.$$

We already know that the Levi factor of $K(e)$ is isomorphic to $\text{GL}(4)$ so that any character is a power of the determinant. We describe now the center of $\text{GL}(4)$ and the determinant as an element of dual lattice of R^\vee . Let $z^* = x_1^* - x_2^* - x_3^* - x_4^* = 2\varepsilon_1^* + \varepsilon_2^* + \varepsilon_3^* + \varepsilon_4^* + \varepsilon_5^*$, then using the description of \mathfrak{h}_0 given in Section 5.5 it is easy to see that z^* is a central cocharacter. In particular, if $R_Z^\vee = \mathbb{Z}z^*$ then

$$T_Z = R_Z^\vee \otimes_{\mathbb{Z}} \mathbb{C}^*$$

is the center of $L \simeq \text{GL}(4)$. Now we compute the character γ_e . We have already noticed that it is enough to compute its restriction to T . Moreover, its restriction to $T_0(e)$ must be trivial since $\text{SL}(4)$ has no characters so we compute its restriction to T_Z . Using the description of the stabilizer $\mathfrak{k}(e)$ in (8.2) and the description of \mathfrak{h}_0 given in Section 5.5 it is easy to prove that $\langle \gamma_e, z^* \rangle = 4$. Hence we see that γ_e restricted to the Levi of $K(e)$, which is isomorphic to $\text{GL}(4)$, equals the determinant and

$$\gamma_e = x_1 - x_2 - x_3 - x_4.$$

In particular, it is not the square of a character. However, on the covering G of K we can consider the character $-\omega_0^D = -\varepsilon_1$, and we denote by χ its restriction to $G(e)$. We have

$$\chi = \frac{1}{2}(x_1 - x_2 - x_3 - x_4) = \frac{1}{2}\gamma'_e.$$

Notice that $\xi_{D'_0} = \chi$. Indeed, $\omega_{D'_0} = \omega_0^D$ and $G(e)$ contains the unipotent radical U_0^- of P_0^- . Hence the only $G(e)$ -semi-invariant in $V(\omega_0^D)$ is the lowest weight vector which has weight $-\omega_0^D$.

We now describe the space of sections $\Gamma(\mathcal{O}_{\mathfrak{p}}, \mathcal{V}_\chi)$. In [1] the same description follows from a vanishing result which is not proved in this case (see Conjecture 8.6 and Theorem 8.10 in [1]).

Theorem 8.10. *We have the following isomorphism of G -modules:*

$$\Gamma(\mathcal{O}_{\mathfrak{p}}, \mathcal{V}_\chi) \simeq \bigoplus_{\lambda \in \Lambda_{\mathbb{E}}^+} V(\omega_0^D + \lambda).$$

Proof. We have seen above that $\xi_{D'_0} = \chi$. Hence we can apply Lemma 8.7 with $D = D'_0$ and $E = D'_8$. We obtain

$$\Gamma(\mathcal{O}_{\mathfrak{p}}, \mathcal{V}_\chi) \simeq \bigoplus_{\substack{n \in \mathbb{Z}, \sigma \in \mathbb{Z}\Sigma \\ D'_0 + nD'_8 - \sigma \in \mathbb{N}\Delta}} V_{D'_0 + nD'_8 - \sigma}.$$

Write $D'_0 + nD'_8 - \sigma = \sum_{i=0}^8 a_i D'_i$. Notice that as in the proof of Lemma 8.9 this implies $a_1, \dots, a_8 \geq 0$. Hence we obtain $n \geq 0$ and $\sigma \in \mathbb{N}\Sigma$. In particular, the condition $a_0 \geq 0$ is automatically satisfied. As already noticed in the proof of Theorem 8.6, $\Lambda_{\mathbb{E}}^+$ consists of the weights of the form $\omega(nD'_8 - \sigma)$ for $n \geq 0$, $\sigma \in \mathbb{N}\Sigma$ and $nD'_8 - \sigma \in \mathbb{N}\Delta$. \square

9. DEGENERACY OF THE MULTIPLICATION

Here we give a counterexample to the surjectivity of the multiplication of sections of line bundles, on wonderful varieties, generated by global sections.

9.1. Counterexample. Let G be $\mathrm{SO}(2r+1)$ and let M be the model wonderful variety for the group G . It is not isomorphic to the model wonderful variety for $\mathrm{Spin}(2r+1)$. Its set of colors $\Delta = \{D_1, \dots, D_r\}$ is in bijection, via ω , with the set $\{\omega_1, \dots, \omega_{r-1}, 2\omega_r\}$. Its set of spherical roots is $\{\alpha_1 + \alpha_2, \dots, \alpha_{r-1} + \alpha_r, 2\alpha_r\}$. If $r = 2$ or $r = 3$, then the multiplication of sections is surjective. Set $r = 4$ and consider the low triple (D_2, D_2, D_1) ; then $V(\omega_1) \not\subset V(\omega_2)^{\otimes 2}$, hence $s^{2D_2 - D_1} V_{D_1} \not\subset V_{D_2}^2$. In particular, the multiplication of sections is not surjective, for all $r \geq 4$.

9.2. Degeneracy of the multiplication. Roughly speaking, in its nature the previous example does not express a lack of the multiplication, but rather a lack of the tensor product. Indeed, $V(\omega_1) \not\subset V(\omega_2)^{\otimes 2}$ but $V(2\omega_1) \subset V(2\omega_2)^{\otimes 2}$, so that it expresses the fact that the saturation property does not hold for groups of type B: similar things cannot happen if G is of type A and, conjecturally, whenever G is simply laced. It is worth noticing that the same situation holds if we consider the multiplication of the wonderful variety considered in a previous example: $s^{2D_2 - D_1} V_{D_1} \not\subset V_{D_2}^2$ but $s^{4D_2 - 2D_1} V_{2D_1} \subset V_{2D_2}^2$.

We briefly recall what the saturation property is. Let G be a simply connected almost simple algebraic group. We say that the *saturation property* holds for G if, whenever $d > 0$ and $\lambda, \mu, \nu \in \Lambda^+$ are such that $\nu \leq \lambda + \mu$ and $V(d\nu) \subset V(d\lambda) \otimes V(d\mu)$, then it also holds that $V(\nu) \subset V(\lambda) \otimes V(\mu)$. In [31] A. Knutson and T. Tao showed that the saturation property holds if G is of type A, while in [28] M. Kapovich and J. Millson conjectured that the saturation property holds whenever G is simply laced.

We want to consider the saturation property in the more general context of the multiplication law attached to a wonderful variety, the classical case corresponding to the wonderful compactification of an adjoint group ([27, Lemma 3.1]). We say that the *saturation property* holds for a wonderful variety M with set of colors Δ and set of spherical roots Σ if, whenever $d > 0$ and $D, E, F \in \mathbb{N}\Delta$ are such that $D \leq_{\Sigma} E + F$ and $s^{d(E+F-D)}V_{dD} \subset V_{dE}V_{dF}$, then the inclusion $s^{E+F-D}V_D \subset V_EV_F$ also holds.

Suppose that M is a wonderful variety and let $E, F \in \mathbb{N}\Delta$. Then the following inclusion holds:

$$V_EV_F \subset \bigoplus_{\substack{D \in \mathbb{N}\Delta : D \leq_{\Sigma} E+F \\ V(\omega_D) \subset V(\omega_E) \otimes V(\omega_F)}} s^{E+F-D}V_D.$$

If the equality holds in the previous inclusion, then we say that the product V_EV_F is *non-degenerate*, while if the equality holds for every $E, F \in \mathbb{N}\Delta$ then we say that the multiplication of M is *non-degenerate*. It is easy to show that if the multiplication of M is non-degenerate and if the saturation property holds for G , then the latter holds for M as well.

Another equivalent description of the multiplication follows by identifying sections of line bundles on M with H -semi-invariant functions on G , H acting on the right. More precisely, given $E, F \in \mathbb{N}\Delta$, we may identify $\Gamma(M, \mathcal{L}_E)$, $\Gamma(M, \mathcal{L}_F)$ and $\Gamma(M, \mathcal{L}_{E+F})$ with submodules of $\mathbb{C}[G]^{(H)}$, and the multiplication of sections $\Gamma(M, \mathcal{L}_E) \otimes \Gamma(M, \mathcal{L}_F) \rightarrow \Gamma(M, \mathcal{L}_{E+F})$ coincides with the restriction of the multiplication defined in $\mathbb{C}[G]^{(H)}$. Therefore we regard the modules V_E, V_F, V_EV_F inside $\mathbb{C}[G]^{(H)}$, and the multiplication is the multiplication of functions.

Example 1. Let $G = \mathrm{SL}(2)$ and consider the basic case of the rank one wonderful variety $M = \mathbb{P}^1 \times \mathbb{P}^1$, whose generic stabilizer is the maximal torus T . Then $\Sigma = \{\alpha\}$ is the unique simple root of G and $\Delta = \{D^+, D^-\}$, where $D^+ = \mathbb{P}^1 \times \{[1, 0]\}$ and $D^- = \{[1, 0]\} \times \mathbb{P}^1$. Moreover $\alpha = D^+ + D^-$ and $\omega_{D^+} = \omega_{D^-} = 1$ equal the fundamental weight of G .

Given $n \in \mathbb{N}$, identify the simple G -module $V(n)$ of highest weight n with $\mathbb{C}[x, y]_n$, the set of homogeneous polynomials of degree n in two variables. Given $aD^+ + bD^- \in \mathbb{N}\Delta$, the corresponding T -eigenvector is $h(a, b) = x^a y^b$.

The projection $\pi_m : V(a+b) \otimes V(c+d) \rightarrow V(a+b+c+d-2m)$ is described on the basis of T -eigenvectors as follows:

$$x^a y^b \otimes x^c y^d \mapsto \sum_{u+v=m} (-1)^u u! v! \binom{a}{u} \binom{b}{v} \binom{c}{v} \binom{d}{u} x^{a+c-m} y^{b+d-m}.$$

As shown by the following two examples, the multiplication of M is degenerate.

- Consider $D^+ + D^- \leq_{\Sigma} 2(D^+ + D^-)$. Then $V(2) \subset V(2)^{\otimes 2}$ but $\pi_1(h(1, 1) \otimes h(1, 1)) = 0$, so that $V_{D^++D^-} \not\subset V_{D^++D^-}^2$. This can also be explained since $V(2)$ is not contained in the second symmetric power of $V(2)$.
- Consider $2D^+ + D^- \leq_{\Sigma} (3D^+ + D^-) + (D^+ + 2D^-)$. Then $V(3) \subset V(4) \otimes V(3)$ but $\pi_2(h(3, 1) \otimes h(1, 2)) = 0$, so that $V_{2D^++D^-} \not\subset V_{3D^++D^-} V_{D^++2D^-}$.

More generally, the multiplication is degenerate whenever $\Sigma \cap S \neq \emptyset$. This can be reduced to the basic case of $\mathrm{SL}(2)/T$ as shown in the following proposition.

Proposition 9.1. *Suppose that $S \cap \Sigma \neq \emptyset$. Then the multiplication is degenerate.*

Proof. Let $\alpha \in S \cap \Sigma$ and suppose that the multiplication is non-degenerate. Consider the rank one wonderful subvariety M' defined by intersecting all the G -stable prime divisors M^σ with $\sigma \in \Sigma \setminus \{\alpha\}$, denote Δ' its set of colors. Then M' is the quotient of a parabolic induction of a wonderful variety for $\mathrm{SL}(2)$ isomorphic to $\mathbb{P}^1 \times \mathbb{P}^1$, whose generic stabilizer is a maximal torus of $\mathrm{SL}(2)$ and whose set of colors is identified with $\Delta'(\alpha) = \{D \in \Delta' : P_\alpha D \neq D\}$, where P_α denotes the minimal parabolic associated to α . It follows then by Corollary 1.7 that, for every $D' \in \mathbb{N}[\Delta' \setminus \Delta'(\alpha)]$, $V_{D'}V_{E'} = V_{D'+E'}$ for all $E' \in \mathbb{N}\Delta'$.

Let $D', E', F' \in \mathbb{N}\Delta'$ be such that $D' \leq_{\Sigma'} E' + F'$ and $V(\omega_{D'}) \subset V(\omega_{E'}) \otimes V(\omega_{F'})$. Denote $\rho: \mathrm{Pic}(M) \rightarrow \mathrm{Pic}(M')$ the restriction. From the combinatorics of colors and spherical roots it follows that there exist $E'_0, F'_0 \in \mathbb{N}[\Delta' \setminus \Delta'(\alpha)]$ and $E, F \in \mathbb{N}\Delta$ such that $E' + E'_0 = \rho(E)$ and $F' + F'_0 = \rho(F)$. Let $D = E + F - (E' + F' - D')$ and notice that $D \in \mathbb{N}\Delta$ and $D' + E'_0 + F'_0 = \rho(D)$. Moreover, the inclusion $V(\omega_{D'}) \subset V(\omega_{E'}) \otimes V(\omega_{F'})$ implies that $V(\omega_D) \subset V(\omega_E) \otimes V(\omega_F)$, hence the non-degeneracy of the multiplication of M implies that $V_D \subset V_E V_F$ and it follows $V_{D'+E'_0+F'_0} \subset V_{E'+E'_0} V_{F'+F'_0}$. On the other hand, by Corollary 1.7, $V_{D'+E'_0+F'_0} \subset V_{E'+E'_0} V_{F'+F'_0} = V_{E'} V_{F'} V_{E'_0+F'_0}$ if and only if $V_{D'} \subset V_{E'} V_{F'}$.

Therefore, we have deduced that the multiplication of M' is non-degenerate, but this is a contradiction by the previous example together with Proposition 1.6. \square

Remark 2. More generally, if M is a wonderful variety whose multiplication is non-degenerate and if $M' \subset M$ is a localization of M , notice that the multiplication of M' is non-degenerate as well. If indeed $\rho: \mathrm{Pic}(M) \rightarrow \mathrm{Pic}(M')$ is the restriction of line bundles and if $D, E \in \mathbb{N}\Delta$, then $\rho(V_D V_E) = V_{\rho(D)} V_{\rho(E)}$, and if $V_F \subset V_D V_E$, then $\rho(V_F) \neq 0$ if and only if $\mathrm{supp}_{\Sigma'}(D + E - F) \subset \Sigma'$ (where $\Sigma' \subset \Sigma$ are the spherical roots of M'). On the other hand, given $E', F' \in \mathbb{N}\Delta'$, by making use of Corollary 1.7, one can reduce the description of the product $V_{E'} V_{F'}$ to the one corresponding to a pair $E'', F'' \in \rho(\mathbb{N}\Delta)$ (where Δ' is the set of colors of M').

As well, if $M \rightarrow M'$ is a quotient of M by a distinguished set of colors and if $H \subset H'$ are the corresponding spherical subgroups, by Corollary 1.4 it follows that the multiplication of M' is non-degenerate if that of M is non-degenerate. Indeed, if we identify the set of colors Δ' of M' with a subset of Δ , then $V_E V_F \subset \mathbb{C}[G]^{(H')} \subset \mathbb{C}[G]^{(H)}$ for all $E, F \in \mathbb{N}\Delta'$.

By using Proposition 1.6, similar statements are easily proved for parabolic inductions and localizations at a parabolic subgroup (see e.g. [8, 1.1.4] for the definition of the latter). Therefore the non-degeneracy property of the multiplication of a wonderful variety is stable under the standard operations of localization, quotient by a distinguished subset of colors, parabolic induction, localization at a parabolic subgroup.

Given a spherical subgroup $H \subset G$, define its *spherical closure* \overline{H} as the kernel of the action of the normalizer $N_G(H)$ on the set of colors of G/H . If H is equal to its spherical closure, then we say that H is *spherically closed*. By a theorem of F. Knop [29, Corollary 7.6], if H is spherically closed, then G/H admits a wonderful compactification. If G is not simply laced, then not spherically closed spherical subgroups H such that G/H admits a wonderful compactification exists. The projection $G/H \rightarrow G/\overline{H}$ canonically identifies the colors of G/\overline{H} with those of G/H .

Generally speaking, if M is the wonderful compactification of G/H where H is not spherically closed, then the multiplication may be degenerate. Indeed, it is easy to show that $\mathbb{C}[G]^{(H)} = \mathbb{C}[G]^{(\overline{H})}$, therefore, if $\overline{\Sigma}$ is the set of spherical roots of G/\overline{H} and if $D \in \mathbb{N}\Delta$ is such that $V_D \subset V_E V_F$, then we must have $D \leq_{\overline{\Sigma}} E + F$. In this way we may construct examples of non-spherically closed spherical subgroups H of G (possessing no simple spherical roots) whose associated multiplication is degenerate.

Example 2. Consider the non-adjoint symmetric wonderful variety M for $\mathrm{Sp}(8)$ with spherical roots $\sigma_1 = \alpha_1 + 2\alpha_2 + \alpha_3, \sigma_2 = \alpha_3 + \alpha_4$. Then M possesses two colors D_2 and D_4 , where $\omega_{D_2} = \omega_2$ and $\omega_{D_4} = \omega_4$. Then we have $D_2 <_{\Sigma} 2D_2$ and $V(\omega_2) \subset V(\omega_2)^{\otimes 2}$, on the other hand $\overline{\Sigma} = \{\sigma_1, 2\sigma_2\}$ and $2D_2 - D_2 = \sigma_1 + \sigma_2 \notin \mathbb{N}\overline{\Sigma}$, therefore it cannot be $V_{D_2} \subset V_{D_2}^2$.

Proposition 9.2. *Let $\mathcal{M} \subset \mathbb{N}\Sigma$ be the semigroup generated by the set of differences $\{E + F - D : V_D \subset V_E V_F\}$. Then $\mathcal{M} = \mathbb{N}\overline{\Sigma}$.*

Proof. As already noticed we have $\mathcal{M} \subset \mathbb{N}\overline{\Sigma}$. Here we show that indeed $\mathcal{M} \supset \overline{\Sigma}$. Since $\mathbb{C}[G]^{(H)} = \mathbb{C}[G]^{(\overline{H})}$ we may assume that H is spherically closed. Let $\sigma \in \Sigma$, proceeding by localization and parabolic induction as in Proposition 9.1 we may assume that M is a cuspidal (i.e., parabolic induction of no other wonderful variety) rank one wonderful variety with spherical root σ (see [44, Table 1] for a complete list of such varieties). With three exceptions (cases 9B, 9C, 15 in [44]) such wonderful varieties are all flag varieties for their automorphism group, so that they are projectively normal. Being H spherically closed, notice that apart from the cases 9B, 7C, 9C, 12, 15 in [44] we have that $\sigma = D + E$ for some $D, E \in \Delta$; then $(0, D, E)$ is a low triple and the projective normality of M implies that $V_0 \subset V_D V_E$, hence $\sigma \in \mathcal{M}$. Moreover, since the case 7C is a quotient of the case 9C, we are reduced to the cases 9B, 7C, 12 and 15, where the following inclusions may be checked directly (we index the colors as the corresponding fundamental weights):

- 9B) G is of type B_r , $\sigma = \alpha_1 + \dots + \alpha_r = D_1$ and we have $V_{D_r} \subset V_{D_1} V_{D_r}$.
- 7C) G is of type C_r , $\sigma = \alpha_1 + 2\alpha_2 + \dots + 2\alpha_{r-1} + \alpha_r = D_2$ and we have $V_{D_2} \subset V_{D_2}^2$.
- 12) G is of type F_4 , $\sigma = \alpha_1 + 2\alpha_2 + 3\alpha_3 + 2\alpha_4 = D_4$ and we have $V_{D_4} \subset V_{D_4}^2$.
- 15) G is of type G_2 , $\sigma = \alpha_1 + \alpha_2 = D_2 - D_1$ and we have $V_{2D_1} \subset V_{D_1} V_{D_2}$.

□

Suppose now that M is a strict wonderful variety and suppose that $E, F \in \mathbb{N}\Delta$ are such that $E + F$ is not faithful. The following example shows that the product $V_E V_F$ may be degenerate, essentially reducing to a not spherically closed case.

Example 3. Let M be the model wonderful variety of type C_4 . Then $D_2 <_{\Sigma} 2D_2$ and $V(\omega_2) \subset V(\omega_2)^{\otimes 2}$ (more precisely, $V(\omega_2)$ is also contained in the second symmetric power of $V(\omega_2)$). Notice that $2D_2$ is not faithful: the maximal distinguished subset of Δ which does not intersect the support of $2D_2$ is $\Delta^* = \{D_1, D_3\}$, and the quotient M/Δ^* is the symmetric wonderful variety M' considered in Example 2, whose colors we still be denoted by D_2 and D_4 . By Corollary 1.4 we may identify $\Gamma(M, \mathcal{L}_{D_2}) = \Gamma(M', \mathcal{L}_{D_2})$ and $\Gamma(M, \mathcal{L}_{2D_2}) = \Gamma(M', \mathcal{L}_{2D_2})$, so that we may regard the product $V_{D_2}^2$ inside $\Gamma(M', \mathcal{L}_{2D_2})$ and by Example 2 it follows that $V_{D_2} \not\subset V_{D_2}^2$.

Question. Let M be a strict wonderful variety and let $E, F \in \mathbb{N}\Delta$ be such that $E + F$ is a faithful divisor. Is the product $V_E V_F$ non-degenerate?

Suppose that G is simply laced: then the class of the strict wonderful varieties is stable with respect to the operation of quotient by a distinguished set of colors, so phenomena as that in Example 3 cannot happen. Therefore, if the answer to the previous question were affirmative, proceeding by induction on the rank of M , it would follow that the multiplication is non-degenerate whenever M is a strict wonderful variety for a simply laced group.

ACKNOWLEDGMENTS

We would like to thank Domingo Luna, who suggested some years ago that we study the normality of spherical nilpotent orbit closures via the projective normality of wonderful varieties.

The paper was partially written during a stay of the second author at Friedrich-Alexander-Universität Erlangen-Nürnberg between March and October 2012, partially supported by a DAAD fellowship. He is grateful to the Emmy Noether Zentrum and especially to Friedrich Knop for their warm hospitality. Also, many thanks are due to Friedrich Knop for fruitful discussions.

REFERENCES

- [1] J. Adams, J.-S. Huang, and D. A. Vogan Jr., *Functions on the model orbit in E_8* , Represent. Theory **2** (1998), 224–263 (electronic), DOI 10.1090/S1088-4165-98-00048-X. MR1628031 (99g:20077)
- [2] D. Ahiezer, *Equivariant completions of homogeneous algebraic varieties by homogeneous divisors*, Ann. Global Anal. Geom. **1** (1983), no. 1, 49–78, DOI 10.1007/BF02329739. MR739893 (85j:32052)
- [3] D. Barbasch and D. A. Vogan Jr., *Unipotent representations of complex semisimple groups*, Ann. of Math. (2) **121** (1985), no. 1, 41–110, DOI 10.2307/1971193. MR782556 (86i:22031)
- [4] I. N. Bernšteĭn, I. M. Gel’fand, and S. I. Gel’fand, *Models of representations of compact Lie groups* (Russian), Funkcional. Anal. i Priložen. **9** (1975), no. 4, 61–62. MR0414792 (54 #2884)
- [5] P. Bravi, *Primitive spherical systems*, Trans. Amer. Math. Soc. **365** (2013), no. 1, 361–407, DOI 10.1090/S0002-9947-2012-05621-2. MR2984062
- [6] P. Bravi and S. Cupit-Foutou, *Classification of strict wonderful varieties* (English, with English and French summaries), Ann. Inst. Fourier (Grenoble) **60** (2010), no. 2, 641–681. MR2667789 (2011m:14085)
- [7] P. Bravi, J. Gandini, A. Maffei, and A. Ruzzi, *Normality and non-normality of group compactifications in simple projective spaces* (English, with English and French summaries), Ann. Inst. Fourier (Grenoble) **61** (2011), no. 6, 2435–2461 (2012), DOI 10.5802/aif.2679. MR2976317
- [8] P. Bravi and D. Luna, *An introduction to wonderful varieties with many examples of type F_4* , J. Algebra **329** (2011), 4–51, DOI 10.1016/j.jalgebra.2010.01.025. MR2769314 (2012f:14102)
- [9] P. Bravi and G. Pezzini, *Wonderful varieties of type D* , Represent. Theory **9** (2005), 578–637 (electronic), DOI 10.1090/S1088-4165-05-00260-8. MR2183057 (2006g:14078)
- [10] M. Brion, *Groupe de Picard et nombres caractéristiques des variétés sphériques* (French), Duke Math. J. **58** (1989), no. 2, 397–424, DOI 10.1215/S0012-7094-89-05818-3. MR1016427 (90i:14048)
- [11] M. Brion, *The total coordinate ring of a wonderful variety*, J. Algebra **313** (2007), no. 1, 61–99, DOI 10.1016/j.jalgebra.2006.12.022. MR2326138 (2008d:14067)
- [12] A. Broer, *Decomposition varieties in semisimple Lie algebras*, Canad. J. Math. **50** (1998), no. 5, 929–971, DOI 10.4153/CJM-1998-048-6. MR1650954 (99k:14077)
- [13] R. Chirivì, C. de Concini, and A. Maffei, *On normality of cones over symmetric varieties*, Tohoku Math. J. (2) **58** (2006), no. 4, 599–616. MR2297202 (2008g:14087)

- [14] R. Chirivì, P. Littelmann, and A. Maffei, *Equations defining symmetric varieties and affine Grassmannians*, Int. Math. Res. Not. IMRN **2** (2009), 291–347, DOI 10.1093/imrn/rnn132. MR2482117 (2010b:14104)
- [15] R. Chirivì and A. Maffei, *Projective normality of complete symmetric varieties*, Duke Math. J. **122** (2004), no. 1, 93–123, DOI 10.1215/S0012-7094-04-12213-4. MR2046808 (2005b:14085)
- [16] R. Chirivì and A. Maffei, *Plücker relations and spherical varieties: application to model varieties*, Transform. Groups **19** (2014), no. 4, 979–997, DOI 10.1007/s00031-014-9285-6. MR3278858
- [17] D. H. Collingwood and W. M. McGovern, *Nilpotent orbits in semisimple Lie algebras*, Van Nostrand Reinhold Mathematics Series, Van Nostrand Reinhold Co., New York, 1993. MR1251060 (94j:17001)
- [18] M. Costantini, *On the coordinate ring of spherical conjugacy classes*, Math. Z. **264** (2010), no. 2, 327–359, DOI 10.1007/s00209-008-0468-5. MR2574980 (2010k:20071)
- [19] C. De Concini, *Normality and non normality of certain semigroups and orbit closures*, Algebraic transformation groups and algebraic varieties, Encyclopaedia Math. Sci., vol. 132, Springer, Berlin, 2004, pp. 15–35, DOI 10.1007/978-3-662-05652-3_3. MR2090668 (2005g:20068)
- [20] C. De Concini and C. Procesi, *Complete symmetric varieties*, Invariant theory (Montecatini, 1982), Lecture Notes in Math., vol. 996, Springer, Berlin, 1983, pp. 1–44, DOI 10.1007/BFb0063234. MR718125 (85e:14070)
- [21] W. A. de Graaf, *Constructing representations of split semisimple Lie algebras*, J. Pure Appl. Algebra **164** (2001), no. 1-2, 87–107, DOI 10.1016/S0022-4049(00)00150-X. Effective methods in algebraic geometry (Bath, 2000). MR1854331 (2002h:17007)
- [22] J. Gandini, *Spherical orbit closures in simple projective spaces and their normalizations*, Transform. Groups **16** (2011), no. 1, 109–136, DOI 10.1007/s00031-011-9120-2. MR2785497 (2012f:14091)
- [23] The GAP Group, Aachen, St Andrews, *GAP—Groups, Algorithms, and Programming - a System for Computational Discrete Algebra*, Version 4.5.4, 2012. <http://www-gap.dcs.st-and.ac.uk/~gap>
- [24] I. M. Gel'fand and A. V. Zelevinskii, *Models of representations of classical groups and their hidden symmetries* (Russian), Funktsional. Anal. i Prilozhen. **18** (1984), no. 3, 14–31. MR757246 (86i:22024)
- [25] I. M. Gel'fand and A. V. Zelevinsky, *Representation models for classical groups and their higher symmetries*, Astérisque **Numero Hors Serie** (1985), 117–128. The mathematical heritage of Élie Cartan (Lyon, 1984). MR837197 (88b:22022)
- [26] W. Hesselink, *The normality of closures of orbits in a Lie algebra*, Comment. Math. Helv. **54** (1979), no. 1, 105–110, DOI 10.1007/BF02566258. MR522033 (80g:14018)
- [27] S. S. Kannan, *Projective normality of the wonderful compactification of semisimple adjoint groups*, Math. Z. **239** (2002), no. 4, 673–682, DOI 10.1007/s002090100319. MR1902056 (2003f:20076)
- [28] M. Kapovich and J. J. Millson, *Structure of the tensor product semigroup*, Asian J. Math. **10** (2006), no. 3, 493–539, DOI 10.4310/AJM.2006.v10.n3.a2. MR2253157 (2007h:22009)
- [29] F. Knop, *Automorphisms, root systems, and compactifications of homogeneous varieties*, J. Amer. Math. Soc. **9** (1996), no. 1, 153–174, DOI 10.1090/S0894-0347-96-00179-8. MR1311823 (96c:14037)
- [30] F. Knop, H. Kraft, and T. Vust, *The Picard group of a G-variety*, Algebraische Transformationsgruppen und Invariantentheorie, DMV Sem., vol. 13, Birkhäuser, Basel, 1989, pp. 77–87. MR1044586
- [31] A. Knutson and T. Tao, *The honeycomb model of $GL_n(\mathbf{C})$ tensor products. I. Proof of the saturation conjecture*, J. Amer. Math. Soc. **12** (1999), no. 4, 1055–1090, DOI 10.1090/S0894-0347-99-00299-4. MR1671451 (2000c:20066)
- [32] B. Kostant and S. Rallis, *Orbits and representations associated with symmetric spaces*, Amer. J. Math. **93** (1971), 753–809. MR0311837 (47 #399)
- [33] S. Kumar, *Tensor product decomposition*, Proceedings of the International Congress of Mathematicians. Volume III, Hindustan Book Agency, New Delhi, 2010, pp. 1226–1261. MR2827839
- [34] D. Luna, *Toute variété magnifique est sphérique* (French, with English summary), Transform. Groups **1** (1996), no. 3, 249–258, DOI 10.1007/BF02549208. MR1417712 (97h:14066)

- [35] D. Luna, *Variétés sphériques de type A* (French), Publ. Math. Inst. Hautes Études Sci. **94** (2001), 161–226, DOI 10.1007/s10240-001-8194-0. MR1896179 (2003f:14056)
- [36] D. Luna, *La variété magnifique modèle* (French, with English and French summaries), J. Algebra **313** (2007), no. 1, 292–319, DOI 10.1016/j.jalgebra.2006.10.042. MR2326148 (2008e:14070)
- [37] A. Maffei, *Orbits in degenerate compactifications of symmetric varieties*, Transform. Groups **14** (2009), no. 1, 183–194, DOI 10.1007/s00031-008-9040-y. MR2480858 (2010f:14055)
- [38] D. I. Panyushev, *Some amazing properties of spherical nilpotent orbits*, Math. Z. **245** (2003), no. 3, 557–580, DOI 10.1007/s00209-003-0555-6. MR2021571 (2004j:14051)
- [39] G. Pezzini, *Simple immersions of wonderful varieties*, Math. Z. **255** (2007), no. 4, 793–812, DOI 10.1007/s00209-006-0050-y. MR2274535 (2007j:14069)
- [40] R. W. Richardson, *On orbits of algebraic groups and Lie groups*, Bull. Austral. Math. Soc. **25** (1982), no. 1, 1–28, DOI 10.1017/S0004972700005013. MR651417 (83i:14041)
- [41] J. R. Stembridge, *The partial order of dominant weights*, Adv. Math. **136** (1998), no. 2, 340–364, DOI 10.1006/aima.1998.1736. MR1626860 (2000a:06038)
- [42] D. A. Timashev, *Homogeneous spaces and equivariant embeddings*, Encyclopaedia of Mathematical Sciences, vol. 138, Springer, Heidelberg, 2011. Invariant Theory and Algebraic Transformation Groups, 8. MR2797018 (2012e:14100)
- [43] D. A. Vogan Jr., *The method of coadjoint orbits for real reductive groups*, Representation theory of Lie groups (Park City, UT, 1998), IAS/Park City Math. Ser., vol. 8, Amer. Math. Soc., Providence, RI, 2000, pp. 179–238. MR1737729 (2001k:22027)
- [44] B. Wasserman, *Wonderful varieties of rank two*, Transform. Groups **1** (1996), no. 4, 375–403, DOI 10.1007/BF02549213. MR1424449 (97k:14051)

DIPARTIMENTO DI MATEMATICA, SAPIENZA UNIVERSITÀ DI ROMA, PIAZZALE ALDO MORO 5,
00185 ROMA, ITALY

E-mail address: `bravi@mat.uniroma1.it`

SCUOLA NORMALE SUPERIORE, PIAZZA DEI CAVALIERI 7, 56126 PISA, ITALY

E-mail address: `jacopo.gandini@sns.it`

DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI PISA, LARGO BRUNO PONTECORVO 5, 56127
PISA, ITALY

E-mail address: `maffei@dm.unipi.it`