

Hsp60 expression in skeletal muscle increases after endurance training

Barone Rosario¹, Macaluso Filippo¹, Sangiorgi Claudia¹, Costa Alessandra², Moresi Viviana², Coletti Dario^{2,3}, Adamo Sergio³, Francesco Cappello¹, Farina Felicia¹, Zummo Giovanni¹, Di Felice Valentina¹

¹ Dipartimento di Biomedicina Sperimentale e Neuroscienze Cliniche, Università degli Studi di Palermo, Italy

² Dipartimento di Scienze Anatomiche, Istologiche, Medico-legali e dell'Apparato Locomotore, SAPIENZA Università di Roma, Italy

³ Laboratory of Genetics and Physio-pathology of Muscle Tissues, UR4 - Ageing, Stress, Inflammation, University Pierre et Marie Curie Paris 6, France

Adult skeletal muscle is remarkably plastic. Increased contractile activity, such as endurance exercise, elicits multiple signals to activate a large set of genes, leading to phenotypic changes in skeletal muscle, including IIB-to-IIa fiber type switching, enhanced mitochondrial biogenesis, and angiogenesis, to match physiologic capability to functional demand.

Heat shock protein (Hsp) 60 is a mitochondrial protein which plays a key role in the translocation of proteins from the cytoplasm to mitochondria. Apart from its mitochondrial localization, Hsp60 has been detected in the cytoplasm, in the cellular membrane and inside exosomes [1]. In the skeletal muscle the expression of Hsp60 is fibre type specific, being expressed more in type IIa and I fibers compared to IIx and IIb. Upon endurance training Hsp60 increases particularly in type I fibers (unpublished data).

To investigate the role of Hsp60 in skeletal muscle plasticity and mitochondrial biogenesis, we used three experimental models: 1) Forty-eight trained young healthy male mice; 2) in vitro C2C12, where Hsp60 was over expressed by plasmids or repressed by siRNAs; 3) in vivo transfected muscles where Hsp60 was over expressed by plasmids and inhibited by siRNAs.

References

- [1] Campanella et al. (2012) The odyssey of Hsp60 from tumor cells to other destinations includes plasma membrane-associated stages and Golgi and exosomal protein-trafficking modalities. *PLoS One* 7: e42008.

Keywords

Heat shock protein 60, skeletal muscle, endurance exercise, mitochondrial biogenesis.

This work was supported by MIUR - PRIN 2012 grant.