

Endocrine
DOI 10.1007/s12020-014-0479-4

ORIGINAL ARTICLE

Risk factors of overweight and obesity among preschool children with different ethnic background

Stefania Toselli · Luciana Zaccagni ·
Francesca Celenza · Augusta Albertini ·
Emanuela Gualdi-Russo

Received: 9 September 2014 / Accepted: 9 November 2014
© Springer Science+Business Media New York 2014

Abstract In this study, we evaluated the risk factors associated with overweight and obesity in 2,640 preschool children in Italy taking into account the ethnic background of the parents. Height and weight were measured and BMI was calculated. Personal and lifestyle data for the children (birth weight, type of breastfeeding, sleep duration, skipping breakfast, snacking, physical activity) and parents (ethnicity, educational level, occupation, weight, height) were collected by means of a questionnaire. Italian and Other European children generally showed the highest percentage of normal weight, while the other ethnic groups presented a greater imbalance. Overweight and obesity were high in African males, who also presented high birth weight. Breastfeeding was most common, although formula feeding was significantly higher in Italians than in immigrants. Immigrants, particularly males, tended to skip breakfast more than Italians. Physical activity was significantly higher in Italians than in immigrants. In the parents,

underweight was particularly high in Italian and Other mothers. African parents had high rates of overweight and obesity and a low educational level. The most common profession was worker for the fathers and housewife for the mothers, with the exception of Italians in which clerical work prevailed. Multiple linear regression analysis showed that the BMI of children was closely related to the BMI of the parents and the birth weight. Hence, these are the most informative parameters in preventing obesity.

Keywords Anthropometry · Nutritional status · Childhood obesity · Immigrants · Preschool children · Ethnicity

Introduction

In the past 25 years, the prevalence of overweight and obese children quadrupled, from roughly four to sixteen percent [1]. It is important to document both the prevalence of overweight and the timing of weight gain in children because those who experience early onset weight gain are more likely to be heavier in adulthood than children with later onset [2]. Overweight children are more likely to become obese adults [1, 3].

The worldwide increase in childhood obesity can be attributed to various factors. The prevalence of overweight and obesity is associated with the parents' BMI: children with weight disorders are more frequently observed in obese families [4, 5]. The relationship between birth weight and obesity has often been called into question in this regard, and previous investigations have produced conflicting results: low birth weight (LBW, i.e., birth weight <2,500 g) or high birth weight (HBW, i.e., birth weight >4,000 g) are both at increased risk [6–8]. The positive

S. Toselli
Department of Biomedical and Neuromotor Science, University of Bologna, Via Selmi 3, 40121 Bologna, Italy
e-mail: stefania.toselli@unibo.it

L. Zaccagni (✉) · E. Gualdi-Russo
Department of Biomedical and Specialty Surgical Sciences, University of Ferrara, Corso Ercole I d'Este 32, 44121 Ferrara, Italy
e-mail: luciana.zaccagni@unife.it

E. Gualdi-Russo
e-mail: emanuela.gualdi@unife.it

F. Celenza · A. Albertini
Public Health Service, Bologna, Italy
e-mail: f.celenza@ausl.bologna.it

A. Albertini
e-mail: albertiniaa@libero.it

association of HBW with childhood obesity has been confirmed in some studies [9, 10], but others have found that LBW is not associated with overweight/obesity in children [11, 12]. Another factor associated with increased risk of obesity is short sleep duration: children who slept for 8 h or less were 2.2 times more likely to be overweight/obese [4, 13, 14].

However, the main causes of obesity seem to be decreased physical activity, increased media use, and unfavorable changes in eating habits [15].

Several systematic reviews and observational studies have suggested an effect of breastfeeding on the risk of childhood overweight/obesity and adiposity [16–25]. Many prospective studies have found a positive association between poor childhood dietary habits and body weight in later life [26, 27]. A high BMI has been associated with an increased intake of high-calorie foods such as high-energy snacks, sweets, and sugary beverages [28, 29]. The risk of being overweight or obese is also higher among children who skip breakfast [30]. Watching television and/or eating and snacking during TV time can contribute to overweight and obesity through an increase in caloric intake (decrease in satiety and/or influence of advertized foods on food choices), inactivity, and possibly a decrease in physical activity [25, 31–33].

The socioeconomic and immigrant status also affects the onset of obesity [34]. The prevalence of overweight and obesity is increased in children of immigrant parents and/or of parents with a low educational level (EL) [35, 36]. Children of recently arrived immigrant parents, particularly boys, tend to weight more and are at greater risk of overweight than native children [37–39]. Conversely, a recent study on children living in Italy [40] found that the BMI was lower in children of immigrant mothers than in those of Italian mothers (significantly lower in daughters).

Accordingly, some unhealthy eating habits and sedentary behaviors were more frequently observed in children of immigrants [35, 41]. However, it remains unclear if parental migrant status (MS) and EL contribute independently to such habits and if this relationship is only observed for some selected eating habits related to childhood obesity. Further gaps include the paucity of data in young children and possible differences according to maternal or paternal MS and EL.

Findings such as these provide insights about the kinds of social and economic contexts that might raise or lower the risk of overweight among children [42].

However, it is unclear at what age children become at-risk for overweight or when the at-risk groups become overweight. For example, if at-risk children tend to be overweight at kindergarten, interventions should focus on the preschool environments (families and childcare facilities) rather than on elementary schools [1].

It is now argued that the prevention of weight gain beginning in the formative preschool years is the most effective means of achieving healthy weight in the population [7, 43]. Hence, a better understanding of the risk factors for obesity during the preschool years is urgently needed to inform prevention strategies.

Our objectives were to assess the differences in adiposity, lifestyle habits and parents' features in preschool children according to their parental MS and to evaluate the influence of behavioral/lifestyle factors on the children's BMI.

Materials and methods

This cross-sectional study was conducted on 2,640 children aged 5–6 years (1,350 males and 1,290 females) attending kindergartens in Emilia-Romagna region (northern Italy) and on 4,874 parents (2,396 fathers and 2,478 mothers). Only children who received parental written consent and agreed to participate were allowed to take part in the study. The response rate was 74.4 %; detailed sampling methodology has been described in a previous publication [44].

The present survey was part of a larger research project (SoNIA—Sorveglianza Nutrizionale Infanzia e Adolescenza) on the nutritional status of children and adolescents [45]. The study was approved by the Italian Ministry of Health and was scientifically supported by the National Institute of Nutrition.

The children's MS was defined on the basis of the origin of their parents [27, 46]. They were classified as Italians ($N = 2,337$) or Immigrants ($N = 263$), with the latter subdivided into the following groups: Other Europeans (45.2 %), Africans (27.4 %), and Others (27.4 %).

We measured height and body weight of the children with a stadiometer and an electronic scale (Seca, Basel, Switzerland) in order to calculate BMI (weight/height²) (kg/m²). During the measurements, the children were dressed in light clothes without shoes. No adjustments were made for clothing. Weight status was defined according to the BMI cutoff values by sex and age established by the International Obesity Task Force (IOTF) in 2000 [47] and by Cole et al. [48].

The parents completed at home a self-administered questionnaire to provide data on the children (birth weight, breastfeeding, sleep duration, breakfast, after dinner snacking, physical activity) and on the family (ethnic group, parents' education, parents' work, weight, height). BMI was calculated from the parents' self-reported height and weight and categorized as underweight (<18.5), normal weight (18.5–24.9), overweight (25–29.9), and obese (≥ 30) [49, 50].

The differences between the Italian and Immigrant children were analyzed by considering the latter group as a whole and divided into the ethnic groups.

To compare the groups, we used Student's *t* test, ANOVA (with Tukey's HSD post-hoc test), and Chi-square (χ^2) test (with Fisher's exact test and Bonferroni correction pairwise comparison when appropriate). A stepwise regression analysis used the children's BMI as the dependent variable and the questionnaire data as independent variables. The probability level was set at 0.05. Statistical analyses were performed using the Package "Statistica" version 8.8 (StatSoft, Vigonza, PD).

Results

Children

Table 1 reports the physical characteristics of the children grouped by sex and origin. There were no significant differences between the Italian and Immigrant children as a whole.

The comparison among children divided into their belonging ethnic groups showed that African males were heavier and taller than children of the other groups. Significant sex differences (*t* test) were observed in Italians for height ($p < 0.001$) and in Africans for weight ($p < 0.05$).

Table 2 presents the prevalence (%) of weight status in the children grouped by sex and origin. Normal weight was most common for both sexes in all ethnic groups. African males had a significantly lower prevalence of normal weight than Italian and Other European males, as well as the highest frequency of overweight and obesity. The Italians showed a significant sex difference in the prevalence of normal weight and overweight children.

The ethnic groups did not differ significantly in birth weight. Males generally weighed more at birth than females (Table 3), with a significant difference in Italians ($p < 0.001$).

Sleep duration (Table 3) was lower in African males than in the other groups, with significant differences in

comparison with Italians ($p < 0.05$) and Other Europeans ($p < 0.001$).

Physical activity (Table 4) was significantly higher in Italian children than in Immigrant children. In all groups, it was generally more frequent in females than in males, with a significant difference in Italians. In those who practiced physical activity, the weekly hours dedicated to it (Table 3) did not differ significantly between males and females.

Breastfeeding (Table 4) was more frequent than formula feeding in all groups. Italian children had the highest percentage of formula feeding (20.9 %), differing significantly from Immigrants and Africans.

Daily breakfast ("every day") (Table 4) was most common in Italians and Other Europeans, with significant differences between Italians and Immigrants of both sexes and between Italian and African females. The "never" response also differed significantly between Italians and Africans and between Italian males and those of the Immigrant and Others groups.

Regarding snacking (Table 4), the percentages of children who never or sometimes consumed food after dinner were generally high. Nevertheless, in males, "never" snacking after dinner was significantly higher in Immigrants than in Italians, while, in females, "sometimes" snacking after dinner was significantly higher in Africans than in Others.

Parents

ANOVA revealed significant differences in mother's BMI by origin ($p < 0.001$). Tukey's HSD test showed that African mothers differed significantly from Italian or Other European ones (Table 5).

In the Immigrant groups, the mothers were most commonly normal weight, while overweight was generally prevalent in the fathers (Table 6). Mothers showed a higher prevalence of underweight than fathers. Underweight was not present in African parents.

Table 1 Anthropometric characteristics of the children (mean, SD) by sex and ethnic background

	Italians (M 1,185, F 1,152)		Immigrants (M 125, F 138)		Other Europeans (M 63, F 56)		Africans (M 36, F 36)		Others (M 26, F 46)	
	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD
Males										
Weight (kg)	22.7	4.3	22.8	4.3	22.2	3.8	24.5	4.9	22.1	4.0
Height (cm)	117.1	5.3	117.3	5.4	116.9	4.9	118.4	6.0	116.7	5.5
BMI (kg/m ²)	16.5	2.3	16.5	2.3	16.2	2.0	17.4	2.8	16.1	2.1
Females										
Weight (kg)	22.4	4.1	22.4	4.1	22.3	3.8	21.9	4.3	22.7	4.3
Height (cm)	115.9	5.5	116.5	5.3	116.8	4.7	115.8	5.8	116.6	5.7
BMI (kg/m ²)	16.6	2.2	16.4	2.2	16.3	2.1	16.2	2.1	16.6	2.3

Table 2 Prevalence (%) of weight status in children by sex and ethnic background

	1-Italians	2-Immigrants	3-Other Europeans	4-Africans	5-Others
Males					
Underweight	7.5	4.0	1.6	5.5	7.7
Normal weight	70.9 ⁴	68.8	81.0 ⁴	50.0 ^{1,3}	65.4
Overweight	13.0	16.8	11.1	25.0	19.2
Obesity	8.6	10.4	6.3	19.4	7.7
Females					
Underweight	5.4	8.1	5.5	13.9	6.7
Normal weight	65.7	64.0	70.9	58.3	60.0
Overweight	19.7	19.9	16.4	22.2	22.2
Obesity	9.3	8.1	7.3	5.6	11.1

¹ Significant difference with Italians

³ Significant difference with Other Europeans

⁴ Significant difference with Africans

Table 3 Birth weight and lifestyle factors of the children by sex and ethnic background

	1-Italians		2-Immigrants		3-Other Europeans		4-Africans		5-Others	
	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD
Males										
Birth weight (kg)	3.44	0.46	3.42	0.53	3.37	0.55	3.54	0.56	3.40	0.44
Sleep duration (h/day)	9.8 ⁴	0.9	9.6	1.1	9.9 ⁴	1.0	9.1 ^{1,3}	1.1	9.7	1.2
PA (h/w)	2.2	1.1	2.6	1.1	2.7	1.3	2.9	1.1	2.3	0.8
Females										
Birth weight (kg)	3.27	0.48	3.33	0.49	3.31	0.55	3.29	0.48	3.40	0.42
Sleep duration (h/day)	9.8	0.9	9.7	1.0	9.8	0.8	9.6	1.1	9.6	1.2
PA (h/w)	2.1	1.0	2.2	1.0	2.3	0.9	2.1	1.0	2.2	1.1

¹ Significant difference with Italians

³ Significant difference with Other Europeans

⁴ Significant difference with Africans

The prevalence of underweight was significantly higher in male Others than in the other groups. Normal weight parents were more frequent in Italians than in Immigrants; moreover, Italian mothers presented a significantly higher prevalence of normal weight than African ones. African mothers and fathers had high rates of overweight and obesity (Table 6), and the difference between Italian and African mothers was significant. Italian mothers showed significantly lower values of overweight than Immigrant mothers.

The number of children and of family members differed significantly between Italians and Immigrants. ANOVA revealed a significant difference between Italians and Other Europeans in the number of family members. Africans had a slightly higher number of children than the other groups (Table 5).

Middle school and high school diploma were the most common ELs (Table 6). Italian parents showed a significant difference with Immigrants in terms of the total absence of education and elementary school diploma. These groups also differed for middle school diploma in the fathers and for high school diploma in the mothers. The African parents had the lowest level of education, with the highest frequency of individuals with a total absence of

education or only an elementary school diploma; the difference was significant with respect to Italians and Other Europeans (mothers only). African mothers also differed significantly from Italian ones for high school diploma.

Laborer was the most common profession of the fathers and housewife of the mothers, with the exception of Italian parents for whom clerical work was predominant (Table 6). Italian mothers were significantly more engaged in clerical work than Immigrant ones, the difference being significant for all the three subgroups (Other Europeans, Africans, and Others). The Italian mothers showed significantly lower values for housewife than the above-mentioned groups. Merchant activity was significantly higher in Italian mothers than in Immigrant ones. Italian mothers showed significantly lower frequencies of laborer than Immigrant mothers (particularly than Africans), while the opposite situation was found for mothers seeking employment.

Among the fathers, Italians showed significantly higher values of entrepreneur than Immigrants (particularly than Africans). Immigrants presented significantly lower values for clerical worker than Italians, with Africans having a significantly lower percentage than all the other groups. African employees were significantly lower than all the

Table 4 Prevalence (%) of lifestyle factors of the children by sex and ethnic background

	1-Italians	2-Immigrants	3-Other Europeans	4-Africans	5-Others
<i>Types of feeding</i>					
Breastfeeding	54.6	61.1	57.0	66.7	62.7
Formula feeding	20.9 ^{2,4}	14.8 ¹	18.4	7.9 ¹	14.9
Mixed feeding	24.5	24.2	24.6	25.4	22.4
<i>Breakfast</i>					
Males					
Never	0.3 ^{2,4,5}	1.7 ¹	0.0	3.1 ¹	3.8 ¹
Sometimes	4.5	6.6	7.9	3.1	7.7
Almost every day	10.5	14.9	11.1	18.8	19.2
Every day	84.7 ²	76.9 ¹	81.0	75.0	69.2
Females					
Never	0.3 ⁴	0.7	0.0	2.9 ¹	0.0
Sometimes	6.1	10.3	7.3	14.3	10.9
Almost every day	10.3	13.2	10.9	20.0	10.9
Every day	83.3 ^{2,4}	75.7 ¹	81.8	62.9 ¹	78.3
<i>After dinner snacking</i>					
Males					
Never	26.9 ²	38.2 ¹	32.1	44.8	44.0
Sometimes	42.3	39.1	41.1	34.5	40.0
Almost every day	12.7	7.3	8.9	3.4	8.0
Every day	18.2	15.5	17.9	17.2	8.0
Females					
Never	31.7	36.7	41.2	20.7	42.5
Sometimes	38.6	38.3	37.3	62.1 ⁵	22.5 ⁴
Almost every day	11.8	9.2	13.7	3.4	7.5
Every day	17.9	15.8	7.8	13.8	27.5
<i>Physical activity</i>					
Males					
	38.6 ²	29.0 ¹	29.0	19.4	42.3
Females					
	50.2 ²	40.6 ¹	46.4	30.6	41.3

¹ Significant difference with Italians

² Significant difference with Immigrants

⁴ Significant difference with Africans

⁵ Significant difference with Others

Table 5 Anthropometric characteristics of the parents and family composition by ethnic background

	1-Italians		2-Immigrants		3-Other Europeans		4-Africans		5-Others	
	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD
Mother's height (cm)	163.9	5.8	164.5	6.2	165.1	6.0	165.4 ⁵	6.3	162.7 ⁴	6.1
Mother's weight (kg)	61.2 ⁴	10.6	63.8	10.6	62.1 ⁴	10.3	68.2 ^{1,3,5}	10.8	62.1 ⁴	9.6
Mother's BMI (kg/m ²)	22.8 ⁴	3.6	23.6	3.7	22.8 ⁴	3.6	24.9 ^{1,3}	3.6	23.5	3.7
Father's height (cm)	176.6 ⁴	6.5	175.2	7.6	177.3 ^{4,5}	6.5	173.3 ^{1,3}	7.8	173.6 ³	8.3
Father's weight (kg)	79.9	11.7	79.4	10.8	80.6	11.2	77.9	9.8	78.7	11.0
Father's BMI (kg/m ²)	25.6	3.3	25.9	3.2	25.6	3.2	26.1	3.0	26.1	3.3
Number of children	1.9 ²	0.8	2.1 ¹	0.8	2.0	0.8	2.2	0.9	2.0	0.7
Number of family members	3.9 ^{2,3}	0.9	4.3 ¹	1.1	4.4 ¹	1.1	4.3	1.2	4.0	0.8

¹ Significant difference with Italians

² Significant difference with Immigrants

³ Significant difference with Other Europeans

⁴ Significant difference with Africans

⁵ Significant difference with Others

Table 6 Prevalence of parents' characteristics by ethnic background

	1-Italians	2-Immigrants	3-Other Europeans	4-Africans	5-Others
Mother's weight status					
Underweight	5.5	2.9	3.6	0.0	4.5
Normal weight	74.2 ^{2,4}	67.1 ¹	71.8	59.1 ¹	67.2
Overweight	15.8 ^{2,4}	23.5 ¹	20.9	28.8 ¹	22.4
Obesity	4.5 ⁴	6.6	3.6	12.1 ¹	6.0
Father's weight status					
Underweight	0.2 ⁵	0.8	0.9 ⁵	0.0 ⁵	1.5 ^{1,3,4}
Normal weight	48.4 ²	41.5 ¹	46.3	34.9	40.0
Overweight	42.5	48.3	44.4	52.4	50.8
Obesity	8.9	9.3	8.3	12.7	7.7
Mother's educational level					
None	0.2 ^{2,4}	3.6 ¹	0.0 ⁴	11.6 ^{1,3}	1.4
Primary school	1.5 ^{2,4,5}	9.1 ¹	3.5 ⁴	18.8 ^{1,3}	8.6 ¹
Middle school	32.0	32.9	39.8	30.4	24.3
High school	52.5 ^{2,4}	43.3 ¹	46.0	33.3 ¹	48.6
University	13.9	11.1	10.6	5.8	17.1
Father's educational level					
None	0.2 ^{2,4}	2.8 ¹	0.0 ⁴	9.0 ^{1,3}	1.5
Primary school	3.4 ^{2,4}	9.8 ¹	8.0	14.9 ¹	7.5
Middle school	41.6 ²	34.6 ¹	32.1	35.8	37.3
High school	41.4	39.8	45.5	31.3	38.8
University	13.4	13.0	14.3	9.0	14.9
Mother's occupation					
Entrepreneur	8.0	5.6	6.2	0.0	9.9
Clerical worker	40.4 ^{2,3,4,5}	15.1 ¹	18.6 ¹	7.5 ¹	16.9 ¹
Merchant	8.3 ²	4.4 ¹	6.2	3.0	2.8
Laborer	17.7 ^{2,4}	24.7 ¹	23.9	31.3 ¹	19.7
Housewife	17.4 ^{2,3,4,5}	39.4 ¹	36.3 ¹	38.8 ¹	45.1 ¹
Seeking employment	2.3 ^{2,4}	5.2 ¹	2.7	11.9 ¹	2.8
Other	5.9	5.6	6.2	7.5	2.8
Father's occupation					
Entrepreneur	22.3 ^{2,4}	15.3 ¹	14.9	10.4 ¹	20.6
Clerical worker	28.8 ^{2,4}	16.9 ¹	19.3 ⁴	4.5 ^{1,3,5}	25.0 ⁴
Merchant	16.7	14.9	15.8	11.9	16.2
Laborer	26.2 ^{2,4}	42.2 ¹	36.8 ⁴	64.2 ^{1,3,5}	29.4 ⁴
Household	0.7 ^{2,3,4}	3.6 ¹	3.5 ¹	4.5 ¹	2.9
Seeking employment	0.6	1.6	0.0	3.0	2.9
Other	4.6	5.6	9.6	1.5	2.9

¹ Significant difference with Italians

² Significant difference with Immigrants

³ Significant difference with Other Europeans

⁴ Significant difference with Africans

⁵ significant difference with Others

other groups. The Italian fathers had a lower frequency of laborer than the other groups, with significant differences with Immigrants and particularly Africans. The latter presented the highest frequency, also differing significantly from Other Europeans and Others. There were significantly less household fathers in Italians than in Immigrants (especially Other Europeans and Africans).

The stepwise regression analysis using the children's BMI as the dependent variable and type of breastfeeding,

mother's BMI, father's BMI, skipping breakfast, number of family members, mother's education, father's education, ethnicity, hours of physical activity per week, birth weight, mother's occupation, father's occupation, gender, sleep duration, practice of physical activity, and after dinner snacking as independent variables yielded a model with the variables reported in Table 7. Children's BMI varied according to birth weight and parents' BMI, which are thus the most informative predictors.

Table 7 Regression model of child's BMI ($R^2 = 0.12$)

	Estimate	Std. error	<i>p</i>
(Intercept)	7.92	0.51	0.0002
Mother's BMI	0.15	0.09	<0.0001
Father's BMI	0.12	0.01	<0.0001
Birth weight	0.49	0.01	<0.0001

Discussion

The purposes of this study were to assess differences in BMI and weight status in preschool children of Emilia-Romagna region (Italy) with different ethnic background and to assess the possible risk factors associated with these conditions.

The main limitation of the study is the use of self-reported data for the parents. In fact, some incorrect answers might result from recall bias, for example regarding characteristics of the child at birth, or from possible confusion between the actual and desired physical characteristics with regard to parental height and weight [40]. Another limitation is the small immigrant samples, with a probable reduction in the power of the statistical tests. Finally this is a cross-sectional study.

However, with regard to the children's anthropometric characteristics, our study considered only data measured by specialized personnel with standardized training.

Although the differences were significant only in some cases, the results indicate noteworthy trends. The most pronounced differences among the groups regarded African children, particularly males, with the prevalence of overweight and obesity being remarkably high in this group. Italian and Other European children generally showed the highest percentage of normal weight subjects, while the other ethnic groups presented a greater imbalance.

African children had the highest incidence of risk factors related to obesity: African males had higher birth weight and shorter sleep than the other groups.

There was a significantly higher frequency of Italian children engaged in physical activity than Immigrant children.

There were no notable differences in the type of feeding, even though formula feeding was a more common practice in Italians than in the other groups.

Africans and male Others Immigrants showed a tendency to skip breakfast with respect to Italians and Other European Immigrants.

The weight status of the children partly reflected what was observed in the parents. Indeed, the African parents had a high frequency of overweight and obesity.

Moreover, the data on parental education showed a particularly low EL in Africans. This confirms literature

reports showing that MS, along with a low EL, is a risk factor for overweight/obesity [35, 36, 39].

The stepwise regression analysis revealed three potentially modifiable factors associated with BMI in preschool children. In particular, our study confirmed that HBW is an important predictor of overweight or obesity in children [5, 8]. Infant birth weight is related to the mother's nutrition and body weight during pregnancy, becoming an important indicator of nutritional and developmental status of the child. The other two risk factors with a significant association with children's BMI were the mother's and father's BMI. According to Ebenegger et al. [27], certain maternal factors, such as maternal obesity, are associated with severe kindergarten obesity. In our study, weight disorders were higher among Immigrants, both children and parents. Particularly promising prevention and treatment components for severely obese immigrant children include integration of traditional dietary customs and preferences, family-centered approaches to physical activity and healthy meals [51].

The effects of ethnicity can be partly explained by cultural beliefs and behavioral norms. There may be a need for culturally appropriate health promotion interventions targeted at high-risk minority ethnic groups [52, 53].

As parents are mainly responsible for the purchase and preparation of food, low-income families tend to buy more energy-dense foods since they provide low-cost calories [27, 36, 54].

According to Potter and Ulijazek [55], the prediction of adult obesity from the obesity risk in childhood depends on an ecological multifactor framework that spans multiple phases of the life course. Numerous studies [4, 7, 8, 14, 15, 25, 27, 56] have sought to highlight the factors playing a major role in this relationship. Understanding these factors will help to devise and implement focused preventive measures. Unfortunately, there are still very few studies on immigrant children, especially in countries such as Italy where immigration is a recent, albeit very important, phenomenon. Our study shows that there are different risks for overweight in 5/6-year-old children according to their ethnic background. Monitoring the health status and associated risk factors in these groups and supporting the parents, particularly in the promotion of healthy behaviors, will allow better protection and integration of these immigrants. In order to reduce health inequalities, culturally specific efforts are required to promote a healthy lifestyle among high-risk ethnic groups.

Acknowledgments This study was supported by the Italian Health Ministry and by the European Union Seventh Framework Programme (FP7/2007-2013, Grant 260715). The authors gratefully acknowledge the contribution of the participating parents and children, and public health services.

References

1. K.S. Balistreri, J. Van Hook, Trajectories of overweight among US school children: a focus on social and economic characteristics. *Matern. Child Health J.* **15**, 610–619 (2011)
2. W. Wisemandle, L.M. Maynard, S.S. Guo, R.M. Siervogel, Childhood weight, stature, and body mass index among never overweight, early-onset overweight, and late-onset overweight groups. *Pediatrics* **106**, E14 (2000)
3. R.C. Whitaker, J.A. Wright, M.S. Pepe, K.D. Seidel, W.H. Dietz, Predicting obesity in young adulthood from childhood and parental obesity. *New Engl. J. Med.* **37**, 869–873 (1997)
4. D.A. Dev, B.A. McBride, B.H. Fiese, B.L. Jones, H. Cho, Behalf Of The Strong Kids Research Team, Risk factors for overweight/obesity in preschool children: an ecological approach. *Child. Obes.* **9**, 399–408 (2013)
5. L. Zhou, G. He, J. Zhang, R. Xie, M. Walker, S.W. Wen, Risk factors of obesity in preschool children in an urban area in China. *Eur. J. Pediatr.* **170**, 1401–1406 (2011)
6. P.J. Surkan, C.C. Hseish, A.L.V. Johansson, P.W. Dickman, S. Cnattingius, Reasons for increasing trends in large for gestational age births. *Obstet. Gynecol.* **104**, 720–726 (2004)
7. J. Oldroyd, A. Renzaho, H. Skouteris, Low and high birth weight as risk factors for obesity among 4 to 5-year-old Australian children: does gender matter? *Eur. J. Pediatr.* **170**, 899–906 (2011)
8. N. Li, E. Liu, S. Sun, J. Guo, L. Pan, P. Wang, J. Liu, L. Tan, G. Liu, G. Hu, Birth weight and overweight or obesity risk in children under 3 years in China. *Am. J. Hum. Biol.* **26**(3), 331–336 (2014). doi:10.1002/ajhb.22506
9. S. Brophy, R. Cooksey, M.B. Gravenor, R. Mistry, N. Thomas, R.A. Lyons, R. Williams, Risk factors for childhood obesity at age 5: analysis of the millennium cohort study. *BMC Public Health* **9**, 467 (2009)
10. L. Dubois, M. Girard, Early determinants of overweight at 4.5 years in a population-based longitudinal study. *Int. J. Obes.* **30**, 610–617 (2006)
11. V. Hirschler, J. Bugna, M. Roque, T. Gilligan, C. Gonzalez, Does low birth weight predict obesity/overweight and metabolic syndrome in elementary school children? *Arch. Med. Res.* **39**, 796–802 (2008)
12. X. Zhang, E. Liu, Z. Tian, W. Wang, T. Ye, G. Liu, Y. Li, P. Wang, X. Yang, Z. Yu, G. Hu, High birth weight and overweight or obesity among Chinese children 3–6 years old. *Prev. Med.* **49**, 172–178 (2009)
13. N. Padilla, M.L. Biason, B. Ledésert, Prevalence and risk factors of overweight among 5- to 11-year-old schoolchildren in Narbonne. France. *Sante Publique* **24**, 317–328 (2012)
14. W.W. Wong, C.L. Ortiz, D. Lathan, L.A. Moore, K.L. Konzelmann, A.L. Adolph, E.O. Smith, N.F. Butte, Sleep duration of underserved minority children in a cross-sectional study. *BMC Public Health* **13**, 648 (2013)
15. U. Ekelund, L.B. Sardinha, S.A. Anderssen, M. Harro, P.W. Franks, S. Brage, A.R. Cooper, L.B. Andersen, C. Riddoch, K. Froberg, Associations between objectively assessed physical activity and indicators of body fatness in 9- to 10-y-old European children: a population-based study from 4 distinct regions in Europe (the European Youth Heart Study). *Am. J. Clin. Nutr.* **80**, 584–590 (2004)
16. M.S. Kramer, Do breast-feeding and delayed introduction of solid foods protect against subsequent obesity? *J. Pediatr.* **98**, 883–887 (1981)
17. S. Arenz, R. Ruckerl, B. Koletzko, R. von Kries, Breast-feeding and childhood obesity—a systematic review. *Int. J. Obes. Relat. Metab. Disord.* **28**, 1247–1256 (2004)
18. V. Burke, L.J. Beilin, K. Simmer, W.H. Oddy, K.V. Blake, D. Doherty, G.E. Kendall, J.P. Newnham, L.I. Landau, F.J. Stanley, Breastfeeding and overweight: longitudinal analysis in an Australian birth cohort. *J. Pediatr.* **147**, 56–61 (2005)
19. T. Harder, R. Bergmann, G. Kallischnigg, A. Plagemann, Duration of breastfeeding and risk of overweight: a meta-analysis. *Am. J. Epidemiol.* **162**, 397–403 (2005)
20. C.G. Owen, R.M. Martin, P.H. Whincup, G.D. Smith, D.G. Cook, Effect of infant feeding on the risk of obesity across the life course: a quantitative review of published evidence. *Pediatrics* **115**, 1367–1377 (2005)
21. E.J. Mayer-Davis, S.L. Rifas-Shiman, L. Zhou, F.B. Hu, G.A. Colditz, M.W. Gillman, Breast-feeding and risk for childhood obesity. *Diabetes Care* **29**, 2231–2237 (2006)
22. L. Shields, M. O'Callaghan, G.M. Williams, J.M. Najman, W. Bor, Breastfeeding and obesity at 14 years: a cohort study. *J. Paediatr. Child Health* **42**, 289–296 (2006)
23. M. Weyermann, D. Rothenbacher, H. Brenner, Duration of breastfeeding and risk of overweight in childhood: a prospective birth cohort study from Germany. *Int. J. Obes.* **30**, 1281–1287 (2006)
24. M.W. Gillman, D. Barker, D. Bier, F. Cagampang, J. Challis, C. Fall, K. Godfrey, P. Gluckman, M. Hanson, D. Kuh, P. Nathanielsz, P. Nestel, K.L. Thornburg, Meeting report on the 3rd International Congress on Developmental Origins of Health and Disease (DOHaD). *Pediatr. Res.* **61**, 625–629 (2007)
25. D.D. Bingham, M.I. Varela-Silva, M.M. Ferrão, G. Augusta, M.I. Mourão, H. Nogueira, V.R. Marques, C. Padez, Socio-demographic and behavioral risk factors associated with the high prevalence of overweight and obesity in Portuguese children. *Am. J. Hum. Biol.* **25**, 733–742 (2013)
26. R.A. Forshee, P.A. Anderson, M.L. Storey, Sugar-sweetened beverages and body mass index in children and adolescents: a meta-analysis. *Am. J. Clin. Nutr.* **87**, 1662–1671 (2008)
27. V. Ebenegger, P.M. Marques-Vidal, A. Nydegger, J. Laimbacher, I. Niederer, F. Bürgi, V. Giusti, P. Bodenmann, S. Kriemler, J.J. Puder, Independent contribution of parental migrant status and educational level to adiposity and eating habits in preschool children. *Eur. J. Clin. Nutr.* **65**, 210–218 (2011)
28. T.A. Nicklas, S.J. Yang, T. Baranowski, I. Zakeri, G. Berenson, Eating patterns and obesity in children. The Bogalusa Heart Study. *Am. J. Prev. Med.* **25**, 9–16 (2003)
29. S.A. Bowman, S.L. Gortmaker, C.B. Ebbeling, M.A. Pereira, D.S. Ludwig, Effects of fast-food consumption on energy intake and diet quality among children in a national household survey. *Pediatrics* **113**, 112–118 (2004)
30. W.S. Wolfe, C.C. Campbell, E.A. Frongillo Jr, J.D. Haas, T.A. Melnik, Overweight schoolchildren in New York State: prevalence and characteristics. *Am. J. Public Health* **84**, 807–813 (1994)
31. K.A. Coon, K.L. Tucker, Television and children's consumption patterns. A review of the literature. *Minerva Pediatr.* **54**, 423–436 (2002)
32. E.M. Blass, D.R. Anderson, H.L. Kirkorian, T.A. Pempek, I. Price, M.F. Kolehini, On the road to obesity: television viewing increases intake of high-density foods. *Physiol. Behav.* **88**, 597–604 (2006)
33. L. Dubois, A. Farmer, M. Girard, K. Peterson, Social factors and television use during meals and snacks is associated with higher BMI among pre-school children. *Public Health Nutr.* **11**, 1267–1279 (2008)
34. E. Gualdi-Russo, L. Zaccagni, V.S. Manzon, S. Masotti, N. Rinaldo, M. Khyatti, Obesity and physical activity in children of immigrants. *Eur. J. Public Health* **24**(suppl 1), 40–46 (2014)

35. A.M. Lasserre, A. Chiolero, F. Cachat, F. Paccaud, P. Bovet, Overweight in Swiss children and associations with children's and parents' characteristics. *Obesity* **15**, 2912–2919 (2007)
36. S. Caprio, S.R. Daniels, A. Drewnowski, F.R. Kaufman, L.A. Palinkas, A.L. Rosenbloom, J.B. Schwimmer, Influence of race, ethnicity, and culture on childhood obesity: implications for prevention and treatment: a consensus statement of Shaping America's Health and the Obesity Society. *Diabetes Care* **31**, 2211–2221 (2008)
37. J. Van Hook, K. Balistreri, Immigrant generation, socioeconomic status and economic development of countries of origin: a longitudinal study of body mass index among children. *Soc. Sci. Med.* **65**, 976–989 (2007)
38. K.S. Balistreri, J. Van Hook, Socioeconomic status and body mass index among hispanic children of immigrants and children of natives. *Am. J. Public Health* **99**, 1–8 (2009)
39. S. Toselli, P. Brasili, F. Spiga, Body image, body dissatisfaction and weight status in children from Emilia-Romagna (Italy): comparison between immigrant and native-born. *Ann. Hum. Biol.* **41**, 23–28 (2014)
40. E. Gualdi-Russo, V.S. Manzon, S. Masotti, S. Toselli, A. Albertini, F. Celenza, L. Zaccagni, Weight status and perception of body image in children: the effect of maternal immigrant status. *Nutr. J.* **11**, 85 (2012)
41. S. Sausenthaler, I. Kompauer, A. Mielck, M. Borte, O. Herbarth, B. Schaaf, A. von Berg, J. Heinrich, Impact of parental education and income inequality on children's food intake. *Public Health Nutr.* **10**, 24–33 (2007)
42. T.A. Glass, M.J. McAtee, Behavioral science at the crossroads in public health: extending horizons, envisioning the future. *Soc. Sci. Med.* **62**, 1650–1670 (2006)
43. A.K. Ventura, L.L. Birch, Does parenting affect children's eating and weight status? *Int. J. Behav. Nutr. Phys. Act* **5**, 15 (2008)
44. A. Tripodi, A. Albertini, R. Cecchetti, A. Fabbri, E. Dalle Donne, C. Saletti, M. Mattioli, S. De Giorgi, G. Cavrini, E. Gualdi-Russo, B. Veronesi, A. Rauti, C. Cortesi, E. Di Martino, M. Nicolucci, F. Celenza, S. Toselli, M.T. Beccari, C. Rizzoli, I. Stefanelli, C. Sardo Cardalano, M. Spano, Sorveglianza Nutrizionale. Determinanti e indicatori di rischio obesità nella popolazione infantile. Emilia Romagna. Indagine 2003 e 2005. In: *Prevenzione nei luoghi di vita e di lavoro*. 49. Servizio Sanitario Emilia-Romagna Bologna: Centro Stampa Giunta (2006)
45. E. Gualdi-Russo, A. Albertini, L. Argnani, F. Celenza, M. Nicolucci, S. Toselli, Weight status and body image perception in Italian children. *J. Hum. Nutr. Diet.* **21**, 39–45 (2008)
46. R. Muckelbauer, L. Libuda, K. Clausen, A.M. Toschke, T. Reinehr, M. Kersting, Immigrational background affects the effectiveness of a school-based overweight prevention program promoting water consumption. *Obesity* **18**, 528–534 (2010)
47. T.J. Cole, M.C. Bellizzi, K.M. Flegal, W.H. Dietz, Establishing a standard definition for child overweight and obesity worldwide: international survey. *Br. Med. J.* **320**, 1240–1245 (2000)
48. T.J. Cole, K.M. Flegal, D. Nicholls, A.A. Jackson, Body mass index cut offs to define thinness in children and adolescents: international survey. *Br. Med. J.* **335**, 194–201 (2007)
49. World Health Organization, Physical status: the use and interpretation of anthropometry. Report of a WHO Expert Committee. *World Health Organ. Tech. Rep. Ser.* **854**, 1–452 (1995)
50. National Institute for Health and Clinical Excellence: Obesity: the prevention, identification, assessment and management of overweight and obesity in adults and children. <http://www.ncbi.nlm.nih.gov/books/NBK63696/pdf/TOC.pdf>. Accessed 3 Sep 2014
51. G. Flores, H. Lin, Factors predicting overweight in US kindergartners. *Am. J. Clin. Nutr.* **97**, 1178–1187 (2013)
52. C.L. Falconer, M.H. Park, H. Croker, A.S. Kessel, S. Saxena, R.M. Viner, S. Kinra, Can the relationship between ethnicity and obesity-related behaviours among school-aged children be explained by deprivation? A cross-sectional study. *BMJ Open* **4**(1), e003949 (2014). doi:10.1136/bmjopen-2013-003949
53. S. Toselli, L. Argnani, E. Gualdi-Russo, Growth of Chinese Italian infants in the first 2 years of life. *Ann. Hum. Biol.* **32**, 15–29 (2005)
54. A. Drewnowski, N. Darmon, A. Briend, Replacing fats and sweets with vegetables and fruits—a question of cost. *Am. J. Public Health* **94**, 1555–1559 (2004)
55. C.M. Potter, S.J. Ulijaszek, Predicting adult obesity from measures in earlier life. *J. Epidemiol. Community Health* **67**, 1032–1037 (2013)
56. E. Gualdi-Russo, S. Toselli, S. Masotti, D. Marzouk, K. Sundquist, J. Sundquist, Health, growth and psychosocial adaptation of immigrant children. *European Journal of Public Health* **24** (suppl 1), 26–30 (2014)