



Developing As and Cu Tissue Residue Thresholds to Attain the Good Ecological Status of Rivers in Mining Areas

Iñigo Moreno-Ocio¹ · Leire Méndez-Fernández² · Maite Martínez-Madrid² · Noemí Costas³ · Isabel Pardo³ · Pilar Rodríguez¹

Received: 13 October 2021 / Accepted: 20 January 2022 / Published online: 4 March 2022
© The Author(s) 2022

Abstract

The study was performed on residue-effects datasets from polluted and unpolluted sites in the Nalón River basin (northern Spain). The effects were measured in terms of alteration of field macroinvertebrate communities, and measured as ecological status scores, and number of families and abundance of Ephemeroptera, Plecoptera and Trichoptera (EPT). Non-linear regression models of the field-measured tissue residues in 10 taxa related to the ecological status of the macroinvertebrate communities were used to derive effective tissue residues (ERs). These were estimated for the good/moderate boundary defined by the ecological quality ratio (EQRs) score and for the 50% reduction of EQR and EPT metrics. As, Cu, Hg and Se ERs were calculated for several macroinvertebrate taxa with different feeding styles. The ER dataset allowed us to estimate As and Cu hazardous concentrations (HC), using species sensitivity distribution models, and were interpreted as community thresholds. Further studies for Hg and Se are needed to complete the database required for HC estimation. The reliability and differences of the several thresholds were tested in a risk assessment using a tissue-residue approach (TRA) conducted with field organisms from Cauxa Creek, a tributary from the same basin exposed to high levels of metals in the sediments due to gold mining activities. This risk assessment identified that As and Cu tissue residues satisfactorily explained the reduction in the ecological status of the macroinvertebrate assemblages. Our results indicate that TRA can help in setting future environmental quality standards for the protection of aquatic biota.


✉ Iñigo Moreno-Ocio
inigo.moreno@ehu.eus

¹ Department Zoology and Animal Cell Biology, University of the Basque Country, UPV/EHU, Box 644, 48080 Bilbao, Spain

² Department Genetics, Physical Anthropology and Animal Physiology, University of the Basque Country, UPV/EHU, Bilbao, Spain

³ Department Ecology and Animal Biology, University of Vigo, Vigo, Spain

Graphical Abstract



Areas having a high level of metals and metalloids (hereinafter, metals) due to their lithology have often, historically been exploited for metal extraction. Mining activities usually result in the disposal of soil tailings in the areas adjacent to the mine, which can leach metals to rivers and cause alterations to aquatic communities in the absence of adequate management (Loredo et al. 2010). Monitoring the levels of contaminants in water is a common strategy in European countries, but this may be not enough to achieve the desired level of protection of field communities (EC 2008). For this reason, European policy also considers an important issue to set environmental quality standards in sediment and biota for some priority metals (Cd, Ni, Hg and Pb and their compounds) (EC 2008, Annex II). The objective of this environmental policy includes monitoring the bioaccumulation potential and bioassessment of impacts and trends for certain chemicals, thus ensuring protection against secondary poisoning at the community level. The measurement of metal tissue residues reflects bioavailability, thus reducing uncertainty about the actual bioavailable fraction of chemicals based on external concentrations (Sappington et al. 2011). There is also a general consensus on the convenience of including bioaccumulation data measured in

indigenous organisms for sediment risk assessment (Chapman 2007; Adams et al. 2011) and for the integration of toxicological and ecological information toward regulatory applications using a tissue residue approach (McCarty et al. 2011). This integrative approach is especially suitable for assessing water quality in mining districts, where aquatic macroinvertebrates can accumulate high levels of metals (Cain et al. 2004; Solà et al. 2004; Méndez-Fernández et al. 2015) and can become a significant source of dietary uptake for their predators (Clements 1991; EC 2011). Furthermore, bioaccumulation can also be the cause of alterations in macroinvertebrate assemblages (Luoma et al. 2010; De Jonge et al. 2013; Bervoets et al. 2016).

The measurement of field alterations in the macroinvertebrate assemblage composition and structure is an essential component of the integrated assessment of ecological status, as established by the Water Framework Directive (EC 2000). In that context, the relationships of metal tissue residues in selected biomonitors to adverse effects on aquatic communities can be critical for developing reliable environmental quality criteria for the protection of biota. In a previous publication, the baseline metal concentration in 10 taxa was calculated from unaltered, field reference sites in the Nalón

River basin (northern Spain) (Rodríguez et al. 2018), providing an estimate of the tissue concentration threshold for nine metals, below which alterations in the macroinvertebrate assemblages of the study region are unlikely (ETTC, ecological threshold tissue concentration). In the present study, we aimed to develop high-threshold tissue concentrations above which the impairment of macroinvertebrate assemblages is likely to occur. The impairment was measured through ecologically relevant metrics to protect field populations in the study region.

The tissue residue thresholds for benthic macroinvertebrates are valuable tools in water quality assessments in areas affected by mining activities since they can improve water and sediment quality standards; they can be used as screening benchmarks and provide necessary information in a weight-of-evidence approach using chemistry, toxicity and bioaccumulation data (Meador et al. 2014). We used two approaches to derive tissue residue thresholds: First, we derived the effective tissue residues (ERs) using nonlinear regression models relating taxa tissue residues against two general benchmarks: (1) the good/moderate boundary for macroinvertebrate ecological status, and (2) the 50% reduction of scores and metrics used to assess biological integrity of the macroinvertebrate assemblages. Second, using taxaspecific ERs estimated for As and Cu, we calculated the metal hazard concentrations (HCs) from a multitaxa risk assessment approach. Finally, the reliability of the proposed metal thresholds was tested in the Cauxa Creek risk assessment using a tissue-residue approach with field organisms.

Material and Methods

Study Area and Sampling

The Nalón River basin is located in northern Spain, and its catchment has experienced intense historical and present mining activity (Ordóñez et al. 2013). Samples of sediment and benthic macroinvertebrate taxa were collected and analyzed in two sampling campaigns, 14 reference sites in July 2014 and September 2015 and 15 test sites in September 2015, in the Nalón River basin. Sites were located at four macroinvertebrate community-based river types in Spain (R-T21, R-T25, R-T28, R-T31: MAGRAMA 2015). Detailed data on geographic information, water physicochemistry, and metal concentration in sediments and macroinvertebrate sampling strategy and composition from the study sites were published by Costas et al. (2018) (Online Appendix A, Table A1). Cauxa Creek is a small branch of the Narcea River, the main tributary of the Nalón River, and it is subject to active gold mining (Online Appendix A, Figure A1). Cauxa Creek was resampled in July 2016 for a followup risk assessment. Four sites were scrutinized, one upstream (P1)

and three downstream from the gold mining effluents (from P2, closer to the mining effluents, to P4).

At each site, a composite sample of the upper sediment layer was obtained from fine deposits in the riverbed to measure the sediment metal concentration. Sampling of macroinvertebrates to evaluate bioaccumulation followed a river transect or multihabitat schema to collect 10 bio-monitor taxa grouped into general functional feeding groups: scrapers (Baetidae and Heptageniidae), filterers (Ephemeroidea, Hydropsychidae and Simuliidae), generalists (Ephemeroidea), predators (Perlidae and Rhyacophilidae) and deposit feeders (Lumbricidae and Microdriles oligochaetes) at each site (Table A1). Three field replicates consisting of 1–20 individuals of the larger size class were taken for tissue residue analysis. Detailed information on macroinvertebrate sampling procedures can be found in Rodríguez et al. (2018) and Table A1.

Macroinvertebrate Metrics and Scores

The field community metrics were EPT richness (number of families of Ephemeroptera, Plecoptera and Trichoptera, EPT Fam), EPT abundance (number of individuals of EPT, EPT Ab), and the alteration in the ecological status of the macroinvertebrate assemblages, assessed through ecological quality ratios (EQRs), which are calculated as a quotient between the observed/reference value of a biological metric or score in a previously defined water body type (EC 2000). The EQRs were calculated for the river-type specific multimetric index (METI: MAGRAMA 2015) and for the scores derived from the NORThern Spain Indicators predictive model (NORTI: Pardo et al. 2014), both used in the study region and called METI-EQR and NORTI-EQR, respectively. Data on the tissue residues and site ecological status from the sampling campaigns of 2014 and 2015 in the Nalón River basin were incorporated into the regression models, while data from 2016 in Cauxa Creek were analyzed separately for a tissue-residue risk assessment.

Metal Analysis in Sediments and Macroinvertebrate Tissue Residues

A total of nine metals (As, Cd, Cr, Cu, Hg, Ni, Pb, Se and Zn) were measured in sediments and biota. Limits of quantitation (LOQ) for biota and sediment are shown in Online Appendix A, Table A2. Values below the LOQ were replaced with $\frac{1}{2}$ of LOQ for statistical analysis (US EPA 2000). In present study, we have assessed only four metals (As, Cu, Hg and Se) in biota that were relevant for the study area. All analytical methods can be examined in Costas et al. (2018) and Rodríguez et al. (2018) and are summarized in Table A1. All data in this study are reported in $\mu\text{g g}^{-1}$ dw, and for data in the literature given on a ww basis, we used

a conversion factor of 0.2 for all insect taxa (Meador 2011) and 0.1 for oligochaetes (Méndez-Fernández et al. 2013). For each taxon and site, tissue residues are given as the mean of 3 field replicates (on a few occasions, a single pooled sample was measured because of the scarcity of specimens).

Data Interpretation and Statistical Analyses

Sediment metal concentration was assessed using the probable effect concentration (PEC: MacDonald et al. 2000) and the sediment pollution score (SedPoll: Costas et al. 2018) calculated from As, Cd, Cu, Hg, Pb and Se sediment concentrations in the Nalón River basin. In our previous contributions we described the baseline concentrations (Ecological Tissue threshold concentration, ETTC of 9 metals in several biomonitors of the Nalón River basin, Rodríguez et al. 2018), and more recently a bioaccumulation risk assessment was done, using the number of times the baseline concentrations in selected biomonitor taxa were exceeded (Rodríguez et al. 2021). In the present contribution, ERs were estimated on the same dataset for each taxon and metal from nonlinear regression models of the tissue residues (abscissa) vs EQRs (ordinate). Using the official cutoff for good/moderate ecological status ($EQR = 0.700$, $EQR-ER_{GM}$: effective tissue residue above which the community status changes from good to moderate), a boundary intercalibrated for METI-EQR by the Central/Baltic group for benthic macroinvertebrate fauna (EC 2013a; MAGRAMA 2015). The effect on ecological status was also measured as a 50% reduction in the maximum EQR values ($EQR-ER_{50}$: effective tissue residue above which a 50% reduction in the maximum EQR value occurs) in the model and a 50% reduction in the EPT richness and abundance metrics ($EPT-ER_{50}$: effective tissue residue above which a 50% reduction in EPT richness or abundance occurs) (Fig. 1). In the regression models, ER_{GM} was calculated as the inverse function of the regression equations and solved numerically using Wolfram Mathematica 12 software from $EQR = 0.700$. The $EQR-ER_{50}$ was estimated from the same selected models.

Nonlinear regression analyses were conducted using R software and the extension package drc (Ritz and Streibig 2005). After a preliminary analysis, the best fitted models were selected from a set of 6 commonly used sigmoid models (log-logistic and Weibull) with 3 and 4 parameters:

Log – logistic (LL4) :

$$y = c + (d - c) / (1 + \exp(b(\log x - \log e)))$$

Weibull 1 (W1.4) :

$$y = c + (d - c) \exp(-\exp(b(\log x - \log e)))$$




Fig. 1 Scheme of the construction of the dose–response models between tissue concentration of the metal and the EQR (circles). The blue line is the ETTC value; the red lines are the $EQR = 0.700$ change point between Good and Moderate ecological status (G/M boundary) and the corresponding effective tissue concentration (ER_{GM}). Superimposed are the black lines that represent the effective tissue concentration (ER_{50}) related to the 50% reduction of the EPT indexes

Weibull 2 (W2.4) :

$$y = c + (d - c)(1 - \exp(-\exp(b(\log x - \log e))))$$

The parameters c and d are the lower and upper asymptotes for the y variable, respectively, and they are in the same units as the y variable; parameter e is the inflection point of the dose–response curve and provides the ER_{50} value in the log-logistic models; and parameter b is proportional to the slope of the dose–response curve at dose e (Ritz 2010). Three-parameter models were obtained from each model when $c = 0$.

The models with a difference in Akaike’s Information Criteria ($AIC < 2$) were selected (Burnham and Anderson 2002) and were validated when: (1) the c parameter was ≥ 0 , the minimum value of the EQR and macroinvertebrate assemblage metrics; (2) the estimated ER was within the field range of tissue residue values; and (3) the standard error of the estimated ER was lower than the ER value. In the case of regression models built with the EQRs, we also considered for the validity of the model that the d parameter was ≤ 1.4 , the maximum of the EQR value (Pardo et al. 2010). Graphical tests of the standardized and studentized residuals for the selected equations were examined, and studentized residuals were always $< |3|$.

For each taxon and metal, several $EQR-ER_{GM}$, $EQR-ER_{50}$ and $EPT-ER_{50}$ values were estimated from the validated models, and then averaged. The effective tissue residues (ER) estimated on the 50% reduction in the scores have the advantage over the ER_{GM} that we have been able to calculate

the 95% confidence limits of the estimates. $EQR-ER_{50}$ and ER_{GM} should be similar since the good/moderate boundary of 0.700 is half the maximum expected EQR. Using species sensitivity distribution (SSD) models (ETX v.2.1 program, Van Vlaardingen et al. 2004), the 5th and 50th percentile hazard concentrations (HC_5 , HC_{50}) for the macroinvertebrate assemblages were calculated using the taxa ER_{GM} , $EQR-ER_{50}$ and $EPT-ER_{50}$ mean values.

Finally, the aforementioned threshold values were used in a risk assessment of Cauxa Creek, using the average ratios of the field tissue residues (TRs) to the $EQR-ER_{GM}$ for all the biomonitor taxa present at each site. Four quality classes were considered for the risk assessment based on a tissue-residue approach: (1) Low risk for the community when $TR/ER \leq 1$, (2) Moderate risk when $TR/ER = 1.1-2.0$, (3) High risk when $TR/ER = 2.1-10.0$, and (4) Very High risk when $TR/ER > 10$. The same classification was used to assess the ratios of TR to the HC_{50} values. No risk was expected only when the mean tissue residues were $< HC_5$ or the ETTC (ecological threshold tissue concentration).

Results

Dose–Response Models and Effective Tissue Residues for As, Cu, Hg and Se in Macroinvertebrates

Dose–response models for the relationship of field taxa tissue residues to the macroinvertebrate assemblage metrics (METI- and NORTI-EQRs and EPT richness and abundance) were built when possible for 10 taxa. Tissue residues uploaded to the models from 15 potentially polluted sites from the Nalón River basin are shown in Online Appendix B, Table B1; data from 14 reference sites were reported by Rodriguez et al. (2018). In the study area, METI-EQR values ranged from 0.50–1.27, and NORTI-EQRs ranged from 0.26–1.27. The maximum EPT richness varied from 6 to 25 families, and the EPT abundance varied from 96 to 7333 individuals per site (2.5 m^2). A total of 254 (out of 960 calculated) dose–response models were validated following the criteria reported in “Material and Methods” Section; 87 models for As, 75 for Cu, 52 for Se, and 40 for Hg. ER_{GM} was calculated from 128 models (64 models using METI-EQR data and 64 models using NORTI-EQRs), and the $EQR-ER_{50}$ was calculated from 86 models (Online Appendix B, Table B2); $EPT \text{ Fam-}ER_{50}$ was estimated from 58 regression models, and $EPT \text{ Ab-}ER_{50}$ was estimated from 68 models (Online Appendix B, Table B3). Overall, ERs were calculated for all the study taxa from several selected models: Microdrile oligochaetes (42 models), Rhyacophilidae (33), Baetidae and Heptageniidae (28 each), Lumbricidae (26), Hydropsychidae (23), Ephemerellidae and Ephemeridae (22 each), Simuliidae (19) and Perlidae (11).

The mean ERs calculated for each metal and taxon relative to the good/moderate ecological status boundary (ER_{GM}) are shown in Table 1. The ER_{GM} concentrations estimated from NORTI-EQRs were generally 1–3 times lower than the ER_{GM} concentrations derived from the METI-EQRs. The highest As ER_{GM} values were for Simuliidae or Microdrile oligochaetes, while the lowest values were found in the predators Rhyacophilidae and Perlidae. The Cu ER_{GM} was higher for Heptageniidae and Ephemerellidae, while the deposit feeders and filterers had lower values. Only a small number of models were validated for Hg and Se.

Regarding the ratios of the ER_{GM} to the baseline ETTC for each metal and taxon (Table 1), the METI- ER_{GM} values were usually 4–20 times the ETTC for As (but up to 60 for Simuliidae); these ratios ranged 2–7 times for Cu, 1–14 for Hg, and approximately 2 for Se. The ratios of NORTI- ER_{GM} to ETTC were generally lower, 2–13 for As, and varied typically from 1–2 for Cu, Hg and Se. In most instances, the $EQR-ER_{50}$ values ranged between 0.4 and 7.1 times (mean = 1.7) the corresponding ER_{GM} .

We found $EPT \text{ Fam-}ER_{50}$ for As (Table 2) to be lower for EPT abundance than for EPT richness. This is interpreted as As tissue residues causing a reduction in abundance of sensitive taxa before having an effect on the number of families. For Cu, the ratios of Cu $EPT-ER_{50}$ to ETTC were 1–3 (except for Rhyacophilidae, with ratios < 1), with similar values for richness and abundance. In the case of Hg and Se, the $EPT-ER_{50}$ estimates were limited to a few taxa and, in most instances, $< 1 \mu\text{g g}^{-1}$ for Hg, which resulted in ratios to ETTC ≤ 1 . The $EPT \text{ Ab-}ER_{50}$ was 3 and 2 times higher than the baseline ETTC for As and Cu, respectively (Table 2; Figs. 2 and 3, for Baetidae, Ephemerellidae, Lumbricidae and Rhyacophilidae). However, $EPT \text{ Fam-}ER_{50}$ values were much higher than ETTC for As (mean ratio 16.9) but only 2 times higher for Cu and equal or lower for Hg and Se (Table 2). The ratios for Se were very variable, although based on a limited number of data.

Last, using the SSD models based on the taxa ERs, we estimated hazard concentrations for the integrity of the whole community. The HC_5 and HC_{50} were estimated for As and Cu (Table 3) but not for Hg and Se due to the limited number of data points. The HC_5 derived from the $EQR-ER_{GM}$ values for both As and Cu were similar to those estimated using $EQR-ER_{50}$ or $EPT-ER_{50}$ data (Table 3). The HC_{50} values estimated from the same SSD models were 11–25 times higher than the corresponding HC_5 for As and 4–5 times higher for Cu. The respective 90% confidence limits of HC_5 and HC_{50} did not overlap, which supports their use as low and high community thresholds, respectively.

Table 1 Mean effective tissue residues (ERs) ($\mu\text{g g}^{-1}$ dw) calculated for As, Cu, Hg and Se from non-linear regression models of the tissue concentration and the EQR for each site and taxon

Taxon	Mean METI–EQR versus TR regression models			Mean NORTI–EQR versus TR regression models		
	EQR–ER ₅₀ (range)	EQR–ER _{GM} (range)	Mean ER _{GM} / ETTC	EQR–ER ₅₀ (range)	EQR–ER _{GM} (range)	Mean ER _{GM} / ETTC
As						
Baetidae	71.0 (32.1–186)	41.6 (37.3–52.2)	13.4	99.5 (51.5–123)	18.9 (16.3–21.7)	6.1
Ephemerellidae	–	67.7 (66.7–68.7)	10.5	–	35.5	5.5
Ephemeridae	–	–	–	–	44.3	6.9
Heptageniidae	–	48.6 (47.6–49.4)	7.7	7.8 (7.6–8.0)	9.0 (8.0–10.8)	2.3
Hydropsychidae	16.1 (2.6–29.5)	–	–	–	25.5 (23.1–27.8)	12.7
Lumbricidae	17.2	19.1	1.4	84.9 (83.5–86.2)	85.7 (84.5–86.9)	6.2
Microdrile	561	243.1 (230–256)	16.8	542.4 (509–570)	109.4 (79.0–126)	7.6
Rhyacophilidae	11.1 (10.7–11.7)	7.1 (6.9–7.4)	4.2	1.1	1.1	1.5
Perlidae	–	–	–	–	2.6	1.5
Simuliidae	120	269 (196–341)	59.8	–	17.8	4.0
Cu						
Baetidae	39.2	104.5	3.6	33.6 (32.8–34.6)	38.1 (36.6–39.6)	1.3
Ephemerellidae	237 (236–240)	194 (191–199)	6.5	–	134.4	1.5
Ephemeridae	37.0 (36.5–37.8)	–	–	19.1 (16.2–21.0)	22.9 (16.5–27.3)	1.4
Heptageniidae	220 (199–241)	409.7 (407–414)	5.1	98.1 (97–100)	158 (156–160)	2.0
Hydropsychidae	–	–	–	15.6 (15.3–15.9)	18.5 (17.9–19.1)	1.2
Lumbricidae	146 (145–147)	82.4 (77.7–86.8)	7.1	208	45.7	4.0
Microdrile	36.0 (35.7–36.4)	37.7 (37.3–38.1)	1.8	31.6 (30.1–33.0)	30.9 (29.6–32.2)	1.5
Perlidae	118 (105–131)	98.5 (97.7–99.3)	2.9	–	–	–
Rhyacophilidae	67.1	51.2 (49.5–52.5)	2.6	32.2 (31.9–32.5)	39.4 (32.9–44.7)	2.0
Simuliidae	–	–	–	15.8 (15.5–16.1)	37.2 (36.4–37.9)	0.7
Hg						
Baetidae	–	3.6	13.7	–	–	–
Ephemeridae	0.05	–	–	0.10	–	–
Heptageniidae	–	–	–	0.06 (0.06–0.07)	–	–
Perlidae	–	–	–	0.07 (0.06–0.07)	–	–
Rhyacophilidae	0.60	0.59	1.2	0.23 (0.21–0.24)	0.35 (0.34–0.36)	0.7
Simuliidae	–	7.9 (7.8–8.0)	13.1	–	0.97 (0.96–0.97)	1.6
Se						
Baetidae	–	20.1 (18.7–22.1)	1.8	15.2 (14.8–15.6)	8.7 (8.6–8.7)	0.7
Heptageniidae	–	–	–	3.4 (3.2–3.5)	–	–
Hydropsychidae	3.0	–	–	1.1 (0.9–1.1)	–	–
Microdrile	7.1 (6.6–7.6)	16.8	2.2	5.8	5.9	0.8

Their ranges are given when $n > 1$. EQR–ER₅₀ was calculated as the tissue residues corresponding to a 50% reduction in the EQR score. EQR–ER_{GM} was estimated from the models for the official EQR value used as the boundary between good and moderate ecological status of the macroinvertebrate assemblages. Ratios of the ER_{GM} to the baseline ETTC are shown. ETTC, ecological threshold tissue concentration; TR, tissue residues

Cauxa Creek Risk Assessment: A Tissue Residue Approach

The sediment metal concentration in the 2016 campaign in Cauxa Creek showed that 4 metals, As, Cu, Hg and Se, exceeded the PEC values in sites downstream of mine effluents (P2–P4) (Online Appendix B, Table B4). At P2, As was

up to 48 times the PEC value, Cu 13 times and Se 4 times, while the Hg concentration barely exceeded the PEC. Additionally, sediment metal pollution assessed by the SedPoll index evaluated the upstream site, P1, as unpolluted or similar to the reference, while sites P2 to P4 were assessed as medium to highly polluted.

Table 2 Effective tissue residues (ER₅₀) of As, Cu, Hg and Se (μg g⁻¹ dw) related to the 50% reduction in the EPT number of families (EPT Fam) and the EPT abundance (EPT Ab)

Taxon	As		Cu		Hg		Se	
	EPT Fam–ER ₅₀	EPT Ab–ER ₅₀	EPT Fam–ER ₅₀	EPT Ab–ER ₅₀	EPT Fam–ER ₅₀	EPT Ab–ER ₅₀	EPT Fam–ER ₅₀	EPT Ab–ER ₅₀
Baetidae	24.1	11.4	–	–	–	–	19.5	5.7
Ephemere- llidae	92.1	8.4	237.2	192.0	0.06	–	–	–
Ephemeri- dae	20.5	–	38.0	–	0.14	0.61	–	–
Heptageni- idae	8.1	18.1	–	–	–	–	–	4.8
Hydropsychi- dae	107.7	–	16.7	–	–	–	2.1	4.3
Lumbricidae	16.7	–	–	32.8	–	–	10.0	14.9
Microdrile oli- gochaetes	318.7	85.5	–	55.1	–	0.72	4.4	8.0
Perlidae	–	–	167.8	77.6	–	0.06	–	–
Rhyacophi- lidae	8.1	1.5	–	42.4	–	–	–	1.1
Simuliidae	171.5	8.4	–	–	–	0.72	–	–
ER ₅₀ /ETTC ratio	1.2–53.9	1.3–5.9	1.1–2.7	0.3–2.8	0.2–0.3	0.0–1.2	0.5–1.6	0.2–1.1
Mean	16.9	2.9	2.2	2.1	0.25	1.0	0.8	0.7

ER₅₀ are calculated as the mean ER₅₀ values estimated per taxon from the validated regression models. Abbreviations: EPT, Ephemeroptera, Plecoptera and Trichoptera; ETTC, Ecological Threshold Tissue Concentration




Fig. 2 Arsenic dose—response models for METI- and NORTI-EQRs calculated for several taxa (Baetidae, Ephemerellidae, Lumbricidae and Rhyacophilidae) selected from sites in the Nalón River basin. Superimposed lines mark the low and high threshold values. The low threshold is in all instances the baseline concentration in the basin (ETTC), in blue. There are several parameters useful as high thresh-

olds: METI- and NORTI-ER_{GM} in red. For comparison, the ER₅₀ values calculated when possible for the 50% reduction of EPT number of families and abundances are also superimposed in black-dashed line and in black-dotted line, respectively. The represented regression models are indicated on the upper right corner of each plot




Fig. 3 Copper dose—response models for METI- and NORTI-EQRs calculated for several taxa (Baetidae, Ephemerellidae, Lumbricidae and Rhyacophilidae) selected from sites in the Nalón River basin. Superimposed lines mark the low and high threshold values. The low threshold is in all instances the baseline concentration in the basin (ETTC), in blue. There are several parameters useful as high thresh-

olds: METI- and NORTI- ER_{GM} in red. For comparison, the ER_{50} values calculated when possible for the 50% reduction of EPT number of families and abundances are also superimposed in black-dashed line and in black-dotted line, respectively. The represented regression models are indicated on the upper right corner of each plot

Table 3 Hazard concentrations (HC_5 , HC_{50}) for As and Cu and their 90% confidence limits (CL) derived from SSD models using several effective tissue concentrations (ER)

Chemical	HC	EQR- ER_{GM} (90% CL)	<i>n</i>	EQR- ER_{50} (90% CL)	<i>n</i>	EPT- ER_{50} (90% CL)	<i>n</i>
As	HC_5	2.41 (0.74–5.27)	17	1.67 (0.20–5.74)	11	2.12 (0.59–4.82)	15
	HC_{50}	27.98 (15.06–51.97)		41.05 (14.62–115.3)		24.13 (12.50–46.58)	
	As ETTC- HC_{50}	4.24 (2.46–7.30)					
Cu	HC_5	15.45 (7.52–24.70)	16	11.63 (5.25–19.51)	16	14.31 (4.32–27.48)	9
	HC_{50}	64.62 (44.47–93.89)		56.40 (37.34–85.20)		67.43 (38.44–118.3)	
	Cu ETTC- HC_{50}	29.16 (19.23–44.21)					

EQR- ER_{GM} : Effective tissue residues related to the EQR boundary between Good and Moderate ecological status of the macroinvertebrate community; EQR- ER_{50} : Effective tissue residues related to the 50% reduction in the EQR; EPT- ER_{50} : Effective tissue residues related to the 50% reduction of the EPT metrics of abundance and richness. For comparison, the HC_{50} derived from the baseline concentrations of As and Cu in unpolluted reference sites is given (ETTC- HC_{50} , Rodríguez et al. 2018). *n*: number of data included in each of the models

At site P1, ten biomonitors used in the Nalón River basin were found, but downstream (P2–P4), only four of them were present at the four study sites: Baetidae, Ephemerellidae, Lumbricidae and Rhyacophilidae. These taxa represent four different feeding styles: scraper, generalist, deposit feeder and predator. Bioaccumulation levels of As and Cu were high, up to a maximum of 49 times the ETTC (Baetidae) at P2, as expected from the high metal concentration

in the sediment. However, the Hg and Se tissue residue to ETTC ratios were usually < 1 , with a maximum of 1.9 for Hg and 1.6 for Se (Lumbricidae) in P2 (Online Appendix B, Table B5).

The Cauxa Creek risk assessment based on the tissue residue approach was performed using the average ratios of tissue residues to their corresponding high thresholds (EQR- ER_{GM}) calculated for each taxon present (Table 4). These

Table 4 Cauxa river risk assessment based on the mean ratios of the tissue residues in biomonitor taxa to metal thresholds

Ratios of tissue residues to effective tissue residues						
Threshold concentration	Chemical	Site				
		P1	P2	P3	P4	
METI-EQR ER _{GM}	As	0.1	4.2	2.4	2.9	
	Cu	0.5	1.4	1.0	1.3	
	Hg	0.1	0.1	0.1	0.2	
	Se	0.2	0.5	0.5	0.6	
NORTI-EQR ER _{GM}	As	0.2	9.3	3.5	2.3	
	Cu	0.9	3.5	3.5	1.9	
	Hg	0.3	0.3	0.4	0.6	
	Se	0.5	1.1	1.1	1.5	
Hazard Concentrations						
EQR-ER _{GM} HC ₅₀	As	0.1	3.3	2.9	2.4	
	Cu	0.9	2.7	2.5	1.7	
Ecological status						
METI-EQR		0.837	0.570	0.592	0.500	
NORTI-EQR		0.779	0.621	0.665	0.582	

The effective tissue residues (EQR-ER_{GM}) and the community hazard concentration (HC₅₀) were used as high thresholds. The ratios are classified within the following classes: Low Risk (≤ 1) in blue, Moderate Risk (1.1–2.0) in yellow, High Risk (2.1–10.0) in orange and Very High Risk (> 10) in red. The ecological status of the benthic macroinvertebrate assemblages in the four sites of the Cauxa River assessed by the METI and NORTI EQRs are shown in blue (Good) and yellow (Moderate)

ratios were also averaged for each feeding style (Online Appendix B, Table B6). Site P1, upstream from the gold mine, showed in all cases low risk due to As, Cu, and Hg tissue residue ratios to EQR-ER_{GM}. In all cases, downstream sites were assessed as High Risk related to As bioaccumulation (Table 4). The Cu bioaccumulation risk assessment result was variable, depending on the EQR-ER_{GM} used. Se tissue residues downstream of the mine were assessed as Low or Moderate Risk. The Hg showed Low Risk related to the METI and NORTI assessments. The importance in this risk assessment of the high ratios obtained for predators for As and Se in downstream sites can be seen in Table B6.

Finally, the risk assessment for As and Cu tissue residues using their ratios to HC₅₀ for the macroinvertebrate assemblage showed a consistent evaluation with the EQR-ER_{GM} assessment (Table 4). The upstream site (P1) was at Low Risk, while sites downstream of the mine effluents (P2–P4) were assessed from High to Moderate Risk, in parallel with a decreasing concentration of pollutants in the sediment and with decreasing ecological status score values, both for METI and NORTI (< 0.700 , good/moderate boundary). When HC₅ was used as the benchmark, the four sites showed some degree of risk due to As and Cu bioaccumulation (tissue residue ratios to HC₅ > 1).

Discussion

The main goal in water quality management is to attain or maintain the ecological integrity of aquatic communities, thus incorporating ecological realism into the regulatory

framework (Kiffney and Clements 2002). In Spain, the evaluation of exposures to contaminants in biota is based on the *standstill* principle, which states that priority substances in the sediments or biota should not significantly increase their concentration on a long-term basis (MAGRAMA 2015). This evaluation is inadequate to prevent loss or to recover the good ecological status of aquatic communities, since bioaccumulation can be one of the causes of adverse effects, hindering the recovery of the good ecological status of the macroinvertebrate assemblages. In that context, the tissue residue approach provides a step in the evaluation of causal agents derived from contaminants (Meador et al. 2014) and is a necessary tool for developing tissue quality criteria that should improve risk assessment and remediation policies.

A comparison of our thresholds with others (calculated using different approaches, but also associated with a reduction in macroinvertebrate community metrics), showed that the similarity among values depends on the biomonitors selected. In other studies, adverse effects on aquatic organisms related to As bioaccumulation were similar to the As HC₅₀ (28 $\mu\text{g g}^{-1}$ dw) estimated in the present study: e.g. 1.3–5 $\mu\text{g As g}^{-1}$ ww (≈ 6.5 –25 $\mu\text{g g}^{-1}$ dw) (Eisler 2000), and 6.6 $\mu\text{g As g}^{-1}$ ww ($\approx 33 \mu\text{g g}^{-1}$ dw) (DEQ 2007). Higher As thresholds were reported by Bervoets et al. (2016), but this probably was related to their selection of relatively tolerant biomonitors (Diptera: Chironomidae: 65–130 $\mu\text{g g}^{-1}$ dw and tubificid oligochaetes: 85–93 $\mu\text{g g}^{-1}$ dw). In our study, the filterer Simuliidae (Diptera) and sediment-feeder Microdrile oligochaetes also showed higher As ERs (Table 1).

The Cu ER_{GM} in the present study attained high values (155.7–414.1 $\mu\text{g g}^{-1}$ dw for Heptageniidae and 134.4–199.3

for Ephemerellidae, Table B2). However, the $ER_{GM} HC_{50}$ ($65 \mu\text{g Cu g}^{-1} \text{dw}$, Table 3) was similar to that proposed by Bervoets et al. (2016) ($57 \mu\text{g g}^{-1} \text{dw}$). Higher Cu ERs have been reported for specific biomonitors. For example, the value for Hydropsychidae ($> 170 \mu\text{g g}^{-1} \text{dw}$) was associated with a reduction or absence of heptageniids and ephemerellid mayflies (Rainbow et al. 2012), and the values for Heptageniidae ($165.2\text{--}349.5 \mu\text{g g}^{-1} \text{dw}$) were associated with a 20–50% loss in macroinvertebrate richness (De Jonge et al. 2013).

The range of values estimated for Hg ER_{GM} in the present study is wide ($0.35\text{--}7.9 \mu\text{g g}^{-1} \text{dw}$), but the higher values estimated for Simuliidae and Baetidae (Table 1) should be viewed with caution, since they are very biased with respect to the median value ($0.97 \mu\text{g g}^{-1} \text{dw}$). The lower range is comparable to the Hg guidelines proposed for the biota (e.g. $0.12\text{--}1.68 \mu\text{g g}^{-1} \text{dw}$: CCME 2000; $20 \text{ ng g}^{-1} \text{ww} \approx 0.1 \mu\text{g g}^{-1} \text{dw}$: EC 2013b).

Only a small number of Se ER_{GM} could be estimated in our study, ranging from $5.9\text{--}20.1 \mu\text{g g}^{-1} \text{dw}$, comparable to the tissue residues associated with sublethal toxic effects reported by DeBruyn and Chapman (2007) ($1\text{--}30 \mu\text{g g}^{-1} \text{dw}$), and dietary Se thresholds for fish (e.g. 3 to $11 \mu\text{g g}^{-1} \text{dw}$, May et al. 2008).

In the selection of suitable bioaccumulation thresholds, it is desirable that there is a clear but not very large gap between low and high thresholds to reduce the probability of false positives or negatives in the risk assessment. This was the case for most selected taxa, which showed an $ER_{GM}/ETTC$ ratios of $> 1\text{--}10$ for As and Cu (Table 1). However, for Hg and Se these ratios were calculated in very few instances. In the case of Se, the low ratios are probably associated with its essential nature for metabolism, and to the fact that most species of aquatic macroinvertebrates are relatively insensitive to Se (Janz et al. 2014). The database for Hg and Se should be completed with supplementary sites to better understand these low ratios and provide a better risk assessment in the future.

The estimated metal thresholds (ERs) vary by one or two orders of magnitude, depending on the selected biomonitors. However, the metal Hazard Concentrations (HC_5 and HC_{50}) calculated from different biological effective tissue residues are much more comparable to each other. The HC values estimated from SSD models using $EQR\text{--}ER_{50}$ and ER_{GM} were very similar, and also were similar to the ER_{50} calculated from the 50% reduction of the abundance and richness of the EPT taxa. This suggests that thresholds estimated from the good/moderate boundary are a reliable measure of ecological status. The HC_5 90% confidence limits of As and Cu calculated from different ERs overlapped with each other and with the confidence limits of the baseline $ETTC\text{--}HC_{50}$ (Table 3), thus making the HC_5 a reliable low threshold for risk assessment. Nevertheless, when tissue concentrations

are close to HC_5 , a comparison with the baseline concentrations of the biomonitor will improve the accuracy of the risk assessment. In addition, the accuracy of the HC thresholds would probably improve if the sensitivity of the biomonitors was within the HC confidence interval of the metal, avoiding false negatives or positives.

The risk assessment exercise in Cauxa Creek largely affected by gold mining clearly pointed toward the influence of As and Cu bioaccumulation by macroinvertebrates on the altered ecological status of sites downstream of the mine (P2-P4). The $EQR\text{--}ER_{GM}$ thresholds are taxon specific, thus it is possible to get different assessments depending on the biomonitors. In this case study, the average of the ratios of $TR/EQR\text{--}ER_{GM}$ from a selection of several biomonitor species comprise a wide range of sensitivity to the metals, which helps getting a weighted assessment of the risk. The risk assessment through $NORTI\text{--}ER_{GM}$ was closer to the ecological status assessment than through $METI\text{--}ER_{GM}$. However, in Cauxa Creek, the ratios of the tissue residues to the community high thresholds (HC_{50}) resulted in a straightforward and consistent risk assessment, comparable to the ecological status assessment. The HC thresholds also have the advantage of being less dependent on the presence of certain biomonitor taxa at the study sites.

Despite the relevance of the interaction of several metals to evaluate the effects on the biota due to bioaccumulation, there are few studies that have addressed the effects of metal mixtures in field organisms (e.g., De Jonge et al. 2013). In the present study, the interactions of Se with other bioaccumulated trace metals (e.g., As, Cu and Hg) must be analyzed in more detail, since interactions have been demonstrated in the literature. In particular, Se is recognized for its potential in reducing the toxicity of Hg compounds (Hamilton 2004), an issue that requires further research in Cauxa Creek. Specific thresholds for the protection of higher levels in the aquatic trophic chain should also be developed in the future for the satisfactory protection of aquatic communities. This problem is complex since the risk of metal transfer from macroinvertebrates to aquatic wildlife depends on the diet specificity, prey availability, accumulation pattern and ability of the organisms to depurate the metals (Rainbow 2018).

Conclusions

This study is the first to derive the effective tissue concentration from the cutoff value of good/moderate ecological status of the macroinvertebrate assemblages using ten biomonitor taxa. The ecological status of the field community is regularly evaluated by the water authorities, following the European water directive; therefore, EQR can be useful to calculate environmental thresholds for macroinvertebrates derived through a tissue residue approach. The models provide a

complementary tool not only to monitor environmental risk due to bioaccumulation, but also to predict alterations in the ecological status of field macroinvertebrate assemblages. The HC₅ and HC₅₀ calculated for As and Cu are promising since they can be readily applicable as low and high thresholds in the mining districts of northern Spain. They can contribute to setting future environmental quality standards for the protection of aquatic biota. The same approach can be implemented in other European river basins to calculate threshold concentrations in the biota related to reductions in intercalibrated metrics of ecological status.

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1007/s00244-022-00915-w>.

Acknowledgements This investigation has been part of the CGL 2013-44655-R research project sponsored by the Spanish Ministry of Economy, Industry and Competitiveness. Iñigo Moreno-Ocio was financed by a Ph.D. scholarship from the University of the Basque Country (UPV/EHU), in 2017–2021 (PIF16/162). We greatly appreciate the support provided by Amanda Miranda and the staff of the Cantabrian Hydrographical Confederation, who assisted in the selection and sampling of the sites in the Nalón River basin. The authors are grateful to Dr. Miguel Escobedo (Dpt. of Mathematics, UPV/EHU) for his assistance with the non-linear regression models and Mathematica software.

Authors Contribution The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper. Credit Author Statement: IMO: Methodology, Formal analysis, Investigation, Writing–Original Draft. LMF: Conceptualization, Methodology, Investigation, Data Curation, Writing–Review & Editing. Maite Martínez-Madrid: Investigation, Writing–Review & Editing, Funding acquisition. NC: Investigation, Data Curation. Isabel Pardo: Investigation, Writing–Review & Editing, Funding acquisition. PR: Methodology, Formal analysis, Investigation, Writing–Original Draft, Funding acquisition.

Funding Open Access funding provided thanks to the CRUE-CSIC agreement with Springer Nature. Funding was provided by Spanish Ministry of Economy, Industry and Competitiveness, (Grant no. CGL 2013-44655-R), University of the Basque Country, (Grant no. PIF16/162)

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Adams WJ, Blust R, Borgman U, Brix KV, DeForest DK, Green AS, Meyer JS, McGeer JC, Paquin PR, Rainbow PS, Wood CM (2011) Utility of tissue residues for predicting effects of metal on aquatic organisms. *Integr Environ Assess Manag* 7:75–98. <https://doi.org/10.1002/ieam.108>
- Bervoets L, De Jonge M, Blust R (2016) Identification of threshold body burdens of metals for the protection of the aquatic ecological status using two benthic invertebrates. *Environ Pollut* 210:76–84. <https://doi.org/10.1016/j.envpol.2015.12.005>
- Burnham K, Anderson D (2002) Model selection and multimodel inference. Springer-Verlag, New York, p 488
- Cain DJ, Luoma SN, Wallace WG (2004) Linking metal bioaccumulation of aquatic insects to their distribution patterns in a mining-impacted river. *Environ Toxicol Chem* 23:1463–1473. <https://doi.org/10.1897/03-291>
- CCME, Canadian Council of Ministers of the Environment (2000) Canadian environmental quality guidelines. Canadian tissue residue guidelines for the protection of wildlife consumers of aquatic biota. Methylmercury. Environment Canada, guidelines and standards division p. 7
- Chapman PM (2007) Do not disregard the benthos in sediment quality assessments! *Marine Pollut Bull* 54:633–635. <https://doi.org/10.1016/j.marpolbul.2007.03.006>
- Clements WH (1991) Community responses of stream organisms to heavy metals: a review of observational and experimental approaches. Chapter 13. In: Newman MC, McIntosh AW (eds) *Metal Ecotoxicology concepts and applications*. CRC Press, Boca Raton, pp 363–390
- Costas N, Pardo I, Méndez-Fernández L, Martínez-Madrid M, Rodríguez P (2018) Sensitivity of macroinvertebrate indicator taxa to metal gradients in mining areas in Northern Spain. *Ecol Indic* 93:207–218. <https://doi.org/10.1016/j.ecolind.2018.04.059>
- De Jonge M, Tipping E, Lofts S, Bervoets L, Blust R (2013) The use of invertebrate body burdens to predict ecological effects of metal mixtures in mining-impacted waters. *Aquat Toxicol* 142:294–302. <https://doi.org/10.1016/j.aquatox.2013.08.018>
- DeBruyn AMH, Chapman PM (2007) Selenium toxicity to invertebrates: will proposed thresholds for toxicity to fish and birds also protect their prey? *Environ Sci Technol* 41:1766–1770. <https://doi.org/10.1021/es062253j>
- DEQ, Oregon Department of Environmental Quality (2007) Guidance for assessing bioaccumulative chemicals of concern in sediment. Environmental cleanup program. 07-LQ-023A
- EC, European Commission (2000) Directive 2000/60/EC. Establishing a framework for community action in the field of water policy. OJEC L 327, pp. 1–71
- EC, European Commission (2008) Directive 2008/105/EC of the European parliament and of the council on environmental quality standards in the field of water policy. OJEC L vol. 348: pp. 84–97
- EC, European Commission (2011) Technical guidance for deriving environmental quality standards. Guidance Document No. 27, p. 204
- EC, European Commission (2013a) 2013a/480/UE. Commission Decision of 20 September 2013a establishing, pursuant to Directive 2000/60/EC of the European Parliament and of the Council, the values of the Member State monitoring system classifications as a result of the intercalibration exercise and repealing Decision 2008/915/EC. CELEX 32013aD0480
- EC, European Commission (2013b) Directive 2013/39/EU of the European parliament and of the council as regards priority substances in the field of water policy. OJEU L226 vol. 17

- Eisler R (2000) Handbook of chemical risk assessment: health hazards to humans, plants, and animals. Lewis Publishers, Boca Raton
- Hamilton SJ (2004) Review of selenium toxicity in the aquatic food chain. *Sci Total Environ* 326:1–31
- Janz DM, Liber K, Pickering IJ, Wiramanaden CIE, Weech SA, Gallego-Gallegos M, Driessnack MK, Franz ED, Goertzen MM, Phibbs J, Tse JJ, Himbeault KT, Robertson EL, Burnett-Seidel Ch, England K, Gent A (2014) Integrative assessment of selenium speciation, biogeochemistry, and distribution in a northern coldwater ecosystem. *Integr Environ Assess Manag* 10:543–554. <https://doi.org/10.1002/ieam.1560>
- Kiffney PM, Clements WH (2002) Ecological effects of metals on benthic invertebrates. Chapter 8. In: Simon TP (ed) Biological response signatures: Indicator patterns using aquatic communities. CRC Press, Boca Raton, pp 135–154
- Loredo J, Petit-Domínguez MD, Ordoñez A, Galán MP, Fernández-Martínez R, Alvarez R, Rucandio MI (2010) Surface water monitoring in the mercury mining district of Asturias (Spain). *J Hazard Mater* 176:323–332. <https://doi.org/10.1016/j.jhazmat.2009.11.031>
- Luoma SN, Cain DJ, Rainbow PS (2010) Calibrating biomonitors to ecological disturbance: a new technique for explaining metal effects in natural waters. *Integr Environ Assess Manag* 6:199–209. https://doi.org/10.1897/IEAM_2009-067.1
- MacDonald DD, Ingersoll CG, Berger TA (2000) Development and evaluation of consensus-based sediment quality guidelines for freshwater ecosystems. *Arch Environ Contam Toxicol* 39:20–31. <https://doi.org/10.1007/s002440010075>
- MAGRAMA (2015) Real decreto 817/2015. Criterios de seguimiento y evaluación del estado de las aguas superficiales y las normas de calidad ambiental. BOE-A-2015–980680582- 80677 p. 96 (in Spanish). <https://www.boe.es/eli/es/rd/2015/09/11/817>
- May TW, Fairchild JF, Petty JD, Walther MJ, Lucero J, Delvaux M, Manring J, Armbruster M (2008) An evaluation of selenium concentrations in water, sediment, invertebrates, and fish from the Solomon River Basin. *Environ Monit Assess* 137:213–232. <https://doi.org/10.1007/s10661-007-9742-y>
- McCarty LS, Landrum PF, Luoma SN, Meador JP, Merten AA, Shephard BK, Van Wezel AP (2011) Advancing environmental toxicology through chemical dosimetry: external exposures versus tissue residues. *Integr Environ Assess Manag* 7:7–27. <https://doi.org/10.1002/ieam.98>
- Meador JP (2011) Organotins in aquatic biota: occurrence in tissue and toxicological significance. In: Beyer WN, Meador JP (eds) Environmental contaminants in biota: interpreting tissue concentrations. CRC Press, Boca Raton, pp 256–284
- Meador JP, Warne MSJ, Chapman PM, Chan KM, Yu S, Leung KM (2014) Tissue-based environmental quality benchmarks and standards. *Environ Sci Pollut Res* 21:28–32. <https://doi.org/10.1007/s11356-013-1714-x>
- Méndez-Fernández L, Martínez-Madrid M, Rodríguez P (2013) Toxicity and critical body residues of Cd, Cu and Cr in the aquatic oligochaete *Tubifex tubifex* (Müller) based on lethal and sublethal effects. *Ecotoxicology* 22:1445–1460. <https://doi.org/10.1007/s10646-013-1131-4>
- Méndez-Fernández L, Rodríguez P, Martínez-Madrid M (2015) Sediment toxicity and bioaccumulation assessment in abandoned copper and mercury mining areas of the Nalón River basin (Spain). *Arch Environ Contam Toxicol* 68:107–123. <https://doi.org/10.1007/s00244-014-0093-8>
- Méndez-Fernández L, Casado-Martínez C, Martínez-Madrid M, Moreno-Ocio I, Costas N, Pardo I, Rodríguez P (2019) Derivation of sediment Hg quality standards based on ecological assessment in river basins. *Environ Pollut* 245:1000–1013. <https://doi.org/10.1016/j.envpol.2018.11.068>
- Ordóñez A, Álvarez R, Loredó J (2013) Asturian mercury mining district (Spain) and the environment: a review. *Environ Sci Pollut Res* 20:7490–7508. <https://doi.org/10.1007/s11356-013-1663-4>
- Pardo I, Gómez-Rodríguez C, Abraín R, García-Roselló E, Reynoldson TB (2014) An invertebrate predictive model (NORTI) for streams and rivers: sensitivity of the model in detecting stress gradients. *Ecol Indic* 45:51–62. <https://doi.org/10.1016/j.ecolind.2014.03.019>
- Pardo I, Abraín R, Gómez-Rodríguez C, García-Roselló E (2010) Aplicación de los sistemas de evaluación del estado ecológico desarrollados para ríos en la aplicación de la Directiva Marco del agua en la Demarcación Hidrográfica del Cantábrico 2010. Version 1. NIPO 282–12–001-X
- Rainbow PS (2018) Trace metals in the environment and living organisms: the British Isles as a case study. Cambridge University Press, Cambridge
- Rainbow PS, Hildrew AG, Smith BD, Geatches T, Luoma SN (2012) Caddisflies as biomonitors identifying thresholds of toxic metal bioavailability that affect the stream benthos. *Environ Pollut* 166:196–207. <https://doi.org/10.1016/j.envpol.2012.03.017>
- Ritz C (2010) Toward a unified approach to dose–response modeling in ecotoxicology. *Environ Toxicol Chem* 29:220–229. <https://doi.org/10.1002/etc.7>
- Ritz C, Streibig JC (2005) Bioassay analysis using R. *J Stat Softw* 12:1–22
- Rodríguez P, Méndez-Fernández L, Pardo I, Costas N, Martínez-Madrid M (2018) Baseline tissue levels of trace metals and metalloids to approach ecological threshold concentrations in aquatic macroinvertebrates. *Ecol Indic* 91:395–409
- Rodríguez P, Moreno-Ocio I, Martínez-Madrid M, Costas N, Pardo I, Méndez-Fernández L (2021) Proposal of integrative scores and biomonitor selection for metal bioaccumulation risk assessment in mine-impacted rivers. *Aquat Toxicol* 238:105918. <https://doi.org/10.1016/j.aquatox.2021.105918>
- Sappington KG, Bridges TS, Bradbury SP, Erickson RJ, Hendriks AJ, Lanno RP, Meador JP, Mount DR, Salazar MH, Spry DJ (2011) Application of the tissue residue approach in ecological risk assessment. *Integr Environ Assess Manag* 7:116–140
- Solà C, Burgos M, Plazuelo Á, Toja J, Plans M, Prat N (2004) Heavy metal bioaccumulation and macroinvertebrate community changes in a Mediterranean stream affected by acid mine drainage and an accidental spill (Guadiamar River, SW Spain). *Sci Total Environ* 333:109–126. <https://doi.org/10.1016/j.scitotenv.2004.05.011>
- US EPA, United States Environmental Protection Agency (2000) Guidance for data quality assessment: practical methods for data analysis. EPA/600/R-96/084, Washington, DC
- Van Vlaardingen PLA, Traas TP, Wintersen AM, Aldenberg T (2004) ETX v.2.0. A program to calculate hazardous concentrations and fraction affected, based on normally distributed toxicity data. Bilthoven, the Netherlands (RIVM). Report no. 601501028/2004 p. 68