

Kent Academic Repository

Full text document (pdf)

Citation for published version

Zhou, Jiandong, Leung, Keith Sai Kit, Kong, Dicken, Lee, Sharen, Liu, Tong, Wai, Abraham Ka Chung, Chang, Carlin, Zhang, Qingpeng and Tse, Gary (2021) Low rates of liver injury in edoxaban users: Evidence from a territorywide observational cohort study. *Clinical Cardiology*, 44 (7). pp. 886-889. ISSN 0160-9289.

DOI

<https://doi.org/10.1002/clc.23570>

Link to record in KAR

<https://kar.kent.ac.uk/98749/>

Document Version

Publisher pdf

Copyright & reuse

Content in the Kent Academic Repository is made available for research purposes. Unless otherwise stated all content is protected by copyright and in the absence of an open licence (eg Creative Commons), permissions for further reuse of content should be sought from the publisher, author or other copyright holder.

Versions of research

The version in the Kent Academic Repository may differ from the final published version.

Users are advised to check <http://kar.kent.ac.uk> for the status of the paper. **Users should always cite the published version of record.**

Enquiries

For any further enquiries regarding the licence status of this document, please contact:

researchsupport@kent.ac.uk

If you believe this document infringes copyright then please contact the KAR admin team with the take-down information provided at <http://kar.kent.ac.uk/contact.html>

Medtronic

Partner with a leader in RDN

Symlicity™ renal
denervation system

[Build your program](#)

Enhance your
hypertension
treatment
strategies

LETTER TO THE EDITOR

Low rates of liver injury in edoxaban users: Evidence from a territory-wide observational cohort study

Dear Editor,

We read the recent meta-analysis by Dai et al. on liver injury caused by oral anticoagulants with great interest.¹ With the widespread use of direct oral anticoagulants (DOACs) in clinical practice for the prevention of thrombotic events,^{2,3} certain concerns have been raised with its risk of inducing liver injury. Such findings were not revealed in clinical trials, with a meta-analysis demonstrating no significant associations between DOAC use and liver injury.⁴ Specifically regarding edoxaban, the landmark ENGAGE AF-TIMI 48 trial found that hepatic events, as defined by AST or ALT $\geq 3\times$ upper limit of normal or above with total bilirubin $\geq 2\times$ upper limit of normal, were comparable among the warfarin, high-dose and low-dose edoxaban groups.⁵ However, the occurrence of drug-induced liver injury for DOACs has been reported by several case reports and series in post-market settings, and has been summarized by Liakoni et al. as early in 2015.⁶ These variations in results have led to investigations of drug-induced liver injury by individual DOACs in observational studies.⁷⁻¹¹ Nevertheless, limited data are available for edoxaban, given its comparative late entry into the international market. Therefore, we conducted this territory-wide study to assess the risk of liver injury in edoxaban users who have no prior history of liver diseases. This study received approvals from Institutional Review Board of the University of Hong Kong—Hospital Authority Hong Kong West Cluster, and the Joint Chinese University of Hong Kong—Hospital Authority New Territories East Cluster Clinical Research Ethics Committee. The detailed methods are shown in the **Supplementary Appendix**.

Among the 2107 edoxaban users from January 01, 2016 to December 31, 2019 initially identified, 894 patients were excluded for the following reasons: (1) prescriptions of other anticoagulants ($n = 543$); a prior diagnosis of liver disease ($n = 16$); prior liver injury ($n = 335$) (Figure 1). Finally, 1213 edoxaban users were included (Table 1, left), with 19 developing liver injury after its prescription, which corresponded to an incidence of 1.5% (Figure 2). Univariate Cox regression identified significant predictors of liver injury (Table 1, right), which were: older age (HR: 1.32, 95% CI: [1.03, 3.29], $p = .006$), prior hemorrhagic stroke (HR: 5.24, 95% CI: [1.53, 17.99], $p = .009$), and higher levels of creatinine (HR: 1.002, 95% CI: [1.000, 1.005], $p = .021$) and urea (HR: 1.10, 95% CI: [1.04, 1.16], $p = .001$).

The main finding of this study is that there is a low rate of liver injury of 1.5% in patients receiving edoxaban therapy in Hong Kong. This extends previous findings by another local study, which reported an incidence of 1.8% for dabigatran, 2.1% for rivaroxaban, 2.5% for apixaban, with a pooled incidence rate of 2.0% for all three DOACs, as compared to 3.7% for warfarin.⁸ Edoxaban, similar to the other DOACs, has demonstrated at least non-inferiority compared to warfarin for stroke prevention in patients with atrial fibrillation.¹²⁻¹⁴ Our data provide further evidence regarding the safety of edoxaban specifically regarding hepatotoxicity. However, significant factors include advancing age, impaired renal function and prior haemorrhagic stroke. Therefore, this medication should be used in caution in such users, who should receive regular blood tests. Our findings should be validated in future observational cohort studies or clinical trials. Moreover, a follow-up analysis of the ENGAGE AF-TIMI 48 trial demonstrated that a history of liver disease did not significantly affect the efficacy of safety of edoxaban compared to warfarin and the rates of hepatic adverse events were similar between edoxaban or warfarin users.¹⁵ Future observational studies should examine the efficacy and safety outcomes in this specific subgroup to complement findings from this important trial.




FIGURE 1 Procedures of data processing

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2021 The Authors. *Clinical Cardiology* published by Wiley Periodicals LLC.

TABLE 1 Baseline characteristics of edoxaban users and Cox regression results for significant predictors of liver injury

Characteristics	Edoxaban (N = 1213) median (IQR); max; N or count (%)	HR (95% CI)	p-value
Demographics			
Baseline age, year	74 (64–82); 100; n = 1213	0.99 (0.96, 1.02)	.584
Male gender	622 (51.27%)	1.32 (1.03, 3.29)	.006**
Past comorbidities			
Respiratory	345 (28.44%)	1.04 (0.30, 3.56)	.955
Endocrine	18 (1.48%)	1.65 (0.38, 7.15)	.502
Diabetes mellitus	84 (6.92%)	0.67 (0.24, 1.87)	.449
Hypertension	440 (36.27%)	0.72 (0.17, 3.11)	.657
Gastrointestinal	170 (14.01%)	0.70 (0.16, 3.02)	.630
Hemorrhagic stroke	15 (1.23%)	5.24 (1.53, 17.99)	.009**
Congestive heart failure	28 (2.30%)	2.72 (0.36, 20.41)	.330
Gastrointestinal bleeding	86 (7.08%)	0.70 (0.09, 5.26)	.731
Medications			
Edoxaban duration, day	313 (236–424); 228; n = 1213	0.69 (0.67, 0.73)	.002**
ACEI	422 (34.78%)	0.93 (0.35, 2.44)	.874
ARB	222 (18.30%)	0.57 (0.13, 2.49)	.459
Calcium channel blockers	665 (54.82%)	0.51 (0.20, 1.30)	.160
Beta blockers	625 (51.52%)	0.75 (0.30, 1.86)	.531
Diuretics for heart failure	512 (42.20%)	1.73 (0.70, 4.28)	.232
Nitrates	272 (22.42%)	0.68 (0.20, 2.32)	.534
Antihypertensive drugs	231 (19.04%)	0.51 (0.12, 2.22)	.372
Statins and fibrates	654 (53.91%)	0.54 (0.21, 1.37)	.191
Antihyperlipidemic/Lipid-lowering drugs	646 (53.25%)	0.55 (0.22, 1.41)	.216
Laboratory tests			
Creatinine, umol/L	87 (71–111); 1192; n = 1085	1.002 (1.000, 1.005)	.021*
Urea, mmol/L	6.1 (5–8); 40.6; n = 1084	1.10 (1.04, 1.16)	.001***
Potassium, mmol/L	4.1 (3.8–4.4); 6; n = 1085	0.70 (0.25, 1.91)	.482
Sodium, mmol/L	140 (138–142); 155; n = 1085	0.93 (0.81, 1.08)	.350
Urate, mmol/L	0.41 (0.3365–0.521); 0.98; n = 207	19.51 (0.06, 6018.00)	.310
Albumin, g/L	39 (34.6–42); 51.7; n = 937	0.94 (0.87, 1.01)	.077
Protein, g/L	72 (67–76); 92.48; n = 882	0.99 (0.93, 1.06)	.817
Alkaline phosphatase, U/L	74.55 (61–92); 454; n = 936	1.00 (0.98, 1.01)	.818

* $p \leq .05$.** $p \leq .01$.*** $p \leq .001$.**CONFLICT OF INTEREST**

None.


DATA AVAILABILITY STATEMENT

The data described in this letter are available upon request to the corresponding authors.


Jiandong Zhou MSc¹
 Keith Sai Kit Leung BSc, MRSPH^{2,3}
 Dicken Kong^{3,4}

¹School of Data Science, City University of Hong Kong, Hong Kong, China

²Aston Medical School, Aston University, Birmingham, UK

Sharen Lee³
 Tong Liu MD, PhD⁵
 Abraham Ka Chung Wai MBChB, FRCEM, FRCP⁶
 Carlin Chang MBChB, MPhil, MRCP⁷
 Qingpeng Zhang PhD¹
 Gary Tse PhD, FESC, FACC, FRCP^{3,5} 

Kaplan-Meier curve: Liver injury

**FIGURE 2** Kaplan-Meier curve of liver injury in edoxaban users

³Cardiovascular Analytics Group, Laboratory of Cardiovascular Physiology, Hong Kong, China

⁴Li Ka Shing Institute of Health Sciences, The Chinese University of Hong Kong, Hong Kong, China

⁵Tianjin Key Laboratory of Ionic-Molecular Function of Cardiovascular Disease, Department of Cardiology, Tianjin Institute of Cardiology, Second Hospital of Tianjin Medical University, Tianjin, China

⁶Emergency Medicine Unit, Li Ka Shing Faculty of Medicine, The University of Hong Kong, Hong Kong, China

⁷Division of Neurology, Department of Medicine, Queen Mary Hospital, Hong Kong, China

Correspondence

Qingpeng Zhang, PhD, School of Data Science, City University of Hong Kong, Hong Kong, China.
Email: qingpeng.zhang@cityu.edu.hk

Gary Tse, PhD FRCP, Tianjin Key Laboratory of Ionic-Molecular Function of Cardiovascular Disease, Department of Cardiology, Tianjin Institute of Cardiology, Second Hospital of Tianjin Medical University, Tianjin 300211, China.
Email: garytse86@gmail.com

Jiandong Zhou and Keith Sai Kit Leung are considered as joint first authors.

ORCID

Gary Tse  <https://orcid.org/0000-0001-5510-1253>

REFERENCES

- Dai Q, Deng X, Zhou L, Zhang L, Xiao X, Liao Y. Real-world use of non-vitamin K antagonist oral anticoagulant in atrial fibrillation patients with liver disease: a meta-analysis. *Clin Cardiol.* 2020;43(7):676-683.
- Li KHC, Jesuthasan A, Kui C, Davies R, Tse G, Lip GYH. Acute ischemic stroke management: concepts and controversies. A narrative review. *Expert Rev Neurother.* 2020;21(1):65-79.
- Wang Y, Lv H, Li D, et al. Efficacy and safety of direct Oral anticoagulants for secondary prevention of cancer-associated thrombosis: a systematic review and meta-analysis of randomized controlled trials and prospective cohort studies. *Front Pharmacol.* 2019;10:773.
- Caldeira D, Barra M, Santos AT, et al. Risk of drug-induced liver injury with the new oral anticoagulants: systematic review and meta-analysis. *Heart.* 2014;100(7):550-556.
- Giugliano RP, Ruff CT, Braunwald E, et al. Edoxaban versus warfarin in patients with atrial fibrillation. *N Engl J Med.* 2013;369(22):2093-2104.
- Liakoni E, Ratz Bravo AE, Krahenbuhl S. Hepatotoxicity of new oral anticoagulants (NOACs). *Drug Saf.* 2015;38(8):711-720.
- Maura G, Bardou M, Billionnet C, Weill A, Drouin J, Neumann A. Oral anticoagulants and risk of acute liver injury in patients with non-valvular atrial fibrillation: a propensity-weighted nationwide cohort study. *Sci Rep.* 2020;10(1):11624.
- Zhao J, Blais JE, Chui CSL, et al. Association between nonvitamin K antagonist oral anticoagulants or warfarin and liver injury: a cohort study. *Am J Gastroenterol.* 2020;115(9):1513-1524.

9. Douros A, Azoulay L, Yin H, Suissa S, Renoux C, Non-Vitamin K. Antagonist oral anticoagulants and risk of serious liver injury. *J Am Coll Cardiol*. 2018;71(10):1105-1113.
10. Alonso A, MacLehose RF, Chen LY, et al. Prospective study of oral anticoagulants and risk of liver injury in patients with atrial fibrillation. *Heart*. 2017;103(11):834-839.
11. Bjornsson HK, Gudmundsson DO, Bjornsson ES. Liver injury caused by oral anticoagulants: a population-based retrospective cohort study. *Liver Int*. 2020;40(8):1895-1900.
12. Hanley CM, Kowey PR. Are the novel anticoagulants better than warfarin for patients with atrial fibrillation? *J Thorac Dis*. 2015;7(2):165-171.
13. Sawant AC, Kumar A, McCray W, et al. Superior safety of direct oral anticoagulants compared to warfarin in patients with atrial fibrillation and underlying cancer: a national veterans affairs database study. *J Geriatr Cardiol*. 2019;16(9):706-709.
14. Mongkhon P, Fanning L, Lau WCY, et al. Oral anticoagulant and reduced risk of dementia in patients with atrial fibrillation: a population-based cohort study. *Heart Rhythm*. 2020;17(5 Pt A):706-713.
15. Qamar A, Antman EM, Ruff CT, et al. Edoxaban versus warfarin in patients with atrial fibrillation and history of liver disease. *J Am Coll Cardiol*. 2019;74(2):179-189.

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section at the end of this article.