

Kent Academic Repository

Full text document (pdf)

Citation for published version

Tamè, Luigi and Holmes, Nicholas P. (2013) The role of primary somatosensory cortex in tactile detection and discrimination: fMRI-guided TMS investigations. In: VII Edition, Concept Action and Objects, 23 May - 26 May 2013, Rovereto, Italy.

DOI

Working Poster No. 10

Link to record in KAR

<https://kar.kent.ac.uk/98842/>

Document Version

Presentation

Copyright & reuse

Content in the Kent Academic Repository is made available for research purposes. Unless otherwise stated all content is protected by copyright and in the absence of an open licence (eg Creative Commons), permissions for further reuse of content should be sought from the publisher, author or other copyright holder.

Versions of research

The version in the Kent Academic Repository may differ from the final published version.

Users are advised to check <http://kar.kent.ac.uk> for the status of the paper. **Users should always cite the published version of record.**

Enquiries

For any further enquiries regarding the licence status of this document, please contact:

researchsupport@kent.ac.uk

If you believe this document infringes copyright then please contact the KAR admin team with the take-down information provided at <http://kar.kent.ac.uk/contact.html>

Role of primary somatosensory cortex in tactile detection and discrimination: fMRI-guided TMS investigations




Introduction

In the present study in **seven experiments** we investigated the contribution of the **primary somatosensory cortex (SI)** to the perception of tactile stimuli at the fingers in **discrimination** and **detection** task. We combined the **QUEST** threshold estimation method with **MRI-guided** single and double pulse **TMS (sp/dpTMS)**.

ZIFC

Design

Yes-no




fMRI-guided TMS


TMS Stimulation conditions

Participants (**N=20**) underwent a series of fMRI scans (localisers) to produce somatotopic maps of SI and SII cortices.

The tactile stimuli were **200Hz sinusoidal vibrations** applied to the **index or middle fingers** for **50 ms**.




The peak activation focus in SI was lateral to the hand motor area, and on the posterior bank of the post-central gyrus was used to position the TMS (~2.5 cm from the target area) during a modified version of the behavioural task.



Results

Data values represent the mean (+/- SE) of the last point of the series of 48 trials used to determine the threshold (82 % correct).



Conclusions

Frequency discrimination thresholds at the fingers are increased by double-pulse TMS over SI in comparison to SMG. By contrast, **detection** thresholds at the fingers are not increased by either single or double pulse TMS over SI. This suggests that **SI is required for discrimination**, but **may not be for detection**, of tactile stimuli at the fingers.