



INSTITUT DE FRANCE
Académie des sciences

Comptes Rendus

Physique

Atac Imamoglu, Ovidiu Cotlet and Richard Schmidt

Exciton–polarons in two-dimensional semiconductors and the Tavis–Cummings model


Volume 22, issue S4 (2021), p. 89-96

<<https://doi.org/10.5802/crphys.47>>

Part of the Special Issue: Recent advances in 2D material physics

Guest editors: Xavier Marie (INSA Toulouse, Université Toulouse III Paul Sabatier, CNRS, France) and Johann Coraux (Institut Néel, Université Grenoble Alpes, CNRS, France)

© Académie des sciences, Paris and the authors, 2021.
Some rights reserved.

 This article is licensed under the
CREATIVE COMMONS ATTRIBUTION 4.0 INTERNATIONAL LICENSE.
<http://creativecommons.org/licenses/by/4.0/>



*Les Comptes Rendus. Physique sont membres du
Centre Mersenne pour l'édition scientifique ouverte*
www.centre-mersenne.org



Recent advances in 2D material physics / *Physique des matériaux bidimensionnels*

Exciton–polarons in two-dimensional semiconductors and the Tavis–Cummings model

Exciton–polarons dans des semi-conducteurs bidimensionnels et le modèle de Tavis–Cummings

Atac Imamoglu^{*, a}, Ovidiu Cotlet^a and Richard Schmidt^{b, c}

^a Institute for Quantum Electronics, ETH Zürich, CH-8093 Zürich, Switzerland

^b Max Planck Institute of Quantum Optics, 85748 Garching, Germany

^c Munich Center for Quantum Science and Technology, Schellingstrasse 4, 80799 München, Germany

E-mails: imamoglu@phys.ethz.ch (A. Imamoglu), ocotlet@gmail.com (O. Cotlet), richard.schmidt@mpq.mpg.de (R. Schmidt)

Abstract. The elementary optical excitations of a two-dimensional electron or hole system have been identified as exciton-Fermi-polarons. Nevertheless, the connection between the bound state of an exciton and an electron, termed trion, and exciton–polarons is subject of ongoing debate. Here, we use an analogy to the Tavis–Cummings model of quantum optics to show that an exciton–polaron can be understood as a hybrid quasiparticle—a coherent superposition of a bare exciton in an unperturbed Fermi sea and a bright collective excitation of many trions. The analogy is valid to the extent that the Chevy Ansatz provides a good description of dynamical screening of excitons and provided the Fermi energy is much smaller than the trion binding energy. We anticipate our results to bring new insight that could help to explain the striking differences between absorption and emission spectra of two-dimensional semiconductors.

Résumé. Les excitations optiques élémentaires d'un système bidimensionnel d'électrons ou de trous ont été identifiées comme des exciton-Fermi-polarons. Néanmoins, la connexion entre l'état lié d'un exciton et d'un électron, appelé trion, et les exciton–polarons fait l'objet d'un débat permanent. Ici, nous utilisons une analogie avec le modèle de Tavis–Cummings de l'optique quantique pour montrer qu'un exciton–polaron peut être compris comme une quasi-particule hybride — une superposition cohérente d'un exciton nu dans une mer de Fermi non perturbée et une excitation collective brillante de nombreux trions. L'analogie est valable dans la mesure où l'Ansatz de Chevy fournit une bonne description de l'écrantage dynamique des excitons et à condition que l'énergie de Fermi soit beaucoup plus petite que l'énergie de liaison des trions. Nous espérons que nos résultats apporteront de nouvelles connaissances qui pourraient aider à expliquer les différences frappantes entre les spectres d'absorption et d'émission des semi-conducteurs bidimensionnels.

* Corresponding author.

Keywords. Exciton–polarons, Two-dimensional semiconductors, Tavis–Cummings model, Quantum optics, Many-body physics.

Mots-clés. Exciton–polarons, Semi-conducteurs bidimensionnels, Modèle de Tavis–Cummings, Optique quantique, Physique des corps multiples.

Available online 8th February 2021

Two-dimensional (2D) semiconductors [1] such as monolayers of transition metal dichalcogenides (TMD) have emerged as an exciting platform for investigating many-body physics and strong correlations [2, 3]. Due to strong Coulomb interactions, the optical excitation spectra of neutral TMDs are dominated by tightly bound excitons. The small Bohr radius a_B of TMD excitons leads to ultra-short radiative decay rates, in turn ensuring that in clean samples the exciton resonance is predominantly radiatively broadened [4, 5]. Introduction of itinerant electrons (holes) into the monolayer dramatically modifies the nature of the optical spectra and leads to the emergence of a new absorption/reflection resonance near the energy of the three-body bound—trion—state of an exciton and an electron (hole) [3]. It has been recently shown that the relevant elementary optical excitations in this limit are excitons that are dynamically dressed by Fermi sea electrons (holes), termed attractive or repulsive exciton–polarons [6–8].

The connection between attractive exciton polaron (AP) and trion excitations has been the subject of ongoing debate. The oscillator strength for optical generation of a single isolated trion by diffraction limited resonant light is $f_T \sim f_x(k_{\text{ph}}a_T)^2$, where f_x is the exciton oscillator strength, a_T is the trion Bohr radius and $k_{\text{ph}} = E_T/(\hbar c)$ is the momentum of a photon resonant with the trion transition (E_T). In the presence of a finite electron density n_e , it was predicted that the oscillator strength would be proportional to n_e [9–11], which is precisely the scaling one gets for AP oscillator strength for low n_e . The goal of this Letter is to shed new light on the relation between AP and trion excitations by making use of the Tavis–Cummings (TC) model of quantum optics [12].

1. Tavis–Cummings model

We start our analysis by recalling that the TC model describes an ensemble of N_a two-level atoms with an energy splitting ω_{eg} between the ground ($|g\rangle$) and excited ($|e\rangle$) states coupled to a single cavity mode [12] of frequency ω_c . The interaction Hamiltonian of this system is given by

$$H_{\text{int}} = \sum_i g_c^i (\sigma_{eg}^i a_c + \text{h.c.}), \quad (1)$$

where a_c^\dagger is the cavity creation operator and σ_{eg}^i denotes the raising operator of the i th two-level atom. The cavity mode and atoms are coupled by the single-atom coupling rate g_c^i which we, for simplicity, assume to be identical for all atoms in the following, $\forall i$ $g_c^i = g_c$.

The lowest energy excitation spectrum of the TC model consists of $N_a - 1$ dark states at energy ω_{eg} and two polariton states that can be expressed as a superposition of bare cavity and atomic excitations. We refer to the lowest energy excited state as the lower polariton (LP) state, which can be expressed as

$$|\Phi_{\text{LP}}\rangle = \left(\alpha a_c^\dagger + \beta \sum_{i=1}^{N_a} \sigma_{eg}^i \right) |0\rangle. \quad (2)$$

Here the state $|0\rangle$ describes the vacuum of the cavity and all atoms in their ground state. We consider the case where the cavity frequency ω_c is blue-detuned with respect to the atomic transition ω_{eg} by a detuning $\Delta = \omega_c - \omega_{eg}$.

In the limit when the detuning Δ is large compared to $g_c\sqrt{N_a}$, as well as the cavity (κ_c) and atomic (Γ_{eg}) decay rates, one finds

$$\alpha = g_c\sqrt{N_a}/\Delta. \quad (3)$$

For this parameter range the LP state is a predominantly bright (symmetric) excitation of N_a atoms, together with a small probability amplitude (α) for a single cavity-photon excitation. The expression for α shows the well-known collective enhancement of cavity-atom coupling from g_c to $g_c\sqrt{N_a}$. As a result of this enhanced coupling, the LP state is red-shifted in energy as compared to the $N_a - 1$ atomic dark states by an energy

$$E_{\text{LP}} = \alpha^2 \Delta = g_c^2 N_a / \Delta, \quad (4)$$

provided that the cavity decay rate $\kappa_c \ll g_c\sqrt{N_a}$ (strong-coupling limit). It is important to emphasize that the LP resonance is insensitive to inhomogeneous broadening of atomic energy levels, provided that this broadening is smaller than E_{LP} .

Our simplified discussion of the TC model did not account for the spontaneous emission of the atoms: if the cavity-mode area is large compared to the square of the cavity-mode wavelength λ_c , the total spontaneous emission rate of the atoms is hardly modified. In the opposite limit where cavity-Purcell enhancement dominates the atomic decay, the atomic decay takes place predominantly through a two-step process where coherent excitation exchange between the atoms and the cavity is followed by cavity decay.

The TC model can be extended to a two-dimensional setting by assuming that the Fabry–Perot cavity consists of two parallel mirrors and the atoms are embedded in a 2D lattice with a period $d \ll \lambda_c$. In this case, the in-plane momentum of the polariton excitations constitutes a good quantum number. One may then define the collective atomic raising operator corresponding to an excitation with momentum \mathbf{k} ,

$$\sigma_{eg}(\mathbf{k}) = \sum_j \sigma_{eg}^j e^{i\mathbf{k}\cdot\mathbf{R}_j}, \quad (5)$$

where the \mathbf{R}_j denote the atomic lattice sites. The ansatz for the lower polariton branch of the two-dimensional TC model is then finally obtained by replacing a_c by $a_c(\mathbf{k})$ and $\sum_j \sigma_{eg}^j$ with $\sigma_{eg}(\mathbf{k})$ in (2).

2. Exciton–polarons

The Hamiltonian describing the interacting exciton–electron system in a TMD monolayer can be written as [7, 8, 13]

$$H_{xe} = \sum_{\mathbf{k}} \omega_{\mathbf{k}} x_{\mathbf{k}}^\dagger x_{\mathbf{k}} + \sum_{\mathbf{k}} \epsilon_{\mathbf{k}} e_{\mathbf{k}}^\dagger e_{\mathbf{k}} + \frac{v}{\mathcal{A}} \sum_{\mathbf{k}, \mathbf{k}', \mathbf{q}} x_{\mathbf{k}+\mathbf{q}}^\dagger x_{\mathbf{k}} e_{\mathbf{k}'-\mathbf{q}}^\dagger e_{\mathbf{k}'}. \quad (6)$$

Here, $e_{\mathbf{k}}$ and $x_{\mathbf{k}}$ denote the annihilation operators of electrons and excitons with momentum \mathbf{k} , respectively. The electronic dispersion is $\epsilon_{\mathbf{k}} = \mathbf{k}^2 / (2m_e)$. The exciton dispersion $\omega_{\mathbf{k}} = \mathbf{k}^2 / (2m_x)$ is defined with respect to the exciton energy E_x which we set to zero.

The contact coupling constant v characterizes the short-range interaction between excitons and electrons and it is related to the trion binding energy by the Lippmann–Schwinger equation

$$v^{-1} = -\frac{1}{\mathcal{A}} \sum_{|\mathbf{k}| < \Lambda} \frac{1}{E_T + \omega_{\mathbf{k}} + \epsilon_{\mathbf{k}}}. \quad (7)$$

Here $E_T = \hbar^2 / (2ma_T^2)$ denotes the trion binding energy, and $1/m = 1/m_x + 1/m_e$ is the reduced mass. As evident from (7), the interaction is regularized by a UV cutoff Λ which can physically be related to the inverse Bohr radius of the exciton. However, assuming that the exciton Bohr radius is the smallest length-scale in the problem, one may take the limit $\Lambda \rightarrow \infty$ at the end of the calculation.

It has been shown [14, 15] that the eigenstates of the interacting polariton–electron system can be accurately described using the variational Chevy ansatz [16]

$$|\Psi_{\text{AP},\mathbf{p}}\rangle = \left(\phi_{\mathbf{p}} x_{\mathbf{p}}^{\dagger} + \sum_{\mathbf{k}\mathbf{q}} \phi_{\mathbf{k}\mathbf{q}}^{\mathbf{p}} x_{\mathbf{p}+\mathbf{q}-\mathbf{k}}^{\dagger} e_{\mathbf{k}}^{\dagger} e_{\mathbf{q}} \right) |\Phi\rangle \quad (8)$$

$$= \left(\phi_{\mathbf{p}} x_{\mathbf{p}}^{\dagger} + \sum_{\nu\mathbf{q}} \eta_{\nu\mathbf{q}}^{\mathbf{p}} t_{\nu\mathbf{p}+\mathbf{q}}^{\dagger} e_{\mathbf{q}} \right) |\Phi\rangle, \quad (9)$$

which expands the wavefunction in excitations of the non-interacting ground state $|\Phi\rangle$ of the electron system in the TMD monolayer. The variational ground state $|\Psi_{\text{AP},\mathbf{p}}\rangle$ is the so-called attractive polaron (AP) of momentum \mathbf{p} which describes the exciton as a quasiparticle dressed by the attractive interactions with the Fermi sea of electrons.

In the second line of (8), we have introduced the creation operator $t_{\nu\mathbf{l}}^{\dagger}$ that generates a composite trion state of center-of-mass momentum \mathbf{l} in an internal state ν :

$$t_{\nu\mathbf{l}}^{\dagger} = \sum_{\mathbf{k}} \chi_{\nu\mathbf{k}}^{\mathbf{l}} x_{\mathbf{l}-\mathbf{k}}^{\dagger} e_{\mathbf{k}}^{\dagger}. \quad (10)$$

Here the states ν denote both bound trion as well as electron–exciton scattering states [13]. Importantly, while the sum over ν in (8) runs over all these composite states, for excitations around the AP resonance the bound trion state is the most relevant one ($\nu = 0$). For a zero-momentum AP, Equation (8) can therefore be expressed as

$$|\Psi_{\text{AP},\mathbf{p}=0}\rangle \approx \left(\phi_0 x_0^{\dagger} + \chi_0 \sum_{\mathbf{q}} \tilde{\eta}_{\mathbf{q}} t_{\mathbf{q}}^{\dagger} e_{\mathbf{q}} \right) |\Phi\rangle, \quad (11)$$

where $\phi_0 \equiv \phi_{\mathbf{p}=0}$, $\chi_0 \tilde{\eta}_{\mathbf{q}} \equiv \eta_{\nu=0,\mathbf{q}}^{\mathbf{p}=0}$. Equation (11) can be interpreted as describing a quasiparticle where an exciton with momentum $\mathbf{p} = 0$ is hybridized with a collective optical excitation of all electrons in the Fermi sea. As we will show below, for low electron densities, where the Fermi momentum k_F satisfies $k_F^2 a_T^2 \ll 1$, an AP excitation has a small probability amplitude (ϕ_0) for a bare exciton excitation.

3. Correspondence of the TC and exciton–polaron models

The forms of the LP and AP wavefunctions given in (2) and (11) already hint at a one-to-one correspondence between the elementary excitations occurring in rather different experimental systems. The equivalence of these two models can be clarified by identifying the correspondence between the operators

$$\begin{aligned} \sigma_{eg}^i &\iff \tilde{\eta}_{\mathbf{q}} t_{\mathbf{q}}^{\dagger} e_{\mathbf{q}} \\ a_c^{\dagger} &\iff x_0^{\dagger} \end{aligned}$$

and key parameters

$$\begin{aligned} \omega_c &\iff E_x \\ \omega_{eg} &\iff E_x - E_T \\ \Delta &\iff E_T \\ N_a &\iff N_e = \mathcal{A} n_e = \mathcal{A} k_F^2 / (4\pi) \\ E_{\text{LP}} &\iff \delta E^{\text{AP}}, \end{aligned}$$

where $n_e = k_F^2/4\pi$ denotes the electron density in a single valley, \mathcal{A} is the area of the TMD monolayer, $E_F = \hbar^2 k_F^2/2m_e$ is the Fermi energy, and $\delta E^{\text{AP}} = -E_T - E_{\mathbf{p}=0}^{\text{AP}}$ is the energy difference between the AP and trion resonances. The wave functions of the LP and AP are related by

$$\begin{aligned}\beta &\iff \chi_0 \\ \alpha &\iff \phi_0.\end{aligned}$$

From this correspondence between the two models one would expect δE^{AP} to satisfy an expression similar to the one for E_{LP} in (4). In fact, without calculation, the correspondence would directly imply that $\delta E^{\text{AP}} = \phi_0^2 E_T$. In the following, we demonstrate that this is indeed the case by an explicit calculation.

To this end, we use the electron–exciton scattering T -matrix that accounts for effects of the finite electron density [6, 17, 18],

$$T(\mathbf{p}, \omega)^{-1} = v^{-1} - \frac{1}{\mathcal{A}} \sum_{|\mathbf{k}| > k_F} \frac{1}{\omega - \epsilon_{\mathbf{k}} - \omega_{\mathbf{p}-\mathbf{k}} + i0^+}, \quad (12)$$

where \mathbf{p} and ω denote the total momentum and energy of the exciton and the electron. The exciton self-energy is obtained from the T -matrix as:

$$\Sigma_x(\mathbf{p}, \omega) = \frac{1}{\mathcal{A}} \sum_{|\mathbf{q}| < k_F} T(\mathbf{p} + \mathbf{q}, \omega + \epsilon_{\mathbf{q}}). \quad (13)$$

The quasi-particle weight $|\phi_{\mathbf{p}}|^2$ in turn is given by

$$|\phi_{\mathbf{p}}|^2 = \left(1 - \frac{\partial}{\partial \omega} [\Sigma_x(\mathbf{p}, \omega)]_{\omega = E_{\mathbf{p}}^{\text{AP}}} \right)^{-1}, \quad (14)$$

where $E_{\mathbf{p}}^{\text{AP}}$ denotes the energy of the AP at momentum \mathbf{p} , as determined by the solution of the Dyson equation

$$[\omega - \omega_{\mathbf{p}} - \Sigma_x(\mathbf{p}, \omega)]|_{\omega = E_{\mathbf{p}}^{\text{AP}}} = 0. \quad (15)$$

We now focus on zero momentum $\mathbf{p} = 0$. To obtain an analytical expression for $\phi_{\mathbf{p}=0}$, we consider the low electron density limit where $E_T \gg E_F$. As shown in Appendix A in this limit the exciton self-energy can be approximated by

$$\Sigma_x(\omega) = \Sigma_x(\mathbf{p} = 0, \omega) \simeq n_e T_{xe}(0, \omega), \quad (16)$$

where the *two-body* T -matrix is given by

$$T_{xe}(0, \omega) = \frac{2\pi\hbar^2}{m} \frac{1}{\ln\left[\frac{-E_T}{\omega+i0^+}\right] + i\pi} \quad (17)$$

$$\simeq \frac{2\pi\hbar^2}{m} \frac{E_T}{E_T + \omega}. \quad (18)$$

In the second line the T -matrix is evaluated for energies ω close to the pole at $\omega = -E_T$, i.e. for the condition $|E_T + \omega| \ll E_T$. Using this approximate expression for $T_{xe}(0, \omega)$ evaluated at $\omega = E_{\mathbf{p}=0}^{\text{AP}}$, we obtain

$$|\phi_{\mathbf{p}=0}|^2 \simeq \frac{(\delta E^{\text{AP}})^2}{E_T} \frac{m}{n_e 2\pi\hbar^2}. \quad (19)$$

To express $\phi_{\mathbf{p}=0}$ in terms of the Fermi momentum k_F and a_T , we use the fact that the AP resonance energy $E_{\mathbf{p}=0}^{\text{AP}}$ is given by the lowest energy pole of the exciton propagator, i.e. the solution of (15). In the limit $|E_T - E_{\mathbf{p}=0}^{\text{AP}}| \ll E_T$, we obtain

$$\delta E^{\text{AP}} = -E_T - E_{\mathbf{p}=0}^{\text{AP}} = n_e \frac{2\pi\hbar^2}{m} = \frac{m_e}{m} E_F. \quad (20)$$

Substituting for δE^{AP} in (19), we thus arrive at the expression,

$$\phi_{\mathbf{p}=0}^2 = k_F^2 a_T^2. \quad (21)$$

Equation (21) shows that the AP resonance has a collectively enhanced oscillator strength $f_{\text{AP}} = k_F^2 a_T^2 f_x$. Finally, using (21) and (20), we find $\delta E^{\text{AP}} = \phi_0^2 E_T$, verifying the perfect correspondence between the LP and AP resonances of the two models.

4. Discussion of limitations

We emphasize that despite the remarkable correspondence between the LP and AP resonances, the analogy between exciton–polarons and the TC model breaks down for the repulsive polaron branch owing to the logarithmic energy dependence of the exciton–electron T -matrix, and consequently of $\Sigma_{xe}(\mathbf{p}, \omega)$. In contrast, the photon self-energy in the TC model is given simply by $g_c^2 N_a / (E - \omega_{eg})$.

The approximation $\Sigma_x(\omega, \mathbf{p} = \mathbf{0}) \simeq n_e T_{xe}(0, \omega)$ we used is valid either in the limit of low electron density n_e or if the electron mass were much larger than the exciton mass; this would be the case if the monolayer is embedded in a 2D cavity with a small cavity length where the elementary excitations are exciton–polaritons with a very light effective mass. In the absence of a cavity, however, the fermionic nature of the electrons leads to a broadening of the trion transition of order E_F , which is in turn comparable to the shift of the AP resonance $\phi_{\mathbf{p}=0}^2 E_T$. Consequently, and unlike in the ideal TC model, the trion–hole pairs that contribute to the AP resonance do not have identical energy. Nevertheless, within the Chevy description, the AP resonance is insensitive to this broadening even for finite n_e and is broadened exclusively by radiative decay arising from its bare exciton character.

The analogy we developed uses the simplest Ansatz for describing correlated exciton–electron states. In particular, this Chevy Ansatz does not capture the screening of trions by the Fermi sea of electrons (for a discussion in the context of ultracold atoms see [19]); indeed, neglecting Coulomb repulsion between electrons, it has been shown theoretically that dynamically screened trions have lower energy than AP provided $k_F a_T \leq 0.1$ [20–23].

Arguably, the most important difference between the exciton–electron system and the 2D TC model is the drastic reduction of coupling of high momentum collective atomic excitations to the corresponding cavity modes in the TC model. Since the effective cavity photon mass is orders of magnitude smaller than that describing the collective atomic excitations, only the bright symmetric atomic excitation couples appreciably to the cavity mode. In the limit of a 0D cavity, this description becomes exact and justifies referring to the $N_a - 1$ antisymmetric excitations with energy ω_{eg} as dark states.

Due to the comparable effective masses of the exciton, electron and the trion, each trion–hole pair state with total momentum \mathbf{p} ($t_{v\mathbf{p}+\mathbf{q}}^\dagger e_{\mathbf{q}}$) hybridizes with the exciton mode $x_{\mathbf{p}}$. Therefore, in the limit $k_F a_T \ll 1$, only polaron states couple to light. Moreover, this coupling is proportional to the quasi-particle weight $|\phi_{\mathbf{p}}|^2$, i.e. it is exclusively due to the bare-exciton character of the polaron. This argument of course does not preclude possible observation of single trion decay in a nonequilibrium experiment such as photoluminescence [24]; if we ignore the dynamical screening of trions, each single trion–hole pair state is a superposition of AP eigenstates. Even in the presence of a finite electron density, an optically generated electron–hole pair could form a trion with a single electron and subsequently decay by emitting a photon before the excitation is spread throughout the sample to form the AP state at $\mathbf{p} \simeq 0$.

In summary, we have developed an analogy between the interacting exciton–electron problem in 2D materials and the TC model. Our work shows that the AP resonance can be described as a hybridization of collective trion–hole pair excitations with excitons, which in turn ensures their

enhanced coupling to external light fields. We emphasize that a simplistic picture describing the total optical absorption strength as having contributions from all $N_e = A_L k_F^2 / (4\pi)$ electrons within the excitation spot with area A_L yields a similar result as what we obtain using the polaron model. However, such a description would erroneously predict line broadening by E_F and misses out on the energy shift of the AP resonance (δE^{AP}) from the single trion energy. Finally, the collective nature of the polaron excitation with minimal disturbance of each electron ensures that polaron excitation constitutes an invaluable nondestructive spectroscopic tool for investigating strongly correlated states of electrons.

Acknowledgments

This work was initiated during an informal meeting with M. Glazov, M. Semina, B. Urbaszek, X. Marie, T. Amand, A. Högele and M. Kroner. We thank J. v. Milczewski for insightful discussions, and we particularly acknowledge many useful discussions with M. Glazov. This work was supported by the Swiss National Science Foundation (SNSF) under Grant No. 200021-178909/1. RS is supported by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany's Excellence Strategy – EXC-2111 – 390814868.

Appendix A. Exciton self-energy at low doping

As shown in [25] the solution to the variational ansatz is equivalent to a non-selfconsistent resummation of ladder diagrams. Thus we may express our results by using the language of many-body field theory in terms of T -matrices and selfenergies. Generalizing [17] to the mass-imbalanced case one finds the expression for the exciton selfenergy

$$\begin{aligned} \Sigma_x(\mathbf{p} = \mathbf{0}, \omega) &= \int_{|\mathbf{q}| < k_F} \frac{d^2 q}{(2\pi)^2} T_{xe} \\ &\times \left[\frac{1}{2} \left(\omega + \epsilon_{\mathbf{q}} + \omega_{\mathbf{q}} - 2\epsilon_{\mathbf{q}}^{\text{Tot}} - \epsilon_F^R + i0^+ - \sqrt{(\omega + \epsilon_{\mathbf{q}} - \omega_{\mathbf{q}} - \epsilon_F^R)^2 - 4\epsilon_F^x \omega_{\mathbf{q}} - i0^+} \right) \right], \end{aligned} \quad (\text{A1})$$

where we define $\epsilon_{\mathbf{q}}^{\text{Tot}} = \mathbf{q}^2 / (m_x + m_e)$, $\epsilon_F^x = k_F^2 / (2m_x)$, and $\epsilon_F^R = k_F^2 / (2m)$. Since the integral extends only up to $|\mathbf{q}| < k_F$ and the pole of the polaron will be in the vicinity of $\omega \sim E_T$ at low electron density we may expand the square root in (A1) to obtain

$$\Sigma_x(\mathbf{0}, \omega) \approx \int_{|\mathbf{q}| < k_F} \frac{d^2 q}{(2\pi)^2} T_{xe} \left[\omega + \epsilon_{\mathbf{q}} - \epsilon_{\mathbf{q}}^{\text{Tot}} - \epsilon_F^R + i0^+ \right]. \quad (\text{A2})$$

Inserting the expression for T_{xe} given by (17) yields

$$\Sigma_x(\mathbf{0}, \omega) \approx \frac{2\pi}{m} \int_0^{k_F} \frac{dq q}{2\pi} \frac{1}{\ln \left[\frac{E_T}{\omega - \epsilon_F^R + \gamma q^2 + i0^+} \right] + i\pi} \quad (\text{A3})$$

where we define $\gamma = 1/2m_e - 1/2m_{\text{Tot}}$ with $m_{\text{Tot}} = m_e + m_x$.

Equation (A3) can be integrated analytically and gives

$$\Sigma_x(\mathbf{0}, \omega) \approx \frac{E_T}{2m\gamma} \left[\text{li} \left(\frac{-\omega + \epsilon_F^R - \gamma k_F^2}{E_T} \right) - \text{li} \left(\frac{-\omega + \epsilon_F^R}{E_T} \right) \right] \quad (\text{A4})$$

where $\text{li}(x)$ is the logarithmic integral function. Expanding to second order in k_F finally yields

$$\Sigma_x(\mathbf{0}, \omega) \approx -\frac{k_F^2}{2m} \frac{1}{\ln \left(\frac{-\omega}{E_T} \right)} = n_e T_{xe}(\mathbf{0}, \omega) \quad (\text{A5})$$

which proves (16).

References

- [1] X. Xu, W. Yao, D. Xiao, T. F. Heinz, “Spin and pseudospins in layered transition metal dichalcogenides”, *Nat. Phys.* **10** (2014), p. 343-350.
- [2] J. R. Schaibley, H. Yu, G. Clark, P. Rivera, J. S. Ross, K. L. Seyler, W. Yao, X. Xu, “Valleytronics in 2d materials”, *Nat. Rev. Mater.* **1** (2016), article no. 16055.
- [3] G. Wang, A. Chernikov, M. M. Glazov, T. F. Heinz, X. Marie, T. Amand, B. Urbaszek, “Colloquium: Excitons in atomically thin transition metal dichalcogenides”, *Rev. Mod. Phys.* **90** (2018), article no. 021001.
- [4] P. Back, S. Zeytinoglu, A. Ijaz, M. Kroner, A. Imamoglu, “Realization of an electrically tunable narrow-bandwidth atomically thin mirror using monolayer MoSe₂”, *Phys. Rev. Lett.* **120** (2018), article no. 037401.
- [5] G. Scuri, Y. Zhou, A. A. High, D. S. Wild, C. Shu, K. De Greve, L. A. Jauregui, T. Taniguchi, K. Watanabe, P. Kim, M. D. Lukin, H. Park, “Large excitonic reflectivity of monolayer MoSe₂ encapsulated in hexagonal boron nitride”, *Phys. Rev. Lett.* **120** (2018), article no. 037402.
- [6] R. A. Suris, “Correlation between trion and hole in Fermi distribution in process of trion photo-excitation in doped QWs”, in *Optical Properties of 2D Systems with Interacting Electrons*, Springer, 2003, p. 111-124.
- [7] M. Sidler, P. Back, O. Cotlet, A. Srivastava, T. Fink, M. Kroner, E. Demler, A. Imamoglu, “Fermi polaron-polaritons in charge-tunable atomically thin semiconductors”, *Nat. Phys.* **13** (2016), no. 10, p. 255-261.
- [8] D. K. Efimkin, A. H. MacDonald, “Many-body theory of trion absorption features in two-dimensional semiconductors”, *Phys. Rev. B* **95** (2017), article no. 035417.
- [9] G. V. Astakhov, V. P. Kochereshko, D. R. Yakovlev, W. Ossau, J. Nürnberger, W. Faschinger, G. Landwehr, “Oscillator strength of trion states in ZnSe-based quantum wells”, *Phys. Rev. B* **62** (2000), p. 10345-10352.
- [10] A. Esser, R. Zimmermann, E. Runge, “Theory of trion spectra in semiconductor nanostructures”, *Phys. Status Solidi (b)* **227** (2001), no. 2, p. 317-330.
- [11] G. V. Astakhov, V. P. Kochereshko, D. R. Yakovlev, W. Ossau, J. Nürnberger, W. Faschinger, G. Landwehr, T. Wojtowicz, G. Karczewski, J. Kossut, “Optical method for the determination of carrier density in modulation-doped quantum wells”, *Phys. Rev. B* **65** (2002), article no. 115310.
- [12] M. Tavis, F. W. Cummings, “Exact solution for an n -molecule—radiation-field Hamiltonian”, *Phys. Rev.* **170** (1968), p. 379-384.
- [13] C. Fey, P. Schmelcher, A. Imamoglu, R. Schmidt, “Theory of exciton-electron scattering in atomically thin semiconductors”, *Phys. Rev. B* **101** (2020), article no. 195417.
- [14] R. Combescot, S. Giraud, “Normal state of highly polarized Fermi gases: Full many-body treatment”, *Phys. Rev. Lett.* **101** (2008), article no. 050404.
- [15] C. Trefzger, Y. Castin, “Impurity in a Fermi sea on a narrow feshbach resonance: A variational study of the polaronic and dimeronic branches”, *Phys. Rev. A* **85** (2012), article no. 053612.
- [16] F. Chevy, “Universal phase diagram of a strongly interacting Fermi gas with unbalanced spin populations”, *Phys. Rev. A* **74** (2006), article no. 063628.
- [17] R. Schmidt, T. Enss, V. Pietilä, E. Demler, “Fermi polarons in two dimensions”, *Phys. Rev. A* **85** (2012), article no. 021602.
- [18] O. Cotlet, F. Pientka, R. Schmidt, G. Zarand, E. Demler, A. Imamoglu, “Transport of neutral optical excitations using electric fields”, *Phys. Rev. X* **9** (2019), article no. 041019.
- [19] M. Punk, P. T. Dumitrescu, W. Zwerger, “Polaron-to-molecule transition in a strongly imbalanced Fermi gas”, *Phys. Rev. A* **80** (2009), no. 5, article no. 053605.
- [20] M. M. Parish, “Polaron-molecule transitions in a two-dimensional Fermi gas”, *Phys. Rev. A* **83** (2011), article no. 051603.
- [21] M. M. Parish, J. Levinsen, “Highly polarized Fermi gases in two dimensions”, *Phys. Rev. A* **87** (2013), article no. 033616.
- [22] P. Kroiss, L. Pollet, “Diagrammatic Monte Carlo study of quasi-two-dimensional Fermi polarons”, *Phys. Rev. B* **90** (2014), article no. 104510.
- [23] J. Vlietinck, J. Ryckebusch, K. Van Houcke, “Diagrammatic Monte Carlo study of the Fermi polaron in two dimensions”, *Phys. Rev. B* **89** (2014), article no. 085119.
- [24] M. M. Glazov, “Optical properties of charged excitons in two-dimensional crystals”, <https://arxiv.org/abs/2004.13484>, 2020.
- [25] R. Combescot, A. Recati, C. Lobo, F. Chevy, “Normal state of highly polarized Fermi gases: Simple many-body approaches”, *Phys. Rev. Lett.* **98** (2007), article no. 180402.