

11-22-2022

Association Between Sacubitril/Valsartan Initiation and Mitral Regurgitation Severity in Heart Failure With Reduced Ejection Fraction: The PROVE-HF Study

James L Januzzi

Alaa Mabrouk Salem Omar

Yuxi Liu

Sean Murphy

Javed Butler

See next page for additional authors

Follow this and additional works at: <https://jdc.jefferson.edu/cardiologyp>



Part of the [Cardiology Commons](#)

[Let us know how access to this document benefits you](#)

This Article is brought to you for free and open access by the Jefferson Digital Commons. The Jefferson Digital Commons is a service of Thomas Jefferson University's [Center for Teaching and Learning \(CTL\)](#). The Commons is a showcase for Jefferson books and journals, peer-reviewed scholarly publications, unique historical collections from the University archives, and teaching tools. The Jefferson Digital Commons allows researchers and interested readers anywhere in the world to learn about and keep up to date with Jefferson scholarship. This article has been accepted for inclusion in Division of Cardiology Faculty Papers by an authorized administrator of the Jefferson Digital Commons. For more information, please contact: JeffersonDigitalCommons@jefferson.edu.

Authors

James L Januzzi, Alaa Mabrouk Salem Omar, Yuxi Liu, Sean Murphy, Javed Butler, G Michael Felker, Iliana L Piña, Jonathan Ward, Scott Solomon, and Johanna Contreras



Association Between Sacubitril/Valsartan Initiation and Mitral Regurgitation Severity in Heart Failure With Reduced Ejection Fraction: The PROVE-HF Study

James L. Januzzi¹, MD; Alaa Mabrouk Salem Omar, MD, MSc, PhD; Yuxi Liu, MS; Sean Murphy¹, MBCh; Javed Butler¹, MD, MBA; G. Michael Felker¹, MD, MHS; Iliana L. Piña¹, MD, MPH; Jonathan Ward, PharmD; Scott Solomon¹, MD; Johanna Contreras, MD

Mitral regurgitation (MR) severity is an important determinant of symptom status and prognosis in heart failure with reduced ejection fraction (HFrEF). Functional MR is frequently progressive, associated with adverse myocardial remodeling, neurohormonal activation, worsened symptoms, and poor outcome.¹ Percutaneous MR repair may improve outcomes in HFrEF;² however, optimizing guideline-directed medical therapy (GDMT) before valve repair may reduce MR severity sufficiently to avoid need for such a procedure. Limited data suggest treatment of HFrEF with sacubitril/valsartan (sac/val) may result in improvement of MR regardless of background GDMT.³

We examined the association between treatment with sac/val on change in MR among participants in the PROVE-HF study (Prospective Study of Biomarkers, Symptom Improvement, and Ventricular Remodeling During Sacubitril/Valsartan Therapy for Heart Failure; URL: <https://www.clinicaltrials.gov>; Unique identifier: NCT02887183).⁴ All study procedures were approved by local institutional review boards, and study participants provided informed consent. All supporting data are available within the article. In this study of 794 participants with HF and left ventricular (LV) ejection fraction (LVEF) <40%, sac/val was initiated and titrated to the maximally tolerated dose. An echocardiogram was performed at baseline, 6 months, and 12 months, and interpreted in a temporally and clinically

blinded fashion. MR severity was graded using a combination of visual assessment of the color Doppler jet and MR/left atrial area and vena contracta width⁵ and categorized on a scale of 0 (none), 1+ (trace), 2+ (mild), 3+ (moderate), and 4+ (severe). Those with previous mitral procedures (N=40) were excluded from the analysis, as were 30 study participants missing baseline MR severity.

Study participants had a mean \pm SD age of 65.0 ± 12.4 years. Most (76.4%) were receiving an ACE inhibitor or angiotensin II receptor blocker at baseline. The median LVEF was 28.3%, with LV end diastolic volume index (LVEDVi) of 87.2 mL/kg², LV end systolic volume index of 61.8 mL/kg², left atrial volume index of 37.6 mL/kg², E/e' (the ratio of the early diastolic transmitral Doppler velocity [E] and the early diastolic septal velocity [e']) of 11.3, and LV mass index of 125.2 g/m². At baseline, 42 (5.8%) and 66 (9.1%) had 3+ and 4+ MR, respectively.

From baseline through 12 months, improvement in MR was observed (Figure); by 6 months, prevalence of 3 to 4+ MR decreased to 8.2% (relative 45.0% reduction), and by 12 months, it was 8.4% (relative 44.7% reduction). Those with 3 to 4+ MR at baseline with reduction to $\leq 2+$ by 12 months ("responders"; N=52) had similar baseline clinical characteristics (including vital signs, baseline GDMT, or previous cardiac resynchronization therapy) to those with persistent MR grade 3 to 4+ at 12 months ("nonresponders"; N=33); baseline LVEF and

Key Words: cardiac remodeling ■ health status ■ heart failure ■ mitral regurgitation ■ natriuretic peptides ■ prognosis ■ sacubitril/valsartan

Correspondence to: James L. Januzzi, MD, Cardiology Division, Massachusetts General Hospital, 55 Fruit St, Boston, MA 02114. Email jjanuzzi@partners.org
For Sources of Funding and Disclosures, see page 1640.

© 2022 The Authors. *Circulation* is published on behalf of the American Heart Association, Inc., by Wolters Kluwer Health, Inc. This is an open access article under the terms of the [Creative Commons Attribution Non-Commercial-NoDerivs](https://creativecommons.org/licenses/by-nc-nd/4.0/) License, which permits use, distribution, and reproduction in any medium, provided that the original work is properly cited, the use is noncommercial, and no modifications or adaptations are made.

Circulation is available at www.ahajournals.org/journal/circ

Nonstandard Abbreviations and Acronyms

GDMT	guideline-directed medical therapy
HFrEF	heart failure with reduced ejection fraction
LV	left ventricular
LVEDVi	left ventricular end diastolic volume index
LVEF	left ventricular ejection fraction
MR	mitral regurgitation
NT-proBNP	N-terminal pro-B-type natriuretic peptide
sac/val	sacubitril/valsartan

LV volumes were similar between groups, but responders had lower baseline left atrial volume index (43.9 versus 49.3 mL/kg²; $P=0.01$) and LV mass index (145.0 versus 166.0 g/m²; $P=0.02$). Neither ratio of vena contracta/LVEDVi or left atrial area/LVEDVi was significantly different between responders and nonresponders at baseline ($P=0.11$ and 0.68, respectively). Between responders and nonresponders, the average sac/val dose during the study was 276 versus 277 mg/d ($P=0.57$). By 12 months, LVEF improvement was greater in responders versus nonresponders (+11.0% versus +7.6%; $P=0.05$), and responders had lower final LVEDVi (85.2 versus 96.9 mL/kg²; $P=0.02$), LV end systolic volume index (56.5 versus 66.0 mL/kg²; $P=0.04$), left atrial volume index (33.4 versus 42.3 mL/kg²; $P<0.001$), E/e' (12.6 versus 15.8; $P=0.04$), and LV mass index (125.7 versus 152.2 g/m²). Last, by 12 months, responders had lower median

NT-proBNP (N-terminal pro-B-type natriuretic peptide; 912 versus 1512 pg/mL; $P=0.01$) and higher median Kansas City Cardiomyopathy Questionnaire Overall Summary Scores (82.3 versus 72.9; $P=0.04$) despite similar results for both measures at baseline.

Although valve repair in those with 3 to 4+ MR may improve symptoms and reduce adverse outcomes in HFrEF, optimizing GDMT is important before such an intervention. In this analysis, despite the majority of study participants previously receiving an ACE inhibitor or angiotensin II receptor blocker, we observed a substantial shift to lesser degrees of MR after 12 months of treatment with sac/val, including a nearly 50% reduction in 3 to 4+ MR. Reduction in MR from 3 to 4+ was associated with considerable reverse cardiac remodeling, reduced NT-proBNP, and improved health status.

The baseline clinical and echocardiographic characteristics of those with 3 to 4+ MR who had a reduction to $\leq 2+$ MR by 12 months were similar to those who had persistent 3 to 4+ MR by 12 months, and both groups had comparable baseline LVEF, LV volumes, and ratio of MR severity to LVEDVi despite comparable degrees of MR at baseline. Thus, predicting presence of MR caused by LV dilation versus intrinsic valve dysfunction without a course of sac/val treatment may be difficult.

Limitations of this study include the single-arm, observational design; thus, improvement in MR might have been due to factors other than sac/val. Nonetheless, the results of this study provide important insights into potential change in MR after initiation of sac/val in usual care.

With growth in use of percutaneous approaches for mitral valve repair for those with HFrEF and 3 to 4+ MR,




Figure. Sankey diagram detailing severity of mitral regurgitation at baseline, 6 months, and 12 months among study participants with heart failure and reduced ejection fraction treated with sacubitril/valsartan.

the results of this study reinforce importance of ensuring optimal GDMT before such decisions.

ARTICLE INFORMATION

This work was presented as an abstract at the Heart Failure Society of America's Annual Scientific Meeting 2022, Washington, DC, September 30 through October 3, 2022.

Affiliations

Massachusetts General Hospital (J.L.J., Y.L., S.M.), Brigham and Women's Hospital (S.S.), Harvard Medical School, Boston. Baim Institute for Clinical Research, Boston, MA (J.L.J.). Mount Sinai Morningside, New York, NY (A.M.S.O.). Icahn School of Medicine at Mount Sinai, New York, NY (A.M.S.O., J.C.). Baylor Scott and White Research Institute, Dallas, TX (J.B.). University of Mississippi, Jackson (J.B.). Duke University School of Medicine, Durham, NC (G.M.F.). Thomas Jefferson University, Philadelphia, PA (I.L.P.). Novartis Pharmaceuticals Corporation, East Hanover, NJ (J.W.).

Sources of Funding

Dr Januzzi is supported in part by the Hutter Family Professorship.

Disclosures

Dr Januzzi is a Trustee of the American College of Cardiology; is a board member of Imbria Pharmaceuticals; has received research support from Abbott, Applied Therapeutics, Innolife, Novartis Pharmaceuticals, and Roche Diagnostics; has received consulting income from Abbott, Beckman, Bristol Myers Squibb (BMS), Boehringer-Ingelheim, Janssen, Novartis, Pfizer, Merck, Roche Diagnostics, and Siemens; and participates in clinical end point committees/data safety monitoring boards for Abbott, AbbVie, Bayer, CVRx, Intercept, Janssen, and Takeda. Dr Butler is a consultant for Abbott, Adrenomed, Amgen, Applied Therapeutics, Array, AstraZeneca, Bayer, Boehringer Ingelheim, Bristol Myers Squibb, CVRx, G3 Pharmaceutical, Impulse Dynamics, Innolife, Janssen, LivaNova, Luitpold, Medtronic, Merck, Novartis, Novo Nordisk, Roche, and Vifor. Dr Felker has received research grants from the National Heart, Lung, and Blood Institute, American Heart Association, Amgen, Bayer, BMS, Merck, Cytokinetics, and CSL-Behring; has acted as a consultant to Novartis, Amgen, BMS, Cytokinetics, Medtronic, Cardionomic, Boehringer-Ingelheim, American Regent, Abbott, AstraZeneca, Reprise, Myovant, Sequana, Windtree Therapeutics, and Whiteswell; and has served on clinical

end point committees/data safety monitoring boards for Amgen, Merck, Medtronic, EBR Systems, V-Wave, LivaNova, Siemens, and Rocket Pharma. Dr Ward is an employee of Novartis Pharmaceuticals Corporation. Dr Solomon has received research grants from Alnylam, Amgen, AstraZeneca, Bellerophon, Bayer, BMS, Celladon, Cytokinetics, Eidos, Gilead, GSK, Ionis, Lone Star Heart, Mesoblast, MyoKardia, the National Institutes of Health/National Heart, Lung, and Blood Institute, Novartis, Sanofi Pasteur, and Theracos; and has consulted for Akros, Alnylam, Amgen, AstraZeneca, Bayer, BMS, Cardior, Corvia, Cytokinetics, Gilead, GSK, Ironwood, Merck, Myokardia, Novartis, Roche, Takeda, Theracos, Quantum Genetics, Cardurion, AoBiome, Janssen, Cardiac Dimensions, and Tenaya. Dr Contreras has received consulting fees from Alnylam Pharmaceuticals, Novo Nordisk, Boehringer Ingelheim, and AstraZeneca. The other authors report no conflicts.

REFERENCES

1. Bartko PE, Pavo N, Perez-Serradilla A, Arfsten H, Neuhold S, Wurm R, Lang IM, Strunk G, Dal-Bianco JP, Levine RA, et al. Evolution of secondary mitral regurgitation. *Eur Heart J Cardiovasc Imaging*. 2018;19:622–629. doi: 10.1093/ehjci/jeu023
2. Stone GW, Lindenfeld J, Abraham WT, Kar S, Lim DS, Mishell JM, Whisenant B, Grayburn PA, Rinaldi M, Kapadia SR, et al. Transcatheter mitral-valve repair in patients with heart failure. *N Engl J Med*. 2018;379:2307–2318. doi: 10.1056/NEJMoa1806640
3. Kang DH, Park SJ, Shin SH, Hong GR, Lee S, Kim MS, Yun SC, Song JM, Park SW, Kim JJ. Angiotensin receptor neprilysin inhibitor for functional mitral regurgitation. *Circulation*. 2019;139:1354–1365. doi: 10.1161/CIRCULATIONAHA.118.037077
4. Januzzi JL Jr, Prescott MF, Butler J, Felker GM, Maisel AS, McCague K, Camacho A, Pina IL, Rocha RA, Shah AM, et al; PROVE-HF Investigators. Association of change in N-terminal pro-B-type natriuretic peptide following initiation of sacubitril-valsartan treatment with cardiac structure and function in patients with heart failure with reduced ejection fraction. *JAMA*. 2019;322:1085–1095. doi: 10.1001/jama.2019.12821
5. Zoghbi WA, Adams D, Bonow RO, Enriquez-Sarano M, Foster E, Grayburn PA, Hahn RT, Han Y, Hung J, Lang RM, et al. Recommendations for noninvasive evaluation of native valvular regurgitation: a report from the American Society of Echocardiography developed in collaboration with the Society for Cardiovascular Magnetic Resonance. *J Am Soc Echocardiogr*. 2017;30:303–371. doi: 10.1016/j.echo.2017.01.007