

# Three new species of *Nigrograna* (Dothideomycetes, Pleosporales) associated with Arabica coffee from Yunnan Province, China

Li Lu<sup>1,2,3</sup>, Samantha C. Karunaratna<sup>1</sup>, Dong-qin Dai<sup>1</sup>,  
Ruvishika S. Jayawardena<sup>2,3</sup>, Nakarin Suwannarach<sup>4</sup>, Saowaluck Tibpromma<sup>1</sup>

**1** Center for Yunnan Plateau Biological Resources Protection and Utilization, College of Biological Resource and Food Engineering, Qujing Normal University, Qujing, Yunnan 655011, China **2** Center of Excellence in Fungal Research, Mae Fah Luang University, Chiang Rai 57100, Thailand **3** School of Science, Mae Fah Luang University, Chiang Rai 57100, Thailand **4** Research Center of Microbial Diversity and Sustainable Utilization, Faculty of Science, Chiang Mai University, Chiang Mai 50200, Thailand

Corresponding authors: Nakarin Suwannarach ([suwan.462@gmail.com](mailto:suwan.462@gmail.com)),  
Saowaluck Tibpromma ([saowaluckfai@gmail.com](mailto:saowaluckfai@gmail.com))

---

Academic editor: C. Bhunjun | Received 28 September 2022 | Accepted 17 November 2022 | Published 12 December 2022

---

**Citation:** Lu L, Karunaratna SC, Dai D-q, Jayawardena RS, Suwannarach N, Tibpromma S (2022) Three new species of *Nigrograna* (Dothideomycetes, Pleosporales) associated with Arabica coffee from Yunnan Province, China. MycoKeys 94: 51–71. <https://doi.org/10.3897/mycokeys.94.95751>

---

## Abstract

Coffee is one of the most important cash crops in Yunnan Province, China. Yunnan is ranked as the biggest producer of high-quality coffee in China. During surveys of microfungi from coffee plantations in Yunnan, six fungal strains that resemble Nigrogranaceae were collected. Multi-gene analyses of a combined SSU-LSU-ITS-*rpb2-tef1- $\alpha$*  sequence data matrix were used to infer the phylogenetic position of the new species in *Nigrograna* while morphological characteristics were used to deduce the taxonomic position of the new species. Six fungal strains isolated from decaying branches of *Coffea arabica* represent three new saprobic species in *Nigrograna*. The three new species, *N. asexualis*, *N. coffeae*, and *N. puerensis*, are described with full (macro and micro characteristics) descriptions, illustrations, and a phylogenetic tree that shows the phylogenetic position of new taxa.

## Keywords

3 new taxa, *Coffea arabica*, Nigrogranaceae, phylogeny, saprobic fungi, taxonomy

## Introduction

Coffee (*Coffea* L.) was first planted in Yunnan Province, China in 1982 (Zhang et al. 2014). To date, about 170 varieties of coffee (Global Biodiversity Information Facility database (GBIF), available at: <https://www.gbif.org/species/2895315> (accessed on 07 November 2022)) are available in the world, of which *Coffea arabica* L. is the most popular coffee accounting for 75% of the world's production, while 25% is provided by *C. canephora* Pierre ex A. Froehner, and less than 1% by *C. liberica* W. Bull and other varieties (Sharma 2020). The coffee production in Yunnan Province is approximately 90% of China's total coffee production (Neilson and Wang 2019), while Pu'er is the largest coffee planting area in Yunnan, in terms of the highest yield and the best quality (Li 2014).

Fungal diversity is highly uncertain; the current estimated numbers are between 1.5 to 12 million, of which about 150,000 species have been named and classified (Hawksworth and Lücking 2017; Hyde et al. 2020; Bhunjun et al. 2022). Fungi are important organisms in terrestrial and aquatic ecosystems that are involved in the decomposition and nutrient cycling of dead plant material (Hyde et al. 2020; Bhunjun et al. 2022; Phukhamsakda et al. 2022). Also, saprobic fungi play vital roles in soil food chains, decomposition of plant, and animal materials, and solubilization of phosphorous (Dighton 2003; Pandey et al. 2008). However, coffee saprobic fungi have been poorly investigated (Arias and Abarca 2014; Lu et al. 2022a). Coffee saprobic fungi are distributed in 15 orders, and among them, Pleosporales Luttr. is the most common order (Lu et al. 2022a).

Pleosporales, belonging to Dothideomycetes O.E. Erikss. & Winka, was first proposed by Luttrell (1955), and later it was formally established by Barr (1987). In 2021, it consists of 91 families and 614 genera as the largest order (Hongsanan et al. 2020a; Wijayawardene et al. 2022). They are distributed in terrestrial and aquatic habitats (Zhang et al. 2008; Jiang et al. 2021). The members of Pleosporales are characterized by perithecioid and ostiolar ascomata, with or without periphyses, presence of cellular pseudoparaphyses, bitunicate, with ocular chambers or apical ring asci, various shapes of ascospores, with pigmentation and septation, and sheath present or absent (Zhang et al. 2012; Tennakoon et al. 2021; Yang et al. 2022).

Nigrogranaceae Jaklitsch & Voglmayr (Pleosporales) was proposed as a new family by Jaklitsch and Voglmayr (2016) to accommodate *Nigrograna* Gruyter, Verkley & Crous as the type genus. Liu et al. (2017) estimated that the divergence time of Nigrogranaceae is around 79 (44–124) Mya in crown age and 131 (86–180) Mya in stem age. Nigrogranaceae is monotypic, and they exist as endophytic, human pathogenic, and saprobic lifestyles (Hongsanan et al. 2020b; Zhang et al. 2020; Boonmee et al. 2021). The sexual morph of Nigrogranaceae is characterized by globose and black, ostiolar, clavate, and fissitunicate ascomata, with a short stipe and asci with a knob-like base, fusoid to narrowly ellipsoid, septate, and smooth or faintly verruculose ascospores (Jaklitsch and Voglmayr 2016). The asexual morph is characterized by pycnidia similar to ascomata, filiform and branched conidiophores, ampulliform or lageniform phialides, rod-like to ellipsoid, and hyaline or sub-hyaline conidia (Jaklitsch and Voglmayr 2016).

*Nigrograna* was introduced by de Gruyter (2012) with *N. mackinnonii* (Borelli) Gruyter, Verkley & Crous (basonym: *Pyrenochaeta mackinnonii* Borelli) as the type species. *Pyrenochaeta mackinnonii* was reported from a mycetoma patient by Borelli (1976), but it was found to be remote from the generic type species *P. nobilis* De Not. (de Gruyter et al. 2010, 2013). Since it was not possible to determine which family in Pleosporales *P. mackinnonii* belongs to, only the new genus *Nigrograna* was introduced to accommodate *P. mackinnonii* and named as *N. mackinnonii* (de Gruyter 2012). Later, *Nigrograna* was used as a synonym of *Biatriospora* K.D. Hyde & Borse, as *N. mackinnonii* is phylogenetically closely related to the type species of *Biatriospora* (*B. marina* K.D. Hyde & Borse) (Ahmed et al. 2014), while Hongsanan et al. (2020a) treated *Biatriospora* and *Nigrograna* as two separate genera. In 2022, *Nigrograna* represents 20 epithets listed in Index Fungorum (2022), and the members have been reported as saprobic, human pathogenic, and endophytic worldwide (Kolařík 2018; Zhao et al. 2018), showing a wide range of hosts (marine and terrestrial habitats) (Hyde et al. 2017; Tibpromma et al. 2017; Dayarathne et al. 2020). The sexual morph of *Nigrograna* is characterized by globose to subglobose and black ascomata, with ostiolar, two-layered peridium, clavate and fissitunicate asci, fusoid to narrowly ellipsoid, straight or curved, septate, and smooth or verruculose ascospores (Jaklitsch and Voglmayr 2016; Zhang et al. 2020). Asexual morph is characterized by globose to subglobose or pyriform pycnidia, filiform and branched conidiophores, hyaline, phialidic, discrete conidiogenous cells, sub-hyaline, aseptate and ellipsoidal conidia (de Gruyter 2012; Jaklitsch and Voglmayr 2016).

In this study, three saprobic *Nigrograna* were collected from *Coffea arabica* branches in Yunnan Province, China. One species was isolated as an asexual morph (*N. asexualis*), while the other two isolated as sexual morphs (*N. coffeae*, *N. puerensis*) are illustrated and described as new species based on morphology and multi-gene phylogenetic analyses and are compared with closely related taxa.

## Materials and methods

### Collection, morphology and isolation

Coffee branch samples were collected from coffee plantations in Pu'er and Xishuangbanna, Yunnan Province, China. Specimens were put in plastic bags and taken to the mycology laboratory at Qujing Normal University. The vertical sections of fruiting structures were made for microscope studies and photomicrography. Micro-morphological characteristics were observed using a Leica DM2500 compound microscope and photographed with a Leica DMC4500 camera fitted onto the microscope. Color codes in the manuscript followed colorhexa (<https://www.colorhexa.com>). The measurements were processed in Tarosoft (R) Image Frame Work v. 0.9.7, and photographic plates were made in Adobe Photoshop CC 2018. Single spore isolation was carried out following Senanayake et al. (2020). Herbarium specimens were deposited at Zhongkai University of Agriculture and Engineering (ZHKU), while the living cultures growing on potato dextrose agar (PDA) were deposited at the culture collection of Zhongkai

University of Agriculture and Engineering (ZHKUCC). Faces of fungi (FoF) numbers and Index Fungorum (IF) numbers were obtained as explained in Jayasiri et al. (2015) and Index Fungorum (2022).

## DNA extraction and PCR amplification

Genomic DNA was extracted from the fresh fungal mycelia which were grown on PDA for about two weeks, using Biospin Fungus Genomic DNA Extraction Kit–BSC14S1 (BioFlux, China) following the manufacturer's instructions. Lu et al. (2021) was followed for the Polymerase Chain Reaction (PCR). Five partial gene regions were used in this study viz. the internal transcribed spacer (ITS) region was amplified with the primers ITS4 and ITS5 (White et al. 1990), the 18 s small subunit (SSU) region was amplified by primers NS1 and NS4 (White et al. 1990), the nuclear ribosomal 28 s large subunit (LSU) region was amplified by the primers LROR and LR5 (Vilgalys and Hester 1990), the partial RNA polymerase II subunit (*rpb2*) region was amplified with the primers RPB2-5F and RPB2-7cR (Liu et al. 1999), and the partial translation elongation factor 1-alpha (*tefl- $\alpha$* ) gene was amplified with the primers EF1-983F and 2218R (Rehner and Buckley 2005). Lu et al. (2022b) was followed for the amplification reactions of different primers. Amplified PCR products were sent to Sango Biotechnology Co., Ltd. (Shanghai, China) for sequencing. All sequences generated in this study were deposited in GenBank (Table 1).

## Phylogenetic analyses

Phylogenetic analyses of the aligned sequences referred to Dissanayake et al. (2020). Newly generated reverse and forward sequences were assembled with Geneious program (9.1.2) and the preliminary identification was done by the BLASTn search in NCBI (<https://www.ncbi.nlm.nih.gov>). Additional highly similar sequences were downloaded from GenBank (<https://www.ncbi.nlm.nih.gov/genbank/>) based on the BLASTn results and recent publications. Single-gene sequence alignments were made in MAFFT v. 7 (<http://mafft.cbrc.jp/alignment/server/>), edited in trimAl v1.2 (<http://trimal.cgenomics.org>), and multi-gene alignments were made by Sequence Matrix program (1.7.8) (Vaidya et al. 2011). The sequence datasets used to build the phylogenetic trees are shown in Table 1.

Phylogenetic analyses were conducted with maximum likelihood (ML) and Bayesian inference (BI) algorithms on the CIPRES Science Gateway portal (<https://www.phylo.org/>) (Miller et al. 2012). The ML tree was run with RAxML-HPC v.8 on XSEDE (Stamatakis 2014), and GTRGAMMA substitution model with 1000 bootstrap iterations. The BI tree was run with MrBayes on XSEDE (3.2.7a) (Ronquist et al. 2012). MrModeltest 2.2 (Nylander 2004) and PAUP v. 4.0b10 (Ronquist and Huelsenbeck 2003) were used to evaluate the best models of evolution, the evolutionary model of SYM+I+G substitution model was selected for LSU, HKY+I+G substitution model was selected for SSU, and GTR+I+G substitution model was selected for ITS, *rpb2* and *tefl- $\alpha$* . Six simultaneous Markov Chains were run for two million generations

**Table 1.** Taxa names, strain numbers, and corresponding GenBank accession numbers of the taxa used in the phylogenetic analyses. Newly generated sequences in this study are indicated in bold. The type species are noted with <sup>T</sup> after the species name, while NA indicates the unavailability of data.

Taxon	Strain numbers	ITS	LSU	<i>rpb2</i>	SSU	<i>tefl-α</i>
<i>Cylothyriella rubronotata</i> (Berk. & Broome) Jaklitsch & Voglmayr <sup>T</sup>	CBS 141486	KX650544	KX650519	NA	KX650507	KX650574
<i>Cylothyriella rubronotata</i>	CBS 419.85	NA	GU349002	GU301875	NA	GU371728
<i>Nigrograna antibiotica</i> (M. Kolařík & A. Kubátová) M. Kolařík <sup>T</sup>	CCF 4378	JX570932	KF925327	NA	KF925328	JX570934
<i>Nigrograna antibiotica</i>	CCF 4998	LT221894	NA	LT221895	NA	NA
<i>Nigrograna aquatica</i> W. Dong, H. Zhang & K.D. Hyde <sup>T</sup>	MFLUCC 14-1178	MF399065	MF415392	NA	MF415394	MF498582
<i>Nigrograna aquatica</i>	MFLUCC 17-2318	MT627705	MN913705	NA	NA	NA
<i>Nigrograna asexualis</i> <sup>T</sup>	<b>ZHKUCC 22-0214</b>	<b>OP450965</b>	<b>OP450971</b>	<b>OP432241</b>	<b>OP450979</b>	<b>OP432245</b>
<i>Nigrograna asexualis</i>	<b>ZHKUCC 22-0215</b>	<b>OP450966</b>	<b>OP450972</b>	<b>OP432242</b>	<b>OP450980</b>	<b>OP432246</b>
<i>Nigrograna cangshanensis</i> Z.L. Luo, H.Y. Su & K.D. Hyde <sup>T</sup>	MFLUCC 15-0253	KY511063	KY511064	NA	KY511065	NA
<i>Nigrograna carollii</i> M. Kolařík <sup>T</sup>	CCF 4484	LN626657	LN626682	LN626662	LN626674	LN626668
<i>Nigrograna chromolaenae</i> Mapook & K.D. Hyde <sup>T</sup>	MFLUCC 17-1437	MT214379	MT214473	NA	NA	MT235801
<i>Nigrograna coffeae</i> <sup>T</sup>	<b>ZHKUCC 22-0210</b>	<b>OP450967</b>	<b>OP450973</b>	<b>OP432243</b>	<b>OP450981</b>	<b>OP432247</b>
<i>Nigrograna coffeae</i>	<b>ZHKUCC 22-0211</b>	<b>OP450968</b>	<b>OP450974</b>	<b>OP432244</b>	<b>OP450982</b>	<b>OP432248</b>
<i>Nigrograna fuscidula</i> (Sacc.) Jaklitsch & Voglmayr <sup>T</sup>	CBS 141556	KX650550	NA	NA	NA	KX650525
<i>Nigrograna fuscidula</i>	CBS 141476	KX650547	NA	KX650576	KX650509	KX650522
<i>Nigrograna fuscidula</i>	MF1a	KX650548	NA	NA	NA	KX650523
<i>Nigrograna fuscidula</i>	MF3	KX650549	NA	NA	NA	KX650524
<i>Nigrograna hydei</i> J.F. Zhang, J.K. Liu & Z.Y. Liu <sup>T</sup>	GZCC 19-0050	MN387225	MN387227	NA	NA	MN389249
<i>Nigrograna impatientis</i> J.F. Zhang, J.K. Liu & Z.Y. Liu <sup>T</sup>	GZCC 19-0042	MN387226	MN387228	NA	NA	MN389250
<i>Nigrograna jingbongensis</i> Wanas. & K.D. Hyde <sup>T</sup>	KUMUCC 21-0035	MZ493303	MZ493317	MZ508421	MZ493289	MZ508412
<i>Nigrograna jingbongensis</i>	KUMUCC 21-0036	MZ493304	MZ493318	MZ508422	MZ493290	MZ508413
<i>Nigrograna kunmingensis</i> T.Y. Du & Tibpromma <sup>T</sup>	ZHKUCC 22-0242	OP456214	OP456379	NA	OP456382	OP471608
<i>Nigrograna kunmingensis</i>	ZHKUCC 22-0243	OP484334	OP456380	NA	OP456383	OP471609
<i>Nigrograna locuta-pollinis</i> F. Liu & L. Cai <sup>T</sup>	CGMCC 3.18784	MF939601	MF939583	MF939610	NA	MF939613
<i>Nigrograna locuta-pollinis</i>	LC11690	MF939603	MF939584	MF939611	NA	MF939614
<i>Nigrograna mackinnonii</i> <sup>T</sup>	CBS 674.75	KF015654	KF015612	KF015703	GQ387552	KF407986
<i>Nigrograna mackinnonii</i>	E5202H	JX264157	KJ605422	JX264156	JX264155	JX264154
<i>Nigrograna mackinnonii</i>	E9303e	JN545759	LN626681	LN626666	LN626678	LN626673
<i>Nigrograna magnoliae</i> Wanas. <sup>T</sup>	MFLUCC 20-0020	MT159628	MT159622	MT159611	MT159634	MT159605
<i>Nigrograna magnoliae</i>	GZCC 17-0057	MF399066	MF415393	NA	MF415395	MF498583
<i>Nigrograna magnoliae</i>	MFLUCC 20-0021	MT159629	MT159623	MT159612	MT159635	MT159606
<i>Nigrograna mycophila</i> Jaklitsch, Friesbe & Voglmayr <sup>T</sup>	CBS 141478	KX650553	NA	NA	NA	KX650526
<i>Nigrograna mycophila</i>	CBS 141483	KX650555	NA	KX650577	KX650510	KX650528
<i>Nigrograna mycophila</i>	MF6	KX650554	NA	NA	NA	KX650527
<i>Nigrograna norvegica</i> Jaklitsch & Voglmayr <sup>T</sup>	CBS 141485	KX650556	NA	KX650578	KX650511	NA
<i>Nigrograna obliqua</i> Jaklitsch & Voglmayr <sup>T</sup>	CBS 141477	KX650560	NA	KX650580	NA	KX650531
<i>Nigrograna obliqua</i>	CBS 141475	KX650558	NA	KX650579	KX650512	KX650530
<i>Nigrograna obliqua</i>	MRP	KX650561	NA	KX650581	NA	KX650532

Taxon	Strain numbers	ITS	LSU	<i>rpb2</i>	SSU	<i>tefl-<math>\alpha</math></i>
<i>Nigrograna peruviana</i> (M. Kolařík & R. Gazis) M. Kolařík <sup>T</sup>	CCF 4485	LN626658	LN626683	LN626665	LN626677	LN626671
<i>Nigrograna puerensis</i> <sup>T</sup>	ZHKUCC 22-0212	OP450969	OP450975	NA	OP450983	OP432249
<i>Nigrograna puerensis</i>	ZHKUCC 22-0213	OP450970	OP450976	NA	OP450984	OP432250
<i>Nigrograna rhizophorae</i> Dayar., E.B.G. Jones & K.D. Hyde <sup>T</sup>	MFLUCC 18-0397	MN047085	NA	MN431489	NA	MN077064
<i>Nigrograna rhizophorae</i>	MFLU 19-1234	NA	MN017845	MN431490	NA	MN077063
<i>Nigrograna samueliana</i> Devadatha, V.V. Sarma & E.B.G. Jones <sup>T</sup>	NFCCI-4383	MK358817	MK358812	MK330939	MK358810	MK330937
<i>Nigrograna thymi</i> Mapook, Camporesi & K.D. Hyde <sup>T</sup>	MFLUCC 14-1096	KY775576	KY775573	NA	KY775574	KY775578
<i>Nigrograna yasuniiana</i> M. Kolařík <sup>T</sup>	YU.101026	HQ108005	LN626684	LN626664	LN626676	LN626670
<i>Occultibambusa bambusae</i> D.Q. Dai & K.D. Hyde <sup>T</sup>	MFLUCC 13-0855	KU940123	KU863112	KU940170	NA	KU940193
<i>Occultibambusa fusispora</i> Phookamsak, D.Q. Dai & K.D. Hyde	MFLUCC 11-0127	MZ329036	MZ325466	MZ329032	MZ329028	MZ325469
<i>Occultibambusa pustula</i> D.Q. Dai & K.D. Hyde <sup>T</sup>	MFLUCC 11-0502	KU940126	KU863115	NA	NA	NA
<i>Paradictyoarthrinium diffractum</i> Matsush.	MFLUCC13-0466	KP744455	NA	KP744498	NA	NA
<i>Paradictyoarthrinium tectonicola</i> Doilom & K.D. Hyde <sup>T</sup>	MFLUCC 13-0465	KP744456	NA	KP744500	KP753961	KX437763
<i>Seriascoma didymosporum</i> Phookamsak, D.Q. Dai, Karun. & K.D. Hyde <sup>T</sup>	MFLUCC 11-0179	KU940127	KU940196	KU863116	NA	KU940173
<i>Seriascoma honghense</i> H.B. Jiang, Phookamsak & K.D. Hyde <sup>T</sup>	KUMCC 21-0021	MZ329039	MZ325468	MZ329035	NA	MZ325470
<i>Versicolorisporium triseptatum</i> Sat. Hatak., Kaz. Tanaka & Y. Harada <sup>T</sup>	HHUF 28815	NR_119392	NA	NG_042318	NG_060995	NA

and trees were sampled at every 200<sup>th</sup> generation (resulting in 10,000 trees), and these chains stopped when all convergences met and the standard deviation fell below 0.01. All resulting trees were plotted using FigTree v. 1.4.0 (Rambaut 2014) and the layout of the trees was made by Microsoft Office PowerPoint 2020.


## Results

### Phylogenetic analyses

Three new species formed a distinct clade in *Nigrograna* with strong statistical support (*N. coffeae* and *N. puerensis* ML = 100%, BIPP = 1.00, and *N. asexualis* ML = 68%, BIPP = 0.97). Multi-locus data (SSU, LSU, ITS, *rpb2* and *tefl- $\alpha$* ) composed of 54 strains (Table 1), and *Cylothyriella rubronotata* strains CBS 141486 and CBS 419.85 were used as the outgroup taxa. A total of 4485 characters were fed to the phylogenetic analysis after alignment, 1–1047 (SSU), 1048–1956 (LSU), 1957–2477 (ITS), 2478–3510 (*rpb2*) and 3511–4485 (*tefl- $\alpha$* ). The topology of the phylogenetic tree generated by the ML method was highly similar to that by BI, and therefore it was chosen to represent the evolutionary history of *Nigrograna*.

The ML analysis of the combined dataset yielded a best-scoring tree with a final ML optimization likelihood value of -23091.568105. The alignment has 1495

distinct alignment patterns, with 33.58% completely undetermined characters and gaps. Parameters for the GTR + I + G model of the combined SSU, LSU, ITS, *rpb2* and *tef1- $\alpha$*  were as follows: estimated base frequencies A = 0.247145, C = 0.250645, G = 0.263985, T = 0.238225; substitution rates AC = 1.810004, AG = 4.475190, AT = 1.758134, CG = 1.340389, CT = 10.583215, GT = 1.000; gamma distribution shape parameter  $\alpha$  = 0.167006. The phylogenetic tree resulting from RAxML analysis is shown in Fig. 1.



**Figure 1.** The maximum-likelihood phylogram of *Nigrograna* based on a combined SSU, LSU, ITS, *rpb2* and *tef1- $\alpha$*  sequence dataset with *Cyclothyriella rubronotata* CBS 141486 and CBS 419.85 as the outgroup taxa (Dayarathne et al. 2020). The maximum-likelihood bootstrap values (ML  $\geq$  60%, left) and Bayesian Inference Posterior Probability values (BIPP  $\geq$  0.90, right) are shown above the nodes. Strains derived from the current study are in blue, while type strains are in bold.

## Taxonomy

### *Nigrograna coffeae* L. Lu & Tibpromma, sp. nov.

Index Fungorum number: IF559425

Facesoffungi Number: FoF12765

Fig. 2

**Etymology.** Species epithet refers to the host genus “*Coffea*” where the fungus was isolated.

**Holotype.** ZHKU 22-0121.


**Description.** *Saprobia* on decaying branch of *Coffea arabica*. **Sexual morph:** *Ascomata* 90–140  $\mu\text{m}$  high, 140–200  $\mu\text{m}$  wide ( $\bar{x}$  = 115  $\times$  168  $\mu\text{m}$ ,  $n$  = 10), immersed, solitary, black spots on substrate, subglobose to oval, sometimes obpyriform, some with ostiolate. *Peridium* 10–15  $\mu\text{m}$  wide, composed of 3–5 layers, hyaline to brown (#937463) cells of *textura angularis*. *Hamathecium* 1.5–3  $\mu\text{m}$  wide, composed of numerous, hyaline, filamentous, septate, branched, pseudoparaphyses. *Asci* 50–70  $\times$  7–11  $\mu\text{m}$  ( $\bar{x}$  = 58  $\times$  9  $\mu\text{m}$ ,  $n$  = 20), 8-spored, bitunicate, fissitunicate, clavate to cylindrical-clavate, short stalked, some with club-shape pedicel, apically rounded, with a small ocular chamber. *Ascospores* 12–16  $\times$  4–5  $\mu\text{m}$ , ( $\bar{x}$  = 14.4  $\times$  4.6  $\mu\text{m}$ ,  $n$  = 30), overlapping uni- to bi-seriately arranged, fusiform, straight or slightly curved, hyaline when immature and become pale brown (#e1af33) to dark-brown (#6e5031) when mature, mostly 1-septate, few 2 or 3-septate, constricted at each septum, with obviously guttulate. **Asexual morph:** Undetermined.

**Culture characteristics.** Ascospores germinated on PDA within 24 h and germ tubes arising from both ends. Colonies on PDA, reaching 4.5 cm diam. after two months of incubation at room temperature (22–26 °C), initially white (#f2f3f4) becoming grey (#bbbeb2) to dark brown (#6e5031) at maturity, dense, circular, slightly raised, smooth surface, radially fimbriate at the edge, reverse dark green (#3a4543) to brown (#937463).

**Material examined.** Pu’wen Town, Xishuangbanna, Yunnan Province, China, on a decaying branch of *Coffea arabica*, (22°31'18"N, 101°2'44"E, 856.89 m), 15 September 2021, LiLu, JHPW16 (ZHKU 22-0121, holotype), ZHKUCC 22-0210 = ZHKUCC 22-0211. GenBank number; ITS: OP450967, LSU: OP450973, *rpb2*: OP432243, SSU: OP450981, *tefl*- $\alpha$ : OP432247 (ZHKUCC 22-0210, ex-type); ITS: OP450968, LSU: OP450974, *rpb2*: OP432244, SSU: OP450982, *tefl*- $\alpha$ : OP432248 (ZHKUCC 22-0211).

**Notes.** Our phylogenetic analyses showed that *Nigrograna coffeae* forms an independent clade (100% ML, 1.00 BIPP, Fig. 1), and is phylogenetically related to *N. yasuniana* and *N. jinghongensis*. *Nigrograna yasuniana* was reported as endophytes from *Conceveiba guianensis* Aubl. in Ecuador, but there were not enough morphological data, the comparison of base pairs in ITS showed 3.4% differences (15/433 bp), LSU showed 1.5% differences (12/812bp), SSU only showed 0.3% differences (3/1028 bp), *rpb2* showed





**Figure 2.** *Nigrograna coffeae* (ZHKU 22-0121, holotype) **a, b** ascmata on the host substrate **c** a vertical section through an ascoma **d** peridium **e** hamathecium **f–k** asci **l** germinated ascospore **m** culture on pda from above and reverse **n–s** ascospores (arrows indicate the septa). Scale bars: 50  $\mu\text{m}$  (**c**); 10  $\mu\text{m}$  (**d–l**); 5  $\mu\text{m}$  (**n–s**).

14% differences (117/829 bp), and *tefl-α* showed 3.2% differences (31/954 bp) (Kolařík et al. 2017). *Nigrograna jinghongensis* was introduced as a saprobic fungus from woody litter in China, and our new isolate shares a similar size (12–16 × 4–5 μm vs 12–15 × 4–5.5 μm) and color (hyaline to dark brown vs yellowish-brown to brown) of ascospores with *N. jinghongensis* (Boonmee et al. 2021), but there are some significant differences in the size of the ascomata (90–140 μm high, 140–200 μm wide vs 300–400 μm high 220–300 μm wide) and the shape of ascospores (fusiform, straight or slightly curved vs ellipsoid) (Boonmee et al. 2021). Based on the sequence blast results, ITS, LSU and *rpb2* gene sequences were similar to *Nigrograna* sp., with 97.5% (MZ270683), 98.4% (MK762716), and 86% (MZ508421) respectively, SSU was similar to *N. mycophila* with 99% (KX650510), and *tefl-α* was similar to *N. yasuniana* with 96.6% (LN626670). Therefore, we introduce our new isolate as a new species *N. coffeae* based on both morphological characteristics and phylogenetic analyses.

***Nigrograna puerensis* L. Lu & Tibpromma, sp. nov.**

Index Fungorum number: IF559426

Facesoffungi Number: FoF12766


Fig. 3

**Etymology.** The specific epithet “*puerensis*” refers to the location Pu’er City, where the type species was collected.

**Holotype.** ZHKU 22-0122.

**Description.** *Saprobic* on decaying branch of *Coffea arabica*. **Sexual morph:** **Ascomata** 90–180 μm high, 90–150 μm wide ( $\bar{x}$  = 138 × 115 μm, n = 10), immersed, with only ostiolar necks visible on the host surface or erumpent, solitary, subglobose to ellipsoid, dark brown (#6e5031). **Peridium** 10–15 μm wide ( $\bar{x}$  = 13 μm, n = 15), outer layer consists of 2–3 layers of *textura prismatica*, brown (#937463) and thick-walled cells, inner layer hyaline with thin-walled cells. **Hamathecium** composed of numerous, 1.5–2 μm wide ( $\bar{x}$  = 1.8 μm, n = 20), filamentous, hyaline, septate, pseudoparaphyse. **Asci** 50–80 × 8–11 μm ( $\bar{x}$  = 66 × 9.5 μm, n = 20), 8-spored, bitunicate, fissitunicate, cylindrical to clavate, short pedicellate, apically rounded, with poorly developed ocular chamber. **Ascospores** 15–18 × 4–5 μm, ( $\bar{x}$  = 16 × 4.5 μm, n = 30), uni- to bi-seriately arranged, fusoid, apical cell and basal cell acute, and apical cell slightly wider than basal cell, straight or slightly curved, 1-septate, constricted at septum, guttulate, hyaline to yellow-brownish (#daceb8) when young, brownish (#937463) when mature. **Asexual morph:** Undetermined.

**Culture characteristics.** On PDA, colonies reached up to 4 cm diam. after two months at room temperature (22–26 °C). Colony dense, circular, slightly raised at the center, surface with white aerial mycelium, fluffy, with a serrate edge, grayish (#c9bfb3) to dark brown (#6e5031) from center to edge, reverse dark green (#3a4543) to dark brown (#6e5031).



**Figure 3.** *Nigrograna puerensis* (ZHKU 22-0122, holotype) **a, b** ascomata observed on host substrate **c** a vertical section through an ascoma **d** peridium **e** hamathecium **f–j** asci **k** germinated ascospore **l–p** ascospores **q** culture on PDA from above and reverse. Scale bars: 50  $\mu\text{m}$  (**c**); 30  $\mu\text{m}$  (**d**); 15  $\mu\text{m}$  (**e–k**); 5  $\mu\text{m}$  (**l–p**).

**Material examined.** Pu'er City, Yunnan Province, China, on a decaying branch of *Coffea arabica*, (22°36'2"N, 101°0'59"E, 1016.43 m), 16 September 2021, LiLu, Puer 1–4 (ZHKU 22-0122, holotype), ZHKUCC 22-0212 = ZHKUCC 22-0213. GenBank number; ITS: OP450969, LSU: OP450975, SSU: OP450983, *tef1- $\alpha$* : OP432249 (ZHKUCC 22-0212, ex-type); ITS: OP450970, LSU: OP450976, SSU: OP450984, *tef1- $\alpha$* : OP432250 (ZHKUCC 22-0213).

**Notes.** *Nigrograna puerensis* clusters with *N. carollii* with significant statistical support from ML 100% and BIPP 1.00. In morphology, our new strains best fit *Nigrograna* by having immersed ascomata, clavate and short pedicellate asci, and pale to brown, fusoid to narrowly ellipsoid, and septate ascospores (Jaklitsch and Voglmayr 2016; Zhang et al. 2020). Blast search results of ITS, LSU and *tef1- $\alpha$*  sequence data revealed that our taxon (ZHKUCC 22-0212) is similar to *N. mackinnonii* (96% MZ270697, 99% KJ605422, and 95% LT797087 respectively), while the similarity of SSU sequence to *N. carollii* is as high as 99%. Based on nucleotide comparisons, our isolate (ZHKUCC 22-0212) differs from *N. carollii* (CCF 4484) by 9/490 bp (1.8%) in ITS, 2/222 bp (1%) in LSU, 2/1306 bp (0.2%) in SSU, and 10/530 bp (2%) in *tef1- $\alpha$* . Unfortunately, for *N. carollii*, sufficient morphological data was not available to compare with our novel taxon which was isolated as an endophyte on living sapwood of wild *Hevea brasiliensis* Müll. Arg., and *N. mackinnonii* which was isolated as a human pathogen (de Gruyter 2012; Kolařík et al. 2017). In addition, the colony morphology of *N. carollii* on PDA is described as colonies plane, effuse, and light gray (Kolařík et al. 2017), while *N. puerensis* colony surface is seen as white aerial mycelium, fluffy, with a serrate edge, and grayish to dark brown from center to edge. Therefore, based on morphological and phylogenetic analyses, we introduce *N. puerensis* as a distinct new species.

### ***Nigrograna asexualis* L. Lu & Tibpromma, sp. nov.**

Index Fungorum number: IF559427


Facesoffungi Number: FoF12767

Fig. 4

**Etymology.** The species epithet '*asexualis*' refers to the asexual morph.

**Holotype.** ZHKU 22-0123.

**Description.** *Saprobic* on decaying branch of *Coffea arabica*. **Sexual morph:** Undetermined. **Asexual morph:** Coelomycetous. **Pycnidia** 100–230  $\mu$ m high, 120–180  $\mu$ m wide ( $\bar{x}$  = 156  $\times$  144  $\mu$ m, n = 10), globose to subglobose, or pyriform, immersed, solitary, unilocular, dark brown, papillate ostiole, appearing as black spots on host surface. **Pycnidial wall** 11–16  $\mu$ m wide ( $\bar{x}$  = 14  $\mu$ m, n = 15), brown (#937463), the wall with pseudoparenchymatous cells. **Conidiophores** arising from the pycnidial wall, up to 46  $\mu$ m long and 3–4.4  $\mu$ m wide ( $\bar{x}$  = 3.4  $\mu$ m, n = 25), filiform, septate, hyaline, simple to sparsely branched, with pegs along one or two sides and solitary phialides terminally. **Phialides** 3–6  $\times$  1–2  $\mu$ m ( $\bar{x}$  = 4.5  $\times$  1.5  $\mu$ m, n = 15), variable



**Figure 4.** *Nigrograna asexualis* (ZHKU 22-0123, holotype) **a, b** conidiomata on the host substrate **c, d** vertical sections of a conidioma **e** peridium **f, g** conidiophores with phialides **h** conidia **i** culture on PDA from above and reverse. Scale bars: 100  $\mu\text{m}$  (**c**); 50  $\mu\text{m}$  (**d**); 15  $\mu\text{m}$  (**e**); 30  $\mu\text{m}$  (**f**); 20  $\mu\text{m}$  (**g**); 10  $\mu\text{m}$  (**h**).

in shape, phialidic, discrete, ampulliform-lageniform-subcylindrical. **Conidia** 5–6.5  $\times$  3–4  $\mu\text{m}$  ( $\bar{x}$  = 5.5  $\times$  3.7  $\mu\text{m}$ ,  $n$  = 30), ellipsoidal, unicellular, aseptate with 1–2 granules, subhyaline, smooth-walled.

**Culture characteristics.** Conidium germinated on PDA within 24 h. Colonies growing on PDA reaching 5 cm diam. after two months at room temperature (22–26  $^{\circ}\text{C}$ ). Colony dense, circular, surface sparsely hairy, radially striate, with a fimbriate edge, yellowish (#eabf83) to pale brown (#e1af33) at the center and dark brown (#6e5031) at the margin, reverse dark brown (#6e5031).

**Material examined.** Pu'er City, Yunnan Province, China, on a decaying branch of *Coffea arabica*, (22°36'2"N, 101°0'59"E, 1016.43 m), 16 September 2021, LiLu, Puer 1-14 (ZHKU 22-0123, holotype), ZHKUCC 22-0214 = ZHKUCC 22-0215. GenBank number; ITS: OP450965, LSU: OP450971, *rpb2*: OP432241, SSU: OP450979, *tef1- $\alpha$* : OP432245 (ZHKUCC 22-0214, ex-type); ITS: OP450966, LSU: OP450972, *rpb2*: OP432242, SSU: OP450980, *tef1- $\alpha$* : OP432246 (ZHKUCC 22-0215).

**Notes.** In multi-gene phylogeny, *Nigrograna asexualis* formed a separate (68% ML, 0.97 BIPP) and distinct clade within *Nigrograna* (Fig. 1). Morphologically, *N. asexualis* conforms to the morphological characteristics of *Nigrograna* by having hyaline or subhyaline, long and branched conidiophores, solitary phialides, and aseptate, ellipsoidal or cylindrical conidia (Jaklitsch and Voglmayr 2016; Dayarathne et al. 2020; Wanasinghe et al. 2020). Blast results of the sequences show that ITS is similar to *N. fuscidula* with 89% (MH856004), and SSU is similar to *N. mycophila* with 99.8% (KX650510). *Nigrograna asexualis* is different from *N. fuscidula* and *N. mycophila* by its ellipsoidal conidia, but the similarities of these three species are hyaline, 1-celled, smooth-walled conidia forming on phialides (Jaklitsch and Voglmayr 2016). The LSU and *rpb2* sequences of our strain blast results are similar to *N. obliqua*, and the similarities are 98.9% (KX650560) and 87% (KX650579) respectively, but *N. obliqua* lacks the asexual morph (Jaklitsch and Voglmayr 2016). The *tef1- $\alpha$*  sequence of our strain is 95.8% (MF939615) similar to *N. locuta-pollinis*, which was isolated from hive-stored pollen of *Brassica campestris* L. that lacks morphology (Zhao et al. 2018). Therefore, we introduce *N. asexualis* as a distinct new species from coffee in China.

## Discussion

Members of *Nigrograna* are distributed worldwide in soil, wood, and other plant debris (Mapook et al. 2020), and the hotspots of *Nigrograna* are reported as Central and South America, where the taxa are also found as human pathogens (Kolařík 2018; Puing et al. 2020). To date, five *Nigrograna* species viz. *N. cangshanensis* (decaying wood, Yunnan), *N. jinghongensis* (dead woody litter, Yunnan), *N. kunmingensis* (dead twigs of *Gleditsia sinensis* Lam., Yunnan), *N. magnoliae* (living branches of *Magnolia denudate* Desr., Yunnan), and *N. locuta-pollinis* (hive-stored pollen, Hubei) have been isolated from different hosts in China (Tibpromma et al. 2017; Zhao et al. 2018; Wanasinghe et al. 2020; Boonmee et al. 2021; Zhou et al. 2022). In this study, three new saprobic fungi were isolated from decaying branches of *Coffea arabica* in Yunnan Province, China, and this is the first report of *Nigrograna* species from coffee.

Species of *Nigrograna* are morphologically very similar and overlapping, hence can be interpreted as cryptic species. Therefore, it is difficult to delimit the species based only on their morphological characteristics (Jaklitsch and Voglmayr 2016; Zhang et al. 2020). In our research, we found that *N. coffeae* and *N. puerensis* have similar morphology, but in phylogeny, they are distributed differently within *Nigrograna*. This confirms the view of Jaklitsch and Voglmayr (2016) that the gene sequences are important and crucial for the identification of taxa at the genus and the species level.

## Acknowledgements

Li Lu thanks Mae Fah Luang University for the award of a fee-less scholarship. The Center for Yunnan Plateau Biological Resources Protection and Utilization, College of Biological Resource and Food Engineering, Qujing Normal University is thanked for the facilities provided for the research work. Dr. Shaun Pennycook is thanked for his advice on new fungi names. Dai Dong-qin thanks the National Natural Science Foundation of China (No. NSFC 31760013, 31950410558), and High-Level Talent Recruitment Plan of Yunnan Provinces (“Young Talents” Program). Samantha C. Karunarathna thanks the National Natural Science Foundation of China grant number 32260004 for the support. Nakarin Suwannarach thanks Chiang Mai University, Thailand for financial support. Saowaluck Tibpromma thanks “the most cited article award” for allowing one free publication in MycoKeys.

## References

- Ahmed SA, van de Sande WWJ, Stevens DA, Fahal A, van Diepeningen AD, Menken SBJ, de Hoog GS (2014) Revision of agents of black-grain eumycetoma in the order of Pleosporales. *Persoonia* 33: 141–154. <https://doi.org/10.3767/003158514X684744>
- Arias RM, Abarca GH (2014) Fungal diversity in coffee plantation systems and in a tropical montane cloud forest in Veracruz, Mexico. *Agroforestry Systems* 88: 921–933. <https://doi.org/10.1007/s10457-014-9736-z>
- Barr ME (1987) *Prodromus to class Loculoascomycetes*. Published by the Author, Amherst, Massachusetts; University of Massachusetts, U.S.A., 168 pp.
- Bhunjun CS, Niskanen T, Suwannarach N, Wannathes N, Chen YJ, McKenzie EH, Maharachchikumbura SS, Buyck B, Zhao CL, Fan YG, Zhang JY, Dissanayake AJ, Marasinghe DS, Jayawardena RS, Kumla J, Padamsee M, Chen YY, Liimatainen K, Ammirati JF, Phukhamsakda C, Liu JK, Phonrob W, Randrianjohany E, Hongsan S, Cheewangkoon R, Bundhun D, Khuna S, Yu WJ, Deng LS, Lu YZ, Hyde KD, Lumyong S (2022) The numbers of fungi: Are the most speciose genera truly diverse? *Fungal Diversity* 27: 1–76. <https://doi.org/10.1007/s13225-022-00501-4>
- Boonmee S, Wanasinghe DN, Calabon MS, Huanraluek N, Chandrasiri SK, Jones GE, Rossi W, Leonardi M, Singh SK, Rana S, Singh PN, Maurya DK, Lagashetti AC, Choudhary D, Dai YC, Zhao CL, Mu YH, Yuan HS, He SH, Phookamsak R, Jiang HB, Martín MP, Dueñas M, Telleria MT, Kaľucka LK, Jagodziński AM, Liimatainen K, Pereira DS, Phillips AJL, Suwannarach N, Kumla J, Khuna S, Lumyong S, Potter TB, Shivas RG, Sparks AH, Vaghefi N, Abdel-Wahab MA, Abdel-Aziz FA, Li GJ, Lin WF, Singh U, Bhatt RP, Lee HB, Nguyen TTT, Kirk PM, Dutta AK, Acharya K, Sarma VV, Niranjana M, Rajeshkumar KC, Ashtekar N, Lad S, Wijayawardene NN, Bhat DJ, Xu RJ, Wijesinghe SN, Shen HW, Luo ZL, Zhang JY, Sysouphanthong P, Thongklang N, Bao DF, Aluthmuhandiram JVS, Abdollahzadeh J, Javadi A, Dovana F, Usman M, Khalid AN, Dissanayake AJ, Telagathoti A, Probst M, Peintner U, Garrido-Benavent I, Bóna L, Merényi Z, Boros L, Zoltán B, Stielow JB, Jiang N, Tian CM, Shams E, Dehghanizadeh F, Pordel A, Javan-Nikkhah M,

- Denchev TT, Denchev CM, Kemler M, Begerow D, Deng CY, Harrower E, Bozorov T, Kholmuradova T, Gafforov Y, Abdurazakov A, Xu JC, Mortimer PE, Ren GC, Jeewon R, Maharachchikumbura SSN, Phukhamsakda C, Mapook A, Hyde KD (2021) Fungal diversity notes 1387–1511: Taxonomic and phylogenetic contributions on genera and species of fungal taxa. *Fungal Diversity* 111: 1–335. <https://doi.org/10.1007/s13225-021-00489-3>
- Borelli D (1976) *Pyrenochaeta mackinnonii* nova species agente de micetoma. *Castellania* 4: 227–234.
- Dayarathne MC, Jones E, Maharachchikumbura S, Devadatha B, Sarma V, Khongphinitbunjong K, Chomnunti P, Hyde KD (2020) Morpho-molecular characterization of microfungi associated with marine based habitats. *Mycosphere* 11: 1–188. <https://doi.org/10.5943/mycosphere/11/1/1>
- de Gruyter JH (2012) Revised taxonomy of *Phoma* and allied genera. Wageningen University and Research, 1–181.
- de Gruyter J, Woudenberg JH, Aveskamp MM, Verkley GJ, Groenewald JZ, Crous PW (2010) Systematic reappraisal of species in *Phoma* section *Paraphoma*, *Pyrenochaeta* and *Pleurophoma*. *Mycologia* 102: 1066–1081. <https://doi.org/10.3852/09-240>
- de Gruyter J, Woudenberg JH, Aveskamp MM, Verkley GJ, Groenewald JZ, Crous PW (2013) Redisposition of *Phoma*-like anamorphs in Pleosporales. *Studies in Mycology* 75: 1–36. <https://doi.org/10.3114/sim0004>
- Dighton J (2003) Fungi in ecosystem processes. Marcel Dekker, New York, 434 pp. <https://doi.org/10.1201/9781315371528>
- Dissanayake AJ, Bhunjun CS, Maharachchikumbura SSN, Liu JK (2020) Applied aspects of methods to infer phylogenetic relationships amongst fungi. *Mycosphere* 11: 2652–2676. <https://doi.org/10.5943/mycosphere/11/1/18>
- Hawksworth DL, Lücking R (2017) Fungal diversity revisited: 2.2 to 3.8 million species. *Microbiology Spectrum* 5(4). <https://doi.org/10.1128/microbiolspec>
- Hongsanan S, Hyde KD, Phookamsak R, Wanasinghe DN, McKenzie EHC, Sarma VV, Boonmee S, Lücking R, Bhat DJ, Liu NG, Tennakoon DS, Pem D, Karunarathna A, Jiang SH, Jones EBG, Phillips AJL, Manawasinghe IS, Tibpromma S, Jayasiri SC, Sandamali DS, Jayawardena RS, Wijayawardene NN, Ekanayaka AH, Jeewon R, Lu Y, Dissanayake AJ, Zeng XY, Luo ZL, Tian Q, Phukhamsakda C, Thambugala KM, Dai DQ, Samarakoon MC, Chethana KWT, Ertz D, Doilom M, Liu JK, Pérez-Ortega S, Suija A, Senwana C, Wijesinghe SN, Niranjana M, Zhang SN, Ariyawansa HA, Jiang HB, Zhang JF, Norphanphoun C, de Silva NI, Thiagaraja V, Zhang H, Bezerra JDP, Miranda-González R, Aptroot A, Kashiwadani H, Harishchandra D, Sérusiaux E, Abeywickrama PD, Bao DF, Devadatha B, Wu HX, Moon KH, Gueidan C, Schumm F, Bundhun D, Mapook A, Monkai J, Bhunjun CS, Chomnunti P, Suetrong S, Chaiwan N, Dayarathne MC, Yang J, Rathnayaka AR, Xu JC, Zheng JS, Liu G, Feng Y, Xie N (2020a) Refined families of Dothideomycetes: Dothideomycetidae and Pleosporomycetidae. *Mycosphere* 11: 1553–2107. <https://doi.org/10.5943/mycosphere/11/1/13>
- Hongsanan S, Hyde KD, Phookamsak R, Wanasinghe DN, McKenzie EHC, Sarma VV, Lücking R, Boonmee S, Bhat DJ, Liu NG, Tennakoon DS, Pem D, Karunarathna A, Jiang SH, Jones EBG, Phillips AJL, Manawasinghe IS, Tibpromma S, Jayasiri SC, Sandamali DS, Jayawardena RS, Wijayawardene NN, Ekanayaka AH, Jeewon R, Lu YZ, Phukhamsakda C, Dissanayake



- AJ, Zeng XY, Luo ZL, Tian Q, Thambugala KM, Dai DQ, Samarakoon MC, Chethana KWT, Ertz D, Doilom M, Liu JK, Pérez-Ortega S, Suija A, Senwana C, Wijesinghe SN, Niranan M, Zhang SN, Ariyawansa HA, Jiang HB, Zhang JF, Norphanphoun C, de Silva NI, Thiyagaraja V, Zhang H, Bezerra JDP, Miranda-González R, Aptroot A, Kashiwadani H, Harishchandra D, Sérusiaux E, Abeywickrama PD, Bao DF, Devadatha B, Wu HX, Moon KH, Gueidan C, Schumm F, Bundhun D, Mapook A, Monkai J, Bhunjun CS, Chomnunti P, Suetrong S, Chaiwan N, Dayarathne MC, Yang J, Rathnayaka AR, Xu JC, Zheng JS, Liu G, Feng Y, Xie N (2020b) Refined families of Dothideomycetes: Orders and families incertae sedis in Dothideomycetes. *Fungal Diversity* 105(1): 17–318. <https://doi.org/10.1007/s13225-020-00462-6>
- Hyde KD, Norphanphoun C, Abreu VP, Bazzicalupo A, Chethana KWT, Clericuzio M, Dayarathne MC, Dissanayake AJ, Ekanayaka AH, He MQ, Hongsan S, Huang SK, Jayasiri SC, Jayawardena RS, Karunarathna A, Konta S, Kušan I, Lee H, Li J, Lin CG, Liu NG, Lu YZ, Luo ZL, Manawasinghe IS, Mapook A, Perera RH, Phookamsak R, Phukhamsakda C, Siedlecki I, Soares AM, Tennakoon DS, Tian Q, Tibpromma S, Wanasinghe DN, Xiao YP, Yang J, Zeng XY, Abdel-Aziz FA, Li WJ, Senanayake IC, Shang QJ, Daranagama DA, De Silva NI, Thambugala KM, Abdel-Wahab MA, Bahkali AH, Berbee ML, Boonmee S, Bhat DJ, Bulgakov TS, Buyck B, Camporesi E, Castañeda-Ruiz RF, Chomnunti P, Doilom M, Dovana F, Gibertoni TB, Jadan M, Jeewon R, Jones GEB, Kang JC, Karunarathna SC, Lim YW, Liu JK, Liu ZY, Plautz Jr HL, Lumyong S, Maharachchikumbura SSN, Matočec N, McKenzie EHC, Mešič A, Miller D, Pawłowska J, Pereira OL, Promputtha I, Romero AI, Ryvarde L, Su HY, Suetrong S, Tkalc̃ec Z, Vizzini A, Wen TC, Wisitrasameewong K, Wrzosek M, Xu JC, Zhao Q, Zhao RL, Mortimer PE (2017) Fungal diversity notes 603–708: Taxonomic and phylogenetic notes on genera and species. *Fungal Diversity* 87: 1–235. <https://doi.org/10.1007/s13225-017-0391-3>
- Hyde KD, Jeewon R, Chen YJ, Bhunjun CS, Calabon MS, Jiang HB, Lin CG, Norphanphoun C, Sysouphanthong P, Pem D, Tibpromma S, Zhang Q, Doilom M, Jayawardena RS, Liu JK, Maharachchikumbura SSN, Phookamsakda C, Phookamsak R, Al-Sadi AM, Thongklang N, Wang Y, Gafarov Y, Jones EBG, Lumyong S (2020) The numbers of fungi: Is the descriptive curve flattening? *Fungal Diversity* 103(1): 219–271. <https://doi.org/10.1007/s13225-020-00458-2>
- Index Fungorum (2022) Index Fungorum. <http://www.indexfungorum.org/Names/Names.asp> [Retrieved 20 September 2022]
- Jaklitsch WM, Voglmayr H (2016) Hidden diversity in *Thyridaria* and a new circumscription of the Thyridariaceae. *Studies in Mycology* 85(1): 35–64. <https://doi.org/10.1016/j.simyco.2016.09.002>
- Jayasiri SC, Hyde KD, Ariyawansa HA, Bhat J, Buyck B, Cai L, Dai YC, Abd-Elsalam KA, Ertz D, Hidayat I, Jeewon R, Jones EBG, Bahkali AH, Karunarathna SC, Liu JK, Luangsa-ard JJ, Lumbsch HT, Maharachchikumbura SSN, McKenzie EHC, Moncalvo JM, Ghobad-Nejhad M, Nilsson H, Pang KA, Pereira OL, Phillips AJL, Raspé O, Rollins AW, Romero AI, Etayo J, Selçuk F, Stephenson SL, Suetrong S, Taylor JE, Tsui CKM, Vizzini A, Abdel-Wahab MA, Wen TC, Boonmee S, Dai DQ, Daranagama DA, Dissanayake AJ, Ekanayaka AH, Fryar SC, Hongsan S, Jayawardena RS, Li WJ, Perera RH, Phookamsak R, de Silva NI, Thambugala KM, Tian Q, Wijayawardena NN, Zhao RL, Zhao Q, Kang JC, Promputtha I (2015) The Faces of Fungi database: Fungal names linked with morphol-

- ogy, phylogeny and human impacts. *Fungal Diversity* 74: 3–18. <https://doi.org/10.1007/s13225-015-0351-8>
- Jiang HB, Phookamsak R, Hyde KD, Mortimer PE, Xu JC, Kakumyan P, Karunarathna SC, Kumla JA (2021) A taxonomic appraisal of bambusicolous fungi in Occultibambusaceae (Pleosporales, Dothideomycetes) with new collections from Yunnan Province, China. *Life* 11(9): 932. <https://doi.org/10.3390/life11090932>
- Kolařík M (2018) New taxonomic combinations in endophytic representatives of the genus *Nigrograna*. *Czech Mycology* 70: 123–126.
- Kolařík M, Spakowicz DJ, Gazis R, Shaw J, Kubátová A, Nováková A, Chudíčková M, Forcina GC, Kang KW, Kelnarová I, Skaltsas D (2017) *Biatriospora* (Ascomycota: Pleosporales) is an ecologically diverse genus including facultative marine fungi and endophytes with biotechnological potential. *Plant Systematics and Evolution* 303(1): 35–50. <https://doi.org/10.1007/s00606-016-1350-2>
- Li ZW (2014) Issues and suggestions on building an international brand of “Pu’er Coffee”. Selected papers on rural agricultural reform, innovation and agricultural modernization 2: 719–721.
- Liu YJ, Whelen S, Hall BD (1999) Phylogenetic relationships among ascomycetes: Evidence from an RNA polymerase II subunit. *Molecular Biology and Evolution* 16: 1799–1808. <https://doi.org/10.1093/oxfordjournals.molbev.a026092>
- Liu JK, Hyde KD, Jeewon R, Phillips AJ, Maharachchikumbura SS, Ryberg M, Liu ZY, Zhao Q (2017) Ranking higher taxa using divergence times: A case study in Dothideomycetes. *Fungal Diversity* 84(1): 75–99. <https://doi.org/10.1007/s13225-017-0385-1>
- Lu L, Tibpromma S, Karunarathna SC, Thiyagaraja V, Xu JC, Jayawardena RS, Lumyong S, Hyde KD (2021) Taxonomic and phylogenetic appraisal of a novel species and a new record of Stictidaceae from coffee in Yunnan Province, China. *Phytotaxa* 528(2): 111–124. <https://doi.org/10.11646/phytotaxa.528.2.4>
- Lu L, Tibpromma S, Karunarathna SC, Jayawardena RS, Lumyong S, Xu JC, Hyde KD (2022a) Comprehensive review of fungi on coffee. *pathogens* 11: 411. <https://doi.org/10.3390/pathogens11040411>
- Lu L, Karunarathna SC, Dai DQ, Xiong YR, Suwannarach N, Stephenson SL, Elgorban AM, Al-Rejaie S, Jayawardena RS, Tibpromma S (2022b) Description of four novel species in Pleosporales associated with coffee in Yunnan, China. *Journal of Fungi* 8(10): 1113. <https://doi.org/10.3390/jof8101113>
- Luttrell ES (1955) The ascostromatic Ascomycetes. *Mycologia* 47: 511–532. <https://doi.org/10.1080/00275514.1955.12024473>
- Mapook A, Hyde KD, McKenzie EH, Jones EB, Bhat DJ, Jeewon R, Stadler M, Samarakoon MC, Malaithong M, Tanunchai B, Buscot F (2020) Taxonomic and phylogenetic contributions to fungi associated with the invasive weed *Chromolaena odorata* (Siam weed). *Fungal Diversity* 101(1): 1–175. <https://doi.org/10.1007/s13225-020-00444-8>
- Miller MA, Pfeiffer W, Schwartz T (2012) The CIPRES science gateway: Enabling high-impact science for phylogenetics researchers with limited resources. *Proceedings of the 1<sup>st</sup> Conference of the Extreme Science and Engineering Discovery Environment: Bridging from the Extreme to the Campus and Beyond*, 1–8. <https://doi.org/10.1145/2335755.2335836>

- Neilson J, Wang JHZ (2019) China and the changing economic geography of coffee value chains. *Singapore Journal of Tropical Geography* 40(3): 429–451. <https://doi.org/10.1111/sjtg.12279>
- Nylander JAA (2004) MrModeltest v2. Program distributed by the author Evolutionary Biology Centre, Uppsala University, Uppsala.
- Pandey A, Das N, Kumar B, Rinu K, Trivedi P (2008) Phosphate solubilization by *Penicillium* spp. isolated from soil samples of Indian Himalayan region. *World Journal of Microbiology & Biotechnology* 24(1): 97–102. <https://doi.org/10.1007/s11274-007-9444-1>
- Phukhamsakda C, Nilsson RH, Bhunjun CS, de Farias AR, Sun YR, Wijesinghe SN, Raza M, Bao DF, Lu L, Tibpromma S, Dong W, Tennakoon DS, Tian XG, Xiong YR, Karunaratna SC, Cai L, Luo ZL, Wang Y, Manawasinghe IS, Camporesi E, Kirk PM, Promputtha I, Kuo CH, Su HY, Doilom M, Li Y, Fu YP, Hyde KD (2022) The numbers of fungi: Contributions from traditional taxonomic studies and challenges of metabarcoding. *Fungal Diversity* 28: 1–60. <https://doi.org/10.1007/s13225-022-00502-3>
- Puing AG, Couture-Cossette A, Wang AX, Zygourakis CC, Cheng X, Stevens BA, Banaei N, Novoa RA, Ho DY, Subramanian AK (2020) Simultaneous coccidioidomycosis and phaeohyphomycosis in a kidney transplant recipient: A case report and literature review. *Transplant Infectious Disease* 22(6): e13365. <https://doi.org/10.1111/tid.13365>
- Rambaut A (2014) FigTree v1. 4.2, a graphical viewer of phylogenetic trees.
- Rehner SA, Buckley EA (2005) *Beauveria* phylogeny inferred from nuclear ITS and *EF1- $\alpha$*  sequences: Evidence for cryptic diversification and links to *Cordyceps* teleomorphs. *Mycologia* 97: 84–98. <https://doi.org/10.1080/15572536.2006.11832842>
- Ronquist F, Huelsenbeck JP (2003) MrBayes 3: Bayesian phylogenetic inference under mixed models. *Bioinformatics* 19: 1572–1574. <https://doi.org/10.1093/bioinformatics/btg180>
- Ronquist F, Teslenko M, Van Der Mark P, Ayres DL, Darling A, Höhna S, Larget B, Liu L, Suchard MA, Huelsenbeck JP (2012) MrBayes 3.2: Efficient Bayesian phylogenetic inference and model choice across a large model space. *Systematic Biology* 61: 539–542. <https://doi.org/10.1093/sysbio/sys029>
- Senanayake IC, Rathnayaka AR, Marasinghe DS, Calabon MS, Gentekaki E, Lee HB, Hurdeal VG, Pem D, Dissanayake LS, Wijesinghe SN, Bundhun D, Nguyen TT, Goonasekara ID, Abeywickrama PD, Bhunjun CS, Jayawardena RS, Wanasinghe DN, Jeewon R, Bhat DJ, Xiang MM (2020) Morphological approaches in studying fungi: Collection, examination, isolation, sporulation and preservation. *Mycosphere* 11: 2678–2754. <https://doi.org/10.5943/mycosphere/11/1/20>
- Sharma H (2020) A detail chemistry of coffee and its analysis. In *Coffee-Production and Research*, 79 pp. <https://doi.org/10.5772/intechopen.91725>
- Stamatakis A (2014) RAXML Version 8: A tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics* 30: 1312–1313. <https://doi.org/10.1093/bioinformatics/btu033>
- Tennakoon DS, Kuo CH, Maharachchikumbura SS, Thambugala KM, Gentekaki E, Phillips AJ, Bhat DJ, Wanasinghe DN, de Silva NI, Promputtha I, Hyde KD (2021) Taxonomic and phylogenetic contributions to *Celtis formosana*, *Ficus ampelas*, *F. septica*, *Macaranga tanarius* and *Morus australis* leaf litter inhabiting microfungi. *Fungal Diversity* 108: 1–215. <https://doi.org/10.1007/s13225-021-00474-w>

- Tibpromma S, Hyde KD, Jeewon R, Maharachchikumbura SSN, Liu JK, Bhat DJ (2017) Fungal diversity notes 491–602: Taxonomic and phylogenetic contributions to fungal taxa. *Fungal Diversity* 83: 1–261. <https://doi.org/10.1007/s13225-017-0378-0>
- Vaidya G, Lohman DJ, Meier R (2011) SequenceMatrix: Concatenation software for the fast assembly of multi-gene datasets with character set and codon information. *Cladistics* 27(2): 171–180. <https://doi.org/10.1111/j.1096-0031.2010.00329.x>
- Vilgalys R, Hester M (1990) Rapid genetic identification and mapping of enzymatically amplified ribosomal DNA from several *Cryptococcus* species. *Journal of Bacteriology* 172: 4238–4246. <https://doi.org/10.1128/jb.172.8.4238-4246.1990>
- Wanasinghe DN, Wijayawardene NN, Xu J, Cheewangkoon R, Mortimer PE (2020) Taxonomic novelties in Magnolia-associated pleosporalean fungi in the Kunming Botanical Gardens (Yunnan, China). *PLoS ONE* 15(7): e0235855. <https://doi.org/10.1371/journal.pone.0235855>
- White TJ, Bruns T, Lee SJWT, Taylor JL (1990) Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. *PCR protocols: a guide to methods and applications* 18: 315–322. <https://doi.org/10.1016/B978-0-12-372180-8.50042-1>
- Wijayawardene NN, Hyde KD, Dai DQ, Sánchez-García M, Goto BT, Saxena RK, Erdođdu M, Selçuk F, Rajeshkumar KC, Aptroot A, Błaszowski J, Boonyuen N, da Silva GA, de Souza FA, Dong W, Ertz D, Haelewaters D, Jones EBG, Karunarathna SC, Kirk PM, Kukwa M, Kumla J, Leontyev DV, Lumbsch HT, Maharachchikumbura SSN, Marguno F, Martínez-Rodríguez P, Mešić A, Monteiro JS, Oehl F, Pawłowska J, Pem D, Pfliegler WP, Phillips AJL, Pošta A, He MQ, Li JX, Raza M, Sruthi OP, Suetrong S, Suwannarach N, Tedersoo L, Thiyagaraja V, Tibpromma S, Tkalčec Z, Tokarev YS, Wanasinghe DN, Wijesundara DSA, Wimalaseana SDMK, Madrid H, Zhang GQ, Gao Y, Sánchez-Castro I, Tang LZ, Stadler M, Yurkov A, Thines M (2022) Outline of fungi and fungus-like taxa–2021. *Mycosphere* 13: 53–453. <https://doi.org/10.5943/mycosphere/13/1/2>
- Yang EF, Tibpromma S, Karunarathna SC, Phookamsak R, Xu JC, Zhao ZX, Karunanayake C, Promputtha I (2022) Taxonomy and phylogeny of novel and extant taxa in Pleosporales associated with *Mangifera indica* from Yunnan, China (Series I). *Journal of Fungi* 8(2): 152. <https://doi.org/10.3390/jof8020152>
- Zhang Y, Schoch CL, Fournier J, Crous PW, De Gruyter J, Woudenberg JH, Hirayama K, Tanaka K, Pointing SB, Spathafora JW, Hyde KD (2008) Multi-locus phylogeny of Pleosporales: A taxonomic, ecological and evolutionary re-evaluation. *Studies in Mycology* 61(1): 111–119. <https://doi.org/10.3114/sim.2009.64.04>
- Zhang Y, Crous PW, Schoch CL, Hyde KD (2012) Pleosporales. *Fungal Diversity* 53(1): 1–221. <https://doi.org/10.1007/s13225-011-0117-x>
- Zhang H, Li J, Zhou H, Chen Z, Song G, Peng Z, Pereira AP, Silva MC, Varzea VM (2014) Arabica coffee production in the Yunnan Province of China. *Proceedings of the 24<sup>th</sup> International Conference on Coffee Science (ASIC)*.
- Zhang JF, Liu JK, Thambugala KM, Yang J, Meng ZH, Liu ZY (2020) Two new species and a new record of *Nigrograna* (Nigrogranaceae, Pleosporales) from China and Thailand. *Mycological Progress* 19: 1365–1375. <https://doi.org/10.1007/s11557-020-01633-0>

- Zhao YZ, Zhang ZF, Cai L, Peng JW, Liu F (2018) Four new filamentous fungal species from newly-collected and hive stored bee pollen. *Mycosphere* 9: 1089–1116. <https://doi.org/10.5943/mycosphere/9/6/3>
- Zhou LW, Wang XW, Liu SL, Li GJ, Deng CY, Rossi W, Leonardi M, Liimatainen K, Kekki T, Niskanen T, Smith ME, Ammirati J, Bojantchev D, Abdel-Wahab MA, Zhang M, Tian E, Lu YZ, Zhang JY, Ma J, Dutta AK, Acharya K, Du TY, Xu JZ, Kim JS, Lim YW, Gerlach A, Zeng NK, Han YX, Razaghi P, Raza M, Cai L, Calabon MS, Jones EBG, Saha R, Kumar TKA, Krishnapriya K, Thomas A, Kaliyaperumal M, Kezo K, Gunaseelan S, Singh SK, Singh PN, Lagashetti AC, Pawar KS, Jiang SH, Zhang C, Zhang H, Qing Y, Bau T, Ramirez NA, Niveiro N, Li MX, Yang ZL, Wu G, Tarafder E, Tennakoon DS, Kuo CH, da Silva TM, Souza-Motta CM, Bezerra JDP, He G, Ji XH, Suwannarach N, Kumla J, Lumyong S, Wannathes N, Rana S, Hyde KD (2022) Fungal diversity notes: taxonomic and phylogenetic contributions on genera and species of fungal taxa. *Fungal Diversity* (in press).