



Brief communication: The hidden labyrinth: deep groundwater in Wright Valley, Antarctica

Hilary A. Dugan¹, Peter T. Doran², Denys Grombacher³, Esben Auken³, Thue Bording³, Nikolaj Foged³, Neil Foley⁴, Jill Mikucki⁵, Ross A. Virginia⁶, and Slawek Tulaczyk⁷

¹Center for Limnology, University of Wisconsin-Madison, 53706 Madison, WI, USA

²Department of Geology and Geophysics, Louisiana State University, 70802 Baton Rouge, LA, USA

³Department of Geoscience, Aarhus University, 8000 Aarhus, Denmark

⁴Environmental Sciences Department, University of Montana Western, 59725 Dillon, MT, USA

⁵Department of Microbiology, University of Tennessee, 37996 Knoxville, TN, USA

⁶Department of Environmental Studies, Dartmouth College, 03755 Hanover, NH, USA

⁷Department of Earth and Planetary Sciences, University of California, Santa Cruz, 95064 Santa Cruz, CA, USA

Correspondence: Hilary A. Dugan (hdugan@wisc.edu)

Received: 27 April 2022 – Discussion started: 4 July 2022

Revised: 14 November 2022 – Accepted: 21 November 2022 – Published: 16 December 2022

Abstract. Since the 1960s, a deep groundwater system in Wright Valley, Antarctica, has been the hypothesized source of brines to hypersaline Don Juan Pond and Lake Vanda, both of which are rich in calcium and chloride. Modeling studies do not support other possible mechanisms, such as evaporative processes, that could have led to the current suite of ions present in both waterbodies. In 2011 and 2018, an airborne electromagnetic survey was flown over Wright Valley to map subsurface resistivity (down to 600 m) in exploration of liquid water. The surveys revealed widespread unfrozen brine in the subsurface near Lake Vanda, Don Juan Pond, and the North Fork of Wright Valley. While our geophysical survey can neither confirm nor deny deep groundwater connectivity between Lake Vanda and Don Juan Pond, it does point to the potential for deep valley-wide brine, likely within the Ferrar Dolerite formation.

1 Introduction

Wright Valley in the McMurdo Dry Valleys of East Antarctica is notable for being home to the Onyx River, the longest river in Antarctica, Lake Vanda, the deepest and warmest lake in Antarctica, and Don Juan Pond, one of the saltiest bodies of water on Earth. It is also notable for having a long-standing debate as to the origin and connectivity of ground-

water brines in the valley. Both Don Juan Pond and Lake Vanda hold salty calcium-chloride-rich brines, distinct from the sodium-chloride brines seen to the south in Taylor Valley (e.g., Blood Falls, Mikucki et al., 2009) and to the north in Victoria Valley (e.g., Lake Vida, Dugan et al., 2015a).

The earliest limnological investigations of Wright Valley in the 1960s revealed saline waters in both closed-basin Don Juan Pond and the bottom waters of closed-basin Lake Vanda (Wilson and Wellman, 1962; Armitage and House, 1962; Meyer et al., 1962), but it was not until the summers of 1973–1974 and 1974–1975 with the undertaking of the Dry Valleys Drilling Project (DVDP) that subsurface hydrology was able to be investigated. Important for our investigation were three boreholes. DVDP4A was drilled in the center of Lake Vanda, DVDP13 was drilled immediately to the west of Don Juan Pond, and DVDP14 was drilled, at the time, to the west of Lake Vanda in the North Fork of Wright Valley (it is now underwater).

In 1973–1974, DVDP4A penetrated 17.4 m beneath the sediment–water interface of the bottom of Lake Vanda. The core consisted of 3.5 m of lacustrine sediments, underlain by 2.2 m of glacial sediments, 6.7 m of marine sediments, and eventually basement rock (Cartwright et al., 1974). Cartwright et al. (1974) hypothesized the presence of a groundwater system flowing from Lake Vanda to the west. They later invalidated this hypothesis based on fluid poten-

tials measured in the sediments. They found that there was no change in total head between the lake bottom and the underlying sand and gravel, indicating that groundwater was flowing laterally at the borehole location, and if there was an area of groundwater discharge or recharge with the lake, the borehole did not intersect it (Cartwright and Harris, 1981).

During the 1974–1975 summer, DVDP14 was drilled 1.5 km west of Lake Vanda. Total penetration depth was 78 m, with basement rock reached at 27.94 m depth (Mudrey Jr. et al., 1975). All water and sediments in the borehole were solidly frozen, which, at the time, was thought to negate the previous summer's hypothesis that water flowed westward from Lake Vanda into the North Fork. Water was later squeezed from the 28 m of sediments and found to be hypersaline (50–180 ppt) (McGinnis et al., 1981).

Also during the 1974–1975 summer, at Don Juan Pond, DVDP13 was drilled 75 m deep, reaching highly fractured basement rock at 12.7 m depth (Chapman-Smith, 1975). Water was encountered in the borehole and consistently rose to a level 0.65–0.80 m higher than the surface of the pond, which indicated that the pond was receiving groundwater discharge. The borehole water had a salinity of 200–225 ppt (McGinnis et al., 1981; Harris, 1981), which is less than the salinity of Don Juan Pond. In December 1978, the borehole was revisited. Once cleared of ice, -15.5°C water flowed from the hole for 3 d as an artesian well before being shut off (McGinnis, 1979). It was hypothesized that the fractured and highly mineralized Ferrar Dolerite might be a confined aquifer, sourcing water from the base of the Antarctic Ice Sheet (McGinnis, 1979; Harris, 1981). The water levels and salinity of the pond vary as well, which suggests a hydrological driver beyond surface conditions (Harris et al., 1979). Geophysical resistivity logs of the DVDP13 borehole showed higher resistivity in the Ferrar Dolerite from 23 to 33 m depth (McGinnis et al., 1981). They suggested that this may be a zone of high permeability and high flow and therefore have less diffusion of salts.

Since the DVDP era, a number of scientists have investigated the origin of salts in the lacustrine systems of the Dry Valleys (Green and Lyons, 2009; Lyons et al., 2005). The oddity of calcium-chloride brines in Wright Valley has led to a special focus on Don Juan Pond. Remote sensing of the surface surrounding Don Juan Pond and laboratory experiments with salts have shown that brines can be generated at the surface via deliquescence in CaCl_2 -rich sediments (Gough et al., 2017). Surface discharge into the pond has been shown to be correlated with meteorological conditions, implying that surface hydrology plays a role in the water balance of the pond (Dickson et al., 2013). However, most mixing models predict that the salts in Lake Vanda and Don Juan Pond could not have originated solely from surface inputs from the Onyx River or the surrounding catchment and must be derived in part from a deep groundwater source (Green and Canfield, 1984; Green and Lyons, 2009; Carlson et al., 1990). Furthermore, if a regional groundwater sys-

tem did exist that connected both waterbodies, water should recharge from higher-elevation Don Juan Pond (113 m a.s.l.) and discharge to lower-elevation Lake Vanda (90 m a.s.l.).

Evidence for deep groundwater flow in Wright Valley is presented by Toner et al. (2017), who developed a model of brine evolution in Don Juan Pond and found that the ionic ratios in Don Juan Pond could only be explained by the upwelling and evaporation of deep groundwater and that the turnover time must be relatively short (<1 year). They concluded the presence of a regional groundwater flow system, but note that their investigation is equivocal on whether a groundwater connection exists between Don Juan Pond and Lake Vanda. Lake Vanda has much higher $\text{Mg} : \text{Cl}$ and $\text{K} : \text{Cl}$ ratios than Don Juan Pond, which suggests that the lake undergoes closed-basin evaporation without a significant outflow, unlike Don Juan Pond, where they hypothesize that groundwater flows both into and out of the pond (Toner et al., 2017).

The aforementioned studies in Wright Valley were limited in their spatial observations by the difficulty in carrying out investigations in a remote and protected area. This study provides the first integrative overview of subsurface brines in Wright Valley using non-destructive geophysical measurements. Our research goals were to map water distribution and hydrological connectivity in order to further our understanding of permafrost hydrogeology.

2 Wright Valley airborne electromagnetic (AEM) survey

Here, we present data from AEM surveys flown in Wright Valley in 2011 and 2018 (Foley et al., 2015). Previous examinations of these data in Victoria Valley and Taylor Valley have revealed both the existence of confined brine pockets (e.g., under Lake Vanda, Dugan et al., 2015b) and expansive groundwater connectivity (e.g., lower Taylor Valley, Mikucki et al., 2015; Foley et al., 2020) and have transformed our understanding of hydrological connectivity in this polar desert ecosystem.

A comprehensive methodology on our 2018 AEM survey can be found in Grombacher et al. (2021). Briefly, a transient electromagnetic (TEM) measurement involves a current in a transmitter loop being turned on and off, which generates a time-varying magnetic field in the subsurface, subsequently leading to electrical eddy currents in the subsurface that generate a secondary time-varying magnetic field. These secondary fields are measured by a receiver coil. Highly conductive regions produce larger amplitude, slowly decaying eddy currents, whereas highly resistive regions give rise to smaller amplitude, quickly decaying currents. Our AEM surveys were conducted by suspending a TEM system beneath a helicopter. Inversion is used to estimate the underlying electrical resistivity structures, consistent with observed TEM signals. Here, standard 1D TEM forward modeling ap-

proaches are employed (Auken et al., 2015), except in the case of Don Juan Pond, where a more complex forward response is required to account for strong induced polarization effects (Fiandaca et al., 2018). In 2017, a ground-based configuration was used to investigate sites prior to the airborne survey. The ground-based TEM had larger transmitter coils and stacking times and therefore could penetrate more deeply with potentially higher accuracy. A comparison of the ground-based and airborne TEM data revealed high consistency between the methods (Madsen et al., 2022).

We focus on three E–W flight lines flown in 2018: Line 1, along the North Fork and Lake Vanda, Line 2, along the South Fork over Don Juan Pond, and Line 3, along the southern edge of Lake Vanda (Fig. 1a). These flight lines were picked to align with the lowest-elevation tracks of Wright Valley, where one might assume brine would pool (although we acknowledge that surface water and groundwater flow potential can be very different). Additional E–W flight lines along higher-elevation tracks returned higher resistivity readings, which confirm that the low-elevation tracks (Lines 1–3) are the most hydrologically interesting. Additional flight lines were flown N–S across Lake Vanda, but many appear to display 3D effects where measured signals are sensitive to features not beneath the sensor (i.e., to the side/front/rear). In these cases, forward modeling that employs a 1D assumption may introduce spurious features in an attempt to attribute observed signals to structures directly beneath the sensor. These spurious features, which sometimes manifest as “pant legs” where a shallow feature is stretched deeper to the sides, can render interpretation of connectivity difficult.

Inversion results are presented as cross-sectional images of resistivity (Ωm) to infer subsurface properties (i.e., highly resistive ice or bedrock or highly conductive water and brines). The cross-track sampling width is on the order of 60–70 m wide at the shallowest depths and extends to several hundred meters wide at the bottom of the profiles. The depth of investigation (DOI), where the signal-to-noise ratio is too low for accurate interpretation, was several hundreds of meters in many areas. In addition, a gap between or within a flight line indicates an area where no earth signal could be reliably detected that exceeded instrumental noise levels (e.g., within Line 1 and between Lines 2 and 3). This does not negate the presence of a deeper conductive unit, only that it is not present within the DOI of the employed AEM system.

3 Wright Valley AEM findings

The returns from the three along-valley flight lines are intriguing (Fig. 1b–d). Firstly, the surface conductors of Lake Vanda and Don Juan Pond are clear, as is the surface of saline Don Quixote Pond at the very western edge of Line 1 (Englert et al., 2014). The top 20 m of Lake Vanda has a resistivity of $\sim 20\ \Omega\text{m}$ (a typical value for freshwater). At 60 m depth, resistivity drops to $3.9\ \Omega\text{m}$ and further drops to

$0.2\ \Omega\text{m}$ at 70 m depth. This is consistent with electrical conductivity (EC) of lake water in Vanda, which shows low EC ($<5\ \text{mS cm}^{-1}$) from the surface to 63.6 m depth (in 2014) followed by a rapid increase to $120\ \text{mS cm}^{-1}$ between the depths of 63.6 and 78 m (Castendyk et al., 2016). The minimum resistivity of $0.14\ \Omega\text{m}$ is recorded from 80 to 92 m, a depth within the lake sediments that overlaps with the highly saline water ($>$ lake water) found in the sediments of DVDP4A.

The surface of Don Juan Pond, from 0 to 4 m, has a resistivity of $1\ \Omega\text{m}$ in Line 2. This might appear high for the second saltiest body of water on Earth, but the pond itself is only 20–30 cm deep, and the AEM inversions are averaged over the top 4 m. The averaged resistivity would take into account the porosity of the sediments as well as the conductivity of the groundwater. The conductivity of CaCl_2 is greater than the conductivity of an equivalent mass of NaCl , and the relationship between concentration (ppt) and specific conductance (mS cm^{-1}) is parabolic, with a maximum SpC of approximately $200\ \text{mS cm}^{-1}$ at $300\ \text{g CaCl}_2\ \text{L}^{-1}$ (Sun and Newman, 1970).

The second depth of low resistivities ($12\text{--}14\ \Omega\text{m}$) is seen from 23 to 36 m. This depth overlaps with the hypothesized aquifer in the Ferrar Dolerite (Harris, 1981). A recent analysis of soluble salts surrounding Don Juan Pond showed that the salts in hillslope surface wet streaks were most likely derived from Ferrar Dolerite bedrock outcrops based on the chemical signatures of Cl^- and NO_3^- (Toner et al., 2022). This is yet another line of evidence that supports a connection between Ferrar Dolerite and Don Juan Pond brine. Based on barometric efficiency calculations, the Ferrar Dolerite is thought to be highly fractured with a porosity upwards of 20 % (Harris, 1981), which could hold a significant volume of brine. Line 2 also reveals a vertical low-resistivity feature approximately 2 km to the east of Don Juan Pond. This vertical feature (possibly a dolerite dyke) would intersect a surface basin which contains the VXE-6 pond. This pond is known to hold shallow saline water and be chemically distinct from other South Fork ponds (Harris, 1981; Toner et al., 2022).

In the Don Juan Pond basin, AEM data provide evidence for regional connectivity. From the Line 2 transect, there is subsurface brine present on either side of the pond and possibly a connection between VXE-6 and Don Juan Pond. The elevation of the brine appears to be higher than the elevation of the pond. This topographic control could be responsible for the artesian well discharge observed during the DVDP drilling.

From our AEM survey, any valley-wide groundwater connectivity is obscure. To the east of Lake Vanda, there is little evidence of subsurface conductors, and it is unlikely that there is deep groundwater. From Lake Vanda to the west, there is no obvious shallow ($<100\ \text{m}$) connectivity between the lake and the North or South Forks. There is evidence of a layer of shallow brines in the North Fork, possibly a




Figure 1. (a) AEM flight lines over the North Fork of Wright Valley and Lake Vanda (Line 1), over Don Juan Pond (Line 2), and over Lake Vanda towards the South Fork (Line 3). White circles show the locations of the DVDP boreholes. Coordinates are in UTM-58S. This contains modified Copernicus Sentinel data 2020. (b–d) Resistivity profiles of the three flight lines. The black line represents the sediment–water interface (bathymetry) of the Lake Vanda basin. Lower data are shaded beyond the depth of investigation. Black vertical rectangles show the approximate location and depth of the DVDP boreholes crossed by Lines 1 and 3.

remnant of paleolakes. DVDP14 did not penetrate the permafrost, and it may be that it did not extend deep enough or that the spatial heterogeneity is such that it did not intersect an unfrozen brine layer. The region between Don Juan Pond and Lake Vanda was surveyed by both airborne and ground-based TEM. Neither survey was able to record a signal that exceeded the noise floor of the system in the area to the SW of Line 3 (see the area of *no signal* in Fig. 1a). If any brine is present here, it would be > 500 m deep. If the key to finding subsurface brine in Wright Valley is to “follow the dolerite”, more knowledge is needed of the subsurface geology in Wright Valley. The dolerite sills generally have a gentle westward dip (McKelvey and Webb, 1962), but evidence of brine-saturated dolerite is not present on the eastern end of the South Fork transect. If there is a dolerite conduit between the two surface waterbodies, it does not follow the surface topography. To the west of Don Juan Pond at higher elevations along the rock glacier, our airborne and ground-based surveys were unable to measure a signal exceeding the noise floor of the system. From this, we cannot draw any conclusions about the source of the brine or possible connections to the East Antarctic Ice Sheet.

4 Conclusions

From Lake Vanda towards Don Juan Pond in the South Fork, we cannot establish a direct subsurface connection from our data; however, a deep (>500 m) connection or a connection outside our study transects may be present. Certainly, resistivities <100 Ω m to either side of Don Juan Pond support this speculation. This depth is in the range at which liquid brines were inferred in Taylor Valley from AEM (Mikucki et al., 2015) and is also similar to the frozen–unfrozen transition at 183 m found in DVDP10 at New Harbour, Taylor Valley, and the inferred frozen–unfrozen transition at 248 m in DVDP11 in front of Commonwealth Glacier, Taylor Valley (Cartwright and Harris, 1981). DVDP geological data established that any deep groundwater connectivity in Wright Valley would be through fractured bedrock (Ferrar Dolerite) and not sediments, as in Taylor Valley.

The formation and spatial extent of Wright Valley brines are relevant to Antarctic hydrological and geochemical processes, including those at subglacial and submarine interfaces (Foley et al., 2019), as well as hydrogeological processes on other icy planets (Toner et al., 2022). Furthermore, these brines may be a refuge for unique microbial life (Campen et al., 2019). Our spatial investigation of Wright Valley did not resolve the potential for valley-wide groundwater connectivity but did confirm the presence of unfrozen brine-saturated regions in the subsurface and importantly highlights regions that deserve further investigation.

Code availability. R code to reproduce the figure in this paper is available at <https://doi.org/10.5281/zenodo.7416021> (Dugan, 2022).

Data availability. Data for the three flight lines presented in this paper are available at <https://doi.org/10.5281/zenodo.7416021> (Dugan, 2022). Data for the project are available via the US Antarctic Program Data Center: <https://doi.org/10.15784/601373> (Tulaczyk, 2020).

Author contributions. HAD led manuscript preparation. DG, EA, NF, and TB oversaw data collection, data processing, inversion, and interpretation. All the authors contributed to the field campaigns and intellectual development of the manuscript.

Competing interests. The contact author has declared that none of the authors has any competing interests.

Disclaimer. Publisher’s note: Copernicus Publications remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Acknowledgements. The thoughtful critiques of two reviewers significantly improved this paper. We would like to thank the many contractors who supported this work based out of McMurdo Station, with special thanks to the pilots and helicopter technicians who made this research possible.

Financial support. This research has been supported by the National Science Foundation (Office of Polar Programs grant nos. 1643536, 1643687, 1643775, and 1644187).

Review statement. This paper was edited by Elizabeth Bagshaw and reviewed by Jonathan Toner and Joseph Levy.

References

- Armitage, K. and House, H.: A limnological reconnaissance in the area of McMurdo Sound, Antarctica, *Limnol. Oceanogr.*, 7, 36–41, <https://doi.org/10.4319/lo.1962.7.1.0036>, 1962.
- Auken, E., Christiansen, A. V., Kirkegaard, C., Fiandaca, G., Schamper, C., Behroozmand, A. A., Binley, A., Nielsen, E., Eftersø, F., Christensen, N. B., Sørensen, K., Foged, N., and Vignoli, G.: An overview of a highly versatile forward and stable inverse algorithm for airborne, ground-based and borehole electromagnetic and electric data, *Explor. Geophys.*, 46, 223–235, <https://doi.org/10.1071/EG13097>, 2015.
- Campen, R., Kowalski, J., Lyons, W. B., Tulaczyk, S., Dachwald, B., Pettit, E., Welch, K. A., and Mikucki, J. A.: Microbial diversity of an Antarctic subglacial community and high-

- resolution replicate sampling inform hydrological connectivity in a polar desert, *Environ. Microbiol.*, 21, 2290–2306, <https://doi.org/10.1111/1462-2920.14607>, 2019.
- Carlson, C. A., Phillips, F. M., Elmore, D., and Bentley, H. W.: Chlorine-36 tracing of salinity sources in the Dry Valleys of Victoria Land, Antarctica, *Geochim. Cosmochim. Ac.*, 54, 311–318, [https://doi.org/10.1016/0016-7037\(90\)90320-K](https://doi.org/10.1016/0016-7037(90)90320-K), 1990.
- Cartwright, K. and Harris, H. J. H.: Hydrogeology of the Dry Valley region, Antarctica, in: *Dry Valley Drilling Project, Antarctic Research Series 33*, edited by: McGinnis, L. D., 33, 193–214 pp., American Geophysical Union, Washington DC, USA, <https://doi.org/10.1029/AR033p0193>, 1981.
- Cartwright, K., Treves, S. B., and Torii, T.: Geology of DVDP 4, Lake Vanda, Wright Valley, Antarctica, *Dry Valley Drilling Project Bulletin*, 1974.
- Castendyk, D. N., Obyrk, M. K., Leidman, S. Z., Gooseff, M., and Hawes, I.: Lake Vanda: A sentinel for climate change in the McMurdo Sound Region of Antarctica, *Global Planet. Change*, 144, 213–227, <https://doi.org/10.1016/j.gloplacha.2016.06.007>, 2016.
- Chapman-Smith, M.: Geological log of DVDP 14, North Fork Basin, Dry Valley Drilling Project Bulletin No. 5, 94–99 pp., 1975.
- Dickson, J. L., Head, J. W., Levy, J. S., and Marchant, D. R.: Don Juan Pond, Antarctica: Near-surface CaCl₂-brine feeding Earth's most saline lake and implications for Mars, *Sci. rep.*, 3, 1166, <https://doi.org/10.1038/srep01166>, 2013.
- Dugan, H. A.: WrightValley_AEM: Cryosphere release (v1.0.0), Zenodo [code and data set], <https://doi.org/10.5281/zenodo.7416021>, 2022.
- Dugan, H. A., Doran, P. T., Wagner, B., Kenig, F., Fritsen, C. H., Arcone, S. A., Kuhn, E., Ostrom, N. E., Warnock, J. P., and Murray, A. E.: Stratigraphy of Lake Vida, Antarctica: hydrologic implications of 27 m of ice, *The Cryosphere*, 9, 439–450, <https://doi.org/10.5194/tc-9-439-2015>, 2015a.
- Dugan, H. A., Doran, P. T., Tulaczyk, S., Mikucki, J. A., Arcone, S. A., Auken, E., Schamper, C., and Virginia, R. A.: Subsurface imaging reveals a confined aquifer beneath an ice-sealed Antarctic lake, *Geophys. Res. Lett.*, 42, 96–103, <https://doi.org/10.1002/2014GL062431>, 2015b.
- Englert, P. A. J., Bishop, J. L., Patel, S., Gibson, E. K., and Koerber, C.: Don Quixote Pond Sediments: Surface and Subsurface Chemistry and Mineralogy, 2014, AGU Fall Meeting Abstracts ADS, 2014AGUFM.P41A3893E, P41A–3893, <https://ui.adsabs.harvard.edu/abs/2014AGUFM.P41A3893E> (last access: 9 February 2022), 2014.
- Fiandaca, G., Madsen, L. M., and Maurya, P. K.: Re-parameterisations of the Cole-Cole model for improved spectral inversion of induced polarization data, *Near Surf. Geophys.*, 16, 385–399, <https://doi.org/10.3997/1873-0604.2017065>, 2018.
- Foley, N., Tulaczyk, S., Auken, E., Schamper, C., Dugan, H. A., Mikucki, J., Virginia, R., and Doran, P.: Helicopter-borne transient electromagnetic in high-latitude environments: An application in the McMurdo Dry Valleys, Antarctica, *Geophysics*, 81, WA87–WA99, <https://doi.org/10.1190/GEO2015-0186.1>, 2015.
- Foley, N., Tulaczyk, S. M., Grombacher, D., Doran, P. T., Mikucki, J., Myers, K. F., Foged, N., Dugan, H., Auken, E., and Virginia, R.: Evidence for Pathways of Concentrated Submarine Groundwater Discharge in East Antarctica from Helicopter-Borne Electrical Resistivity Measurements, *Hydrology*, 6, 54, <https://doi.org/10.3390/hydrology6020054>, 2019.
- Foley, N., Tulaczyk, S., Auken, E., Grombacher, D., Mikucki, J., Foged, N., Myers, K., Dugan, H., Doran, P. T., and Virginia, R. A.: Mapping geothermal heat flux using permafrost thickness constrained by airborne electromagnetic surveys on the western coast of Ross Island, Antarctica, *Explor. Geophys.*, 51, 84–93, <https://doi.org/10.1080/08123985.2019.1651618>, 2020.
- Gough, R. V., Wong, J., Dickson, J. L., Levy, J. S., Head, J. W., Marchant, D. R., and Tolbert, M. A.: Brine formation via deliquescence by salts found near Don Juan Pond, Antarctica: Laboratory experiments and field observational results, *Earth Planet. Sci. Lett.*, 476, 189–198, <https://doi.org/10.1016/j.epsl.2017.08.003>, 2017.
- Green, W. J. and Canfield, D. E.: Geochemistry of the Onyx River (Wright Valley, Antarctica) and its role in the chemical evolution of Lake Vanda, *Geochim. Cosmochim. Ac.*, 48, 2457–2467, 1984.
- Green, W. J. and Lyons, W. B.: The Saline Lakes of the McMurdo Dry Valleys, Antarctica, *Aquat. Geochem.*, 15, 321–348, <https://doi.org/10.1007/s10498-008-9052-1>, 2009.
- Grombacher, D., Auken, E., Foged, N., Bording, T., Foley, N., Doran, P. T., Mikucki, J., Dugan, H. A., Garza-Giron, R., Myers, K., Virginia, R. A., and Tulaczyk, S.: Induced polarization effects in airborne transient electromagnetic data collected in the McMurdo Dry Valleys, Antarctica, *Geophys. J. Int.*, 226, 1574–1583, <https://doi.org/10.1093/gji/ggab148>, 2021.
- Harris, H. J. H., Cartwright, K., and Torii, T.: Dynamic Chemical Equilibrium in a Polar Desert Pond: A Sensitive Index of Meteorological Cycles, *Science*, 204, 301–303, 1979.
- Harris, H. J. H.: Hydrology and Hydrogeochemistry of the South Fork, Wright Valley, Southern Victoria Land, Antarctica, PhD, University of Illinois at Urbana-Champaign, USA, p. 359, ISBN 9798662247758, 1981.
- Lyons, W. B., Welch, K. a., Snyder, G., Olesik, J., Graham, E. Y., Marion, G. M., and Poreda, R. J.: Halogen geochemistry of the McMurdo dry valleys lakes, Antarctica: Clues to the origin of solutes and lake evolution, *Geochim. Cosmochim. Ac.*, 69, 305–323, <https://doi.org/10.1016/j.gca.2004.06.040>, 2005.
- Madsen, L. M., Bording, T., Grombacher, D., Foged, N., Foley, N., Dugan, H. A., Doran, P. T., Mikucki, J., Tulaczyk, S., and Auken, E.: Comparison of ground-based and airborne transient electromagnetic methods for mapping glacial and permafrost environments: Cases from McMurdo Dry Valleys, Antarctica, *Cold Reg. Sci. Technol.*, 199, 103578, <https://doi.org/10.1016/j.coldregions.2022.103578>, 2022.
- McGinnis, L. D.: Artesian well at Don Juan Pond, *Antarct. J. US*, 14, 26–27, 1979.
- McGinnis, L. D., Struckless, J., Osby, D. R., and Kyle, P.: Gamma ray, salinity, and electric log of DVDP boreholes, *Antar. Res. S.-Dry Valley Drilling Project*, 33, 95–108, 1981.
- McKelvey, B. C. and Webb, P. N.: Geological investigations in southern Victoria Land, Antarctica, *New Zeal. J. Geol. Geop.*, 5, 143–162, <https://doi.org/10.1080/00288306.1962.10420116>, 1962.
- Meyer, G. H., Morrow, M. B., Wyss, O., Berg, T. E., and Littlepage, J. L.: Antarctica: The Microbiology of an Unfrozen Saline Pond, *Science*, 138, 1103–1104, <https://doi.org/10.1126/science.138.3545.1103>, 1962.

- Mikucki, J. A., Pearson, A., Johnston, D. T., Turchyn, A. V., Farquhar, J., Schrag, D. P., Anbar, A. D., Priscu, J. C., and Lee, P. A.: A contemporary microbially maintained subglacial ferrous “ocean”, *Science*, 663, 397–400, <https://doi.org/10.1126/science.1167350>, 2009.
- Mikucki, J. A., Auken, E., Tulaczyk, S., Virginia, R. A., Schamper, C., Sørensen, K. I., Doran, P. T., Dugan, H. A., and Foley, N.: Deep groundwater and potential subsurface habitats beneath an Antarctic dry valley, *Nat. Commun.*, 6, 6831, <https://doi.org/10.1038/ncomms7831>, 2015.
- Mudrey Jr., M. G., Torii, T., and Harris, H. J. H.: Geological log of DVD 13 – Don Juan Pond, Dry Valley Drilling Project Bulletin No. 5, 78–93, 1975.
- Sun, I. and Newman, J.: The electrical conductivity of aqueous solutions of calcium chloride, Tech. Rep. UCRL-19150, AEC Contract No. W-740S-eng-48, Inorganic Materials Research Division, Lawrence Radiation Laboratory, <https://escholarship.org/uc/item/5v01s3c6> (last access: 23 October 2022), 1970.
- Toner, J., Catling, D., and Sletten, R. S.: The geochemistry of Don Juan Pond: Evidence for a deep groundwater flow system in Wright Valley, Antarctica, *Earth Planet. Sci. Lett.*, 474, 190–197, <https://doi.org/10.1016/j.epsl.2017.06.039>, 2017.
- Toner, J. D., Sletten, R. S., Liu, L., Catling, D. C., Ming, D. W., Mushkin, A., and Lin, P. C.: Wet streaks in the McMurdo Dry Valleys, Antarctica: Implications for Recurring Slope Lineae on Mars, *Earth Planet. Sci. Lett.*, 589, 117582, <https://doi.org/10.1016/j.epsl.2022.117582>, 2022.
- Tulaczyk, S.: ANTAEM project airborne EM resistivity data from McMurdo Region, U.S. Antarctic Program (USAP) Data Center [data set], <https://doi.org/10.15784/601373>, 2020.
- Wilson, A. T. and Wellman, H. W.: Lake Vanda: An Antarctic Lake: Lake Vanda as a Solar Energy Trap, *Nature*, 196, 1171–1173, <https://doi.org/10.1038/1961171a0>, 1962.