

# Antimicrobial peptides from plants: stabilization of the $\gamma$ core of a tomato defensin by intramolecular disulfide bond<sup>†</sup>

C. Avitabile,<sup>a</sup> R. Capparelli,<sup>b</sup> M. M. Rigano,<sup>b</sup> A. Fulgione,<sup>b</sup> A. Barone,<sup>b</sup> C. Pedone<sup>a</sup> and A. Romanelli<sup>a\*</sup>

**Cysteine-containing antimicrobial peptides of diverse phylogeny share a common structural signature, the  $\gamma$  core, characterized by a strong polarization of charges in two antiparallel  $\beta$  sheets. In this work, we analyzed peptides derived from the tomato defensin SolyC07g007760 corresponding to the protein  $\gamma$  core and demonstrated that cyclization of the peptides, which results in segregation of positive charges to the turn region, produces peptides very active against Gram negative bacteria, such as *Salmonella enterica* and *Helicobacter pylori*. Interestingly, these peptides show very low hemolytic activity and thus represent a scaffold for the design of new antimicrobial peptides. Copyright © 2013 European Peptide Society and John Wiley & Sons, Ltd.**

**Keywords:** tomato defensin; cyclization; antimicrobial

## Introduction

A tremendous effort has been recently devoted to the discovery of biologically active peptides from natural sources [1–3]. Plants produce a wide variety of antimicrobial peptides and proteins showing activity against either plant or human pathogens [4]. Many of these, such as thionins and knottins, are rich in cysteines and are stabilized by a number of disulfide bonds [5,6]. Interestingly, some of the structural features displayed by peptides produced in plants are well conserved also within proteins produced by evolutionarily diverse organisms, and in fact, the existence of multidimensional signatures for antimicrobial peptides has been hypothesized [7–9]. The  $\gamma$  core motif has recently been indicated as a three dimensional signature for antimicrobial peptides or proteins and, generally, as a feature shown by membrane-active proteins. In particular, the  $\gamma$  core motif, characterized by a conserved CXG sequence and two antiparallel  $\beta$  sheets, is common to host defence polypeptides showing multiple disulfide bonds such as toxins, kinocidins, and thionins [10,11]. This structural motif can be generated by different amino acid sequences and can exist in the dextrameric or levomeric form, depending on whether it is derived by the amino acid forward or reverse sequence (Figure 1). In all cases,  $\gamma$  core motifs show polarization of positive charge and segregation of hydrophobic amino acids. There are some cases in which the only  $\gamma$  core is sufficient for the peptide antimicrobial activity, as observed for protegrins,  $\theta$  defensin RTD-1, and other cases in which the  $\gamma$  core represents the scaffold around which helices and sheets packs, as for kinocidins [12,13].

In a recent paper, we identified, after a bioinformatics analysis of the tomato genome, the tomato defensin SolyC07g007760 (iTAG v.2.3) and demonstrated that a synthetic peptide (SolyC) corresponding to its  $\gamma$  core exerts antimicrobial activity against both Gram positive and Gram negative bacteria, being particularly active against different strains of the Gram negative bacterium *Helicobacter pylori* [14,15]. Interestingly, the peptide SolyC was able to discriminate between pathogens and probiotic

bacteria, killing only the minority of the probiotic tested. In addition, SolyC downregulated the proinflammatory cytokines to an extent comparable with the known anti-inflammatory drug acetyl salicylic acid and did not cause red blood cell hemolysis and was not toxic toward eukaryotic cells.


SolyC is a peptide composed by 17 amino acids and contains three cysteines in positions 6, 13, and 15. The results described so far were obtained employing the peptide in its linear form. The presence of three cysteines that might be subjected to spontaneous oxidation and the observation that in typical  $\gamma$  cores those cysteines are involved in disulfide bonds urged us to investigate how formation of disulfide bonds in a controlled fashion could affect the secondary structure and the antimicrobial activity of the peptide. Two peptides in which Cys13 or Cys15 was mutated into a serine were obtained in the linear and oxidized form. Disulfide bonds were formed between C6 and C13 (SolyC1) or C6 and C15 (SolyC2); in both cases, one cysteine was missing. Truncated peptides, starting from the conserved GXC motif, were also obtained in the linear and oxidized form. The secondary structure of all peptides was analyzed by circular dichroism (CD); the antimicrobial activity of the peptides was tested against Gram positive and Gram negative bacteria. All the oxidized peptides

\* Correspondence to: Alessandra Romanelli, University of Naples 'Federico II', School of Biotechnological Sciences, Department of Biological Sciences, Via Mezzocannone 16, 80134, Naples, Italy. E-mail: [alessandra.romanelli@unina.it](mailto:alessandra.romanelli@unina.it)

† Special issue devoted to contributions presented at the 13<sup>th</sup> Naples Workshop on Bioactive Peptides, June 7–10, 2012, Naples.

a University of Naples 'Federico II', School of Biotechnological Sciences, Department of Biological Sciences, Via Mezzocannone 16, 80134, Naples Italy

b University of Naples 'Federico II', School of Biotechnological Sciences, Department of Soil, Plant, Environmental and Animal Production Sciences, Via Università 100, 80055, Portici Italy



**Figure 1.** Schematic representation of the gamma core isoforms.

are highly active against Gram negative bacteria; the linear peptide SolyC2 shows antimicrobial activity against Gram negative bacteria comparable with that of the reference peptide SolyC, whereas linear SolyC1 exhibits an antimicrobial activity against Gram negative bacteria at higher concentrations. The activity of modified peptides against Gram positive bacteria was found in all cases lower as compared with that of the reference peptide. These results suggest that the interaction of the peptides with the membranes of Gram negative bacteria is stronger when the positive charges are exposed, and this likely occurs when the peptides are in the oxidized form, in which the antiparallel  $\beta$  sheets are constrained by the disulfide bond.

## Materials and Methods

### Reagents

The amino acids used for the peptide synthesis, Fmoc-Arg(Pbf)-OH, Fmoc-Asn(Trt)-OH, Fmoc-Cys(Trt)-OH, Fmoc-Gly-OH, Fmoc-Lys(Boc)-OH, Fmoc-Phe-OH, Fmoc-Ser(OtBu)-OH, Fmoc-Thr(tBu)-OH, the Rink amide MBHA resin and the activators *N*-hydroxybenzotriazole (HOBT), and *O*-benzotriazole-*N,N,N,N*-tetramethyl-uronium-hexafluoro-phosphate (HBTU), were purchased from Novabiochem (Gibbstown, NJ, USA). Acetonitrile was from Reidel-deHaën (Seelze, Germany) and dry *N,N*-dimethylformamide (DMF) from LabScan (Dublin, Ireland). All other reagents were from Sigma Aldrich (Milan, Italy). LC-MS analyses were performed on a LC-MS Thermo Finnigan with an electrospray source (MSQ) on a Phenomenex Jupiter 4 $\mu$  Proteo (50  $\times$  4.6 mm) column for all peptides except for SolyC-t that was analyzed on a Jupiter 4 $\mu$  Proteo (150  $\times$  4.6 mm) column. Purification was carried out on an Onyx monolithic semi-prep C18 (100  $\times$  10 mm) column.

### Synthesis

Peptides were synthesized on solid phase by Fmoc chemistry on the MBHA (0.54 mmol/g) resin by consecutive deprotection, coupling, and capping cycles [16]. Deprotection: 30% piperidine in DMF, 5 min (2 $\times$ ). Coupling: 2.5 equivalents of amino acid + 2.49 equivalent of HOBT/HBTU (0.45 M in DMF/DMSO 75/25 v/v) + 3.5 equivalents NMM, 40 min. Capping: acetic anhydride/DIPEA/DMF 15/15/70 v/v/v, 5 min.

Peptides were cleaved off from the resin and deprotected by treatment of the resin with a solution of TFA/TIS/EDT/H<sub>2</sub>O 94/1/2.5/2.5 v/v/v/v, 3 h, rt. TFA was concentrated, and peptides were precipitated in cold ethyl ether. Analysis of the crudes was performed by LC-MS using a gradient of acetonitrile (0.1% TFA) in water (0.1% TFA) from 5% to 70% in 15 min. Purification was performed by semipreparative RP-HPLC using a gradient of acetonitrile (0.1% TFA) in water (0.1% TFA) from 5% to 70% in 15 min.

SolyC (Da): calculated 1994.30; [M + 2H]<sup>2+</sup>: 998.16; [M + 3H]<sup>3+</sup>: 665.77; found: [M + 2H]<sup>2+</sup>: 997.55; [M + 3H]<sup>3+</sup>: 665.64; retention time: 5.40 min.

SolyC-t (Da): calculated 1703.01; [M + 2H]<sup>2+</sup>: 853.50; [M + 3H]<sup>3+</sup>: 568.67; found: [M + 1H]<sup>1+</sup>: 1702.58; [M + 2H]<sup>2+</sup>: 852.29; [M + 3H]<sup>3+</sup>: 568.52; retention time: 12.0 min.

SolyC1 (Da): calculated 1978.26; [M + 2H]<sup>2+</sup>: 990.13; [M + 3H]<sup>3+</sup>: 660.42; found: [M + 2H]<sup>2+</sup>: 990.13; [M + 3H]<sup>3+</sup>: 660.41; retention time: 5.12 min.

SolyC1-t (Da): calculated 1686.95; [M + 2H]<sup>2+</sup>: 844.47; [M + 3H]<sup>3+</sup>: 563.31; found: [M + 2H]<sup>2+</sup>: 844.40; [M + 3H]<sup>3+</sup>: 563.21; retention time: 5.12 min.

SolyC2 (Da): calculated 1978.26; [M + 2H]<sup>2+</sup>: 990.13; [M + 3H]<sup>3+</sup>: 660.42; found: [M + 2H]<sup>2+</sup>: 989.90; [M + 3H]<sup>3+</sup>: 660.21; retention time: 5.33 min.

SolyC2-t (Da): calculated 1686.95; [M + 2H]<sup>2+</sup>: 844.47; [M + 3H]<sup>3+</sup>: 563.31; found: [M + 2H]<sup>2+</sup>: 844.03; [M + 3H]<sup>3+</sup>: 563.23; retention time: 4.93 min.

### Oxidation Reaction

Disulfide-bridged analogs were obtained from the purified linear peptides after an oxidation reaction. Linear peptides were dissolved in a mixture of H<sub>2</sub>O/acetic acid (95/5, v/v) at a final concentration of 2.5 mM (9 mL total volume); pH was adjusted to 6 with ammonium hydrogen carbonate and then DMSO (1 mL) was added. The reaction was complete after 24 h. Analysis of the crudes was performed by LC-MS using a gradient of acetonitrile (0.1% TFA) in water (0.1% TFA) from 5% to 70% in 15 min. Purification was performed by semipreparative RP-HPLC using a gradient of acetonitrile (0.1% TFA) in water (0.1% TFA) from 5% to 70% in 15 min. Peptides were characterized by ES mass spectrometry.

SolyC1-ox (Da): calculated 1976.26; [M + 2H]<sup>2+</sup>: 989.13; [M + 3H]<sup>3+</sup>: 659.75; found: [M + 2H]<sup>2+</sup>: 988.70; [M + 3H]<sup>3+</sup>: 659.60; retention time: 5.35 min.

SolyC1-t-ox (Da): calculated 1684.94; [M + 2H]<sup>2+</sup>: 843.47; [M + 3H]<sup>3+</sup>: 562.64; found: [M + 2H]<sup>2+</sup>: 843.23; [M + 3H]<sup>3+</sup>: 562.53; retention time: 4.96 min.

SolyC2-ox (Da): calculated 1976.26; [M + 2H]<sup>2+</sup>: 989.13; [M + 3H]<sup>3+</sup>: 659.75; found: [M + 2H]<sup>2+</sup>: 988.60; [M + 3H]<sup>3+</sup>: 659.60; retention time: 5.10 min.

SolyC2-t-ox (Da): calculated 1684.94; [M + 2H]<sup>2+</sup>: 843.47; [M + 3H]<sup>3+</sup>: 562.64; found: [M + 2H]<sup>2+</sup>: 843.26; [M + 3H]<sup>3+</sup>: 562.50; retention time: 5.15 min.

### Bacteria

Bacterial isolates were from patients hospitalized at the Medical School of the University of Naples 'Federico II' and at the 'Villa Betania' hospital (Naples, Italy). Species identification was carried out by PCR [17–21]. Bacteria were grown at 37 °C in Tryptic Soy Broth (TSB) or Luria Broth (LB) medium, harvested in the exponential phase (OD 600 nm 0.6–0.8), centrifuged (10 min at 8  $\times$  10<sup>3</sup> g), and resuspended in Muller Hinton broth at the concentration of  $\sim$ 10<sup>5</sup> CFU/ml. The resistance/susceptibility of the different strains used in this study to conventional antibiotics was also determined using the disk diffusion method on Mueller–Hinton agar according to the NCCLS guidelines 2002 (data not shown).

### Antibacterial and Hemolytic Activity

Bacteria were distributed in triplicate into plates (60  $\mu$ l/well), mixed with dilutions of the peptides (5–100  $\mu$ g/ml; 40  $\mu$ l/well),

and incubated at 37 °C for 20 h. The minimal concentration of peptides causing 100% growth inhibition (MIC100) was determined by measuring the absorbance at 600 nm (Biorad microplate reader model 680, CA) [22]. The test was performed in triplicate. The hemolytic activity of the peptides was tested using mouse red blood cells. The hemolytic activity was measured according to the formula  $OD_{\text{peptide}} - OD_{\text{negative control}} / OD_{\text{positive control}} - OD_{\text{negative control}} \times 100$ , where the negative control (0% hemolysis) was represented by erythrocytes suspended in saline and the positive control (100% hemolysis) was represented by the erythrocytes lysed with 1% triton X100 [22].

The LC50 value relative to the SolyC was calculated as described [23].

### Circular dichroism

Circular dichroism spectra were recorded at 25 °C using a 1 cm quartz cell with the Jasco-810 spectropolarimeter using a 260–200 nm measurement range, 100 nm/min scanning speed, 1 nm bandwidth, 4 s response time, 0.5 nm data pitch.

Peptides concentration for CD measurement was 25  $\mu\text{M}$ . CD spectra were registered in 10 mM sodium phosphate buffer, pH 7 and in 10 mM sodium phosphate, 20 mM SDS buffer, pH 7.

## Results


### Peptide Design and Synthesis

The SolyC peptide (Table 1) contains the  $\gamma$  core sequence of the tomato defensin SolyC07g007760 and shows strong similarity in the primary sequence with the levomeric form of porcine protegrin PG-1. In PG-1, four cysteines form two disulfide bonds, one involving C6 and C15 and the other C8 and C13, which force the peptide to fold into two antiparallel beta sheets; the oxidation of cysteines induces the formation of a turn characterized by a strong concentration of positive charges [24,25].

SolyC contains only three cysteines; therefore, to induce the peptide to assume a PG-1-like structure, only one disulfide bond can be formed. We designed two peptides with a different disulfide pattern, with the aim to explore the effect of cyclization on the antimicrobial activity of the peptide, to understand whether


there is a preference for the formation of one of the two disulfides and also whether there is a difference in the activity of the peptides with cycles of different size (Table 1). Because in  $\gamma$  cores cysteines contained in sequences CXC are rarely connected by disulfide bonds [26], we explored only two of the possible combinations of disulfides, namely C6–C13 (SolyC1) and C6–C15 (SolyC2) and mutated the cysteine not involved in the disulfide into a serine. To investigate the effect of reducing the hydrophobicity of the peptide, we also obtained truncated peptides, lacking three amino acids at the N-terminus. In the truncated peptides, the residues that characterize the  $\gamma$  core motif GXC were conserved (Table 1). HPLC retention times for truncated peptides are similar or lower as compared with those of the full length peptides, as expected comparing the mean hydrophobicity calculated for each peptide (Table 1) [27].

All peptides were obtained in the linear and oxidized form. Oxidation was carried out incubating linear peptides with DMSO (10%) at pH 6.0. LC-MS analysis of the peptides, showing for the majority of the peptides a decrease of the retention time and a reduction of 2 units in the mass, confirmed that the reaction occurred (Figure 2).



**Figure 2.** Comparison of the LC profiles for SolyC2 in the linear (A) and oxidized form (B). Zoom between 4 and 6 min of the LC run.

Table 1. Peptide sequences, names, reference number, and mean hydrophobicity			
Peptide sequence	Name	Number	Mean hydrophobicity
FSGGNCRGFRRRCFCTK	SolyC	1	−0.71
GNCRGFRRRCFCTK	SolyC-t	2	−0.98
FSGGNCRGFRRRCFSTK	SolyC1	3	−0.91
GNCRGFRRRCFSTK	SolyC1-t	4	−1.22
FSGGNCRGFRRRSFCTK	SolyC2	5	−0.91
GNCRGFRRRSFCTK	SolyC2-t	6	−1.22
FSGGNCRGFRRRCFSTK	SolyC1-ox	7	−0.91
GNCRGFRRRCFSTK	SolyC1-t-ox	8	−1.22
FSGGNCRGFRRRSFCTK	SolyC2-ox	9	−0.91
GNCRGFRRRSFCTK	SolyC2-t-ox	10	−1.22




**Figure 3.** CD spectra of SolyC2 (blue) and SolyC2-ox (red) (50 μM) recorded in 10 mM phosphate buffer pH 7 (continuous line) and in 10 mM phosphate buffer, 30 mM SDS pH 7 (dotted lines).

### Secondary Structure Studies

Linear and oxidized peptides were analyzed by CD at a 50 μM concentration in phosphate buffer, pH7. Peptides are mostly in an unordered form; spectra for the linear peptides, in fact, show one minimum around 204 nm and a very shallow minimum around 230 nm; upon oxidation, the minimum shifts toward 207 nm. Spectra recorded in the presence of SDS 30 mM are in most cases very similar to those recorded in buffer but more intense (Figure 3).

### Antimicrobial Activity Test

The antimicrobial activity of the peptides was tested against the Gram positive bacteria *Staphylococcus aureus*, *Staphylococcus epidermidis*, *Listeria monocytogenes* and against the Gram negative bacteria *H. pylori* and *Salmonella enterica* (Table 2). For the standard peptide SolyC, truncation results in loss of activity against the Gram positive *Staphilococci*, whereas no difference was found when the peptides SolyC and SolyC-t were tested against Gram negative bacteria. When tested against Gram positive bacteria, the analogs containing two cysteines (SolyC1 and SolyC2) show a reduced antimicrobial ability as compared with the reference peptide SolyC; MIC values are in all cases higher than that of SolyC, and there is no large difference between the activity of linear and oxidized peptides. Peptides are in general more active against Gram negative bacteria; linear peptides show lower MIC in the full length version as compared with the truncated form. Noteworthy, SolyC2, the peptide having the disulfide between C6 and C15, shows activity identical to that of the reference peptide. Oxidation results in an increase of the



**Figure 4.** Hemolytic activity of the peptides measured at different concentrations.

antimicrobial activity of the modified peptides against Gram negative bacteria, either in the extended or in the truncated form: all oxidized peptides exhibit activity identical or comparable with that of the reference peptide.

The hemolytic activity of the peptides was tested for all peptides at several concentrations (Figure 4). Up to 90 μg/ml, all peptides have hemolytic activity below 30%. At low

**Table 2.** Antimicrobial activity of the peptides against Gram positive and Gram negative bacteria

Strains	MIC <sub>100</sub> (μg/ml)									
	Peptide number									
	1	2	3	4	5	6	7	8	9	10
<i>Staphylococcus aureus</i>	40	50	50	50	80	80	80	50	100	80
<i>Staphylococcus epidermidis</i>	40	80	100	100	100	100	100	100	100	100
<i>Listeria monocytogenes</i>	80	80	80	80	80	80	80	80	80	80
<i>Salmonella serovar paratyphi B</i>	15	15	40	80	15	40	15	15	15	20
<i>Helicobacter pylori</i>	15	15	15	15	15	20	15	15	20	20

**Table 3.** LC50 for the SolyC and SolyC analogs

	Peptide number									
	1	2	3	4	5	6	7	8	9	10
LC50 (µg/ml)	275.2	266.7	212.3	228.5	226.5	213.0	229.6	231.0	197.4	235.1

concentrations, all long linear peptides and the oxidized form of SolyC1 show hemolytic activity comparable with that of truncated peptides. Interestingly, SolyC2 has in the long oxidized form (SolyC2-ox) very low hemolytic activity, comparable with that of the standard peptide.


LC50 measured for all peptides was found lower for the peptides containing only two cysteines as compared with SolyC (Table 3). The lowest LC50 value was observed for SolyC2 in the oxidized form (SolyC2-ox).

## Discussion

Peptides containing a  $\gamma$  core motif and, in particular, analogs of protegrins have widely been studied as they are capable to exert antimicrobial activity even at high salt concentrations, unlike other antimicrobial peptides that form sheets such as  $\beta$  defensins [28,29]. Interestingly, the PG-1 analog IB-367 has been tested in phase II clinical trials for the treatment of oral mucositis, and PG-1-based peptidomimetics were found to be active at nanomolar concentrations against *Pseudomonas aeruginosa* [30,31]. It has been reported that the stabilization of the hairpin is necessary for the activity: PG-1 analogs lacking cysteines, in which the hairpin is stabilized by a tryptophan zipper motif, possess antimicrobial activity comparable with PG-1 [32]. In PG-1, the number of disulfide bonds and their position affect the activity; analogs of PG-1 with a single disulfide bond possess antimicrobial activity comparable with PG-1 in the linear form.

The  $\gamma$  core of the tomato defensin family, to which SolyC peptides belong, has a strong similarity in the amino acid sequence with the  $\gamma$  core of other plant defensins as the *Medicago sativa* defensin 4 (MtDef4), and also with the levomeric form of PG-1 [9,33]. The three-dimensional structures of plant defensins, which are composed of about 50–60 amino acids, are very complex, and cysteines belonging to the  $\gamma$  core form disulfide bonds with cysteines located outside of the  $\gamma$  core [34–36]. In self-consistent  $\gamma$  cores as PG-1 and RTD-1, instead, the beta-hairpin structure is stabilized by two disulfide bonds between cysteines inside the  $\gamma$  core. SolyC is a fragment of a tomato defensin and, as most of the plant defensins, has a very conserved GXCRG motif within the  $\gamma$  core with three cysteines. It is reasonable to think that SolyC has a structural organization similar to that of plant defensin in which cysteines in the  $\gamma$  core are not connected between them. As we have demonstrated in a previous paper that the linear fragment SolyC possess high antimicrobial activity, we now explored the possibility of stabilizing the peptide structure introducing in a controlled fashion one disulfide bond and evaluated the effects of cyclization on the peptide antimicrobial activity and secondary structure. Furthermore, similar studies were carried out on truncated peptides.

In SolyC, cysteines are located in position  $i$ ,  $i+7$ , and  $i+9$ ; peptides with disulfide bonds between Cys  $i-i+7$  (SolyC1) and  $i-i+9$  (SolyC2) were obtained. The secondary structure of all peptides was analyzed by CD in phosphate buffer and in phosphate



**Figure 5.** Schematic representation of the peptide SolyC2-ox; in red is the positively charged amino acids in the turn, in yellow is the cysteines involved in the disulfide bond, and in green is the cysteine residue mutated into serine.

buffer containing SDS: both linear and oxidized peptides seem to be in an unordered form (Figure 3). All the analogs are more active against Gram negative than against Gram positive bacteria; these results are in line to what has been reported for PG-1 and PG-1 analogs with a single disulfide bond [28]. Interestingly, cyclic peptides are generally more active than linear ones against Gram negative bacteria. Cyclization induces a strong polarization of the peptides, confining the positive charges in the turn and the hydrophobic residues on the 'tails' and reasonably allows the peptide to interact with the bacterial LPS, neutralizing and permeating it (Figure 5). Recent studies aimed to characterize the interaction of rationally designed peptides with the LPS demonstrate that amphipatic peptides are active antimicrobials as they assume the structure of a  $\beta$  boomerang upon interaction with LPS, exerting their activity by disaggregating it [37]. These peptides are mimics of the outer membrane, as they possess a polar head and hydrophobic tails and easily intercalate into LPS. The SolyC cyclic analogs are already in the active conformation with the segregation of charges, and therefore, it is likely that they exert their activity as the  $\beta$  boomerang peptides do.

SolyC2 in the full length and truncated version shows antimicrobial activity identical to that of the reference peptide SolyC. These result may be interpreted hypothesizing that, in the experimental conditions employed for the activity test, SolyC spontaneously oxidizes, and the formation of the disulfide bond between Cys in position  $i$  and  $i+9$  is favored. As in SolyC2, only the disulfide bridge between Cys  $i$  and  $i+9$  can be formed, it is reasonable to think that the linear peptide spontaneously converts into the oxidized form that corresponds also to the active form. This hypothesis is consistent with the observation that full length SolyC2 in the linear and oxidized form has the same antimicrobial activity as the reference SolyC. Interestingly, in SolyC2, the cysteines involved in the disulfide bond are located in positions identical to external cysteines of PG-1, reproducing the cyclization pattern of the highly active 'bullet' peptide proposed by Harwig [28].

The observation that there is no difference in the activity against Gram negative bacteria of cyclic SolyC1 and SolyC2 derivatives, which differ in the number of amino acids included in the cycle, suggests that the charge segregation and not the dimension of the cycle is crucial for activity.

All peptides at concentrations up to threefold that of MIC show very low hemolytic activity, indicating that these peptides are selective for bacterial outer membrane. The lowest percentage of hemolytic activity was found for the oxidized SolyC2. LC50 calculated for all peptides on murine red blood cells is lower for SolyC2, confirming the selectivity for negatively charged membrane (as the bacterial LPS) versus zwitterionic membranes (as the erythrocytes).

In conclusion, we have demonstrated that isolated  $\gamma$  cores of plant defensins possess strong antimicrobial activity against Gram negative bacteria; formation of an intramolecular disulfide bond stabilizes the peptides in the 'active' conformation. We found that the peptide SolyC2, which reproduces the disulfide bond pattern observed in PG-1, shows a high selectivity towards bacterial outer membrane and possess very low hemolytic activity at concentrations that are threefold the MIC. These results encourage future studies on the antimicrobial activity of isolated  $\gamma$  core peptides from plants due to their strength and specificity of action.

## References

- 1 Ganz T. Defensins: antimicrobial peptides of innate immunity. *Nat. Rev. Immunol.* 2003; **3**: 710–720.
- 2 Zanetti M, Gennaro R, Romeo D. Cathelicidins: a novel protein family with a common proregion and a variable C-terminal antimicrobial domain. *FEBS Lett.* 1995; **374**: 1–5.
- 3 Schitteck B, Hipfel R, Sauer B, Bauer J, Kalbacher H, Stevanovic S, Schirle M, Schroeder K, Blin N, Meier F, Rassner G, Garbe C. Dermcidin: a novel human antibiotic peptide secreted by sweat glands. *Nat. Immunol.* 2001; **2**: 1133–1137.
- 4 Castro MS, Fontes W. Plant defense and antimicrobial peptides. *Protein Pept. Lett.* 2005; **12**: 13–18.
- 5 Stec B. Plant thionins—the structural perspective. *Cell. Mol. Life Sci.* 2006; **63**: 1370–1385.
- 6 Chiche L, Heitz A, Gelly JC, Gracy J, Chau PT, Ha PT, Hernandez JF, Le-Nguyen D. Squash inhibitors: from structural motifs to macrocyclic knottins. *Curr. Protein Pept. Sci.* 2004; **5**: 341–349.
- 7 Yeaman MR, Yount NY. Unifying themes in host defence effector polypeptides. *Nat. Rev. Microbiol.* 2007; **5**: 727–740.
- 8 Yount NY, Andres MT, Fierro JF, Yeaman MR. The gamma-core motif correlates with antimicrobial activity in cysteine-containing kaliocin-1 originating from transferrins. *Biochim. Biophys. Acta* 2007; **1768**: 2862–2872.
- 9 Yount NY, Yeaman MR. Multidimensional signatures in antimicrobial peptides. *Proc. Natl. Acad. Sci. U.S.A.* 2004; **101**: 7363–7368.
- 10 Yeaman MR, Yount NY, Waring AJ, Gank KD, Kupferwasser D, Wiese R, Bayer AS, Welch WH. Modular determinants of antimicrobial activity in platelet factor-4 family kinocidins. *Biochim. Biophys. Acta* 2007; **1768**: 609–619.
- 11 Pelegrini PB, Franco OL. Plant gamma-thionins: novel insights on the mechanism of action of a multi-functional class of defense proteins. *Int. J. Biochem. Cell Biol.* 2005; **37**: 2239–2253.
- 12 Steinberg DA, Hurst MA, Fujii CA, Kung AH, Ho JF, Cheng FC, Loury DJ, Fiddes JC. Protegrin-1: a broad-spectrum, rapidly microbicidal peptide with in vivo activity. *Antimicrob. Agents Chemother.* 1997; **41**: 1738–1742.
- 13 Tran D, Tran PA, Tang YQ, Yuan J, Cole T, Selsted ME. Homodimeric theta-defensins from rhesus macaque leukocytes: isolation, synthesis, antimicrobial activities, and bacterial binding properties of the cyclic peptides. *J. Biol. Chem.* 2002; **277**: 3079–3084.
- 14 The tomato genome sequence provides insights into fleshy fruit evolution. *Nature* 2012; **485**: 635–641.
- 15 Rigano MM, Romanelli A, Fulgione A, Nocerino N, D'Agostino N, Avitabile C, Frusciantè L, Barone A, Capuano F, Capparelli R. A novel synthetic peptide from a tomato defensin exhibits antibacterial activities against *Helicobacter pylori*. *J. Pept. Sci.* 2012; **12**: 755–762.
- 16 Romanelli A, Moggio L, Montella RC, Campiglia P, Iannaccone M, Capuano F, Pedone C, Capparelli R. Peptides from Royal Jelly: studies on the antimicrobial activity of jelleins, jelleins analogs and synergy with temporins. *J. Pept. Sci.* 2011; **17**: 348–352.
- 17 Becker K, Roth R, Peters G. Rapid and specific detection of toxigenic *Staphylococcus aureus*: use of two multiplex PCR enzyme immunoassays for amplification and hybridization of staphylococcal enterotoxin genes, exfoliative toxin genes, and toxic shock syndrome toxin 1 gene. *J. Clin. Microbiol.* 1998; **36**: 2548–2553.
- 18 Hong Y, Liu T, Lee MD, Hofacre CL, Maier M, White DG, Ayers S, Wang L, Berghaus R, Maurer JJ. Rapid screening of *Salmonella enterica* serovars Enteritidis, Hadar, Heidelberg and Typhimurium using a serologically-correlative allelotyping PCR targeting the O and H antigen alleles. *BMC Microbiol.* 2008; **8**: 178.
- 19 Tomasini ML, Zanussi S, Sozzi M, Tedeschi R, Basaglia G, De Paoli P. Heterogeneity of cag genotypes in *Helicobacter pylori* isolates from human biopsy specimens. *J. Clin. Microbiol.* 2003; **41**: 976–980.
- 20 Bubert A, Hein I, Rauch M, Lehner A, Yoon B, Goebel W, Wagner M. Detection and differentiation of *Listeria* spp. by a single reaction based on multiplex PCR. *Appl. Environ. Microbiol.* 1999; **65**: 4688–4692.
- 21 Martineau F, Picard FJ, Roy PH, Ouellette M, Bergeron MG. Species-specific and ubiquitous DNA-based assays for rapid identification of *Staphylococcus epidermidis*. *J. Clin. Microbiol.* 1996; **34**: 2888–2893.
- 22 Capparelli R, Romanelli A, Iannaccone M, Nocerino N, Ripa R, Pensato S, Pedone C, Iannelli D. Synergistic antibacterial and anti-inflammatory activity of temporin A and modified temporin B in vivo. *PLoS One* 2009; **4**: e7191.
- 23 Vinhal Costa Orsine J, Vinhal da Costa R, Carvalho da Silva R, de Fatima Menezes Almeida Santos M, Carvalho Garbi Novaes MR. The acute cytotoxicity and lethal concentration (LC50) of *Agaricus sylvaticus* through hemolytic activity on human erythrocyte. *Intl. J. Nutr. Metabol.* 2012; **4**: 19–23.
- 24 Fahrner RL, Dieckmann T, Harwig SS, Lehrer RI, Eisenberg D, Feigon J. Solution structure of protegrin-1, a broad-spectrum antimicrobial peptide from porcine leukocytes. *Chem. Biol.* 1996; **3**: 543–550.
- 25 Mani R, Cady SD, Tang M, Waring AJ, Lehrer RI, Hong M. Membrane-dependent oligomeric structure and pore formation of a beta-hairpin antimicrobial peptide in lipid bilayers from solid-state NMR. *Proc. Natl. Acad. Sci. U.S.A.* 2006; **103**: 16242–16247.
- 26 Yount NY, Yeaman MR. Emerging themes and therapeutic prospects for anti-infective peptides. *Annu. Rev. Pharmacol. Toxicol.* 2012; **52**: 337–360.
- 27 Kyte J, Doolittle RF. A simple method for displaying the hydropathic character of a protein. *J. Mol. Biol.* 1982; **157**: 105–132.
- 28 Harwig SS, Waring A, Yang HJ, Cho Y, Tan L, Lehrer RI. Intramolecular disulfide bonds enhance the antimicrobial and lytic activities of protegrins at physiological sodium chloride concentrations. *Eur. J. Biochem.* 1996; **240**: 352–357.
- 29 Goldman MJ, Anderson GM, Stolzenberg ED, Kari UP, Zasloff M, Wilson JM. Human beta-defensin-1 is a salt-sensitive antibiotic in lung that is inactivated in cystic fibrosis. *Cell* 1997; **88**: 553–560.
- 30 Mosca DA, Hurst MA, So W, Viajar BS, Fujii CA, Falla TJ. IB-367, a protegrin peptide with in vitro and in vivo activities against the microflora associated with oral mucositis. *Antimicrob. Agents Chemother.* 2000; **44**: 1803–1808.
- 31 Srinivas N, Jetter P, Ueberbacher BJ, Werneburg M, Zerbe K, Steinmann J, Van der Meijden B, Bernardini F, Lederer A, Dias RL, Misson PE, Henze H, Zumbunn J, Gombert FO, Obrecht D, Hunziker P, Schauer S, Ziegler U, Kach A, Eberl L, Riedel K, DeMarco SJ, Robinson JA. Peptidomimetic antibiotics target outer-membrane biogenesis in *Pseudomonas aeruginosa*. *Science* 2010; **327**: 1010–1013.
- 32 Lai JR, Huck BR, Weisblum B, Gellman SH. Design of non-cysteine-containing antimicrobial beta-hairpins: structure-activity relationship studies with linear protegrin-1 analogues. *Biochemistry* 2002; **41**: 12835–12842.
- 33 Sagaram US, Pandurangi R, Kaur J, Smith TJ, Shah DM. Structure-activity determinants in antifungal plant defensins MsDef1 and MtDef4 with different modes of action against *Fusarium graminearum*. *PLoS One* 2011; **6**: e18550.
- 34 Janssen BJ, Schirra HJ, Lay FT, Anderson MA, Craik DJ. Structure of *Petunia hybrida* defensin 1, a novel plant defensin with five disulfide bonds. *Biochemistry* 2003; **42**: 8214–8222.
- 35 Fant F, Vranken W, Broekaert W, Borremans F. Determination of the three-dimensional solution structure of *Raphanus sativus* antifungal protein 1 by 1H NMR. *J. Mol. Biol.* 1998; **279**: 257–270.
- 36 Fant F, Vranken WF, Borremans FA. The three-dimensional solution structure of *Aesculus hippocastanum* antimicrobial protein 1 determined by 1H nuclear magnetic resonance. *Proteins* 1999; **37**: 388–403.
- 37 Bhunia A, Mohanram H, Domadia PN, Torres J, Bhattacharjya S. Designed beta-boomerang antiendotoxic and antimicrobial peptides: structures and activities in lipopolysaccharide. *J. Biol. Chem.* 2009; **284**: 21991–22004.