

Contents lists available at [SciVerse ScienceDirect](http://SciVerse.Sciencedirect.com)

Biochimica et Biophysica Acta

journal homepage: www.elsevier.com/locate/bbamcr

Role of GALNT2 in the modulation of ENPP1 expression, and insulin signaling and action GALNT2: A novel modulator of insulin signaling



Antonella Marucci ^a, Flora Cozzolino ^b, Claudia Dimatteo ^a, Maria Monti ^b, Piero Pucci ^b,
Vincenzo Trischitta ^{a,c,*}, Rosa Di Paola ^{a,**,1}

^a Research Unit of Diabetes and Endocrine Diseases, IRCCS “Casa Sollievo della Sofferenza”, San Giovanni Rotondo, Italy

^b CEINGE Biotecnologie Avanzate, Naples and Dipartimento di Scienze Chimiche, Università Federico II, Naples, Italy

^c Department of Experimental Medicine, Sapienza University, Rome, Italy

ARTICLE INFO

Article history:

Received 18 October 2012

Received in revised form 25 February 2013

Accepted 27 February 2013

Available online 14 March 2013

Keywords:

3'UTR binding proteins

GALNT2

ENPP1

Insulin signaling

ABSTRACT

Ectonucleotide pyrophosphatase phosphodiesterase 1 (ENPP1) inhibits insulin signaling and action. Understanding the mechanisms underlying *ENPP1* expression may help unravel molecular mechanisms of insulin resistance. Recent data suggest a role of *ENPP1*-3'UTR untranslated region (UTR), in controlling *ENPP1* expression. We sought to identify *trans*-acting *ENPP1*-3'UTR binding proteins, and investigate their role on insulin signaling.

By RNA pull-down, 49 proteins bound to *ENPP1*-3'UTR RNA were identified by mass spectrometry (MS). Among these, *in silico* analysis of genome wide association studies and expression profile datasets pointed to N-acetylgalactosaminyltransferase 2 gene (*GALNT2*) for subsequent investigations. Gene expression levels were evaluated by RT-PCR. Protein expression levels, IRS-1 and Akt phosphorylation were evaluated by Western blot. Insulin receptor (IR) autophosphorylation was evaluated by ELISA.

GALNT2 down-regulation increased while *GALNT2* over-expression reduced ENPP1 expression levels. In addition, *GALNT2* down-regulation reduced insulin stimulation of IR, IRS-1 and Akt phosphorylation and insulin inhibition of phosphoenolpyruvate carboxykinase (*PEPCK*) expression, a key neoglucogenetic enzyme.

Our data point to *GALNT2* as a novel factor involved in the modulation of ENPP1 expression as well as insulin signaling and action in human liver HepG2 cells.

© 2013 Elsevier B.V. All rights reserved.

1. Introduction

Insulin resistance plays a major role in the pathogenesis of type 2 diabetes, pro-atherogenic dyslipidemia and cardiovascular disease, thus imposing a tremendous burden to morbidity and mortality in developed countries [1].

The molecular mechanisms underlying insulin resistance are mostly unraveled [2]. Among several molecules involved in this process, the class II transmembrane glycoprotein ectonucleotide pyrophosphatase phosphodiesterase 1 (ENPP1) was identified as a putative candidate. ENPP1 binds to [3] and inhibits the insulin receptor (IR) and subsequent downstream insulin signaling and action in both cultured cells [4–10] and animal models [9,11,12]. In addition, ENPP1 is over-expressed in

several tissues of insulin-resistant subjects [13–17]. Finally, *in vitro* and *in vivo* studies on a gain of function amino acid substitution (*i.e.* K121Q polymorphism; rs1805101) provided further support to the notion that ENPP1 affects insulin action [7,10,18–22]. Thus, unraveling the mechanisms involved in the modulation of ENPP1 expression might help developing strategies aimed at counteracting and possibly reversing some forms of insulin resistance.

Recent data indicated that the 3' untranslated region (UTR) of *ENPP1* mRNA plays a role in controlling ENPP1 expression, suggesting the existence of *trans*-acting proteins that affect *ENPP1* mRNA stability [23,24]. The aim of this study was to identify specific *ENPP1*-3' UTR binding proteins by means of RNA pull-down experiments and to investigate their potential role on affecting ENPP1 expression and modulating insulin signaling and action in human liver cells, the ultimate target of insulin action on glucose homeostasis.

2. Materials and methods

2.1. RNA preparation

The most conserved region (*i.e.* nucleotide (nt) 2750 to nt 3176 of the ENPP1 cDNA) of *ENPP1*-3'UTR (either wild type or polymorphic-“P” [25])

Abbreviations: ENPP1, ectonucleotide pyrophosphatase phosphodiesterase 1; UTR, untranslated region; GALNT2, N-acetylgalactosaminyltransferase 2 gene; IR, insulin receptor; PEPCK, phosphoenolpyruvate carboxykinase; GWAS, genome wide association studies; HEK293, human embryonic kidney; HepG2, human hepatoma cell line; GAPDH, glyceraldehyde 3-phosphate dehydrogenase

* Corresponding author. Tel.: +39 0882 416259; fax: +39 0882 416266.

** Corresponding author. Tel.: +39 0882 416276; fax: +39 0882 416266.

E-mail addresses: vincenzo.trischitta@uniroma1.it (V. Trischitta),

r.dipaola@operapadrepio.it (R. Di Paola).

¹ These two authors equally supervised the entire study.

was obtained as previously described [24] and then used as template. The antisense sequence of the same region was used as control RNA.

2.2. RNA pull-down

In order to identify those proteins which specifically bind the *ENPP1*-3'UTR, the RNA pull-down with adipic acid dehydrazide bead method was used. Briefly, 15 µg of *ENPP1*-3'UTR, "P" *ENPP1*-3'UTR or control RNA, obtained as previously described [24], was placed in a 400 µl reaction mixture containing 100 mmol/l NaOAc, pH 5.2, and 5 mmol/l sodium m-periodate (Sigma-Aldrich, St. Louis, MO), incubated for 1 h in the dark at room temperature, ethanol precipitated, and resuspended in 100 µl of 100 mmol/l NaOAc, pH 5.2. Then 300 µl of adipic acid dehydrazide agarose beads 50% slurry (Sigma-Aldrich, St. Louis, MO) equilibrated in 100 mmol/l NaOAc, pH 5.2, was added to this mixture and incubated for 12 h at 4 °C on a rotator. The RNA that covalently immobilized onto agarose beads was pelleted, washed twice with 1 ml of 2 mol/l NaCl, equilibrated in washing buffer (5 mmol/l HEPES, pH 7.9, 1 mmol/l MgCl₂, 0.8 mmol/l magnesium acetate) and, finally, incubated with 3 mg of cellular protein extract for 30 min at room temperature in 0.6 ml final volume. Heparin was also added to a final concentration of 7 µg/µl. After further washing (*i.e.* four times in 1.5 ml of washing buffer), proteins retained by RNA sequences were eluted in SDS sample buffer and fractionated by a 12.5% SDS-PAGE. Gels were stained with colloidal blue staining kit (Invitrogen Life Technologies, Carlsbad, CA) [26,27].

2.3. Identification and characterizations of *ENPP1*-3'UTR binding proteins

Protein bands were excised from the destained gel, reduced, alkylated and digested with trypsin [28]. Peptide mixtures were extracted from the gel and analyzed by nano-chromatography tandem MS (nanoLC-MS/MS) on a CHIP MS Ion Trap XCT Ultra equipped with a capillary 1100 HPLC system and a chip cube (Agilent Technologies, Palo Alto, CA). Peptide analysis was performed using data-dependent acquisition of one MS scan (mass range from 400 to 2000 m/z) followed by MS/MS scans of the three most abundant ions in each MS scan. Raw data from nanoLC-MS/MS analyses were employed to query a non-redundant protein database using in house MASCOT software (Matrix Science, Boston, MA). Proteins identified in both the control and the sample lane were discarded, whereas those proteins solely identified in the sample and absent in the control were selected as putative *ENPP1*-3'UTR interactors.

2.4. In silico analysis

In order to prioritize proteins that were specifically bound to *ENPP1*-3'UTR and identified by mass spectrometry analysis, we interrogated and cross-checked genome wide association studies (GWAS) and expression profile datasets. Only proteins coded by genes that satisfied the following criteria were selected for further analyses:

1. Genes containing or being nearby single nucleotide polymorphisms (SNPs) showing genome-wide association with one or more of the following traits: adiposity measures (*i.e.* BMI and waist circumference), insulin resistance (*i.e.* homeostatic model assessment of insulin resistance – HOMA-IR index – and fasting insulin), glucose homeostasis (*i.e.* fasting and 2 h glucose at OGTT and type 2 diabetes), and pro-atherogenic dyslipidemia (*i.e.* triglycerides and HDL-cholesterol).
2. Genes whose expression in human and animal tissues is relevant for glucose homeostasis (*i.e.* liver, skeletal muscle and adipose tissue) was significantly ($p < 0.05$) associated with at least one of the above-mentioned trait.

The *in silico* analysis was carried out by browsing:

- HuGENavigator/GWAS Integrator (www.hugenavigator.net/HuGENavigator/AHitStartPage.do), a bioinformatics tool that provides robust lookup and analytic functionalities for all published GWAS.
- NextBio (NextBio, www.nextbio.com) that allows exploring experimental data from an extensive set of public sources, covering a broad range of techniques and therapeutic areas.

2.5. Cell culture

HEK293 (human embryo kidney) or HepG2 (human hepatoma cell line) cells (ECACC, Salisbury, UK) were maintained at 37 °C and 5% CO₂ in DMEM/F12 containing 10% FBS (EuroClone S.p.A., Milano, Italy). Before experiments, HepG2 cells were seeded in six-well plates and grown in DMEM/F12 complete medium for 48 h.

2.6. *GALNT2* siRNA and cDNA transfections

HepG2 cells were either transfected with 10 nmol/l of small interfering RNA (siRNA) targeted against *GALNT2* mRNA (ON-TARGETplus SMARTpool Human *GALNT2* L-011865-01-0005 Thermo-Scientific Dharmacon Lafayette, CO) or with 10 nmol/l of scrambled siRNA (ON-TARGETplus non-targeting Pool D-001810-10-20 Thermo-Scientific Dharmacon Lafayette, CO) by using Interferin™ Transfection Reagents (Polyplus, Illkirch, France), according to the manufacturer's instructions. After 48 or 96 h treatment, followed by cell lysis, equal amount of proteins was analyzed by Western blot as described below.

Otherwise, HepG2 were transiently transfected either with empty vector (mock cells) or with *GALNT2* cDNA Myc/DDK tagged, cloned in pCMV6 Entry vector TrueORF Gold (OriGene Technologies, Rockville, MD) (*GALNT2* cells), by using TransIT Reagent (Mirus, Madison, WI), according to the manufacturer's instructions.

2.7. RNA extraction, cDNA synthesis, and gene expression analysis

Total RNA was isolated from cells using RNeasy Mini kit (Qiagen S.r.l., Milan, Italy). cDNA was generated by reverse transcription with iScript™ Reverse Transcription (Biorad, Hercules, CA) according to the manufacturer's instructions and used as template in the subsequent analyses. Gene Expression Assay on Demand Kit Reagents (Applied Life Technologies, Carlsbad, CA) or PrimeTime Std qPCR Assay (IDT, San Jose, CA) were used to quantify relative gene expression levels of *ENPP1*, *PEPCK* (phosphoenolpyruvate carboxykinase) and *GAPDH* (glyceraldehyde 3-phosphate dehydrogenase) on ABI-PRISM 7500 (Applied Life Technologies, Carlsbad, CA). Expression levels of *ENPP1* and *PEPCK* were calculated by using the comparative $\Delta\Delta\text{CT}$ method. Briefly, the amount of *ENPP1* and *PEPCK* were normalized to *GAPDH* as endogenous reference ($2^{-\Delta\text{CT}}$) in experiments run in triplicate and expressed as percentage of control cells of the first of several experiments.

2.8. Antibodies

Antibody directed against *GALNT2* was obtained from Abcam (Cambridge, UK). Anti-*ENPP1* anti-*GAPDH* and anti-IR β -subunit antibodies were from Santa Cruz Biotechnology (Santa Cruz, CA). Anti-Phospho-IRS-1 Tyr⁸⁹⁵, anti-IRS-1, anti-Phospho-Akt Ser⁴⁷³ and anti-Akt, antibodies were purchased from Cell signaling (Boston, MA).

2.9. Western blot analysis

Cell lysates were separated by SDS-PAGE and transferred to nitrocellulose membrane (Amersham Pharmacia Biotech, Piscataway, NJ). After blotting with specific antibodies described above, immunocomplexes were detected with the Super Signal West Pico (Thermo Fisher

Scientific Pierce, Rockford, IL). Gel images were acquired by Molecular Imager ChemiDoc XRS (Biorad, Hercules, CA) and band intensities were measured by Kodak Molecular Imaging Software 4.0 as Optical Density (OD) values. ENPP1 protein expression was evaluated by ENPP1/GAPDH OD ratio and expressed as means \pm SD. IRS-1-Tyr⁸⁹⁵ and Akt-Ser⁴⁷³ phosphorylations were calculated as IRS-1-Tyr⁸⁹⁵ phosphorylation/IRS-1 or Akt-Ser⁴⁷³ phosphorylation/Akt OD ratio respectively and expressed as means \pm SD.

2.10. IR β -subunit autophosphorylation

After 18 h starvation, both GALNT2 silenced and GALNT2 transfected as well as their respective control HepG2 cells were stimulated with 100 nmol/l insulin for 5 min and lysed. IR β -subunit autophosphorylation was analyzed by using CST's PathScan Phospho-Insulin Receptor β (Tyr1150/1151) solid phase sandwich enzyme-linked immunosorbent assay (ELISA) Kit according to manufacturer's instruction (Cell signaling, Boston, MA). Briefly, equal amount of protein was incubated with a coated anti-Insulin Receptor β mouse antibody in order to capture both phospho- and nonphospho-insulin receptor proteins. Following extensive washing, anti-phospho Insulin Receptor β (Tyr1150/1151) rabbit antibody is added to detect the captured phospho-insulin receptor protein. Anti-rabbit IgG, HRP-linked antibody is then used to recognize the bound detection antibody. OD values were used as a measure of IR β -subunit autophosphorylation. Data are means \pm SD.

2.11. IRS-1-Tyr⁸⁹⁵ and Akt-Ser⁴⁷³ phosphorylation

After 18 h starvation, GALNT2 silenced or not silenced HepG2 cells were stimulated with 100 nmol/l insulin for 5 min and lysed; then, equal amount of proteins was analyzed by Western blot and probed with anti-Phospho-IRS-1 Tyr⁸⁹⁵, anti-Phospho-Akt Ser⁴⁷³, anti IRS-1 or anti-Akt. IRS-1 and Akt phosphorylation levels were normalized against IRS-1 or Akt content and expressed as described above.

2.12. PEPCK mRNA expression levels

Insulin action on glucose metabolism was assessed by studying mRNA level of the gluconeogenic enzyme PEPCK in HepG2 cells. After 18 h starvation, silenced or not silenced HepG2 cells were stimulated with 100 nmol/l insulin for 10 h. Total RNA extraction and cDNA synthesis and PEPCK mRNA level measurement were performed as previously reported [10] and calculated as described above.

2.13. Statistical analyses

Differences between the mean values were evaluated by Student's *t* test. Data are presented as means \pm SD. SPSS 13 software package was used for all analyses.

3. Results

3.1. Identification and characterization of GALNT2 as an ENPP1-3'UTR binding proteins

Proteins that specifically bind the ENPP1-3'UTR were identified by a functional proteomic approach using affinity capture and tandem MS. Fig. 1 shows the corresponding gel stained with colloidal blue Coomassie. The entire sample and control lanes of the gel were cut in slices and proteins occurring in each slice were identified by tandem-MS. Common proteins identified in both the sample and the control gel slices were eliminated, thus greatly decreasing the number of false positives. A total of forty-nine putative candidate proteins were identified (Supplementary Table 1).

A screening procedure was applied to the list of identified proteins to select the most promising 3'UTR interactors by interrogating

GWAS and expression profile datasets using the selection criteria described in Materials and methods. Among the forty-nine putative protein interactors, ten matched the inclusion criteria at the GWAS dataset, whereas only two proteins were selected when using the expression profile datasets. When data from the two screening procedures were cross-checked, only GALNT2 matched inclusion criteria at both datasets and was, then, selected for subsequent investigation. The GALNT2 gene codes for an UDP-N-acetyl-alpha-D-galactosamine polypeptide N-acetylgalactosaminyltransferase (O-GalNAc) which contributes to the initiation of mucin-type O-linked glycosylation [29].

3.2. Confirmation of GALNT2-ENPP1-3'UTR interaction

The interaction of GALNT2 with ENPP1-3'UTR RNA was confirmed by an additional RNA pull-down assay from HEK293 total protein extract followed by Western-blot analysis (Fig. 2). By using anti-GALNT2 specific antibody, a clear band was detected in RNA pulled-down human HEK293 cells (Fig. 2 panel A lane 2, *n* = 2), thus confirming that GALNT2 binds to ENPP1-3'UTR. The interaction between ENPP1 and GALNT2 was further confirmed in typical human insulin target cells, namely liver HepG2 cells (Fig. 2 panel B). Such binding was not influenced by the polymorphic "P" 3'UTR haplotype (Supplementary Fig. 1), reported to affect ENPP1 mRNA expression [25].

3.3. Role of GALNT2 downregulation on ENPP1 expression and on insulin signaling and action

Exposure of HepG2 cells to GALNT2 siRNA for both 48 and 96 h almost abolished GALNT2 expression as compared to scrambled siRNA transfected cells (Fig. 3, panel A: lanes 2 vs. 1, *n* = 3 and lanes 4 vs. 3, *n* = 5, respectively). As compared to scrambled siRNA transfected cells, in GALNT2 downregulated cells (Fig. 3, panel B), ENPP1 expression tended to be increased at 48 h (*i.e.* 43% increase, *p* = 0.05–0.1), and was significantly increased at 96 h (116% increase, *p* < 0.05). Coherently, although to a lesser extent, also ENPP1 protein level was increased at 96 h in GALNT2 downregulated as compared to control cells (27% increase, *p* < 0.01, Fig. 3, panel C).

Given the well established inhibitory role of ENPP1 on insulin signaling and action [10], we tested the effect of GALNT2 downregulation on such pathways. After GALNT2 downregulation, insulin-induced IR




Fig. 1. SDS-PAGE of RNA protein pulled-down. HEK293 total protein extract was incubated with either agarose beads alone (lane 1) or ENPP1-3'UTR RNA (lane 2) or control RNA (lane 3) covalently linked to agarose beads. The complex were eluted with SDS sample buffer, loaded on 12.5% SDS-PAGE, separated by electrophoresis and stained with colloidal blue staining kit. Entire lanes observed in the gel were cut in slices and submitted to the identification procedure as described in Materials and methods.




Fig. 2. GALNT2/ENPP1-3'UTR interaction in HEK293 and HepG2 cells. RNA pull-down was performed as described in [Materials and methods](#). Following incubation with 3 mg of HEK293 (panel A), or HepG2 (panel B) total cell lysates, pulled-down proteins with either ENPP1-3'UTR, or control RNA were analyzed by Western blot with anti-GALNT2 specific antibody. A clear band was evident in ENPP1-3'UTR (lane 2, panels A and B) pulled-down proteins while no band was detectable in control RNA pulled-down proteins (lane 3, panels A and B). GALNT2 expression in 50 µg of total cell lysate is shown in lane 1, panels A and B.

β-subunit autophosphorylation (Fig. 4 panel B) was significantly reduced as compared to insulin-stimulated scrambled siRNA cells (23% reduction, $p < 0.05$). IR protein content was superimposable across all different experimental conditions (Fig. 4 panel A). In GALNT2 siRNA cells also IRS-1 Tyr⁸⁹⁵ (Fig. 5, panel A) and Akt-Ser⁴⁷³ (Fig. 5, panel B) phosphorylations were significantly reduced as compared to what

observed in insulin-stimulated scrambled siRNA cells (25% reduction $p < 0.01$ and 57% reduction $p < 0.05$ respectively).

Insulin significantly suppressed gluconeogenic enzyme *PEPCK* mRNA levels in scrambled siRNA cells (54% reduction, $p < 0.01$). This effect was partially lost in GALNT2 siRNA HepG2 cells in which only a 33% reduction of *PEPCK* mRNA levels was found ($p =$ not significant vs. non insulin stimulated cells; Fig. 6). In addition, *PEPCK* mRNA levels in scrambled siRNA insulin-stimulated cells were significantly lower than those in GALNT2 siRNA insulin-stimulated cells ($p < 0.05$, Fig. 6). Thus, the observed deleterious effect of GALNT2 downregulation on *ENPP1* expression is paralleled by a similar effect on downstream insulin signaling and action on glucose metabolism.

3.4. Role of GALNT2 upregulation on ENPP1 expression and on insulin signaling

Exposure of HepG2 cells to GALNT2 cDNA for 96 h, which almost doubled GALNT2 expression (Fig. 7, panel A), reduced *ENPP1* mRNA expression levels by 30% as compared to mock HepG2 control cells ($p < 0.01$, Fig. 7, panel B), thus mirroring the data obtained by GALNT2 downregulation. In contrast, GALNT2 upregulation did not affect *ENPP1* protein content as shown in Fig. 7, panel C. This lack of effect is observed despite the maintained ability of transfected GALNT2 to bind *ENPP1*-3'UTR (Supplementary Fig. 2). Conversely, these apparently discordant results may be due to a high *ENPP1* protein stability in HepG2 cells which makes it difficult to detect any effect of *ENPP1* mRNA reduction due to GALNT2 overexpression. To test this hypothesis, we exposed cells to *ENPP1* siRNA for 96 h and then measured both *ENPP1* mRNA and protein content. While *ENPP1* mRNA expression levels were decreased by $39 \pm 12\%$ as compared to scrambled siRNA HepG2 transfected cells ($n = 3$ experiments,




Fig. 3. Effect of GALNT2 siRNA on *ENPP1* expression. A. HepG2 cells were transfected with either scrambled siRNA or GALNT2 siRNA for 48 or 96 h, as described in [Materials and methods](#). Expression of GALNT2 (upper blot) and GAPDH (lower blot) was evaluated by Western blot by using anti-GALNT2 or anti-GAPDH specific antibody respectively. A representative blot for each condition is shown. B. HepG2 cells were either treated with scrambled siRNA (black bar) or GALNT2 siRNA (gray bar) and *ENPP1* mRNA expression levels were measured by Real-Time PCR as described in [Materials and methods](#). Data are expressed as percentage of scrambled siRNA cells of the first experiment at 48 h ($n = 3$) and 96 h ($n = 5$) respectively. Data are means \pm SD. * $p < 0.05$. C. HepG2 cells were either transfected with scrambled siRNA or GALNT2 siRNA for 96 h, as described in [Materials and methods](#). Equal amount of protein from cell lysates was separated by SDS-PAGE and probed with anti-*ENPP1* and anti-GAPDH specific antibody. Bars (upper panel) represent *ENPP1*/GAPDH OD ratio (representative immunoblots are shown in lower panel). Data are means \pm SD of 6 experiments in separate times. § $p < 0.01$.




Fig. 6. Effect of *GALNT2* siRNA on *PEPCK* expression. HepG2 cells were either transfected with scrambled siRNA or *GALNT2* siRNA for 96 h and stimulated with insulin as described in [Materials and methods](#). *PEPCK* mRNA expression levels were measured by Real-Time PCR, as described in [Materials and methods](#) and expressed as percentage of unstimulated scrambled siRNA cells of the first experiment. Data are means \pm SD of 4 experiments in separate times. * $p < 0.05$ and § $p < 0.01$.

Fig. 4. Effect of *GALNT2* siRNA on IR β -subunit autophosphorylation. HepG2 cells were either transfected with scrambled siRNA or *GALNT2* siRNA for 96 h and then stimulated with insulin as described in [Materials and methods](#). A. Equal amount of protein from cell lysates was separated by SDS-PAGE. IR and GAPDH protein content was evaluated by Western blot by using anti-IR β -subunit or anti-GAPDH specific antibody, respectively (a representative blot out of 3 experiments for each condition is shown). B. Bars represent OD values of IR β -subunit autophosphorylation as assessed by ELISA (see [Materials and methods](#)). Data are means \pm SD of 3 experiments in separate times. * $p < 0.05$.

$p < 0.05$), ENPP1 protein content was totally unaffected ($110 \pm 15\%$ as compared to scrambled siRNA cells, $n = 3$ experiments, $p < 0.05$), thus reinforcing the hypothesis of a high ENPP1 protein stability, which does not allow to appreciate variation in protein level, at least under these experimental conditions.




Fig. 5. Effect of *GALNT2* siRNA on IRS-1-Y⁸⁹⁵ and Akt-S⁴⁷³ phosphorylation. HepG2 cells were either transfected with scrambled siRNA or *GALNT2* siRNA for 96 h and then stimulated with insulin as described in [Materials and methods](#). Equal amount of protein from cell lysates was separated by SDS-PAGE. A. Insulin stimulated IRS-1-Y⁸⁹⁵ phosphorylation and both IRS-1 and GAPDH protein content were evaluated by Western blot analysis by using anti phospho-IRS-1-Y⁸⁹⁵, anti-IRS-1 or anti-GAPDH specific antibody, respectively. Bars (upper panel) represent quantitative analysis of IRS-1-Y⁸⁹⁵ phosphorylation calculated as IRS-1-Y⁸⁹⁵ phosphorylation/IRS-1 OD ratio. Data are means \pm SD of 4 experiments in separate times. § $p < 0.01$. B. Insulin stimulated Akt-Ser⁴⁷³ phosphorylation and both Akt and GAPDH protein content were evaluated by Western blot analysis by using anti phospho-Akt-Ser⁴⁷³, anti-Akt or anti-GAPDH specific antibody, respectively. Bars (upper panel) represent quantitative analysis of Akt-Ser⁴⁷³ phosphorylation calculated as Akt-Ser⁴⁷³ phosphorylation/Akt OD ratio. Data are means \pm SD of 3 experiments in separate times. * $p < 0.05$.

Coherently with the unchanged ENPP1 protein content, insulin-induced IR β -subunit autophosphorylation was similar in GALNT2 overexpression and mock HepG2 control cells (Fig. 8 panel B). IR protein content was superimposable across all different experimental conditions (Fig. 8 panel A). Based upon these negative results no further experiments on downstream insulin signaling were carried out.

4. Discussion

Functional [4,7,10] metabolic [13,14,16] and genetic [18,23,25,30–38] studies suggest that ENPP1 acts as a negative modulator of insulin signaling and action. Thus, addressing the mechanisms underlying ENPP1 expression may help unravel novel molecular mechanisms of insulin resistance.

Our study on human liver HepG2 cells shows that GALNT2 binds to ENPP1-3'UTR mRNA and inhibits ENPP1 transcript and protein levels. This finding adds to previous functional [24,25] and genetic [23,25] reports suggesting that, in fact, the 3'UTR is involved in the regulation of ENPP1 expression in human liver cells, the utmost target of insulin action on glucose homeostasis. Of note, a previously reported 3'UTR polymorphic haplotype did not impair GALNT2 binding affinity, thus suggesting that its effect on ENPP1 mRNA levels [25] is not mediated by GALNT2. Whether other SNPs associated with metabolic traits [23] which are located outside the most conserved 3'UTR we used for our experiments, affect GALNT2 binding may be interested to be addressed by further investigations. Also of note is that HSP70 was not identified as a binding protein of ENPP1-3'UTR, a finding which is in contrast with our previous reports [24]. We like to speculate that the different methods utilized in our present and previous studies (RNA-pull-down vs. RNA Mobility Shift Analysis) underline the different results we obtained.

A second important finding of this study is that GALNT2 downregulation affects insulin-induced IR β -subunit autophosphorylation, IRS-1 Tyr⁸⁹⁵ and Akt Ser⁴⁷³ phosphorylations and PEPCK expression, four key steps of insulin signaling and gluconeogenic




Fig. 7. Effect of GALNT2 transfection on ENPP1 expression. A. HepG2 cells were either transfected with mock or with GALNT2 cDNA as described in Materials and methods. Equal amount of protein from cell lysates was separated by SDS-PAGE and probed with anti-GALNT2 (upper blot) or anti-GAPDH (lower blot) specific antibody. A representative blot out of 3 experiments for each condition is shown. B. ENPP1 expression levels were measured by Real-Time PCR as described in Materials and methods and expressed as percentage of mock HepG2 cells of the first experiment. Data are means \pm SD of 3 experiments in separate times. § $p < 0.01$. C. HepG2 cells were either transfected with mock or with GALNT2 cDNA as described in Materials and methods. Equal amount of protein from cell lysates was separated by SDS-PAGE and probed with anti-ENPP1 and anti-GAPDH specific antibody. Bars (upper panel) represent ENPP1/GAPDH OD ratio (representative immunoblots are shown in lower panel). Data are means \pm SD of 3 experiments in separate times.




Fig. 8. Effect of GALNT2 transfection on IR β -subunit autophosphorylation. HepG2 cells were either transfected with mock or with GALNT2 cDNA and then stimulated with insulin as described in Materials and methods. A. Equal amount of protein from cell lysates was separated by SDS-PAGE. IR and GAPDH protein content was evaluated by Western blot by using anti-IR β -subunit or anti-GAPDH specific antibody, respectively (a representative blot out of 3 experiments for each condition is shown). B. Bars represent OD values of IR β -subunit autophosphorylation as assessed by ELISA (see Materials and methods). Data are means \pm SD of 3 experiments in separate times.

activity. These effects are very likely mediated, by ENPP1 protein up-regulation. In fact, GALNT2 overexpression, which down-regulates ENPP1 mRNA levels but not ENPP1 protein content, is not paralleled by any change of insulin-induced IR β -subunit autophosphorylation. This finding reinforces the hypothesis that ENPP1 up-regulation, mediates, at least partly, GALNT2 down-regulation effects on insulin signaling and action. We acknowledge that further experiments using cellular models lacking both ENPP1 alleles, are needed to definitively confirm such hypothesis.

It is entirely possible that other mechanisms also underlie such GALNT2 effects. As a matter of fact, GalNAc-T2 coded by GALNT2 is responsible for the O-linked glycosylation, allowing the transfer of *N*-acetylgalactosamine from UDP-GalNAc to the hydroxyl group of a serine or threonine residue [29]. Such glycosylation has been reported to play an important role on insulin resistance and diabetes, either by competing for phosphorylation on insulin-stimulated sites on effector molecules, or by directly regulating components of insulin signaling, including IRS1 and Akt [39–42].

Regardless the mechanism of action, also data from the few studies so far published on GALNT2 expression levels in several models of metabolic abnormalities related to insulin resistance suggests that it modulates insulin resistance traits. While GALNT2 expression is reduced in liver of insulin resistant Goto-Kakizaki diabetic rats [43], it is increased in adipose tissue of obese, insulin resistant non-diabetic PIMA Indians [44]. In addition, GALNT2 expression in mice causes an inverse modulation of serum HDL-c levels, an established marker of insulin sensitivity [45]. All in all, expression data in both humans and rodents, although somehow variable, clearly point to GALNT2 as a potential mediator of abnormalities related to insulin resistance. Further support to this hypothesis comes from studies showing that genetic variability at the GALNT2 locus (SNP rs4846914) is associated with decreased HDL-c and increased triglyceride levels [46–48], two main components of the insulin resistance/metabolic syndrome.

5. Conclusions

In conclusion, our data identified GALNT2 as a new potential modulator of insulin signaling and action in human liver cells through the modulation of ENPP1 expression. Additional experiments in other cell types and tissues relevant to glucose metabolism are needed to confirm this finding, thus deeper addressing the role of GALNT2 as a potential target of novel treatments of insulin resistance and abnormal glucose homeostasis. Finally, because of the very restrictive prioritizing criteria we used to identify GALNT2, the list of ENPP1-3'UTR binding proteins may, in fact, contain additional important candidates to be investigated in further studies.

Supplementary data to this article can be found online at <http://dx.doi.org/10.1016/j.bbamcr.2013.02.032>.

Acknowledgements

This research was supported by the Italian Ministry of Health Grants: RC2010, RC2011 (R.D.P.), and RC2011 (V.T.); the Fondazione Roma “Sostegno alla ricerca scientifica biomedica 2008” (V.T.), and the FIRB Project Rete Nazionale per lo Studio della Proteomica Umana (Italian Human ProteomeNet) from the MIUR (P.P). F.C. was supported by P.O.R. Campania FSE 2007–2013, Project “CREMe”.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Disclosure statement

The authors declare that they do not have any actual or potential conflict of interest.

References

- [1] G.M. Reaven, Banting lecture 1988. Role of insulin resistance in human disease, *Diabetes* 37 (1988) 1595–1607.
- [2] C.M. Taniguchi, B. Emanuelli, C.R. Kahn, Critical nodes in signalling pathways: insights into insulin action, *Nat. Rev. Mol. Cell Biol.* 7 (2006) 85–96.
- [3] B.A. Maddux, I.D. Goldfine, Membrane glycoprotein PC-1 inhibition of insulin receptor function occurs via direct interaction with the receptor alpha-subunit, *Diabetes* 49 (2000) 13–19.
- [4] B.A. Maddux, P. Sbraccia, S. Kumakura, S. Sasson, J. Youngren, A. Fisher, S. Spencer, A. Grupe, W. Henzel, T.A. Stewart, et al., Membrane glycoprotein PC-1 and insulin resistance in non-insulin-dependent diabetes mellitus, *Nature* 373 (1995) 448–451.
- [5] A. Belfiore, A. Costantino, F. Frasca, G. Pandini, R. Mineo, P. Vigneri, B. Maddux, I.D. Goldfine, R. Vigneri, Overexpression of membrane glycoprotein PC-1 in MDA-MB231 breast cancer cells is associated with inhibition of insulin receptor tyrosine kinase activity, *Mol. Endocrinol.* 10 (1996) 1318–1326.
- [6] C.N. Chin, Q. Dallas-Yang, F. Liu, T. Ho, K. Ellsworth, P. Fischer, T. Natasha, C. Ireland, P. Lu, C. Li, I.M. Wang, W. Strohl, J.P. Berger, Z. An, B.B. Zhang, G. Jiang, Evidence that inhibition of insulin receptor signaling activity by PC-1/ENPP1 is dependent on its enzyme activity, *Eur. J. Pharmacol.* 606 (2009) 17–24.
- [7] B.V. Costanzo, V. Trischitta, R. Di Paola, D. Spampinato, A. Pizzuti, R. Vigneri, L. Frittitta, The Q allele variant (GLN121) of membrane glycoprotein PC-1 interacts with the insulin receptor and inhibits insulin signaling more effectively than the common K allele variant (LYS121), *Diabetes* 50 (2001) 831–836.
- [8] J. Liang, M. Fu, E. Ciociola, M. Chandalia, N. Abate, Role of ENPP1 on adipocyte maturation, *PLoS One* 2 (2007) e882.
- [9] H.H. Zhou, C.N. Chin, M. Wu, W. Ni, S. Quan, F. Liu, Q. Dallas-Yang, K. Ellsworth, T. Ho, A. Zhang, T. Natasha, J. Li, K. Chapman, W. Strohl, C. Li, I.M. Wang, J. Berger, Z. An, B.B. Zhang, G. Jiang, Suppression of PC-1/ENPP-1 expression improves insulin sensitivity *in vitro* and *in vivo*, *Eur. J. Pharmacol.* 616 (2009).
- [10] R. Di Paola, N. Caporarello, A. Marucci, C. Dimatteo, C. Iadicicco, S. Del Guerra, S. Prudente, D. Sudano, C. Miele, C. Parrino, S. Piro, F. Beguinot, P. Marchetti, V. Trischitta, L. Frittitta, ENPP1 affects insulin action and secretion: evidences from *in vitro* studies, *PLoS One* 6 (2011) e19462.
- [11] H. Dong, B.A. Maddux, J. Altomonte, M. Meseck, D. Accili, R. Terkeltaub, K. Johnson, J.F. Youngren, I.D. Goldfine, Increased hepatic levels of the insulin receptor inhibitor, PC-1/NPP1, induce insulin resistance and glucose intolerance, *Diabetes* 54 (2005) 367–372.
- [12] B.A. Maddux, Y.N. Chang, D. Accili, O.P. McGuinness, J.F. Youngren, I.D. Goldfine, Overexpression of the insulin receptor inhibitor PC-1/ENPP1 induces insulin resistance and hyperglycemia, *Am. J. Physiol. Endocrinol. Metab.* 290 (2006) E746–E749.
- [13] L. Frittitta, J.F. Youngren, P. Sbraccia, M. D'Adamo, A. Buongiorno, R. Vigneri, I.D. Goldfine, V. Trischitta, Increased adipose tissue PC-1 protein content, but not tumour necrosis factor- α gene expression, is associated with a reduction of both whole body insulin sensitivity and insulin receptor tyrosine-kinase activity, *Diabetologia* 40 (1997) 282–289.
- [14] L. Frittitta, D. Spampinato, A. Solini, R. Nosadini, I.D. Goldfine, R. Vigneri, V. Trischitta, Elevated PC-1 content in cultured skin fibroblasts correlates with decreased *in vivo* and *in vitro* insulin action in nondiabetic subjects: evidence that PC-1 may be an intrinsic factor in impaired insulin receptor signaling, *Diabetes* 47 (1998) 1095–1100.
- [15] S. Teno, H. Kanno, S. Oga, S. Kumakura, R. Kanamuro, Y. Iwamoto, Increased activity of membrane glycoprotein PC-1 in the fibroblasts from non-insulin-dependent diabetes mellitus patients with insulin resistance, *Diabetes Res. Clin. Pract.* 45 (1999) 25–30.
- [16] F.B. Stentz, A.E. Kitabchi, Transcriptome and proteome expressions involved in insulin resistance in muscle and activated T-lymphocytes of patients with type 2 diabetes, *Genomics Proteomics Bioinformatics* 5 (2007) 216–235.
- [17] M. Chandalia, H. Davila, W. Pan, M. Szuszkiewicz, D. Tuvdendorj, E.H. Livingston, N. Abate, Adipose tissue dysfunction in humans: a potential role for the transmembrane protein ENPP1, *J. Clin. Endocrinol. Metab.* 97 (2012).
- [18] A. Pizzuti, L. Frittitta, A. Argiolas, R. Baratta, I.D. Goldfine, M. Bozzali, T. Ercolino, G. Scarlato, L. Iacoviello, R. Vigneri, V. Tassi, V. Trischitta, A polymorphism (K121Q) of the human glycoprotein PC-1 gene coding region is strongly associated with insulin resistance, *Diabetes* 48 (1999) 1881–1884.
- [19] N. Abate, L. Carulli, A. Cabo-Chan Jr., M. Chandalia, P.G. Snell, S.M. Grundy, Genetic polymorphism PC-1 K121Q and ethnic susceptibility to insulin resistance, *J. Clin. Endocrinol. Metab.* 88 (2003) 5927–5934.
- [20] N. Grarup, S.A. Urhammer, J. Ek, A. Albrechtsen, C. Glumer, K. Borch-Johnsen, T. Jorgensen, T. Hansen, O. Pedersen, Studies of the relationship between the ENPP1 K121Q polymorphism and type 2 diabetes, insulin resistance and obesity in 7,333 Danish white subjects, *Diabetologia* 49 (2006) 2097–2104.
- [21] R. Baratta, P. Rossetti, S. Prudente, F. Barbetti, D. Sudano, A. Nigro, M.G. Farina, F. Pellegrini, V. Trischitta, L. Frittitta, Role of the ENPP1 K121Q polymorphism in glucose homeostasis, *Diabetes* 57 (2008) 3360–3364.
- [22] E.S. Stolerman, A.K. Manning, J.B. McAteer, J. Dupuis, C.S. Fox, L.A. Cupples, J.B. Meigs, J.C. Florez, Haplotype structure of the ENPP1 gene and nominal association of the K121Q missense single nucleotide polymorphism with glycemic traits in the Framingham Heart Study, *Diabetes* 57 (2008) 1971–1977.
- [23] D. Meyre, N. Bouatia-Naji, A. Tounian, C. Samson, C. Lecoeur, V. Vatin, M. Ghossaini, C. Wachter, S. Hercberg, G. Charpentier, W. Patsch, F. Pattou, M.A. Charles, P. Tounian, K. Clement, B. Jouret, J. Weill, B.A. Maddux, I.D. Goldfine, A. Walley, P. Boutin, C. Dina, P. Froguel, Variants of ENPP1 are associated with childhood and adult obesity and increase the risk of glucose intolerance and type 2 diabetes, *Nat. Genet.* 37 (2005) 863–867.

- [24] A. Marucci, G. Miscio, L. Padovano, W. Boonyasrisawat, J.C. Florez, A. Doria, V. Trischitta, R. Di Paola, The role of HSP70 on ENPP1 expression and insulin-receptor activation, *J. Mol. Med.* 87 (2009) 139–144.
- [25] L. Frittitta, T. Ercolino, M. Bozzali, A. Argiolas, S. Graci, M.G. Santagati, D. Spampinato, R. Di Paola, C. Cisternino, V. Tassi, R. Vigneri, A. Pizzuti, V. Trischitta, A cluster of three single nucleotide polymorphisms in the 3'-untranslated region of human glycoprotein PC-1 gene stabilizes PC-1 mRNA and is associated with increased PC-1 protein content and insulin resistance-related abnormalities, *Diabetes* 50 (2001) 1952–1955.
- [26] A. Russo, C. Cirulli, A. Amoresano, P. Pucci, C. Pietropaolo, G. Russo, cis-Acting sequences and trans-acting factors in the localization of mRNA for mitochondrial ribosomal proteins, *Biochim. Biophys. Acta, Gene Regul. Mech.* 1779 (2008) 820–829.
- [27] V. Pisa, M. Cozzolino, S. Gargiulo, C. Ottone, F. Piccioni, M. Monti, S. Gigliotti, F. Talamo, F. Graziani, P. Pucci, A.C. Verrotti, The molecular chaperone Hsp90 is a component of the cap-binding complex and interacts with the translational repressor Cup during *Drosophila* oogenesis, *Gene* 432 (2009) 67–74.
- [28] E. Zito, M. Buono, S. Pepe, C. Settembre, I. Annunziata, E.M. Surace, T. Dierks, M. Monti, M. Cozzolino, P. Pucci, A. Ballabio, M.P. Cosma, Sulfatase modifying factor 1 trafficking through the cells: from endoplasmic reticulum to the endoplasmic reticulum, *EMBO J.* 26 (2007) 2443–2453.
- [29] K.G. Ten Hagen, T.A. Fritz, L.A. Tabak, All in the family: the UDP-GalNAc:polypeptide N-acetylgalactosaminyltransferases, *Glycobiology* 13 (2003) 1R–16R.
- [30] H.F. Gu, P. Almgren, E. Lindholm, L. Frittitta, A. Pizzuti, V. Trischitta, L.C. Groop, Association between the human glycoprotein PC-1 gene and elevated glucose and insulin levels in a paired-sibling analysis, *Diabetes* 49 (2000) 1601–1603.
- [31] L. Frittitta, R. Baratta, D. Spampinato, R. Di Paola, A. Pizzuti, R. Vigneri, V. Trischitta, The Q121 PC-1 variant and obesity have additive and independent effects in causing insulin resistance, *J. Clin. Endocrinol. Metab.* 86 (2001) 5888–5891.
- [32] R. Baratta, R. Di Paola, D. Spampinato, G. Fini, A. Marucci, A. Coco, R. Vigneri, L. Frittitta, V. Trischitta, Evidence for genetic epistasis in human insulin resistance: the combined effect of PC-1 (K121Q) and PPARgamma2 (P12A) polymorphisms, *J. Mol. Med.* 81 (2003) 718–723.
- [33] A. Kubaszek, J. Pihlajamaki, P. Karhapa, I. Vauhkonen, M. Laakso, The K121Q polymorphism of the PC-1 gene is associated with insulin resistance but not with dyslipidemia, *Diabetes Care* 26 (2003) 464–467.
- [34] K. Hamaguchi, H. Terao, Y. Kusuda, T. Yamashita, J.A. Hazoury Bahles, L.M. Cruz, V.L. Brugal, W.B. Jongchong, H. Yoshimatsu, T. Sakata, The PC-1 Q121 allele is exceptionally prevalent in the Dominican Republic and is associated with type 2 diabetes, *J. Clin. Endocrinol. Metab.* 89 (2004) 1359–1364.
- [35] N. Abate, M. Chandalia, P. Satija, B. Adams-Huet, S.M. Grundy, S. Sandeep, V. Radha, R. Deepa, V. Mohan, ENPP1/PC-1 K121Q polymorphism and genetic susceptibility to type 2 diabetes, *Diabetes* 54 (2005) 1207–1213.
- [36] S. Bacci, O. Ludovico, S. Prudente, Y.Y. Zhang, R. Di Paola, D. Mangiacotti, A. Raueo, D. Nolan, J. Duffy, G. Fini, L. Salvemini, C. Amico, C. Vigna, F. Pellegrini, C. Menzaghi, A. Doria, V. Trischitta, The K121Q polymorphism of the ENPP1/PC-1 gene is associated with insulin resistance/atherogenic phenotypes, including earlier onset of type 2 diabetes and myocardial infarction, *Diabetes* 54 (2005) 3021–3025.
- [37] I. Tasic, M. Milojkovic, R. Sunder-Plassmann, G. Lazarevic, N.M. Tasic, V. Stefanovic, The association of PC-1 (ENPP1) K121Q polymorphism with metabolic syndrome in patients with coronary heart disease, *Clin. Chim. Acta* 377 (2007) 237–242.
- [38] J.B. McAtteer, S. Prudente, S. Bacci, H.N. Lyon, J.N. Hirschhorn, V. Trischitta, J.C. Florez, The ENPP1 K121Q polymorphism is associated with type 2 diabetes in European populations: evidence from an updated meta-analysis in 42,042 subjects, *Diabetes* 57 (2008) 1125–1130.
- [39] L. Wells, K. Vosseller, G.W. Hart, Glycosylation of nucleocytoplasmic proteins: signal transduction and O-GlcNAc, *Science* 291 (2001) 2376–2378.
- [40] S.Y. Park, J. Ryu, W. Lee, O-GlcNAc modification on IRS-1 and Akt2 by PUGNAc inhibits their phosphorylation and induces insulin resistance in rat primary adipocytes, *Exp. Mol. Med.* 37 (2005) 220–229.
- [41] Y.A. Soesanto, B. Luo, D. Jones, R. Taylor, J.S. Gabrielsen, G. Parker, D.A. McClain, Regulation of Akt signaling by O-GlcNAc in euglycemia, *Am. J. Physiol. Endocrinol. Metab.* 295 (2008) E974–E980.
- [42] X. Yang, P.P. Ongusaha, P.D. Miles, J.C. Havstad, F. Zhang, W.V. So, J.E. Kudlow, R.H. Michell, J.M. Olefsky, S.J. Field, R.M. Evans, Phosphoinositide signalling links O-GlcNAc transferase to insulin resistance, *Nature* 451 (2008) 964–969.
- [43] R.R. Almon, D.C. DuBois, W. Lai, B. Xue, J. Nie, W.J. Jusko, Gene expression analysis of hepatic roles in cause and development of diabetes in Goto-Kakizaki rats, *J. Endocrinol.* 200 (2009) 331–346.
- [44] Y.H. Lee, S. Nair, E. Rousseau, D.B. Allison, G.P. Page, P.A. Tataranni, C. Bogardus, P.A. Permana, Microarray profiling of isolated abdominal subcutaneous adipocytes from obese vs non-obese Pima Indians: increased expression of inflammation-related genes, *Diabetologia* 48 (2005) 1776–1783.
- [45] T.M. Teslovich, K. Musunuru, A.V. Smith, A.C. Edmondson, I.M. Stylianou, M. Koseki, J.P. Pirruccello, S. Ripatti, D.I. Chasman, C.J. Willer, C.T. Johansen, S.W. Fouchier, A. Isaacs, G.M. Peloso, M. Barbalic, S.L. Ricketts, J.C. Bis, Y.S. Aulchenko, G. Thorleifsson, M.F. Feitosa, J. Chambers, M. Orho-Melander, O. Melander, T. Johnson, X. Li, X. Guo, M. Li, Y. Shin Cho, M. Jin Go, Y. Jin Kim, J.Y. Lee, T. Park, K. Kim, X. Sim, R. Twee-Hee Ong, D.C. Croteau-Chonka, L.A. Lange, J.D. Smith, K. Song, J. Hua Zhao, X. Yuan, J. Luan, C. Lamina, A. Ziegler, W. Zhang, R.Y. Zee, A.F. Wright, J.C. Witteman, J.F. Wilson, G. Willemsen, H.E. Wichmann, J.B. Whitfield, D.M. Waterworth, N.J. Wareham, G. Waeber, P. Vollenweider, B.F. Voight, V. Vitart, A.G. Uitterlinden, M. Uda, J. Tuomilehto, J.R. Thompson, T. Tanaka, I. Surakka, H.M. Stringham, T.D. Spector, N. Soranzo, J.H. Smit, J. Sinisalo, K. Silander, E.J. Sijbrands, A. Scuteri, J. Scott, D. Schlessinger, S. Sanna, V. Salomaa, J. Saharinen, C. Sabatti, A. Ruokonen, I. Rudan, L.M. Rose, R. Roberts, M. Rieder, B.M. Psaty, P.P. Pramstaller, I. Pichler, M. Perola, B.W. Penninx, N.L. Pedersen, C. Pattaro, A.N. Parker, G. Pare, B.A. Oostra, C.J. O'Donnell, M.S. Nieminen, D.A. Nickerson, G.W. Montgomery, T. Meitinger, R. McPherson, M.I. McCarthy, et al., Biological, clinical and population relevance of 95 loci for blood lipids, *Nature* 466 (2010) 707–713.
- [46] S. Kathiresan, O. Melander, C. Guiducci, A. Surti, N.P. Burt, M.J. Rieder, G.M. Cooper, C. Roos, B.F. Voight, A.S. Havulinna, B. Wahlstrand, T. Hedner, D. Corella, E.S. Tai, J.M. Ordovas, G. Berglund, E. Vartiainen, P. Jousilahti, B. Hedblad, M.R. Taskinen, C. Newton-Cheh, V. Salomaa, L. Peltonen, L. Groop, D.M. Altshuler, M. Orho-Melander, Six new loci associated with blood low-density lipoprotein cholesterol, high-density lipoprotein cholesterol or triglycerides in humans, *Nat. Genet.* 40 (2008) 189–197.
- [47] C.J. Willer, S. Sanna, A.U. Jackson, A. Scuteri, L.L. Bonnycastle, R. Clarke, S.C. Heath, N.J. Timpson, S.S. Najjar, H.M. Stringham, J. Strait, W.L. Duren, A. Maschio, F. Busonero, A. Mulas, G. Albai, A.J. Swift, M.A. Morken, N. Narisu, D. Bennett, S. Parish, H. Shen, P. Galan, P. Meneton, S. Hercberg, D. Zelenika, W.M. Chen, Y. Li, L.J. Scott, P.A. Scheet, J. Sundvall, R.M. Watanabe, R. Nagaraja, S. Ebrahim, D.A. Lawlor, Y. Ben-Shlomo, G. Davey-Smith, A.R. Shuldiner, R. Collins, R.N. Bergman, M. Uda, J. Tuomilehto, A. Cao, F.S. Collins, E. Lakatta, G.M. Lathrop, M. Boehnke, D. Schlessinger, K.L. Mohlke, G.R. Abecasis, Newly identified loci that influence lipid concentrations and risk of coronary artery disease, *Nat. Genet.* 40 (2008) 161–169.
- [48] A.G. Holleboom, H. Karlsson, R.S. Lin, T.M. Beres, J.A. Sierts, D.S. Herman, E.S. Stroes, J.M. Aerts, J.J. Kastelein, M.M. Motazacker, G.M. Dallinga-Thie, J.H. Levels, A.H. Zwinderman, J.G. Seidman, C.E. Seidman, S. Lundgren, D.J. Lefebvre, E. Morava, R.A. Wevers, T.A. Fritz, L.A. Tabak, M. Lindahl, G.K. Hovingh, J.A. Kuivenhoven, Heterozygosity for a loss-of-function mutation in GALNT2 improves plasma triglyceride clearance in man, *Cell Metab.* 14 (2011) 811–818.