

RESEARCH ARTICLE

Open Access

Day-surgery inguinal hernia repair in the elderly: single centre experience

Bruno Amato^{1*}, Rita Compagna¹, Francesca Fappiano¹, Roberto Rossi¹, Tommaso Bianco¹, Michele Danzi¹, Antonello Accurso¹, Raffaele Serra², Giovanni Aprea¹, Salvatore Massa¹

From 26th National Congress of the Italian Society of Geriatric Surgery
Naples, Italy. 19-22 June 2013

Abstract

Background: Inguinal hernioplasty is well established as a day-surgery procedure, our purpose is to assess the safeness of this approach in elderly patients.

Methods: A total of 292 inguinal hernioplasty were performed between June 2009 and February 2013. Patients were divided into 3 groups depending on the age and postoperative complications were compared in these groups.

Results: Despite of a large number of higher risk (ASA 3-4) patients and a higher rate of comorbidity in older patients, unplanned admission postoperative, symptoms and complications were comparable with those for the younger patients.

Conclusions: Ambulatory surgery is feasible also in older patients. Age, comorbidity and higher ASA risk should not be a barrier to elective day surgery.

Background

Inguinal hernia repair is one of the commonly surgical procedure with good results and minimal morbidity and it is usually performed in an ambulatory surgery setting unless coexisting medical conditions merit hospitalization for specialized monitoring or care [1-3]. Inguinal hernia is more frequent in elderly than in younger patients because of loss of strength of the abdominal wall and conditions which increase intraabdominal pressure [4,10,11]. Also the ageing population is increasing, with a corresponding increasing demand for surgical services [5]. For this reason there is a growing interest in the use of this procedure in elderly patients and recent studies suggest that it is quite safe [7-9] despite the previous idea that is better a conservative approach [5,6]. In our study we evaluate the impact of day surgery approach in octogenarians.

Methods

In the study we included patients with inguinal hernia that reached our department in the period between June 2009 and February 2013. Selection criteria used were: absence of unstable medical condition, availability of a carer overnight, access to a telephone, BMI < 32. Both unilateral and bilateral hernias were included, recurrent and complicated hernias were excluded. A total of 292 surgeries were performed by resident surgeons. Patients were divided into three age classes: a) under 75; b) between 75 and 85; c) over 85. All procedures were performed as described by Rutkow and Robbins [12] or with Lichtenstein techniques with standard-weight polypropylene mesh [13]. Prophylactic antibiotics were not used [14]. American Society of Anaesthesiologists (ASA) grade 1 and 2 patients underwent deep sedation (midazolam, fentanyl) combined with a field block of local anaesthesia (lidocaine without adrenaline) and were monitored by pulse oximetry. For ASA grade 3 and 4 patients procedures were performed in the presence of the anaesthetist and deep sedation was not used [8,15-18]. Patients were discharged within 2 h of completion of the operation

* Correspondence: bruno.amato@unina.it

¹Department of Clinical Medicine and Surgery University of Naples Federico II Via S. Pansini, 5 - 801311 Napoli, Italy
Full list of author information is available at the end of the article

after a clinical control with detailed postoperative instructions, analgesics and a 24 h contact number. Patients were telephoned by a nurse at 24\48 h post-operatively and they were asked informations about swelling, haematoma, possible wound infections. Follow up were made after 1 week, and second time after 1 month.

Results

Comorbidity found are described in table 1.

Although most of the patients in both groups were successfully discharged, a small proportion required overnight admission caused by a variety of complications and the number of unplanned admission was lightly higher in older patients (Table 2) but overall not particularly high. Main complications occurred were recurrence, wound infections (most in over 85 patients), wound hematoma. There were no deaths or major complications and all admitted patients were discharged the next day. Occurrence of unplanned contact with healthcare services was low and not significantly different between the three groups: 7 patients (5,7%) in the under 75 group; 6 patients (4,5%) in the 75/85 group; 2 patients (5,2%) in the over 85 group. 3 The main reasons for calling the healthcare service were nausea, vomiting and pain. Sometimes patients only needed a psychological comfort.

Conclusions

Data collected in this study show that hernia repair can be safely offered to elderly patients with no significant increase in complications and unplanned admissions compared with younger patient, also in over 85 patients, despite the higher percentage of comorbidity found. We also note that the telephone healthcare service helped to reduce the anxiety and ensure the patient safety. Whereas day surgery has many advantages, including avoidance of stress, reduction of the risk of contracting nosocomial infection and cost savings for the hospital, we conclude that age, comorbidity and higher ASA risk should not be a barrier to elective day surgery for inguinal hernioplasty.

Table 1 Patient Characteristics

N = 292			
Age	Under 75 121 (41.4%)	75 - 85 133 (45.5%)	Over 85 38 (13%)
ASA 1-2	90 (30.8%)	73 (25%)	9 (3%)
ASA 3-4	31 (10.6%)	60 (20.5%)	29 (10%)
Hypertension	110 (90%)	125 (94%)	37 (97%)
CD	85 (70.2%)	105 (79%)	36 (95%)
Diabetes	73 (60.3%)	40 (30%)	4 (10.5%)
Renal Failure	2 (1.6%)	3 (2.2%)	2 (5.2%)
COPD	18 (14.9%)	27 (20.3%)	11 (29%)

Table 2 Patient Complications

N = 292			
Age	Under 75 121 (41.4%)	75 - 85 133 (45.5%)	Over 85 38 (13%)
Recurrence	4 (3.3%)	9 (6.7%)	1 (2.6%)
Wound infection	1 (0.8%)	5 (3.7%)	2 (5.2%)
Wound hematoma	1 (0.8%)	2 (1.5%)	1 (2.6%)
Nerve entrapment	3 (2.5%)	2 (1.5%)	1 (2.6%)
Mesh infection	2 (1.6%)	3(2.2%)	1 (2.6%)
Severe bleeding	1 (0.8%)	3(2.2%)	1 (2.6%)
Arrhythmia	0	1 (0.75%)	2 (5.2%)

List of Abbreviations

ASA: American Society of Anaesthesiologists
 COPD: chronic obstructive pulmonary disease
 BPH: benign prostatic hypertrophy
 CD: cardiovascular disease
 BMI: body mass index

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

BA: conception and design, interpretation of data, given final approval of the version to be published.
 RC, FF, RR, TB, MD, AA, RS, GA: acquisition of data, drafting the manuscript, given final approval of the version to be published
 SM: conception and design, given final approval of the version to be published

Authors' information

BA: Associate Professor of Surgery at University "Federico II" of Naples, Italy.
 RC: Post-Graduate Doctorate in Vascular Surgery at University "Federico II" of Naples.
 FF, RR, TB: Resident in General Surgery Training Programme at University "Federico II" of Naples.
 MD, AA, GA: Aggregate Professor of Surgery at University "Federico II" of Naples, Italy.
 RS: Assistant Professor of Surgery at University Magna Graecia of Catanzaro,
 SM: Full Professor of Surgery at University "Federico II" of Naples, Italy.

Declarations

Funding of this supplement has come from University funds.
 This article has been published as part of *BMC Surgery* Volume 13 Supplement 2, 2013: Proceedings from the 26th National Congress of the Italian Society of Geriatric Surgery. The full contents of the supplement are available online at <http://www.biomedcentral.com/bmcscrg/supplements/13/S2>

Authors' details

¹Department of Clinical Medicine and Surgery University of Naples Federico II Via S. Pansini, 5 - 801311 Napoli, Italy. ²Department of Medical and Surgical Science -University Magna Graecia of Catanzaro - Viale Europa, Località Germaneto - 88100 Catanzaro, Italy.

Published: 8 October 2013

References

- Stephenson BM: **Complication of open groin hernia repairs.** *Surg Clin North Am* 2003, **83**:1255-1278.
- Collaboration EUHT: **Repair of groin hernia with synthetic mesh: meta-analysis of randomized controlled trials.** *Ann Surg* 2002, **235**:322-332.

3. Kingsnorth A, LeBlanc K: **Hernias: inguinal and incisional.** *Lancet* 2003, **362**:1561-1571.
4. Chow A, Purkayastha S, Athanasiou T, Tekkis P, Darzi A: **Inguinal Hernia.** *BMJ Clin Evid* 2007, **4**:1-20.
5. Turrentine FE, Wang H, Simpson VB, et al: **Surgical risk factors, morbidity, and mortality in elderly patients.** *J Am Coll Surg* 2006, **203**:865-877.
6. Allen PIM, Zager M, Goldman M: **Elective repair of groin hernias in the elderly.** *Br J Surg* 1987, **74**:987-991.
7. Nilsson H, Stylianidis G, Haapamaki M, Nilsson E, Nordin P: **Mortality after groin hernia surgery.** *Ann Surg* 2007, **245**:656-660.
8. Nienhuijs SW, Remijn EEG, Rosman C: **Hernia repair in elderly patients under unmonitored local anaesthesia is feasible.** *Hernia* 2005, **9**:218-222.
9. Amato B, Compagna R, et al: **Feasibility of inguinal hernioplasty under local anesthesia in elderly patients.** *BMC Surg* 2012, **12**(Suppl 1):S2.
10. Wagh PV, Leverich AP, Sun CN, White HJ: **Direct inguinal herniation in men: a disease of collagen.** *J Surg Res* 1974, **17**:425-433.
11. Rosenthal RA: **Small-bowel disorders and abdominal wall hernia in the elderly patient.** *Surg Clin North Am* 1994, **74**:261-291.
12. Robbins AW, Rutkow IM: **Mesh plug repair and groin hernia surgery.** *Surg Clin North Am* 1998, **78**:1007-1023.
13. Kurzer M, Belsham PA, Kark AE: **The Lichtenstein repair for groin hernias.** *Surg Clin North Am* 2003, **83**:1099-1117.
14. Simons MP, Aufenacker T, Bouillot JL, Campanelli G, et al: **European Hernia Society guidelines on the treatment of inguinal hernia in adult patient.** *Hernia* 2009, **13**:343-403.
15. Rispoli C, Rocco N, Iannone L, Amato B: **Developing guidelines in geriatric surgery:role of the grade system.** *BMC Geriatrics* 2009, **9**(Suppl 1):A99.
16. Amato B, Moja L, Panico S: **Shouldice technique versus other open techniques for inguinal hernia repair (Review).** *Cochrane database of systematic reviews (online)* 2012, **4**:CD001543.
17. Compagna R, Vigliotti G, Coretti G, Amato M, Aprea G, Puzziello A, Militello C, Amato B: **Comparative study between levobupivacaine and bupivacaine for hernia surgery in the elderly.** *BMC Surgery* 2012, **12**(Suppl 1):S12.
18. Rea R, Falco P, Izzo D, Leongito M, Amato B: **Laparoscopic ventral hernia repair with primary transperitoneal closure of the hernia defect.** *BMC Surgery* 2012, **12**(Suppl 1):S33.

doi:10.1186/1471-2482-13-S2-S28

Cite this article as: Amato et al: Day-surgery inguinal hernia repair in the elderly: single centre experience. *BMC Surgery* 2013 **13**(Suppl 2):S28.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

