


Contents lists available at [ScienceDirect](http://www.sciencedirect.com)

The Breast

journal homepage: www.elsevier.com/brst

Viewpoints and debate

Neoadjuvant treatment of HER2 and hormone-receptor positive breast cancer – Moving beyond pathological complete response

Serena Di Cosimo^{a,*}, Grazia Arpino^b, Daniele Generali^c^aFondazione IRCCS – Istituto Nazionale Tumori, Milano, Italy^bDipartimento di Medicina Clinica e Chirurgia, Università di Napoli Federico II, Italy^cU.O. Multidisciplinare di Patologia Mammaria/U.S. Terapia Molecolare e Farmacogenomica, AZ. Istituti Ospitalieri di Cremona, Italy

ARTICLE INFO

Article history:

Received 4 August 2013

Received in revised form

18 November 2013

Accepted 9 December 2013

Keywords:

Breast cancer
Neoadjuvant therapy
HER2
Hormone receptor
Targeted therapy
Trastuzumab

ABSTRACT

Pathologic complete response (pCR) was noted to be prognostic in all but hormone receptor-positive (HR) breast cancer cases even when HER2 is overexpressed. Evocative data suggest that HER2-positive breast cancer patients are a heterogeneous population and a subset of HER2-positive and HR-positive tumors biologically behave more like HER2-negative. Identification and targeted monitoring of these patients is crucial to consolidate data aiming to optimize combination treatment with new agents, thereby avoiding overtreatment with chemotherapy. The questions surrounding HER2-positive and HR-positive breast cancer patients treatment as well as the potential direction towards development of surrogate markers alternative to pCR are discussed.

© 2013 Elsevier Ltd. All rights reserved.

Introduction

Neoadjuvant trials are a new and reliable approach to testing new agents for breast cancer (BC). Aside from the clinical benefit derived by tumor downstaging, neoadjuvant therapy may provide in fact valuable prognostic information, considering that patients achieving pathologic complete response (pCR) show improved long-term outcome as compared to those left with residual disease after neoadjuvant systemic therapy. Based on this evidence, the FDA recently released specific guidelines to use pCR as a surrogate endpoint to support accelerated drug approval for BC neoadjuvant treatment [1]. To date, however, there has not been a uniform definition of pCR, which has made reporting and interpretation of data from neoadjuvant trials challenging. The FDA has proposed the following definition for regulatory purposes: pCR is defined as the absence of any residual invasive cancer on hematoxylin and eosin evaluation of the resected breast specimen and all sampled ipsilateral lymph nodes following completion of neoadjuvant systemic therapy. Adoption of a single term with a standard definition would facilitate discussion of proposed trial designs and interpretation of

clinical trial data to support accelerated approval. Importantly, according to FDA, the definition of pCR proposed in their guidance would have the greatest likelihood of predicting clinical benefit for regulatory purposes in patients with early-stage breast cancer who achieve pCR following neoadjuvant systemic therapy. However, pCR rates significantly vary depending on BC subtypes, ranging from 8.3% in hormone receptor-positive (HR-positive) to 31.1% in triple negative cases, with aggressive subtypes more highly associated with event-free survival than smaller, less aggressive tumors [2]. In patients with HER2 overexpressing (HER2-positive) BC, trastuzumab dramatically changed the disease's natural history and improved outcome [3,4]. However, pCR rates of neoadjuvant trastuzumab-containing regimes are still in the range of 30–60% with 3-year relapse-free survival (RFS) of 71–78% [5], clearly showing that a substantial number of HER2-positive BC patients undergoing surgery still present residual disease notwithstanding prior neoadjuvant therapy.

Significance of pCR in HER2-positive and HR-positive tumors

Growing evidence indicates that response to anti-HER2 agents and the prognostic impact of pCR after anti-HER2-based therapy depend on HR status. In the retrospective study by Guarneri et al., pCR rates among patients with HER2-positive BC treated with anthracycline/taxane-based chemotherapy was 15% and 29%

* Corresponding author. Department of Oncology, Fondazione IRCCS – Istituto Nazionale Tumori, Via G. Venezian 1, 20100 Milano, Italy. Tel.: +39 02 2390 30 66; fax: +39 02 2390 21 49.

E-mail address: serena.dicosimo@istitutotumori.mi.it (S. Di Cosimo).

($p < 0.001$) in HR-positive and HR-negative cases, respectively [6]. The addition of trastuzumab produced similar differences according to HR status, with pCR rates 1.5–2 fold lower in HR-positive than in HR-negative cases [6]. More recently, data from the NeoSphere [7], NeoALTT0 [8] and TBCRC006 [9] trials further confirmed these findings, consistently showing that pCR rates were significantly lower in HR-positive than HR-negative tumors, regardless the type of HER2-targeted therapy with single or combined HER2 blockade (Table 1) [7–19]. Therefore, new neoadjuvant strategies are urgently needed to increase pCR rates and to improve clinical outcome of HER2-positive/HR-positive BC patients.

Nevertheless, the significance of pCR in this subset of patients needs further considerations. The pooled analysis of the German neoadjuvant studies reported that pCR is a suitable surrogate endpoint for HER2-positive/HR-negative but not for HER2-positive/HR-positive BC patients [20]. Indeed, pCR in HER2-positive/HR-positive cases was relatively uncommon (11%–22%), even with the use of trastuzumab, and achievement of pCR in this subgroup was not prognostic for survival. Although the reasons for these findings are not entirely clear, a possible explanation is that lack of pCR to chemotherapy in less proliferative HR-positive tumors predicts greater benefit to future endocrine therapy. Thus patients with HER2-positive/HR-positive BC not achieving a pCR are not necessarily at poor prognosis, as adjuvant endocrine therapy may further improve clinical outcome. It is important to note, however, that among patients with pCR, those initially diagnosed with HER2-positive/HR-positive BC tend to report the lowest 5-year disease-free survival (DFS) as if, when present, pCR in HER2-positive/HR-positive cases is less beneficial. Although no analysis on biological

and clinical characteristics of these patients relapsing after pCR has been done, it is likely that the poorer outcome in this subgroup is due to either decreased benefit from adjuvant therapy or rapid onset of molecular resistance to anti-HER2 agents and/or endocrine therapy. In support of this hypothesis, evidence exists that relapses after pCR generally occur in young patients and/or in patients with locally advanced BC at presentation [21]. Therefore, in patients with HER2-positive/HR-positive, pCR is infrequent and when reported it does not yield the same good outcome as compared to the rest of patients treated with trastuzumab-based neoadjuvant regimens. Data from adjuvant studies further confirm these findings [22]. In the HERA trial, the hazard ratio for DFS among patients with HER2-positive/HR-positive early BC had the widest confidence intervals as compared with the other subtypes. Although the limitations of the *post-hoc* analysis permit no definitive conclusion on the differential effect of adjuvant trastuzumab according to BC subtypes, these data demonstrate that HER2-positive BC patients are a heterogeneous population. The identification and characterization of the HER2-positive/HR-positive BC subset is essential to avoid overtreatment especially in patients with small tumors, who may benefit from endocrine therapy alone or combined with a HER2 targeted therapy without chemotherapy. Indeed, a crosstalk between the HR and HER2 pathways exists, which plays a major role in the development of both intrinsic and acquired resistance to endocrine agents. Emerging data suggests that this crosstalk is bidirectional and is involved in resistance to HER2-directed agents [23]. Therefore, targeting both HR and HER2 pathways may be beneficial in overcoming resistance to endocrine and anti-HER therapies and optimize clinical outcome.

Table 1

Pathological complete response (pCR)^a rate according to hormone-receptor (HR) status in neoadjuvant trials with anti-HER2 agents.

Study [Ref] phase	No. of patients	Clinical stage	Neoadjuvant regimen	pCR rate	<i>p</i>	pCR rate HR-positive	pCR rate HR-negative		
Randomized trials with trastuzumab	MD Anderson [10] Phase III	42	II–III	P → FEC P + T → FEC + T	26% 65%	0.016	27.2% 61.5%	25% 70%	
	NOAH [11] Phase III	235	III	AP → P → CMF AP + T → P + T → CMF + T	19% 38%	0.001	17% 18%	22% 48%	
	ABCSG-24 [12] Phase III	90	II–III	EC ± Cap EC ± Cap + T	26% (breast only) 40% (breast only)	0.369	NR	NR	
	REMAGUS 02 [13] Phase II	120	II–III	EC → D EC → D + T	19% 26%	NR	20.5% 20.5%	19% 32%	
	H2269s [14] Phase II	30	II–III	D + CBDCA D + CBDCA + T	7.1% (breast only) 40% (breast only)	0.0801	NR	NR	
	Randomized trials comparing HER2-targeting approaches	NeoALTT0 [8] Phase III	455	II–III	Weekly P + T Weekly P + L Weekly P + T + L	28% 20% 47%	0.0007 ^b	22.7% 16.1% 41.6%	36.5% 33.7% 61.3%
		NeoSphere [7] Phase III	417	II–III	D + T D + Pert D + Pert + T Pert + T	23% 17% 42% 12%	0.0141 ^b	20% 17.4% 26% 5.9%	36.8% 30% 63.2% 27.3%
		NSABP B-41 [15] Phase III	519	II–III	AC → weekly P + T AC → weekly P + L AC → weekly P + T + L	49.4% 47.4% 60.2%	0.056 ^b	46.7% 48% 55.6%	65.5% 60.6% 73%
		GeparQuinto [16] Phase III	615	II–III	EC + T → D + T EC + L → D + L	44% 30%	0.04	NR	NR
		CHERLOB [17] Phase II	121	II–III	P + T → FEC + T P + L → FEC + L P + T + L → FEC + T + L	25% 26.3% 46.7%	0.019 ^b	25% 22.7% 35.7%	26.6% 35.7% 56.2%
GEICAM 2006–14 [18] Phase II		99	II–III	EC → D + T EC → D + L	48% 24%	0.01	NR	NR	
Holmes et al. [19] Phase II		100	II–III	FEC → weekly P + T FEC → weekly P + L FEC → weekly P + T + L	54% 45% 74%	NR	NR	NR	

A = doxorubicin; Cap = capecitabine; CBDCA = carboplatin; CMF = cyclophosphamide, methotrexate, 5-fluorouracil; D = docetaxel; E = epirubicin; FEC = 5-fluorouracil, epirubicin, and cyclophosphamide; L = lapatinib; NR = not reported; P = paclitaxel; Pert = pertuzumab; T = trastuzumab.

^a pCR = defined as no invasive tumor in the breast and axilla, if not otherwise specified.

^b Comparison between dual HER2-targeted therapy + chemotherapy versus trastuzumab + chemotherapy.

Table 2
Summary of studies assessing biomarkers in HER2 positive breast cancer.

Biological markers	Type of study [Ref]	Patient population (n)	Results	Conclusion/remarks	
Proteins	Basal p95/HER2	Retrospective [37]	MBC (93)	Decreased PFS (HR 1.9; $p = 0.017$) and OS (HR 2.2; $p = 0.012$) as compared with p95/HER2 negative	p95/HER2 is a marker of poor prognosis (TBC in RCT)
		Analysis from RCT [38,17]	NeoAdj (153) NeoAdj (88)	pCR 58.2% vs 32.6% in p95/HER2-negative vs positive pts (0.009) in the GeparQuattro study No difference in the CHERLOB study	Inconclusive data on p95/HER2 predictive value
	Basal Ki67	Analysis from RCT [39]	NeoAdj (1166)	Ki67 significantly linked to prognosis in uni- and multivariate analysis Ki67 $\leq 15\%$ vs 15.1%–35% vs $> 35\%$ had pCR% of 4.2%, 12.8%, and 29.0% ($p < 0.0005$)	Ki67 is associated with poor prognosis and increased pCR (TBC in RCT)
	Post-therapy Ki67	Retrospective [28,40]	NeoAdj (102) NeoAdj (64)	High Ki67 expression correlated with poor DFS and OS Multivariate analysis showed Ki67 significantly ($p = 0.05$) associated with prognosis	Prognostic value of post treatment Ki67 to be further investigated
	Basal and post-therapy TOPOII α	Retrospective [41]	NeoAdj (283)	Decreased DFS ($p = 0.010$) and OS ($p < 0.001$) in TOPOII α cases Significant prognostic value of post-treatment TOPOII α ($p = 0.002$)	Prognostic value of TOPOII α to be further investigated
Immunological-related	Loss of PTEN and p110 expression	Retrospective [42]	MBC, EBC (155)	Decreased PFS correlated with basal expression of p110 α ($p = 0.024$)	Prognostic value of PI3K activation to be further investigated
		Retrospective [34]	MBC, EBC (54)	158 V/V genotype correlated with ORR (HR, 8.7; $p = 0.02$) and PFS (HR 5.3; $p = 0.005$)	FcYR-mediated ADCC to be further investigated
	Lymphocyte Infiltration	Retrospective [32] Retrospective [43]	NeoAdj [25] EBC (1250)	Higher infiltration of CD8+ T cells, FOXP3(+) regulatory T cells (Tregs) and CD8(+) cytotoxic T lymphocytes as compared to HER2 negative cases ($p < 0.005$)	Value of reduced tumor-related immune suppression to be further investigated
Geneprofile	Blueprint PAM50	Retrospective [31] Retrospective [30]	NeoAdj (996) NeoAdj (437)	Increased predictive accuracy for pCR by adding the immune module to clinicopathologic characteristics (IDI, 0.093; $p = 0.004$) 40% of HER2-positive cases are luminal-type by BluePrint and present low response rates to therapy	Immune module scores and molecular subtyping improve estimation of pCR (to be further investigated)
		Prospective [44]	MBC [19]	High expression of 17q12-21 amplicon genes <i>HER2</i> and <i>GRB7</i> correlated with response to treatment ($p = 0.0015$)	PAM50-enriched profile associated with clinical benefit (to be further investigated)
Functional pathway	<i>PIK3CA</i>	Retrospective [36]	NeoAdj [46]	Alterations of PI3K genes are more frequent in HER2 positive cases without HER2 amplification ($p = 0.009$)	PTEN loss and <i>PIK3CA</i> mutations confer worse outcome and resistance to trastuzumab (TBC in RCT)
		Retrospective [45] Retrospective [46]	NeoAdj, EBC, MBC (1889) NeoAdj (80)	PTEN loss correlated with poorer efficacy of trastuzumab (RR = 0.682, 95% CI: 0.550–0.846, $p = 0.000$) <i>PIK3CA</i> mutations correlated to decreased DFS ($p = 0.0013$)	
Metabolic profile	¹⁸ FDG PET	Analysis from RCT [47]	NeoAdj (86)	pCR rates twice as high for ¹⁸ FDG PET responders than non-responders (44% vs 19%, $p = 0.05$)	Metabolic responders have increased likelihood of pCR (to be further investigated)
Soluble	Circulating Tumor Cells	Prospective [48]	EBC (88)	CTCs are related to reduced DFS ($p < 0.023$) and OS ($p < 0.04$)	CTCs predict poor clinical outcome (to be further investigated)
		Retrospective [49]	MBC (53)	CTC count is a significant predictive factor for OS ($p < 0.01$)	
	Circulating Endothelial Cells	Retrospective [50]	NeoAdj (53)	Low baseline CEC associated with pCR ($p = 0.046$)	Value of CEC to be assessed in extensive studies
		Retrospective [51]	MBC [52]	Clinical responses correlated with low CEC ($p = 0.023$)	
	HER2	Review [52]	MBC, EBC	No clear correlation with response to treatment or OS	Prospective trials to investigate the clinical use of HER2 ECD test are needed

DFS = Disease Free Survival; EBC = early breast cancer; ECD = extracellular domain; ¹⁸FDG PET = 18-Fluoro-deoxyglucose positron emission tomography; HR = hazard ratio; Integrated discrimination index (IDI); MBC = metastatic breast cancer; NeoAdj = neoadjuvant; ORR = overall response rate; OS = Overall Survival; pCR = pathological complete response, defined as no invasive tumor in the breast and axilla, if not otherwise specified; PFS = Progression free survival; RCT = randomized controlled trial; RR = relative risk; TBC = to be confirmed; vs = versus.

Searching for new surrogate markers to predict long-term clinical outcome from neoadjuvant therapies

Understanding which patients are eligible for new treatments is critical, especially in the neoadjuvant setting, where the lack of prognostic value of pCR makes the identification of alternative markers a priority for cancer research. The proliferation index Ki67 is a feasible marker widely studied as surrogate of disease relapse in patients undergoing neoadjuvant hormonal therapy. Results from a randomized trial comparing tamoxifen versus anastrozole, showed that differences in Ki67 suppression after 2 and 12 weeks of treatment mirrored those in RFS in companion adjuvant trials [24]. Further evidence for the importance of Ki67 suppression were provided by higher 2-week values of Ki67 predicting a significantly worse RFS [25]. However as HER2 positive tumors show poor reduction in Ki67 upon treatment with aromatase inhibitors, this marker may be of limited utility when used in the setting of neoadjuvant hormonal therapy [26]. Nevertheless, in BC patients receiving neoadjuvant chemotherapy, basal Ki67 values predict treatment response and prognosis, as the higher the basal Ki67 the higher the pCR rate and RFS [27]; and, even more importantly, Ki67 values after treatment predict prognosis in patients not achieving a pCR [28]. The use of Ki67 as a guide for neoadjuvant treatment has been the focus of the Z1031B study [29]. All patients enrolled in this study were initially treated with hormonal therapy during 2–4 weeks. A tumor biopsy was performed thereafter and, according to the Ki67 results, patients who had tumors with a high proliferation rate were recommended to switch to chemotherapy/surgery. The study reported a pCR rate of 6% among those patients who switched to neoadjuvant chemotherapy, which did not meet the study's predefined criteria for adequate activity. Nevertheless, the value of Ki67 in HER2-positive/HR-positive BC continues to be investigated in different trials with anti-HER2 targeted agents ± hormonal therapy, i.e. NeoPhoebe, PAMELA, PERELISA, with the ultimate goal of finding a valid strategy for patients unlikely to benefit from chemotherapy. In addition, clinical trials have already started to evaluate the utility of using molecular profiles for breast cancer management in the preoperative setting. Among these, the I-SPY I study tested the MammaPrint and the Blueprint 80-gene expression profiles to stratify response to neoadjuvant therapy according to molecular subtypes. In detail, the pCR rate was 5% in the luminal-A and 10% in luminal-B, 39% in HER2-type and 33% in the basal-like subtypes. These data support the use of MammaPrint and Blueprint to define the clinical correlation between molecular subtyping and treatment outcomes [30]. Furthermore, the pooled analysis, focusing on the associations between pCR and gene modules to discover biologically relevant and potentially druggable subtype-specific oncogenic pathways, showed a strong link between immune modules and pCR in HER2-positive tumors [31]. Both HER2 and estrogens have immunomodulatory effects on BC development by increasing intratumoral Tregs levels [32] and immunomodulation is a critical factor for trastuzumab antitumoral activity. Several studies have shown that in BC patients treated with trastuzumab-based therapy [33] the antibody-dependent cell-mediated cytotoxicity (ADCC) and the FcγR polymorphisms can reliably predict clinical response [34]. Enhancing antitumor immunity may therefore increase therapy efficacy in HER2-positive/HR-positive BC, a strategy that deserves further investigations in clinical trials. Finally, sequencing analysis in BC reported that dysregulation of the PI3K pathway is associated with poor response to HER2-targeted therapy [35]. Furthermore, alterations in the PI3K and HR signaling are prognostic for patient outcome in the HR-positive BC subset [36]. The crosstalk between PI3K and HR pathways is involved in resistance to chemotherapy with trastuzumab, thus leading to the conclusion that HR status influenced the prognostic significance of genomic alterations in

HER2-positive tumors. If these data will be further confirmed, mutation analysis of PI3K status should become part of the routine biomarkers panel as it may be helpful in selecting the proper treatment of patients with HER2-positive/HR-positive tumors. Studies assessing biomarkers in HER2 positive breast cancer are summarized in Table 2 [17,28,30–32,34,36–52].

Conclusions

The addition of trastuzumab to neoadjuvant chemotherapy regimen significantly increases the pCR rates in all HER2-positive tumors. However, the benefit of trastuzumab is highest in HR-negative tumors and progressively decreases with increase in tumor HR expression. This information prompts the search of new strategies and/or new combination of therapies associated to an increased pCR but also novel surrogate markers of long-term outcome. Studies using surrogate markers as Ki67 and gene amplification and/or mutation are currently ongoing and promise to assist not only the choice of best neoadjuvant therapeutic approach but also to address the clinician in scheduling the proper treatment in the adjuvant setting in the next future.

Conflict of interest

Serena Di Cosimo is member of the speakers' bureau for GlaxoSmithKline S.p.A. and is a Consultant for TEVA pharmaceuticals. Grazia Arpino is member of the speakers' bureau for Roche S.p.A. and GlaxoSmithKline S.p.A. Daniele Generali has no conflict of interest to declare.

References

- [1] Guidance for industry pathologic complete response in neoadjuvant treatment of high-risk early-stage breast cancer: use as an endpoint to support accelerated approval <http://www.fda.gov/Drugs/GuidanceComplianceRegulatoryInformation/Guidances/default.htm>.
- [2] Houssami N, Macaskill P, von Minckwitz G, Marinovich ML, Mamounas E. Meta-analysis of the association of breast cancer subtype and pathologic complete response to neoadjuvant chemotherapy. *Eur J Cancer* 2012;48(18):3342–54.
- [3] Yin W, Jiang Y, Shen Z, Shao Z, Lu J. Trastuzumab in the adjuvant treatment of HER2-positive early breast cancer patients: a meta-analysis of published randomized controlled trials. *Plos One* 2011;6(6):e21030.
- [4] De Mattos-Arruda L, Cortes J. Advances in first-line treatment for patients with HER-2+ metastatic breast cancer. *Oncologist* 2012;17(5):631–44.
- [5] Valachis A, Mauri D, Polyzos NP, Chlouverakis G, Mavroudis D, Georgoulas V. Trastuzumab combined to neoadjuvant chemotherapy in patients with HER2-positive breast cancer: a systematic review and meta-analysis. *Breast* 2011;20(6):485–90.
- [6] Guarneri V, Broglio K, Kau SW, Cristofanilli M, Buzdar AU, Valero V, et al. Prognostic value of pathologic complete response after primary chemotherapy in relation to hormone receptor status and other factors. *J Clin Oncol* 2006;24(7):1037–44.
- [7] Gianni L, Pienkowski T, Im YH, Roman L, Tseng LM, Liu MC, et al. Efficacy and safety of neoadjuvant pertuzumab and trastuzumab in women with locally advanced, inflammatory, or early HER2-positive breast cancer (NeoSphere): a randomised multicentre, open-label, phase 2 trial. *Lancet Oncol* 2012;13:25–32.
- [8] Baselga J, Bradbury I, Eidtmann H, Di Cosimo S, de Azambuja E, Aura C, et al. Lapatinib with trastuzumab for HER2-positive early breast cancer (Neo-ALTTO): a randomised, open-label, multicentre, phase 3 trial. *Lancet* 2012;379:633–40.
- [9] Rimawi MF, Mayer IA, Forero A, Nanda R, Goetz MP, Rodríguez AA, et al. Multicenter phase II study of neoadjuvant lapatinib and trastuzumab with hormonal therapy and without chemotherapy in patients with human epidermal growth factor receptor 2-overexpressing breast cancer: TBCRC 006. *J Clin Oncol* 2013;31(14):1726–1733.
- [10] Buzdar AU, Ibrahim NK, Francis D, Booser DJ, Thomas ES, Theriault RL, et al. Significantly higher pathologic complete remission rate after neoadjuvant therapy with trastuzumab, paclitaxel, and epirubicin chemotherapy: results of a randomized trial in human epidermal growth factor receptor 2-positive operable breast cancer. *J Clin Oncol* 2005;23:3676–85.
- [11] Gianni L, Eiermann W, Semiglazov V, Manikhas A, Lluch A, Tjulandin S, et al. Neoadjuvant chemotherapy with trastuzumab followed by adjuvant trastuzumab versus neoadjuvant chemotherapy alone, in patients with HER2-

- positive locally advanced breast cancer (the NOAH trial): a randomised controlled superiority trial with a parallel HER2-negative cohort. *Lancet* 2010;375:377–84.
- [12] Steger G, Greil R, Jakesz R, Lang A, Mlineritsch B, Melbinger-Zeinitzer E, et al. Final results of ABCSG-24, a randomized phase III study comparing epirubicin, docetaxel, and capecitabine (EDC) to epirubicin and docetaxel (ED) as neoadjuvant treatment for early breast cancer and comparing ED/EDC plus trastuzumab (T) to ED/EDC as neoadjuvant treatment for early HER-2 positive breast cancer. *Cancer Res* 2009;69(S24). S64s, Abstract 1081.
 - [13] Pierga JY, Delaloge S, Espié M, Brain E, Sigal-Zafrani B, Mathieu MC, et al. A multicenter randomized phase II study of sequential epirubicin/cyclophosphamide followed by docetaxel with or without celecoxib or trastuzumab according to HER2 status, as primary chemotherapy for localized invasive breast cancer patients. *Breast Cancer Res Treat* 2010;122:429–37.
 - [14] Chang HR, Glaspy J, Allison MA, Kass FC, Elashoff R, Chung DU, et al. Differential response of triple-negative breast cancer to a docetaxel and carboplatin-based neoadjuvant treatment. *Cancer* 2010;116:4227–37.
 - [15] Robidoux A, Tang G, Rastogi P, Geyer Jr CE, Azar CA, Atkins JN, et al. Lapatinib as a component of neoadjuvant therapy for HER2-positive operable breast cancer (NSABP protocol B-41): an open-label, randomised phase 3 trial. *Lancet Oncol* 2013;14:1183–92.
 - [16] Untch M, Loibl S, Bischoff J, Eidtmann H, Kaufmann M, Blohmer JU, et al. Lapatinib versus trastuzumab in combination with neoadjuvant anthracycline-taxane-based chemotherapy (GeparQuinto, GBG 44): a randomised phase 3 trial. *Lancet Oncol* 2012;13:135–44.
 - [17] Guarneri V, Frassoldati A, Bottini A, Cagossi K, Bisagni G, Sarti S, et al. Preoperative chemotherapy plus trastuzumab, lapatinib, or both in human epidermal growth factor receptor 2-positive operable breast cancer: results of the randomized phase II CHER-LOB study. *J Clin Oncol* 2012;30:1989–95.
 - [18] Alba E, Albanell J, de la Haba J, Barnadas A, Calvo L, Sanchez P, et al. PD07-04: lapatinib vs trastuzumab in combination with standard EC-D chemotherapy in the neoadjuvant treatment of HER2+ patients. Results from the GEICAM 2006–14 phase II randomized trial. *Cancer Res* December 15, 2011;71(24). Supplement 3.
 - [19] Holmes FA, Nagarwala YM, Espina VA, Liotta LA, Danso MA, Gallagher RI, et al. Correlation of molecular effects and pathologic complete response to preoperative lapatinib and trastuzumab, separately and combined prior to neoadjuvant breast cancer chemotherapy. *J Clin Oncol* 2011;29(S15). 46s, Abstract 506.
 - [20] von Minckwitz G, Untch M, Blohmer JU, Costa SD, Eidtmann H, Fasching PA, et al. Definition and impact of pathologic complete response on prognosis after neoadjuvant chemotherapy in various intrinsic breast cancer subtypes. *J Clin Oncol* 2012;30(15):1796–804.
 - [21] Ju NR, Jeffe DB, Keune J, aft R. Patient and tumor characteristics associated with breast cancer recurrence after complete pathological response to neoadjuvant chemotherapy. *Breast Cancer Res Treat* 2013;137(1):195–201.
 - [22] Untch M, Gelber RD, Jackisch C, Procter M, Baselga J, Bell R, et al. Estimating the magnitude of trastuzumab effects within patient subgroups in the HERA trial. *Ann Oncol* 2008 Jun;19(6):1090–6.
 - [23] Arpino G, Gutierrez C, Weiss H, Rimawi M, Massarweh S, Bharwani L, et al. Treatment of human epidermal growth factor receptor 2-overexpressing breast cancer xenografts with multiagent HER-targeted therapy. *J Natl Cancer Inst* 2007;99:694–705.
 - [24] Dowsett M, Smith IE, Ebbs SR, Dixon JM, Skene A, A'Hern R, et al. Prognostic value of Ki67 expression after short-term presurgical endocrine therapy for primary breast cancer. *J Natl Cancer Inst* 2007 Jan 17;99(2):167–70.
 - [25] Ellis MJ, Tao Y, Luo J, A'Hern R, Evans DB, Bhatnagar AS, et al. Outcome prediction for estrogen receptor-positive breast cancer based on postneoadjuvant endocrine therapy tumor characteristics. *J Natl Cancer Inst* 2008;100(19):1380–8.
 - [26] Dunbier AK, Anderson H, Ghazoui Z, Folkard EJ, A'hern R, Crowder RJ, et al. Relationship between plasma estradiol levels and estrogen-responsive gene expression in estrogen receptor-positive breast cancer in postmenopausal women. *J Clin Oncol* 2010 Mar 1;28(7):1161–7.
 - [27] Fasching PA, Heusinger K, Haerberle L, Niklos M, Hein A, Bayer CM, et al. Ki67, chemotherapy response, and prognosis in breast cancer patients receiving neoadjuvant treatment. *BMC Cancer* 2011;11:486.
 - [28] Tanei T, Shimomura A, Shimazu K, Nakayama T, Kim SJ, Iwamoto T, et al. Prognostic significance of Ki67 index after neoadjuvant chemotherapy in breast cancer. *Eur J Surg Oncol* 2011;37:155–61.
 - [29] Ellis MJ, Suman V, McCall L, Luo R, Hoog J, Brink A, et al. Z1031B neoadjuvant aromatase inhibitor trial: a phase 2 study of triage to chemotherapy based on 2 to 4 week Ki67 level >10%. *Cancer Res* 2012;72(Suppl. 24). Abstract nr PD07-01.
 - [30] Glück S, de Snoo F, Peeters J, Stork-Sloots L, Somlo G, et al. Molecular subtyping of early-stage breast cancer identifies a group of patients who do not benefit from neoadjuvant chemotherapy. *Breast Cancer Res Treat* 2013 Jun;139(3):759–67.
 - [31] Ignatiadis M, Singhal SK, Desmedt C, Haibe-Kains B, Criscitiello C, Andre F, et al. Gene modules and response to neoadjuvant chemotherapy in breast cancer subtypes: a pooled analysis. *J Clin Oncol* 2012;30(16):1996–2004.
 - [32] Muraro E, Martorelli D, Turchet E, Miolo G, Scalone S, Comaro E, et al. A different immunologic profile characterizes patients with HER-2-overexpressing and HER-2-negative locally advanced breast cancer: implications for immune-based therapies. *Breast Cancer Res* 2011;13(6). R117.
 - [33] Tagliabue E, Campiglio M, Pupa SM, Ménard S, Balsari A. Activity and resistance of trastuzumab according to different clinical settings. *Cancer Treat Rev* 2012 May;38(3):212–7.
 - [34] Musolino A, Naldi N, Bortesi B, Pezzuolo D, Capelletti M, Missale G, et al. Immunoglobulin G fragment C receptor polymorphisms and clinical efficacy of trastuzumab-based therapy in patients with HER-2/neu-positive metastatic breast cancer. *J Clin Oncol* 2008;26(11):1789–96.
 - [35] Jensen JD, Knoop A, Laenkhölm AV, Grauslund M, Jensen MB, Santoni-Rugiu E, et al. PIK3CA mutations, PTEN, and pHER2 expression and impact on outcome in HER2-positive early-stage breast cancer patients treated with adjuvant chemotherapy and trastuzumab. *Ann Oncol* 2000;23:2034–42.
 - [36] Takada M, Higuchi T, Tozuka K, Takei H, Haruta M, Watanabe J, et al. Alterations of the genes involved in the PI3K and estrogen-receptor pathways influence outcome in human epidermal growth factor receptor 2-positive and hormone receptor-positive breast cancer patients treated with trastuzumab-containing neoadjuvant chemotherapy. *BMC Cancer* 2013;13:241.
 - [37] Sperinde J, Jin X, Banerjee J, Penuel E, Saha A, Diedrich G, et al. Quantitation of p95HER2 in paraffin sections by using a p95-specific antibody and correlation with outcome in a cohort of trastuzumab-treated breast cancer patients. *Clin Cancer Res* 2010 Aug 15;16(16):4226–35.
 - [38] Loibl S, Bruey J, von Minckwitz G, Huober JB, Press MF, Darb-Esfahani S, et al. Validation of p95 as a predictive marker for trastuzumab-based therapy in primary Her2-positive breast cancer: a translational investigation from the neoadjuvant GeparQuattro study. *J Clin Oncol* 2011;29(suppl; abstr 530).
 - [39] Denkert C, Loibl S, Müller BM, Eidtmann H, Schmitt WD, Eiermann W, et al. Ki67 levels as predictive and prognostic parameters in pretherapeutic breast cancer core biopsies: a translational investigation in the neoadjuvant Gepar-Trio trial. *Ann Oncol* 2013 Nov;24(11):2786–93.
 - [40] Yoshioka T, Hosoda M, Yamamoto M, Taguchi K, Hatanaka KC, Takakuwa E, et al. Prognostic significance of pathologic complete response and Ki67 expression after neoadjuvant chemotherapy in breast cancer. *Breast Cancer* 2013 May 5.
 - [41] Chen S, Huang L, Liu Y, Chen CM, Wu J, Shao ZM. The predictive and prognostic significance of pre- and post-treatment topoisomerase II alpha in anthracycline-based neoadjuvant chemotherapy for local advanced breast cancer. *Eur J Surg Oncol* 2013;39(6):619–26.
 - [42] Gallardo A, Lerma E, Escuin D, Tibau A, Muñoz J, Ojeda B, et al. Increased signalling of EGFR and IGF1R, and deregulation of PTEN/PI3K/Akt pathway are related with trastuzumab resistance in HER2 breast carcinomas. *Br J Cancer* 2012;106(8):1367–73.
 - [43] Liu F, Lang R, Zhao J, Zhang X, Pringle GA, Fan Y, et al. CD8(+) cytotoxic T cell and FOXP3(+) regulatory T cell infiltration in relation to breast cancer survival and molecular subtypes. *Breast Cancer Res Treat* 2011;130(2):645–55.
 - [44] Montemurro F, Prat A, Rossi V, Valabrega G, Sperinde J, Peraldo-Neia C, et al. Potential biomarkers of long-term benefit from single-agent trastuzumab or lapatinib in HER2-positive metastatic breast cancer. *Mol Oncol* 2013.
 - [45] Wang CL, MacDonald LR, Rogers JV, Aravkin A, Haseley DR, Beatty JD. Positron emission mammography: correlation of estrogen receptor, progesterone receptor, and human epidermal growth factor receptor 2 status and 18F-FDG. *AJR Am J Roentgenol* 2011;197(2):W247–55.
 - [46] Cizkova M, Dujaric ME, Lehmann-Che J, Scott V, Tembo O, Asselain B, et al. Outcome impact of PIK3CA mutations in HER2-positive breast cancer patients treated with trastuzumab. *Br J Cancer* 2013;108(9):1807–9.
 - [47] Gebhart G, Gámez C, Holmes E, Robles J, Garcia C, Cortés M, et al. 18F-FDG PET/CT for early prediction of response to neoadjuvant lapatinib, trastuzumab, and their combination in HER2-positive breast cancer: results from Neo-ALTO. *J Nucl Med* 2013 Nov, 1862–1868;54(11).
 - [48] Ignatiadis M, Xenidis N, Perraki M, Apostolaki S, Politaki E, Kafousi M, et al. Different prognostic value of cytokeratin-19 mRNA positive circulating tumor cells according to estrogen receptor and HER2 status in early-stage breast cancer. *J Clin Oncol* 2007;25(33):5194–202.
 - [49] Munzone E, Botteri E, Sandri MT, Esposito A, Adamoli L, Zorzino L, et al. Prognostic value of circulating tumor cells according to immunohistochemically defined molecular subtypes in advanced breast cancer. *Clin Breast Cancer* 2012;12(5):340–6.
 - [50] Ali AM, Ueno T, Tanaka S, Takada M, Ishiguro H, Abdellah AZ, et al. Determining circulating endothelial cells using CellSearch system during preoperative systemic chemotherapy in breast cancer patients. *Eur J Cancer* 2011;47(15):2265–72.
 - [51] Rugo HS, Chien AJ, Franco SX, Stopeck AT, Glencer A, Lahiri S, et al. A phase II study of lapatinib and bevacizumab as treatment for HER2-overexpressing metastatic breast cancer. *Breast Cancer Res Treat* 2012;134(1):13–20.
 - [52] Leyland-Jones B, Smith BR. Serum HER2 testing in patients with HER2-positive breast cancer: the death knell tolls. *Lancet Oncol* 2011;12(3):286–95.