

[Oncotarget](#). 2015 Jun 20;6(17):15628-38.

# Ccdc6 knock-in mice develop thyroid hyperplasia associated to an enhanced CREB1 activity.

[Leone V](#)<sup>1</sup>, [Langella C](#)<sup>1</sup>, [Esposito F](#)<sup>1</sup>, [Arra C](#)<sup>2</sup>, [Palma G](#)<sup>2</sup>, [Rea D](#)<sup>2</sup>, [Paciello O](#)<sup>3</sup>, [Merolla F](#)<sup>1</sup>, [De Biase D](#)<sup>3</sup>, [Papparella S](#)<sup>3</sup>, [Celetti A](#)<sup>1</sup>, [Fusco A](#)<sup>1,4</sup>.

## [Author information](#)

### Abstract

CCDC6 was originally identified upon rearrangement with RET in human thyroid papillary carcinomas generating the RET/PTC1 oncogene. We have previously reported that CCDC6 interacts with CREB1 and represses its transcriptional activity. Since the function of at least one allele of CCDC6 is lost following RET/PTC1 rearrangements, we aimed at the generation of mice, carrying a CCDC6 mutant gene. Previous studies suggested that the coiled-coil domain of CCDC6, mainly encoded by human exon 2, is required for the protein function. Therefore, we engineered a murine Ccdc6 construct, carrying a deletion of the exon 2, that was able to exert only a mild repression on CREB1 transcriptional activity, with respect to the wild type Ccdc6. Subsequently, we generated Ccdc6-ex2 knock-in mice. These mice developed thyroid hyperplasia associated with an enhanced CREB1 activity and an increased expression of the CREB-1 regulated genes. These results strongly support a CCDC6 promoting role, ascribed to its functional impairment, in the development of thyroid papillary carcinomas harboring the RET/PTC1 oncogene.

### KEYWORDS:

CREB1; Ccdc6; knock-in mice; thyroid