

This is a provisional PDF only. Copyedited and fully formatted version will be made available soon.



P O L I S H G Y N E C O L O G Y

# GINEKOLOGIA

## POLSKA

ORGAN POLSKIEGO TOWARZYSTWA GINEKOLOGICZNEGO  
THE OFFICIAL JOURNAL OF THE POLISH GYNECOLOGICAL SOCIETY

ISSN: 0017-0011

e-ISSN: 2543-6767

## Sonographic imaging and differential diagnosis of fetal perineal masses

**Authors:** Sharon Perlman, Katarzyna Zych-Krekora, Michal Krekora, Ron Bardin, Yinon Gilboa

**DOI:** 10.5603/GP.a2022.0157

**Article type:** Research paper

**Submitted:** 2022-11-09

**Accepted:** 2022-11-09

**Published online:** 2022-12-19

This article has been peer reviewed and published immediately upon acceptance. It is an open access article, which means that it can be downloaded, printed, and distributed freely, provided the work is properly cited.

Articles in "Ginekologia Polska" are listed in PubMed.

**Sonographic imaging and differential diagnosis of fetal perineal masses**

Sharon Perlman<sup>1, 2\*</sup>, Katarzyna Zych-Krekora<sup>3\*</sup>, Michal Krekora<sup>4</sup>, Ron Bardin<sup>1, 2</sup>, Yinon Gilboa<sup>1, 2, 5</sup>

\*These authors contributed equally to this work

<sup>1</sup>*Ultrasound Unit, The Helen Schneider Women's Hospital, Rabin Medical Center, Beilinson Campus, Petach Tikva, Israel*

<sup>2</sup>*Sackler School of Medicine, Tel Aviv University, Tel Aviv, Israel*

<sup>3</sup>*Polish Mother's Memorial Hospital Research Institute in Lodz, Department of Perinatology, Obstetrics and Gynaecology, Lodz, Poland*

<sup>4</sup>*Polish Mother's Memorial Hospital Research Institute in Lodz, Department of Obstetrics and Gynaecology, Lodz, Poland*

<sup>5</sup>*Ultrasound Unit, Department of Obstetrics and Gynecology, The Chaim Sheba Medical Center, Tel-HaShomer, Israel*

**Corresponding author:**

Katarzyna Zych-Krekora

Polish Mother's Memorial Hospital Research Institute, Perinatology,  
Obstetrics and Gynecology Department, 281/289 Rzgowska St, 93-338  
Lodz, tel. +48 783134859, e-mail [zychkrekora@gmail.com](mailto:zychkrekora@gmail.com)

## **ABSTRACT**

This case series describes our experience with a prenatal diagnosis of fetal perineal masses. We propose a clinical-sonographic approach for prenatal workup in cases presenting perineal findings based on imaging the target sign at the posterior perineal triangle, the fetal genitalia, and sacrum. Targeted, structured prenatal anatomical scan in fetuses presenting with perineal masses may aid in the prenatal differential diagnosis and enable appropriate genetic analysis, prenatal counseling, and postnatal treatment.

Key words: perineal masses; fetus; fetal diagnosis; ultrasound

## **INTRODUCTION**

Congenital perineal masses are a rare finding that may originate from the soft tissues of the perineum itself (perineal lipomas, lipoblastomas, hemangiomas, hamartomas), the sacrum (sacroccygeal teratomas), the external genitalia (accessory scrotum or bifid scrotum, penoscrotal transposition). Findings associated with anorectal malformations and persistent cloaca may resemble a perineal mass, and therefore the differential diagnosis is essential. Most of the literature cases, specifically those diagnosed prenatally, are case reports [1–11]. Although the fetal perineum is not routinely assessed as a part of the fetal anatomical scan, we have implemented it to all fetal examinations referred for targeted scans in recent years. The present study describes our experience with fetal perineal masses and presents a clinical-sonographic flowchart for prenatal workup.

## **MATERIAL AND METHODS**

The local Ethical Committee approved the study. The committee provided a waiver of informed consent, as this was a retrospective study.

Perineal masses were diagnosed in fifteen cases throughout five years. All cases underwent a thorough sonographic morphologic fetal evaluation, and the parents were counseled by a multidisciplinary team including a neonatologist, a geneticist, and a pediatric surgeon and urologist. The fetal perineum was assessed in the axial plane to locate the anal

mucosa and sphincter. A midsagittal plane assessed the spine, the conus medullaris, and the sacral vertebra.

Examinations were performed with E8 or E10 expert machines (GE Healthcare, Kretz Ultrasound, Zipf, Austria), equipped with an abdominal 4–8-MHz curvilinear transducer. Transvaginal and transabdominal transducers were used according to the gestational age and fetal presentation.

## **RESULTS**

A total of 16 cases were referred for a targeted scan for a perineal finding between 20 and 26 weeks of gestation. Definitive diagnosis was provided after 20 weeks, following visualization of the target sign. All couples were offered genetic counseling and amniocentesis. Genetic analysis revealed abnormal results in three cases — one case of down syndrome and two consecutive cases of Townes–Brocks syndrome in the same couple. Postnatal clinical examination of the neonate or the abortus in case of termination of pregnancy (4 cases) up confirmed the diagnosis in all cases.

Prenatal sonographic findings, genetic workup, and final postnatal diagnoses are presented in Table 1 and Figures 1–3.

## **DISCUSSION**

This study reports our experience with the prenatal diagnosis of fetal perineal masses. We propose a clinical, sonographic assessment flow chart based on prenatal and postnatal findings (Figure 4).

The target sign, representing the fetal anal musculature and mucosa, plays a critical role in the clinical-sonographic differential diagnosis approach to fetal perineal lumps. According to our results, non-visualization of the target sign associated with a midline perineal lump located below the scrotum seems to represent poor development of the perineum and is related to high anorectal malformations in male fetuses. It should be noted that perineal masses can be diagnosed during the early second trimester; however, the target sign is not visualized before 20 weeks of gestation [12].

Assessment of the genitalia should follow the visualization of the target sign. As reported by others and demonstrated in the present series, perineal findings may resemble a male phallus. Application of Doppler flow and imaging of fetal urination through the normal penis with no urination through the perineal mass may aid in the differential diagnosis between rare cases of penoscrotal transposition, double penis, and perineal lipomas or other masses originating from the perineal soft tissues (Fig. 5). Sacrococcygeal teratomas originate from the coccyx and sacrum, and assessment of the sacrum is the third step in the prenatal workup of perineal masses, as they may be associated with sacral malformations and sacral agenesis.

Accurate prenatal diagnosis of perineal lumps is essential for proper prenatal counseling regarding the postnatal course and targeted genetic analysis. Our series demonstrates that anal atresia is associated with trisomy 21 and other syndromes. Similarly, urogenital malformations, precisely a penoscrotal inversion, were associated with a distal deletion of 13q [13]. Genetic syndromes associated with sacrococcygeal teratomas include autosomal dominant Currarino syndrome (OMIM #176450). The genotype-phenotype correlation is variable, although traditionally reported as a triad of a presacral mass, sacral agenesis, and anal atresia. For example, in a series by Kochling et al. [14], ten out of 23 patients from 9 families with mutations in the homeobox gene HLXB9 were asymptomatic. The radiologic investigation did reveal characteristic phenotypic features in all patients; however, the complete triad was found only in eight of the 23 patients.

## **CONCLUSIONS**

To summarize, the diversity of fetal perineal malformations requires an accurate diagnosis based on a targeted, structured anatomical scan of the perineum to provide counseling regarding the postnatal outcome.

### ***Conflict of interest***

All authors have declared all potential conflicts of interest

## **REFERENCES**

1. Ito M, Hashizume K, Kanamori Y. New phenotype of accessory scrotum with perineal lipoblastoma: coexistence of midperineal and lateral accessory scrotums. *Int J Urol*. 2004; 11(2): 125–127, doi: [10.1111/j.1442-2042.2004.00744.x](https://doi.org/10.1111/j.1442-2042.2004.00744.x), indexed in Pubmed: [14706020](https://pubmed.ncbi.nlm.nih.gov/14706020/).
2. Nakamura Y, Jennings RW, Connolly S, et al. Fetal diagnosis of penoscrotal transposition associated with perineal lipoma in one twin. *Fetal Diagn Ther*. 2010; 27(3): 164–167, doi: [10.1159/000295807](https://doi.org/10.1159/000295807), indexed in Pubmed: [20215739](https://pubmed.ncbi.nlm.nih.gov/20215739/).
3. Inde Y, Terada Y, Ikegami Ei, et al. Bifid scrotum and anocutaneous fistula associated with a perineal lipomatous tumor complicated by temporary bilateral cryptorchidism in utero mimicking ambiguous genitalia: 2-D/3-D fetal ultrasonography. *J Obstet Gynaecol Res*. 2014; 40(3): 843–848, doi: [10.1111/jog.12232](https://doi.org/10.1111/jog.12232), indexed in Pubmed: [24738127](https://pubmed.ncbi.nlm.nih.gov/24738127/).
4. Ito M, Hashizume K, Kanamori Y. New phenotype of accessory scrotum with perineal lipoblastoma: coexistence of midperineal and lateral accessory scrotums. *Int J Urol*. 2004; 11(2): 125–127, doi: [10.1111/j.1442-2042.2004.00744.x](https://doi.org/10.1111/j.1442-2042.2004.00744.x), indexed in Pubmed: [14706020](https://pubmed.ncbi.nlm.nih.gov/14706020/).
5. Ahn KiH, Boo YJ, Seol HJ, et al. Prenatally detected congenital perineal mass using 3D ultrasound which was diagnosed as lipoblastoma combined with anorectal malformation: case report. *J Korean Med Sci*. 2010; 25(7): 1093–1096, doi: [10.3346/jkms.2010.25.7.1093](https://doi.org/10.3346/jkms.2010.25.7.1093), indexed in Pubmed: [20592907](https://pubmed.ncbi.nlm.nih.gov/20592907/).
6. Wax JR, Pinette MG, Mallory B, et al. Prenatal sonographic diagnosis of a perineal lipoma. *J Ultrasound Med*. 2010; 29(8): 1257–1259, doi: [10.7863/jum.2010.29.8.1257](https://doi.org/10.7863/jum.2010.29.8.1257), indexed in Pubmed: [20660463](https://pubmed.ncbi.nlm.nih.gov/20660463/).
7. Murase N, Uchida H, Hiramatsu K. Accessory scrotum with perineal lipoma diagnosed prenatally: case report and review of the literature. *Nagoya J Med Sci*. 2015; 77(3): 501–506, indexed in Pubmed: [26412897](https://pubmed.ncbi.nlm.nih.gov/26412897/).
8. Yamaçake KG, Giron AM, Tannuri U, et al. A rare case of perineal hamartoma associated with cryptorchidism and imperforate anus: case report. *Einstein (Sao Paulo)*. 2014; 12(2): 234–236, doi: [10.1590/s1679-45082014rc2746](https://doi.org/10.1590/s1679-45082014rc2746), indexed in Pubmed: [25003932](https://pubmed.ncbi.nlm.nih.gov/25003932/).
9. Guerra-Junior G, Aun AME, Miranda ML, et al. Congenital perineal lipoma presenting as ambiguous genitalia. *Eur J Pediatr Surg*. 2008; 18(4): 269–271, doi: [10.1055/s-2008-1038366](https://doi.org/10.1055/s-2008-1038366), indexed in Pubmed: [18704895](https://pubmed.ncbi.nlm.nih.gov/18704895/).
10. Chen CP, Chang TY, Hsu CY, et al. Persistent cloaca presenting with a perineal cyst: Prenatal ultrasound and magnetic resonance imaging findings. *J Chin Med Assoc*. 2012; 75(4): 190–193, doi: [10.1016/j.jcma.2012.02.007](https://doi.org/10.1016/j.jcma.2012.02.007), indexed in Pubmed: [22541150](https://pubmed.ncbi.nlm.nih.gov/22541150/).
11. Achiron R, Frydman M, Lipitz S, et al. Urorectal septum malformation sequence: prenatal sonographic diagnosis in two sets of discordant twins. *Ultrasound in Obstetrics and Gynecology*. 2002; 16(6): 571–574, doi: [10.1046/j.1469-0705.2000.00233.x](https://doi.org/10.1046/j.1469-0705.2000.00233.x).
12. Perlman S, Bilik R, Leibovitch L, et al. More than a gut feeling - sonographic prenatal diagnosis of imperforate anus in a high-risk population. *Prenat Diagn*. 2014; 34(13): 1307–1311, doi: [10.1002/pd.4472](https://doi.org/10.1002/pd.4472), indexed in Pubmed: [25088160](https://pubmed.ncbi.nlm.nih.gov/25088160/).

13. Bartsch O, Kuhnle U, Wu LL, et al. Evidence for a critical region for penoscrotal inversion, hypospadias, and imperforate anus within chromosomal region 13q32.2q34. *Am J Med Genet.* 1996; 65(3): 218–221, doi: [10.1002/\(SICI\)1096-8628\(19961028\)65:3<218::AID-AJMG9>3.0.CO;2-L](https://doi.org/10.1002/(SICI)1096-8628(19961028)65:3<218::AID-AJMG9>3.0.CO;2-L), indexed in Pubmed: [9240747](https://pubmed.ncbi.nlm.nih.gov/9240747/).
14. Köchling J, Karbasiyan M, Reis A. Spectrum of mutations and genotype-phenotype analysis in Currarino syndrome. *Eur J Hum Genet.* 2001; 9(8): 599–605, doi: [10.1038/sj.ejhg.5200683](https://doi.org/10.1038/sj.ejhg.5200683), indexed in Pubmed: [11528505](https://pubmed.ncbi.nlm.nih.gov/11528505/).


**Table 1.**

Case	Prenatal sonographic findings	Genetic workup	Postnatal final diagnosis
1	Non-visualization of the target sign Perineal lump Tethered cord	Normal CMA	Anal atresia
2	Non-visualization of the target sign Perineal lump Single kidney	Normal CMA	Anal atresia
3	Non-visualization of the target sign Perineal lump Single kidney, single umbilical artery. Tetralogy of Fallot with absent pulmonic valve	Normal CMA	Anal atresia
4	Non-visualization of the target sign Perineal lump	Normal CMA	Anal atresia
5	Non-visualization of the target sign Perineal lump	Trisomy 21	Anal atresia
6 + 7	Non-visualization of the target sign Perineal lump hypoplastic-dysplastic kidneys bifid thumb, accessory tragus (Two consequent pregnancies)	Townes Brocks syndrome (OMIM #107480)	Anal atresia
8	Sacrococcygeal teratoma (type 1)	Normal CMA	Sacrococcygeal teratoma (type 1)
9	Sacrococcygeal teratoma (type 1)	Normal CMA	Sacrococcygeal teratoma (type 1)
10	Sacrococcygeal teratoma (type 4)	Normal CMA	Sacrococcygeal teratoma (type 3)
11	Double penis	Normal CMA	Double penis
12	Infra scrotal, perineal mass	Normal CMA	Perineal lipoma




13	Infra scrotal, perineal mass	Normal CMA	Perineal lipoma
14	Infra scrotal, perineal mass	Normal CMA	Perineal lipoma
15	Infra scrotal, perineal mass	Normal CMA	Perineal lipoma
16	Infra scrotal, perineal mass	Normal CMA	Perineal lipoma


CMA — ??




**Figure 1.** Pre- and postnatal imaging of a fetus at 30 weeks of gestation. The target sign was not visualized, consistent with anal atresia (**A**); 3D rendering of the fetal perineum demonstrating a perineal mass caudal to the scrotum and its corresponding postnatal image (**B**)




**Figure 2.** 3D rendering of the fetal perineum obtained at 22 weeks of gestation and its corresponding postnatal image in a case of a double penis



**Figure 3.** 3D rendering of the fetal perineum and its corresponding postnatal image of perineal lipomas. **A, B** — two lipomas located below the scrotum (arrows); **C, D** — one lipoma located below the scrotum (arrow)



**Figure 4.** A flowchart for prenatal sonographic workup of a perineal mass



**Figure 5.** A clinical, sonographic approach to a fetal perineal lump. The target sign was visualized at the posterior perineal triangle. (A) 2D midsagittal view demonstrating the scrotum and upper and lower findings that resemble a male phallus (\*); (B) 3D rendering of the fetal perineum demonstrating the findings described in A; (C) Doppler flow demonstrates urination from the upper mass, i.e., the penis, while no flow is demonstrated originating from the lower mass; (D) 3D rendering demonstrating fetal urination as described in C; The final diagnosis was a perineal lipoma