

---

12-1-2006

## Conformality And Q-harmonicity In Carnot Groups

Luca Capogna

*University of Arkansas, lcapogna@smith.edu*

Michael Cowling

*UNSW Sydney*

Follow this and additional works at: [https://scholarworks.smith.edu/mth\\_facpubs](https://scholarworks.smith.edu/mth_facpubs)



Part of the [Mathematics Commons](#)

---

### Recommended Citation

Capogna, Luca and Cowling, Michael, "Conformality And Q-harmonicity In Carnot Groups" (2006).

Mathematics and Statistics: Faculty Publications, Smith College, Northampton, MA.

[https://scholarworks.smith.edu/mth\\_facpubs/143](https://scholarworks.smith.edu/mth_facpubs/143)

This Article has been accepted for inclusion in Mathematics and Statistics: Faculty Publications by an authorized administrator of Smith ScholarWorks. For more information, please contact [scholarworks@smith.edu](mailto:scholarworks@smith.edu)

# CONFORMALITY AND $Q$ -HARMONICITY IN CARNOT GROUPS

---

LUCA CAPOGNA and MICHAEL COWLING

## Abstract

We show that if  $f$  is a 1-quasiconformal map defined on an open subset of a Carnot group  $G$ , then composition with  $f$  preserves  $Q$ -harmonic functions. We combine this with a regularity theorem for  $Q$ -harmonic functions and an algebraic regularity theorem for maps between Carnot groups to show that  $f$  is smooth. We give some applications to the study of rigidity.

## 1. Introduction

The theory of quasiconformal maps weaves together analysis and geometry, revealing fundamental aspects of both disciplines and links between them. Quasiconformal maps may be studied in very general settings (see, e.g., [18] and [24]) and with minimal smoothness assumptions. For instance, if  $\Omega_1$  and  $\Omega_2$  are open subsets of a metric space  $(\Omega, \varrho)$ , then the distortion of a homeomorphism  $f: \Omega_1 \rightarrow \Omega_2$  is defined by

$$H_f(p, t) = \frac{\sup\{\varrho(f(q), f(p)) \mid \varrho(q, p) \leq t\}}{\inf\{\varrho(f(q), f(p)) \mid \varrho(q, p) \geq t\}}; \quad (1.1)$$

$f$  is said to be  $\lambda$ -quasiconformal if  $\limsup_{t \rightarrow 0} H_f(p, t) \leq \lambda$  for all  $p \in \Omega_1$ .

Typically, constraints on the distortion of the map or the geometry of the underlying space yield extra regularity and rigidity properties. Liouville's 1850 theorem, which states that  $C^4$ -conformal maps between domains in  $\mathbb{R}^3$  are products of translations, dilations, and inversions, is the prototype. In modern terms, the group  $O(1, 4)$  acts conformally on the sphere  $S^3$  and hence locally on  $\mathbb{R}^3$  by stereographic projection, and any conformal map between domains in  $\mathbb{R}^3$  is the restriction of the action of an element of  $O(1, 4)$ . The same result also holds in  $\mathbb{R}^n$  when  $n > 3$  (see, e.g., R. Nevanlinna [27]). A major advance in the theory was the passage from smoothness assumptions to metric assumptions (see F. W. Gehring [14] and Yu. G. Reshetnyak [32]): the conclusion of Liouville's theorem holds for 1-quasiconformal maps. A useful reference for the classical theory is the book by J. Väisälä [36].

DUKE MATHEMATICAL JOURNAL

Vol. 135, No. 3, © 2006

Received 17 December 2004. Revision received 6 March 2006.

2000 *Mathematics Subject Classification*. Primary 30C65, 35H20; Secondary 22E25.

Capogna's work partially supported by a National Science Foundation Career Grant.

Cowling's work partially supported by an Australian Research Council Discovery Grant.

Similar rigidity theorems hold if the ambient space is not Riemannian. Analogous results with the Heisenberg group instead of Euclidean space and the sphere in  $\mathbb{C}^n$  with its CR structure instead of  $S^n$  have been long known (see, e.g., É. Cartan [4], S.-S. Chern and J. K. Moser [5], N. Tanaka [34], and A. Korányi and H. M. Reimann [22], [23]). In this work, the notion of conformality involves the Levi metric: a conformal map is a differentiable contact map whose differential, restricted to the contact plane, is a multiple of a unitary map. The relevant result for this article is that sufficiently smooth conformal maps come from the action of the group  $SU(1, n)$ . Capogna [2, Corollary 1.4, page 869] and P. Tang [35, Theorem 2.3] extended this to 1-quasiconformal maps by establishing that 1-quasiconformal maps are smooth and appealing to previous results; their regularity theorems are immediate ancestors of this article.

An important generalization of the results for CR manifolds is the study of quasiconformal maps in Carnot groups. These are nilpotent Lie groups with a left-invariant sub-Riemannian metric on a left-invariant subbundle of the tangent bundle, called the horizontal bundle, and they admit a group  $\{\delta_t \mid t \in \mathbb{R}^+\}$  of automorphisms which expand when  $t > 1$  and contract when  $t < 1$ . *We define Carnot groups to exclude  $\mathbb{R}$ .*

In his celebrated work [25], G. D. Mostow introduced quasiconformal maps on certain Carnot groups to prove rigidity theorems for rank-one symmetric spaces. He showed that certain quasi-isometries extend to quasiconformal maps of the ideal boundary and then that these are conformal.

P. Pansu [28] made the next breakthrough: he proved that quasiconformal maps\* on general Carnot groups are differentiable in a suitable sense and applied this to study rigidity. A map  $f: \Omega_1 \rightarrow \Omega_2$  between open subsets of Carnot groups is said to be Pansu differentiable at  $p$  in  $\Omega_1$  if  $\delta_t^{-1}[f(p)^{-1}f(p\delta_t x)]$  converges locally uniformly in  $x$  as  $t \rightarrow 0$ . The limit  $Df(p)$  is called the Pansu differential at  $p$ . In [28] Pansu showed that a quasiconformal map  $f$  is Pansu differentiable almost everywhere and that  $Df(p)$  is a group homomorphism that intertwines the dilations  $\delta_t$ . In particular,  $Df(p)$  induces a Lie algebra homomorphism  $df(p)$  that sends horizontal vectors to horizontal vectors. Pansu used this form of differentiability and a study of Lie algebra automorphisms to prove that in the quaternionic and octonionic Heisenberg groups, there is a natural generalized contact structure of codimension greater than one, and the analogue of Liouville's theorem holds for global maps that preserve the contact structure.

These ideas have been extended to many more Carnot groups. For example, the space of smooth, locally defined quasiconformal maps is finite-dimensional for the so-called groups of Heisenberg type (or,  $H$ -type groups; see [21], [8]) when the center has dimension at least three (see [30]; see also the work of Reimann and F. Ricci [31],

\*Actually, Pansu's definition of quasiconformality was different but equivalent (see Definition 4.5).

Cowling and Reimann [9], and Cowling, F. De Mari, A. Korányi, and Reimann [7]). These results, however, need some smoothness. Capogna [3] followed the approach of Gehring [14] and proved that 1-quasiconformal maps in  $H$ -type groups are smooth. The point of this article is that 1-quasiconformal maps are smooth in *all* Carnot groups.

#### THEOREM 1.1

*Suppose that  $\Omega_1$  and  $\Omega_2$  are open subsets of Carnot groups  $G_1$  and  $G_2$ , and suppose that  $f: \Omega_1 \rightarrow \Omega_2$  is 1-quasiconformal. Then  $f$  is smooth.*

In  $\mathbb{R}^n$ , our proof is simpler than Gehring's [14], as we do not need such a careful study of the capacity of rings. It may be very hard to generalize Gehring's approach to all Carnot groups since there is no explicit Green's function for the  $Q$ -Laplacian (see (2.3)). However, Gehring's method does extend to some noncommutative groups (see [1], [3], [6], [10], and [38]).

The first main step of our proof is to use the deep results of Pansu [28] and Heinonen and Koskela [18] to prove that the components of the first layer of  $f$  are  $Q$ -harmonic (see (2.3)). The regularity theory in [3, Main Theorem, page 264, Theorem 4.6, page 282] then implies that the first layer is smooth. The second main step is an algebraic regularity argument (see Theorem 6.1) to show that the higher layers are smooth; we study how the Pansu differential of the map links the layers of its components.

Theorem 1.1 can be used to prove rigidity theorems for quasiconformal maps between open subsets in certain classes of groups without any a priori smoothness assumption. For instance, in Corollary 7.4 we show that if  $G$  is a two-step Carnot group whose only dilation-preserving automorphisms are dilations (see [28, Proposition 13.1] for a large class of examples), then the only quasiconformal maps between open subsets of  $G$  are translations composed with dilations. Combining Theorem 1.1 with the results in [30], we can show a local version of one of the main results in [28].

#### THEOREM 1.2

*Suppose that  $G$  is the nilpotent group in the Iwasawa decomposition of the isometry group of the quaternionic or Cayley hyperbolic space. Quasiconformal maps between two open subsets of  $G$  form a finite-dimensional subset of the space of smooth, generalized contact maps.*

In conclusion, two more of our contributions should be highlighted. First, the metric definition and Pansu's definition of 1-quasiconformal maps are equivalent (see Corollary 7.2), and second, the morphism property of 1-quasiconformal maps needs no extra regularity assumptions (see Theorem 5.7).

## 2. Preliminary results

In this section we recall the relevant definitions, fix notation, and recall some results from the literature on which our arguments rest.

A grading of step  $R$  of a Lie algebra  $\mathfrak{g}$  is a decomposition of  $\mathfrak{g}$  as a vector-space direct sum  $\sum_{k=1}^R V_k$  such that  $[V_i, V_j] \subseteq V_{i+j}$  when  $1 \leq i, j \leq R$ ; here we interpret  $V_i$  as  $\{0\}$  when  $i > R$  and require that  $V_R \neq \{0\}$ .

A stratified Lie group is a simply connected, necessarily nilpotent group  $G$  whose Lie algebra  $\mathfrak{g}$  has a grading with the additional property that  $V_1$  generates  $\mathfrak{g}$ . The map on  $\mathfrak{g}$  which multiplies the elements of the  $k$ th layer  $V_k$  by  $k$  is a derivation. This gives rise to a group of automorphic dilations  $\{\delta_t \mid t \in \mathbb{R}^+\}$  of  $\mathfrak{g}$  defined by

$$\delta_t(X_1 + \dots + X_R) = tX_1 + \dots + t^R X_R,$$

where  $X_k \in V_k$ , such that  $\delta_s \delta_t = \delta_{st}$  and  $\delta_1$  is the identity map (thus  $\delta_{t^{-1}} = \delta_t^{-1}$ ). We also write  $\delta_t$  for the corresponding automorphisms  $\exp \circ \delta_t \circ \log$  of  $G$ . (Here,  $\exp$  denotes the exponential map, which is bijective, and  $\log$  denotes its inverse.) We identify the Lie algebra with the tangent space  $TG_e$  to  $G$  at the identity  $e$ , and for  $X$  in  $\mathfrak{g}$ , we write  $\tilde{X}$  for the left-invariant vector field that agrees with  $X$  at  $e$ . The vector fields in  $\tilde{V}_k$  determine a subspace  $T_k G_p$  of the tangent space at each point  $p$ . In particular, when  $k = 1$ , we obtain the *horizontal subspace*. These subspaces vary smoothly from point to point and give rise to subbundles  $T_k G$  of the tangent bundle. The hypothesis of stratification implies that the sections of the horizontal tangent bundle, called the horizontal vector fields, generate all possible vector fields with linear combinations of commutators of order up to  $R$ .

A *Carnot group* is a stratified group  $G$ , other than  $\mathbb{R}$ , whose Lie algebra  $\mathfrak{g}$  carries an inner product for which the different layers  $V_k$  are orthogonal. Henceforth,  $G$  denotes a Carnot group with Lie algebra  $\mathfrak{g}$ . Denote by  $\pi_k$  the orthogonal projection onto  $V_k$  by  $m_k$ , the vector space dimension of the layer  $V_k$ , when  $k = 1, \dots, R$ , and denote by  $m$  the dimension  $m_1 + \dots + m_R$ . We identify  $V_k$  isometrically with Euclidean space  $\mathbb{R}^{m_k}$ , and  $\mathfrak{g}$  with  $\mathbb{R}^m$ . We write  $Q$  for the *homogeneous dimension*  $\sum_{k=1}^R km_k$  of  $G$  (see [13]). Since  $\exp$  is a bijection, we may parametrize  $G$  by  $\mathfrak{g}$ . The measure on  $G$  obtained by pushing forward the Lebesgue measure on  $\mathfrak{g}$  is translation invariant. We denote by  $|E|$  the measure of a set  $E$ ; then  $|\delta_t E| = t^Q |E|$ .

The simplest noncommutative Carnot group is the Heisenberg group  $\mathbb{H}^1$ , for which  $R = 2$ ,  $m_1 = 2$ , and  $m_2 = 1$ . We use it as an example to illustrate some of the main ideas in a simple setting. A basis of its Lie algebra is  $\{X_1, X_2, X_3\}$ , where  $[X_1, X_2] = X_3$  and  $[X_1, X_3] = [X_2, X_3] = 0$ . Then  $V_1 = \text{span}\{X_1, X_2\}$ , while  $V_2 = \text{span}\{X_3\}$ . We identify  $\mathbb{H}^1$  as a set with  $\mathbb{R}^3$ , writing  $x$  or  $(x_1, x_2, x_3)$  for  $\exp(x_1 X_1 + x_2 X_2 + x_3 X_3)$ . The group law is

$$xy = \left(x_1 + y_1, x_2 + y_2, x_3 + y_3 + \frac{1}{2}(x_1 y_2 - x_2 y_1)\right).$$

In coordinates, the left-invariant vector fields  $\tilde{X}_1, \tilde{X}_2,$  and  $\tilde{X}_3$  are given by

$$\tilde{X}_1 = \frac{\partial}{\partial x_1} - \frac{x_2}{2} \frac{\partial}{\partial x_3}, \quad \tilde{X}_2 = \frac{\partial}{\partial x_2} + \frac{x_1}{2} \frac{\partial}{\partial x_3}, \quad \text{and} \quad \tilde{X}_3 = \frac{\partial}{\partial x_3}.$$

Note that  $[\tilde{X}_1, \tilde{X}_2]$  is  $\tilde{X}_3$  and that  $\tilde{X}_1, \tilde{X}_2,$  and  $\tilde{X}_3$  span the Lie algebra of all left-invariant vector fields.

The inner product on  $V_1$  induces a left-invariant inner product on each horizontal space. This inner product allows us to define a (left-invariant) Carnot-Carathéodory metric  $\varrho$  on  $G$ , as follows. A smooth curve is said to be *horizontal* if its tangent vectors are horizontal, that is, they lie in the horizontal tangent space, and the length of a horizontal curve is the integral of the lengths of its tangent vectors. The distance between two points is then the infimum of the lengths of the horizontal curves joining them. The metric is left-translation-invariant and is related to the dilations by the formula  $\varrho(\delta_t x, \delta_t y) = t \varrho(x, y)$  for all  $t \in \mathbb{R}^+$  and all  $x, y \in G$ . We denote by  $B(x, t)$  the open ball  $\{y \in G \mid \varrho(y, x) < t\}$ ; then  $|B(x, t)| = C_G t^Q$ . We write  $S(x, t)$  for the *sphere*  $\{y \in G \mid \varrho(y, x) = t\}$ . If  $X \in V_1$ , then the curve  $s \mapsto \exp(sX)$  is horizontal and minimizes lengths, so that

$$\varrho(\exp X, e) = |X|, \quad \forall X \in V_1. \tag{2.1}$$

There is a natural pseudonorm  $\|\cdot\|$  on the stratified Lie algebra  $\mathfrak{g}$ , given by  $\|X\|^{2R_1} = \sum_{k=1}^R |\pi_k X|^{2R_1/k}$ , and an associated left-invariant pseudometric  $\varrho_{\text{NSW}}$ , given by  $\varrho_{\text{NSW}}(x, y) = \|\log(y^{-1}x)\|$  for all  $x, y \in G$ . According to A. Nagel, E. M. Stein, and S. Wainger [26], this pseudometric is equivalent to the Carnot-Carathéodory metric, in that

$$C_1 \varrho(x, y) \leq \varrho_{\text{NSW}}(x, y) \leq C_2 \varrho(x, y), \quad \forall x, y \in G \tag{2.2}$$

(for suitable constants  $C_1$  and  $C_2$ ).

We use the symbol  $\simeq$  in the following sense. For functions  $A$  and  $B$  defined on  $S$ , we write  $A(s) \simeq B(s)$  for all  $s$  in  $S$  if and only if there exist constants  $C_1$  and  $C_2$  (all constants are taken to be positive) such that  $C_1 A(s) \leq B(s) \leq C_2 A(s)$  for all  $s \in S$ . In particular,  $\varrho \simeq \varrho_{\text{NSW}}$ . Constants are denoted by  $C, C', C'', C_1, C_2, \dots$ ; their values may vary from formula to formula and may depend on anything not explicitly quantified.

We take the standard basis  $\{X_1, \dots, X_m\}$  for  $\mathfrak{g}$ , so that  $\{X_1, \dots, X_{m_1}\}$  is a basis for  $V_1$ , while  $\{X_{m_1+1}, \dots, X_{m_1+m_2}\}$  is a basis for  $V_2$ , and so on. Then  $\{\tilde{X}_1, \dots, \tilde{X}_m\}$  is a basis for the left-invariant vector fields on  $G$ , and  $\{\tilde{X}_1, \dots, \tilde{X}_{m_1}\}$  is a basis for the left-invariant horizontal vector fields. The functions  $X_i \cdot \log: G \rightarrow \mathbb{R}$  are called *exponential coordinates* on  $G$ . The Lie algebra element  $X_i$  is in the first layer when

$1 \leq i \leq m_1$ , and accordingly, we call  $(X_1 \cdot \log, \dots, X_{m_1} \cdot \log): G \rightarrow \mathbb{R}^{m_1}$  the *first layer* of coordinates.

Suppose that  $V$  is a finite-dimensional inner product space. For a function  $u: G \rightarrow V$ , we write  $\nabla_H u$  for the vector  $(\tilde{X}_1 u, \dots, \tilde{X}_{m_1} u)$ ; then  $\nabla_H^h u$  is a  $V$ -valued tensor when  $h = 2, 3, \dots$ . We consider a multi-index  $(i_1, i_2, \dots, i_l)$  for which  $1 \leq i_1, i_2, \dots, i_l \leq m$ ; we abbreviate this to  $I$  and define its homogeneous length  $|I|$  to be  $\sum_{j=1}^l \kappa_{i_j}$ , where the integer  $\kappa_i$  is specified by requiring that  $m_1 + \dots + m_{\kappa_i - 1} < i \leq m_1 + \dots + m_{\kappa_i}$  when  $1 \leq i \leq m$ . We write  $\nabla^I u$  for  $\tilde{X}_{i_1} \tilde{X}_{i_2} \dots \tilde{X}_{i_l} u$  and consider this to be a derivative of  $u$  of order  $|I|$  since  $\nabla^I(u \circ \delta_s)(p) = s^{|I|}(\nabla^I u)(\delta_s(p))$ .

If  $0 \leq \alpha \leq 1$ , then the space  $C^{0,\alpha}(\Omega, V)$  is defined to be the set of all continuous functions  $u: \Omega \rightarrow V$  such that  $|u(p) - u(q)| \leq C \varrho(p, q)^\alpha$  for all  $p, q \in \Omega$ , and when  $k \geq 1$ , the spaces  $C^{k,\alpha}(\Omega, V)$  are defined by

$$C^{k,\alpha}(\Omega, V) = \{u: \Omega \rightarrow V \mid \tilde{X}_i u \in C^{k-1,\alpha}(\Omega, V) \text{ when } 1 \leq i \leq m_1\}.$$

Because every vector field in  $\tilde{\mathfrak{g}}$  arises as a commutator of vector fields in  $\tilde{V}_1$ , if  $u \in C^{k,\alpha}(\Omega, V)$ , then  $\nabla^I u \in C^{0,\alpha}(\Omega, V)$  when  $0 \leq |I| \leq k$ . In particular, if  $\tilde{X}_i u \in C^\infty(\Omega, V)$  whenever  $1 \leq i \leq m_1$ , then  $u \in C^\infty(\Omega, V)$ .

The local versions of these spaces  $C_{\text{loc}}^{k,\alpha}(\Omega, V)$  are defined in the standard way. Similarly, the subscript 0 on a function space indicates that the functions are compactly supported. When  $V$  is  $\mathbb{R}$  or  $\mathbb{C}$ , we omit  $V$  in the notation.

By definition, the Lie bracket is bilinear. Clearly, there exists a constant  $C$  such that  $\|[X, Y]\| \leq C\|X\|\|Y\|$  for all  $X, Y \in \mathfrak{g}$ . It follows that if  $U$  and  $V$  lie in  $C_{\text{loc}}^{k,\alpha}(\Omega, \mathfrak{g})$ , then  $[U, V] \in C_{\text{loc}}^{k,\alpha}(\Omega, \mathfrak{g})$ .

Let  $\Omega$  be an open subset of  $G$ . When  $k \in \mathbb{N}$  and  $1 \leq P < \infty$ , we define the *horizontal Sobolev spaces* (see [12] and [13]) by

$$S^{k,P}(\Omega) = \{u: \Omega \rightarrow \mathbb{C} \mid |\nabla_H^h u| \in L^P(\Omega) \text{ when } 0 \leq h \leq k\}.$$

The differentiation here must be interpreted in the distributional sense.

A real-valued function  $u$  in  $S_{\text{loc}}^{1,Q}(\Omega)$  is said to be a weak solution of the  $Q$ -Laplace equation ( $Q$  denotes the homogeneous dimension of  $G$ )

$$\nabla_H \cdot (|\nabla_H u|^{Q-2} \nabla_H u) = 0 \tag{2.3}$$

if  $\int_\Omega |\nabla_H u(x)|^{Q-2} \nabla_H u(x) \cdot \nabla_H \phi(x) dx = 0$  for all  $\phi \in C_0^\infty(\Omega)$ ; such functions are called  $Q$ -harmonic. By a standard density argument, we can change the requirement that  $\phi \in C_0^\infty(\Omega)$  to the requirement that  $\phi \in C_0^{0,1}(\Omega)$ .

The following regularity theorem, proved in [3, Corollary, page 264], plays a crucial role in the proof of Theorem 1.1.

**THEOREM 2.1 (Regularity theorem for the  $Q$ -Laplacian)**

Suppose that  $u$  is  $Q$ -harmonic, and suppose that  $|\nabla_H u| \simeq 1$  in a subset of  $\Omega$  of full measure. Then, after changing  $u$  on a null set,  $u \in C^\infty(\Omega)$ .

The Baker-Campbell-Hausdorff (BCH) formula (see [20]) is useful for computations in Carnot groups. For  $X$  and  $Y$  in  $\mathfrak{g}$ , the product  $e^X e^Y$  is equal to  $\exp(\text{BCH}(X, Y))$ , where  $\text{BCH}(X, Y)$  is a polynomial in  $X$  and  $Y$ ; more precisely,  $\text{BCH}(X, Y) = \text{BCH}_1(X, Y) + \text{BCH}_2(X, Y) + \text{BCH}_3(X, Y) + \dots$ , where each term  $\text{BCH}_i$  is a homogeneous Lie algebra polynomial of degree  $i$ , that is, a weighted sum of commutators of order  $i - 1$ . For instance,

$$\begin{aligned} \text{BCH}_1(X, Y) &= X + Y, \\ \text{BCH}_2(X, Y) &= \frac{1}{2} [X, Y], \\ \text{BCH}_3(X, Y) &= \frac{1}{12} ([X, [X, Y]] - [Y, [X, Y]]), \\ \text{BCH}_4(X, Y) &= -\frac{1}{48} ([Y, [X, [X, Y]]] + [X, [Y, [X, Y]]]), \end{aligned}$$

and so on. This series terminates with  $\text{BCH}_R$ , as  $G$  is nilpotent of step  $R$ .

Suppose that  $T : G_1 \rightarrow G_2$  is a continuous (hence, smooth; see [20, Theorem 2.6, page 117]) homomorphism of Carnot groups with dilations  $\delta_t^1$  and  $\delta_t^2$ . We say that  $T$  *intertwines dilations* (and call  $T$  a *Carnot homomorphism*) if  $\delta_t^2 T = T \delta_t^1$ . Then the (Riemannian) differential  $T_*(e)$  of  $T$  at the identity is a Lie homomorphism from  $\mathfrak{g}_1$  to  $\mathfrak{g}_2$  which respects the layers of the two groups. From the definition of the exponential map,  $\exp T_*(e)W = T \exp W$  for all  $W \in V_1$ . Similarly, if  $\gamma : [0, 1] \rightarrow G_1$  is a horizontal curve, then  $\exp T_*(\gamma(s))(\dot{\gamma}(s)) = T \exp \dot{\gamma}(s)$ . In the rest of this article, we abuse notation by writing  $\delta_t$  for the dilations on all Carnot groups.

**LEMMA 2.2**

Suppose that  $T : G_1 \rightarrow G_2$  is a Carnot homomorphism. Then  $T$  is Lipschitz; that is,  $\varrho(Tx, Ty) \leq C \varrho(x, y)$  for all  $x, y \in G$ , where

$$\begin{aligned} C &= \max\{\varrho(Tz, e) \mid z \in S(e, 1)\} \\ &= \max\{\varrho(Ty, e) \mid y \in \exp(V_1) \cap S(e, 1)\}. \end{aligned}$$

*Proof*

First, since  $\varrho(Tx, Ty) = \varrho(T(y^{-1}x), e)$  and  $\varrho(T\delta_t z, e) = t \varrho(z, e)$  for all  $x, y$ , and  $z$  in  $G_1$ ,

$$\varrho(Tx, Ty) \leq \max\{\varrho(Tz, e) \mid z \in S(e, 1)\} \varrho(x, y).$$



Next, if  $z \in S(e, 1)$  and  $\epsilon > 0$ , then there exists a horizontal curve  $\gamma : [0, 1] \rightarrow G_1$  such that  $|\dot{\gamma}(s)| \leq 1 + \epsilon$  for all  $s$  in  $[0, 1]$ , and  $\gamma(0) = e$  and  $\gamma(1) = z$ . Then  $T\gamma(0) = e$  and  $T\gamma(1) = Tz$ , while

$$\begin{aligned} |(T\gamma)'(s)| &= |T_*(\gamma(s))(\dot{\gamma}(s))| \\ &\leq \max\{|T_*(\gamma(s))(W)| \mid W \in V_1, |W| \leq 1 + \epsilon\} \\ &= (1 + \epsilon) \max\{\varrho(T \exp W, e) \mid W \in V_1, |W| \leq 1\}, \end{aligned}$$

from (2.1), whence  $\varrho(Tz, e) \leq \max\{\varrho(Tw, e) \mid w \in \exp(V_1) \cap S(e, 1)\}$ .  $\square$

### 3. Calculus on Carnot groups

In Sections 3–6, we show that the first layer of a 1-quasiconformal map is smooth. Some of the material is often described as well known, but we had difficulty finding detailed references for all the results. We therefore review some features of calculus on Carnot groups and quasiconformal maps.

Throughout this section, we suppose that  $\Omega_1$ ,  $\Omega_2$ , and  $\Omega_3$  are open subsets of Carnot groups  $G_1$ ,  $G_2$ , and  $G_3$ , and we suppose that  $f$  is a map from  $\Omega_1$  to  $\Omega_2$ . We write  $f \in C^{0,\alpha}(\Omega_1, \Omega_2)$  if  $\varrho(f(p), f(q)) \leq C \varrho(p, q)^\alpha$  for all  $p, q \in \Omega_1$ .

There is an important notational issue: for functions whose values lie in multiplicative groups, expressions such as  $f^{-1}(p)$  are ambiguous. We always write  $f^{-1}(p)$  for the value of the inverse function  $f^{-1}$  at  $p$ , and we write  $f(p)^{-1}$  for  $[f(p)]^{-1}$ , the multiplicative inverse of the group element  $f(p)$ .

#### Definition 3.1

Given  $f : \Omega_1 \rightarrow \Omega_2$  and  $p$  in  $\Omega_1$ , we define

$$\begin{aligned} \text{Lip}_f(p) &= \limsup_{q \rightarrow p} \frac{\varrho(f(q), f(p))}{\varrho(q, p)}, \\ \text{lip}_f(p) &= \liminf_{q \rightarrow p} \frac{\varrho(f(q), f(p))}{\varrho(q, p)}. \end{aligned}$$

Suppose that  $f : \Omega_1 \rightarrow \Omega_2$  is a homeomorphism, and suppose that  $p \in \Omega_1$ . Then

$$\text{lip}_f(p) = [\text{Lip}_{f^{-1}}(f(p))]^{-1}, \tag{3.1}$$

and for small  $t$ , the distortion (1.1) is given by

$$H_f(p, t) = \frac{\max\{\varrho(f(q), f(p)) \mid \varrho(q, p) = t\}}{\min\{\varrho(f(q), f(p)) \mid \varrho(q, p) = t\}} \tag{3.2}$$

because  $G_1$  and  $G_2$  are manifolds.

*Definition 3.2*

A continuous function  $f : \Omega_1 \rightarrow \Omega_2$  is said to be Pansu differentiable at a point  $p$  in  $\Omega_1$  if the functions  $x \mapsto \delta_t^{-1}[f(p)^{-1}f(p\delta_t x)]$  converge locally uniformly to a Lie group homomorphism  $Df(p)$  as  $t \rightarrow 0$ .

The homomorphism  $Df(p) : G_1 \rightarrow G_2$ , called the Pansu differential, is a Carnot homomorphism. The Lie derivative of  $Df(p)$ , written  $df(p)$ , is a grading-preserving Lie homomorphism from  $\mathfrak{g}_1$  to  $\mathfrak{g}_2$ .

LEMMA 3.3

If  $f : \Omega_1 \rightarrow \Omega_2$  is Pansu differentiable at  $p$ , then

$$\begin{aligned} \text{Lip}_f(p) &= \max\{\varrho(Df(p)x, e) \mid x \in S(e, 1)\} \\ &= \max\{|df(p)X| \mid X \in V_1, |X| = 1\}, \end{aligned}$$

and

$$\text{lip}_f(p) = \min\{\varrho(Df(p)x, e) \mid x \in S(e, 1)\}.$$

If, in addition,  $\text{Lip}_f(p) \neq 0$ , then

$$\limsup_{t \rightarrow 0} H_f(p, t) = \frac{\text{Lip}_f(p)}{\text{lip}_f(p)}.$$

*Proof*

Since  $f$  is Pansu differentiable at  $p$ ,

$$\begin{aligned} \lim_{t \rightarrow 0} \frac{\varrho(f(p\delta_t x), f(p))}{t} &= \lim_{t \rightarrow 0} \varrho(\delta_t^{-1}[f(p)^{-1}f(p\delta_t x)], e) \\ &= \varrho(Df(p)x, e) \end{aligned}$$

uniformly for  $x$  in  $S(e, 1)$ . Since the function  $x \mapsto Df(p)x$  is continuous, it has a maximum and minimum on  $S(e, 1)$ . Therefore

$$\begin{aligned} \text{Lip}_f(p) &= \limsup_{q \rightarrow p} \frac{\varrho(f(q), f(p))}{\varrho(q, p)} \\ &= \limsup_{t \rightarrow 0} \max_{x \in S(e, 1)} \frac{\varrho(f(p\delta_t x), f(p))}{t} \\ &= \max_{x \in S(e, 1)} \varrho(Df(p)x, e). \end{aligned}$$

The second equality for  $\text{Lip}_f(p)$  follows from Lemma 2.2. To prove the equality for  $\text{lip}_f(p)$ , we exchange suprema and maxima with infima and minima in the argument

above. Finally, from (3.2),

$$\begin{aligned} \limsup_{t \rightarrow 0} H_f(p, t) &= \limsup_{t \rightarrow 0} \frac{\max\{\varrho(f(q), f(p)) \mid \varrho(q, p) = t\}}{\min\{\varrho(f(q), f(p)) \mid \varrho(q, p) = t\}} \\ &= \limsup_{t \rightarrow 0} \frac{\max\{\varrho(f(p\delta_t x), f(p)) \mid x \in S(e, 1)\}}{\min\{\varrho(f(p\delta_t x), f(p)) \mid x \in S(e, 1)\}} \\ &= \frac{\lim_{t \rightarrow 0} \max\{\varrho(f(p\delta_t x), f(p)) \mid x \in S(e, 1)\}/t}{\lim_{t \rightarrow 0} \min\{\varrho(f(p\delta_t x), f(p)) \mid x \in S(e, 1)\}/t} \\ &= \frac{\text{Lip}_f(p)}{\text{lip}_f(p)}, \end{aligned}$$

unless the last expression is of the form  $0/0$ . □

Since  $df(p)$  respects the gradings, the matrix representing  $df(p)$  is block diagonal, and since it is a Lie algebra homomorphism, the top block, which represents the action on  $V_1$ , determines the other blocks. We now show that the restrictions to  $V_1$  of the Pansu differential  $df$  and of the Riemannian differential coincide when they both exist (cf. [17, Section 3]).

#### LEMMA 3.4

Suppose that  $f: \Omega_1 \rightarrow \Omega_2$  is Pansu differentiable at  $p$  in  $\Omega_1$ . Write  $f(p)^{-1}f(px)$  as  $\exp(F(x))$ , and write  $F(x)$  as  $F_1(x) + \dots + F_R(x)$ , where  $F_j(x) \in V_j$ . If  $1 \leq k \leq j \leq R$  and  $X \in V_k$ , then  $\tilde{X}F_j(e)$  exists, and

$$\tilde{X}F_j(e) = \begin{cases} df(p)X & \text{if } k = j, \\ 0 & \text{if } k < j. \end{cases}$$

In addition,  $F_k \circ \exp|_{V_k}$  is differentiable in the usual sense at the origin, with differential  $(F_k \circ \exp|_{V_k})_*(0)$ , which is equal to  $df(p)|_{V_k}$ .

*Proof*

Since  $f$  is Pansu differentiable at  $p$ ,

$$\lim_{t \rightarrow 0} \delta_t^{-1}[f(p)^{-1}f(p\delta_t \exp X)] = Df(p)(\exp X)$$

uniformly for  $X$  in  $V_k$  of norm at most 1. Thus there exists a function  $\epsilon: \mathbb{R}^+ \rightarrow \mathbb{R}^+$  such that  $\lim_{t \rightarrow 0} \epsilon(t) = 0$  and

$$\varrho(\delta_t^{-1}[f(p)^{-1}f(p\delta_t \exp X)], Df(p)(\exp X)) \leq \epsilon(t)$$

for all  $X \in V_k$  such that  $|X| \leq 1$ . Now  $df(p)X \in V_k$ , and thus

$$\begin{aligned} & \varrho \left( \delta_t^{-1} [f(p)^{-1} f(p \delta_t \exp X)], Df(p)(\exp X) \right) \\ &= t^{-1} \varrho \left( [f(p)^{-1} f(p \delta_t \exp X)], \delta_t Df(p)(\exp X) \right) \\ &= t^{-1} \varrho \left( \exp(F(\delta_t \exp X)), \exp(\delta_t df(p)X) \right) \\ &= t^{-1} \varrho \left( \exp(-df(p)t^k X) \exp(F(\exp t^k X)), e \right) \\ &= t^{-1} \varrho \left( \exp(\text{BCH}(-df(p)t^k X, F(\exp t^k X))), e \right) \\ &\simeq t^{-1} \left( \sum_{j=1}^R |\pi_j \text{BCH}(-df(p)t^k X, F(\exp t^k X))|^{2R1/j} \right)^{1/2R1}, \end{aligned}$$

whence  $|\pi_j \text{BCH}(-df(p)t^k X, F(\exp t^k X))| \leq C [t\epsilon(t)]^j$  uniformly for  $X$  in  $V_k$  of norm at most 1; or equivalently,

$$|\pi_j \text{BCH}(-df(p)Y, F(\exp Y))| \leq C (|Y|^{1/k} \epsilon(|Y|^{1/k}))^j \tag{3.3}$$

uniformly for  $Y$  in  $V_k$ , by (2.2).

If  $1 \leq j < k$  and  $Y \in V_k$ , then  $\text{BCH}(-df(p)Y, F(\exp Y)) = F_j(\exp Y) + R$ , where  $R \in V_{j+1} \oplus \dots \oplus V_R$ , so  $\pi_j \text{BCH}(-df(p)Y, F(\exp Y)) = F_j(\exp Y)$ , and hence,

$$|F_j(\exp Y)| \leq C (|Y|^{1/k} \epsilon(|Y|^{1/k}))^j = C_j |Y|^{j/k} \epsilon_j(|Y|),$$

where  $\epsilon_j(t) \rightarrow 0$  as  $t \rightarrow 0$ .

Next, we consider the case where  $k = j$ . For  $Y$  in  $V_k$ , direct computation shows that  $\pi_k \text{BCH}(-df(p)Y, F(\exp Y)) = F_k(\exp Y) - df(p)Y$ , whence

$$|F_k(\exp Y) - df(p)Y| \leq C |Y| \epsilon(|Y|^{1/k})^k = C_k |Y| \epsilon_k(|Y|),$$

where  $\epsilon_k(t) \rightarrow 0$  as  $t \rightarrow 0$ . Thus  $F_k \circ \exp|_{V_k}$  is differentiable with derivative  $df(p)|_{V_k}$ , as required. Consequently,  $|F_k(\exp Y)| \leq C_k |Y| \epsilon'_k(|Y|)$ , where  $\epsilon'_k(t)$  remains bounded as  $t \rightarrow 0$ .

Now, we consider the case where  $j > k$ . For  $Y$  in  $V_k$ ,

$$\pi_{k+1} \text{BCH}(-df(p)Y, F(\exp Y)) = F_{k+1}(\exp Y) - \frac{1}{2} [df(p)Y, F_1(\exp Y)],$$

and from the estimates above, if  $|Y| \leq 1$ , then

$$\begin{aligned} |[df(p)Y, F_1(\exp Y)]| &\leq C |df(p)Y| |F_1(\exp Y)| \\ &\leq \begin{cases} C' |Y|^{1+1/k} \epsilon(|Y|^{1/k}) & \text{if } k > 1, \\ C'' |Y|^2 & \text{if } k = 1. \end{cases} \end{aligned}$$

It follows that  $|F_{k+1}(\exp Y)| \leq C_{k+1}|Y|^{(k+1)/k}\epsilon_{k+1}(|Y|)$ , where  $\epsilon_{k+1}(t)$  remains bounded as  $t \rightarrow 0$ . Continuing inductively, it is easy to show that if  $j > k$ , then  $|F_j(\exp Y)| \leq C_j|Y|^{j/k}\epsilon_j(|Y|)$ , where  $\epsilon_j(t)$  remains bounded as  $t \rightarrow 0$ , whence  $F_j \circ \exp|_{V_k}$  is differentiable, and its derivative is zero.  $\square$

COROLLARY 3.5

Suppose that  $u : \Omega_1 \rightarrow V$  is a function from an open subset  $\Omega_1$  of the Carnot group  $G_1$  to a vector space  $V$ . Then if  $Du(p)$  exists, so does  $\nabla_H u(p)$ , and  $\tilde{X}u(p) = Du(p) \exp X$  for all  $X \in V_1$ . Consequently,  $\text{Lip}_u(p) = |\nabla_H u(p)|$ .

LEMMA 3.6

Suppose that  $u : \Omega_1 \rightarrow \mathbb{R}$  is  $Q$ -harmonic. Then (after correction on a null set)  $u$  is continuous and Pansu differentiable almost everywhere in  $\Omega_1$ .

Proof

The proof has two steps. First, one shows that for any bounded ball  $\Omega_0$  such that  $\tilde{\Omega}_0 \subset \Omega_1$ , there exists  $P$  in  $(Q, \infty)$  such that  $\nabla_H u$  is in  $L^P(\Omega_0)$ . This requires a standard argument based on Gehring’s reverse Hölder inequality (see [17, (6.2)]). It follows from a Sobolev-type embedding theorem that (after correction on a null set)  $u$  is continuous.

Next, one shows that  $S^{1,P}(\Omega_0)$ -functions are Pansu differentiable almost everywhere. By the  $L^P$ -version of Lebesgue’s differentiation theorem,

$$\lim_{t \rightarrow 0} \frac{1}{|B(p, 2t)|} \int_{B(p, 2t)} |\nabla_H u(x) - \nabla_H u(p)|^P dx = 0 \tag{3.4}$$

for almost every  $p \in \Omega_0$ . Fix any such point  $p$ , and define the function  $v$  by  $v(q) = u(q) - u(p) - \langle \nabla_H u(p), \pi_1(\log p^{-1}q) \rangle$ . (We write  $\pi_1$  for the orthogonal projection of  $\mathfrak{g}$  onto  $V_1$ .) To show that  $u$  is Pansu differentiable at  $p$ , we must show that  $|v(q)| = o(\varrho(q, p))$  as  $q \rightarrow p$ . For this, we use a Morrey-type estimate (see, e.g., [17, Lemma 6.10]):

$$\begin{aligned} |v(q)| &= |v(q) - v(p)| \\ &\leq C \varrho(q, p)^{1-Q/P} \left( \int_{B(p, 2\varrho(q, p))} |\nabla_H v(x)|^P dx \right)^{1/P} \\ &= C \varrho(q, p)^{1-Q/P} \left( \int_{B(p, 2\varrho(q, p))} |\nabla_H u(x) - \nabla_H u(p)|^P dx \right)^{1/P} \\ &= C' \varrho(q, p) \left( \frac{1}{|B(p, 2\varrho(q, p))|} \int_{B(p, 2\varrho(q, p))} |\nabla_H u(x) - \nabla_H u(p)|^P dx \right)^{1/P} \\ &= o(\varrho(q, p)), \end{aligned}$$

by (3.4), uniformly in  $q$  such that  $B(p, 2\varrho(q, p)) \subset \Omega_0$ . □

**LEMMA 3.7 (Chain rule)**

Suppose that  $f: \Omega_1 \rightarrow \Omega_2$  is Pansu differentiable at  $p$ , and suppose that  $g: \Omega_2 \rightarrow \Omega_3$  is Pansu differentiable at  $f(p)$ . Then  $g \circ f$  is Pansu differentiable at  $p$ , and  $D(g \circ f)(p) = (Dg)(f(p))Df(p)$ .

*Proof*

Fix a compact set  $K_1$  in  $G_1$ . Since  $f$  is Pansu differentiable at  $p$ , if  $t \leq 1$  and  $x \in K_1$ , then  $\delta_t^{-1}[f(p)^{-1}f(p\delta_t x)]$  lies in a compact set  $K_2$  in  $G_2$ . Further,  $(Dg)(f(p))y = \lim_{t \rightarrow 0} \delta_t^{-1}[g(f(p))^{-1}g(f(p)\delta_t y)]$  uniformly for  $y \in K_2$ , as  $g$  is Pansu differentiable at  $f(p)$ . Take  $y$  to be  $\delta_t^{-1}[f(p)^{-1}f(p\delta_t x)]$ . Then if  $t \leq 1$ ,

$$\begin{aligned} &\varrho(\delta_t^{-1}[g(f(p))^{-1}g(f(p\delta_t x))], (Dg)(f(p))Df(p)x) \\ &\leq \varrho(\delta_t^{-1}[g(f(p))^{-1}g(f(p\delta_t x))], (Dg)(f(p))\delta_t^{-1}[f(p)^{-1}f(p\delta_t x)]) \\ &\quad + \varrho((Dg)(f(p))\delta_t^{-1}[f(p)^{-1}f(p\delta_t x)], (Dg)(f(p))Df(p)x) \\ &\leq \sup_{y \in K_2} \varrho(\delta_t^{-1}[g(f(p))^{-1}g(f(p)\delta_t y)], (Dg)(f(p))y) \\ &\quad + C\varrho(\delta_t^{-1}[f(p)^{-1}f(p\delta_t x)], Df(p)x) \end{aligned}$$

since  $(Dg)(f(p))$  is Lipschitz, and this converges as needed. □

**4. Quasiconformal maps**

In this section, we suppose that  $\Omega_1$  and  $\Omega_2$  are open subsets of Carnot groups  $G_1$  and  $G_2$  and that  $f: \Omega_1 \rightarrow \Omega_2$  is a homeomorphism. We define the distortion  $H_f(p, t)$  of  $f$  by (1.1) or (3.2).

*Definition 4.1*

The map  $f: \Omega_1 \rightarrow \Omega_2$  is said to be *quasiconformal* if there exists a constant  $\lambda$  such that  $\limsup_{t \rightarrow 0} H_f(p, t) \leq \lambda$  for all  $p \in \Omega_1$ .

Next,  $f$  is said to be *weakly quasisymmetric* if there exists a constant  $H$  such that  $H_f(p, t) \leq H$  for all  $p \in \Omega_1$  and  $t \in \mathbb{R}^+$ .

Finally,  $f$  is said to be *quasisymmetric* if there exists an increasing homeomorphism  $\eta: \mathbb{R}^+ \rightarrow \mathbb{R}^+$  such that  $\varrho(f(p), f(r)) \leq \eta(s)\varrho(f(p), f(q))$  whenever  $p, q, r \in \Omega_1$  and  $\varrho(p, r) \leq s\varrho(p, q)$ .

Each of these definitions has a local version; for instance,  $f$  is locally quasiconformal if it is quasiconformal in a neighborhood of each point.

## THEOREM 4.2

Suppose that  $f: \Omega_1 \rightarrow \Omega_2$  is a quasiconformal map. Then  $f$  is locally quasisymmetric.

For quasiconformal maps on  $\mathbb{R}^n$ , this result is now classical. Heinonen and Koskela proved it for Carnot groups (see [18, Theorem 1.3]) and later for more general manifolds (see [19, Theorem 4.7]). Väisälä [37, Theorem 2.9] had already shown the equivalence of weak quasisymmetry and quasisymmetry.

Obviously, quasisymmetry implies weak quasisymmetry (and  $H = \eta(1)$ ), which in turn implies quasiconformality, so that the local versions of these properties are equivalent. The next result is elementary.

## LEMMA 4.3

Suppose that  $f: \Omega_1 \rightarrow \Omega_2$  is an  $\eta$ -quasisymmetric map. Then  $f^{-1}: \Omega_2 \rightarrow \Omega_1$  is  $\tilde{\eta}$ -quasisymmetric, where  $\tilde{\eta}(s) = [\eta^{-1}(s^{-1})]^{-1}$ .

## COROLLARY 4.4

Suppose that  $f: \Omega_1 \rightarrow \Omega_2$  is a quasiconformal map. Then  $f^{-1}: \Omega_2 \rightarrow \Omega_1$  is locally quasiconformal.

Pansu [28] uses a different definition of quasiconformality.

## Definition 4.5

A homeomorphism  $f: \Omega_1 \rightarrow \Omega_2$  is said to be *Pansu quasiconformal* if  $f$  and  $f^{-1}$  are locally quasisymmetric.\*

In light of Theorem 4.2, Lemma 4.3, and Corollary 4.4, Pansu quasiconformality is equivalent to quasiconformality, at least locally. We now focus on the connections between Pansu differentiability and quasiconformality. The next fundamental result was proved by Pansu [28, Théorème 2, Proposition 7.3].

## THEOREM 4.6

Suppose that  $f: \Omega_1 \rightarrow \Omega_2$  is locally quasiconformal. Then  $f$  is Pansu differentiable almost everywhere. Moreover, the maps  $f$  and  $f^{-1}$  are absolutely continuous; that is,  $|f(E)| = 0$  if and only if  $|E| = 0$ .

Suppose that  $f: \Omega_1 \rightarrow \Omega_2$  is quasiconformal. Then  $f$  and  $f^{-1}$  are absolutely continuous, and we may define a measure  $\mu$  on  $\Omega_1$  by the formula  $\mu(E) = |f(E)|$ . Clearly,  $\mu$  is absolutely continuous, and so it has a Radon-Nikodym derivative, say,  $f'$ . By

\*Pansu apparently requires a uniform choice of  $\eta$ ; this uniformity does not seem to play a significant role.

Lebesgue’s differentiation theorem,

$$f'(p) = \lim_{t \rightarrow 0} \frac{|f(B(p, t))|}{|B(p, t)|}$$

for almost all  $p$  in  $\Omega_1$ . The change of variables formula

$$\int_{\Omega_1} u(f(x))f'(x) dx = \int_{f(\Omega_1)} u(y) dy$$

follows (see Federer [11, Section 2.9]). If  $Df(p)$  and  $f'(p)$  exist, then

$$f'(p) \leq \lim_{t \rightarrow 0} \frac{|(B(f(p), \text{Lip}_f(p)t))|}{|B(p, t)|} = \text{Lip}_f(p)^Q,$$

and similarly,  $f'(p) \geq \text{lip}_f(p)^Q$ . From Lemma 3.3,

$$\begin{aligned} \text{Lip}_f(p)^Q &\geq f'(p) \geq \limsup_{t \rightarrow 0} H_f(p, t)^{-Q} \text{Lip}_f(p)^Q, \\ \text{lip}_f(p)^Q &\leq f'(p) \leq \limsup_{t \rightarrow 0} H_f(p, t)^Q \text{lip}_f(p)^Q. \end{aligned} \tag{4.1}$$

Since  $f'$  is locally integrable,  $\text{Lip}_f \in L^Q_{\text{loc}}(\Omega_1)$ .

### 5. 1-Quasiconformal maps

Again, we assume throughout this section that  $\Omega_1$  and  $\Omega_2$  are open subsets of Carnot groups  $G_1$  and  $G_2$ .

#### Definition 5.1

A homeomorphism  $f : \Omega_1 \rightarrow \Omega_2$  is said to be 1-quasiconformal if  $\lim_{t \rightarrow 0} H_f(p, t) = 1$  for all  $p \in \Omega_1$ .

Further,  $f$  is said to be 1-quasiconformal in the sense of Pansu (which we shorten to  $P1$ -quasiconformal) if  $f$  is locally quasiconformal, and  $Df(p)$  is a similarity (i.e., a product of a dilation and an isometry) for almost all  $p$ .

#### LEMMA 5.2

Suppose that  $f : \Omega_1 \rightarrow \Omega_2$  is Pansu differentiable at  $p$ , and suppose that  $Df(p)$  is a similarity. Then  $df(p)|_{V_1}$  is a similarity. More precisely, for all  $X \in V_1$ , we have  $|df(p)X| = \text{Lip}_f(p)|X|$ .

#### Proof

Since the Pansu derivative  $Df(p)$  is a similarity, from Lemma 3.3 we infer that  $Q(Df(p)x, e) = \text{Lip}_f(p)Q(x, e)$  for all  $x$  in  $G$ . From Lemma 2.2, if  $X \in V_1$ , then  $Q(\exp(df(p)X), e) = \text{Lip}_f(p)Q(\exp X, e)$ . □



## LEMMA 5.3

Suppose that  $f: \Omega_1 \rightarrow \Omega_2$  is 1-quasiconformal. Then  $f$  is also P1-quasiconformal.

*Proof*

Take  $p$  in  $\Omega_1$ , where  $f$  is Pansu differentiable. In view of Lemma 3.3 and of the fact that  $\limsup_{t \rightarrow 0} H_f(p, t) = 1$ ,

$$\min\{\varrho(Df(p)x, e) \mid x \in S(e, 1)\} = \max\{\varrho(Df(p)x, e) \mid x \in S(e, 1)\}.$$

Consequently, for all  $y$  such that  $\varrho(y, e) = t$ ,

$$\varrho(Df(p)y, e) = t \varrho(Df(p)x, e) = t \text{Lip}_f(p) \varrho(x, e) = \text{Lip}_f(p) \varrho(y, e),$$

where  $x = \delta_t^{-1}y$ , which shows that  $Df(p)$  is indeed a similarity.  $\square$

The converse is less obvious. If  $f$  is not smooth, then we may lack information about  $H_f(p, t)$  for  $p$  in a null set. In Corollary 7.2, we show that the two definitions are in fact equivalent.

## LEMMA 5.4

Suppose that  $f: \Omega_1 \rightarrow \Omega_2$  is P1-quasiconformal. Then  $f^{-1}$  is also P1-quasiconformal. Furthermore,  $\text{Lip}_f^{\varrho} = f'$  almost everywhere.

*Proof*

Compare with [28, Lemme 7.11]. By Theorem 4.6,  $f$  and  $f^{-1}$  are absolutely continuous. By the chain rule,  $D(f^{-1})(f(p)) Df(p) = I$  for almost all  $p$ , and the inverse of a similarity is a similarity. Hence  $D(f^{-1})$  is a similarity almost everywhere. The second statement follows from (4.1).  $\square$

The following crucial result is due to Pansu [28, Lemme 11.4] (see the more recent article [1, Theorem 6.6] for a different proof).

## THEOREM 5.5

Suppose that  $f: \Omega_1 \rightarrow \Omega_2$  is P1-quasiconformal. Then  $f$  and  $f^{-1}$  are locally Lipschitz regular; that is, for any compact subset  $\Omega_0$  of  $\Omega_1$ ,

$$\varrho(p, q) \simeq \varrho(f(p), f(q)), \quad \forall p, q \in \Omega_0.$$

Before our next result, recall that  $(X_1 \cdot \log f, \dots, X_{m_1} \cdot \log f)$  is the first layer of  $f$  in exponential coordinates.

PROPOSITION 5.6

Suppose that  $f: \Omega_1 \rightarrow \Omega_2$  is  $P1$ -quasiconformal. For each compact subset  $\Omega_0$  of  $\Omega_1$ , and for all  $i = 1, \dots, m_1$ ,

$$|\nabla_H(X_i \cdot \log f)(p)| \simeq 1 \quad \text{for a.a. } p \text{ in } \Omega_0. \tag{5.1}$$

*Proof*

From (3.1) and Theorem 5.5 applied to  $f$  and  $f^{-1}$ ,  $\text{Lip}_f$  and  $\text{Lip}_f^{-1}$  are locally bounded. The map  $df(p)|_{V_1}$  is an orthogonal map, multiplied by dilation by a factor of  $\text{Lip}_f(p)$ . The norm of any row (or column) of a positive multiple of an orthogonal matrix is the multiplying factor. Thus if we represent  $df(p)$  as a matrix, the norm of any column is  $\text{Lip}_f(p)$ , which is bounded and bounded away from zero.  $\square$

The following result was known (see [17]) for quasiregular maps  $f$ , satisfying the further regularity condition that the distributional derivatives  $\tilde{X}_i f$  (where  $1 \leq i \leq m$ , not just  $1 \leq i \leq m_1$ ) lie locally in  $L^m(G_1)$ . A priori,  $1$ -quasiconformal maps need not satisfy this condition. N. S. Dairbekov [10] found a clever way to eliminate the regularity assumption in the step-two case.

THEOREM 5.7 (Morphism property)

Suppose that  $f: \Omega_1 \rightarrow \Omega_2$  is  $P1$ -quasiconformal. If  $u$  is  $Q$ -harmonic in  $\Omega_2$ , then  $u \circ f$  is  $Q$ -harmonic in  $\Omega_1$ .

*Proof*

For any  $p$  for which  $Df(p)$  is a similarity and any function  $w$  which is Pansu differentiable at  $f(p)$ , we have  $\text{Lip}_{w \circ f}(p) = \text{Lip}_w(f(p))\text{Lip}_f(p)$  and, by Corollary 3.5,

$$\begin{aligned} \text{Lip}_{w \circ f}(p)^2 &= \nabla_H(w \circ f)(p) \cdot \nabla_H(w \circ f)(p), \\ \text{Lip}_w(f(p))^2 &= (\nabla_H w) \circ f(p) \cdot (\nabla_H w) \circ f(p), \end{aligned}$$

whence  $\nabla_H(w \circ f) \cdot \nabla_H(w \circ f) = \text{Lip}_f^2 (\nabla_H w) \circ f \cdot (\nabla_H w) \circ f$  almost everywhere. Polarization of this identity shows that for almost everywhere Pansu differentiable functions  $u$  and  $v$ ,

$$\nabla_H(u \circ f) \cdot \nabla_H(v \circ f) = \text{Lip}_f^2 (\nabla_H u) \circ f \cdot (\nabla_H v) \circ f$$

almost everywhere.

Now we change variables. For any  $\psi \in C_0^{0,1}(\Omega_1)$ , we write  $\phi$  for  $\psi \circ f^{-1}$ ; then  $\phi \in C_0^{0,1}(\Omega_2)$ . By Lemma 3.6, every  $Q$ -harmonic function  $u$  is Pansu differentiable

almost everywhere From Lemma 5.4,  $f' = \text{Lip}_f^\varrho$ , and

$$\begin{aligned} & \int_{\Omega_1} |\nabla_H(u \circ f)(x)|^{\varrho-2} \nabla_H(u \circ f)(x) \cdot \nabla_H \psi(x) \, dx \\ &= \int_{\Omega_1} |\nabla_H(u \circ f)(x)|^{\varrho-2} \nabla_H(u \circ f)(x) \cdot \nabla_H(\phi \circ f)(x) \, dx \\ &= \int_{\Omega_1} \text{Lip}_f^\varrho(x)^\varrho |(\nabla_H u)(f(x))|^{\varrho-2} (\nabla_H u)(f(x)) \cdot (\nabla_H \phi)(f(x)) \, dx \\ &= \int_{f(\Omega_1)} |(\nabla_H u)(y)|^{\varrho-2} (\nabla_H u)(y) \cdot (\nabla_H \phi)(y) \, dy \\ &= 0 \end{aligned}$$

since  $u$  is a weak solution of (2.3). □

**COROLLARY 5.8** (Regularity of the first layer of 1-quasiconformal maps)  
*Suppose that  $f : \Omega_1 \rightarrow \Omega_2$  is P1-quasiconformal. Then  $\pi_1 \log f \in C^\infty(\Omega_1)$ .*

*Proof*

When  $1 \leq i \leq m_1$ , the coordinate functions  $X_i \cdot \log$  are  $Q$ -harmonic. Hence the functions  $X_i \cdot \log f$  are too, by Theorem 5.7. The conclusion follows from (5.1) and Theorem 2.1. □

### 6. Algebraic regularity

**THEOREM 6.1**

*Suppose that  $\Omega_1$  and  $\Omega_2$  are open subsets of Carnot groups  $G_1$  and  $G_2$ ; that  $\pi_1 \log f \in C_{\text{loc}}^{k,\beta}(\Omega_1, \mathfrak{g}_2)$ , where  $k \geq 1$  and  $\beta \in [0, 1]$ ; and that  $f \in C_{\text{loc}}^{0,\alpha}(\Omega_1, \Omega_2)$ , where  $\alpha > 1/2$ . Then  $\log f \in C_{\text{loc}}^{k,\beta}(\Omega_1, \mathfrak{g}_2)$ .*

Note that the theorem does not hold if  $\alpha = 1/2$ ; to see this, consider the map  $f : (x_1, x_2, x_3) \mapsto (x_3, x_2, |x_1|)$  in the Heisenberg group  $\mathbb{H}^1$ . This map is not contact but is locally  $C^{0,1/2}$ .

To explain our ideas, we first present the proof of Theorem 6.1 in the setting of the Heisenberg group. Take open subsets  $\Omega_1$  and  $\Omega_2$  of  $\mathbb{H}^1$  and  $f : \Omega_1 \rightarrow \Omega_2$ , as in the statement of the theorem. Now  $f \in C_{\text{loc}}^{0,\alpha}(\Omega_1, \Omega_2)$ , so  $\varrho(f(p \exp t X_1), f(p)) \leq Ct^\alpha$  for small  $t$ . In coordinates, we write  $f(p)$  as  $(f_1(p), f_2(p), f_3(p))$ . Using the Nagel-Stein-Wainger pseudometric and the group law, as in the proof of Lemma 3.4, we see that

$$\left| f_3(p \exp t X_1) - f_3(p) + \frac{f_1(p \exp t X_1)f_2(p) - f_2(p \exp t X_1)f_1(p)}{2} \right| \leq Ct^{2\alpha}.$$

We approximate the  $C^{1,\beta}$ -numerator of the fraction, using its first-order Taylor polynomial in  $t$ , and find that

$$\left| f_3(p \exp t X_1) - f_3(p) + \frac{t}{2} (f_2(p) \tilde{X}_1 f_1(p) - f_1(p) \tilde{X}_1 f_2(p)) \right| = o(t).$$

Thus  $\tilde{X}_1 f_3(p)$  exists and is equal to  $(f_1(p) \tilde{X}_1 f_2(p) - f_2(p) \tilde{X}_1 f_1(p))/2$ . By hypothesis,  $f_1, f_2 \in C^{k,\beta}(\Omega_1)$ , and so  $\tilde{X}_1 f_3 \in C^{k-1,\beta}(\Omega_1)$ . The same argument is applied to the vector field  $\tilde{X}_2$  to complete the proof.

*Proof of Theorem 6.1*

We show by induction that if  $\pi_j \log f \in C^{k,\beta}(\Omega_1, \mathfrak{g}_2)$  when  $j = 1, 2, \dots, l-1$ , where  $l \geq 2$ , then  $\pi_l \log f \in C^{k,\beta}(\Omega_1, \mathfrak{g}_2)$ .

Fix a point  $p$  in  $\Omega_1$ , and fix  $X$  in  $V_1$ . Since  $f \in C_{\text{loc}}^{0,\alpha}(\Omega_1, \Omega_2)$ , there exists an interval  $I$  containing zero, and a constant  $C$ , such that

$$\varrho(f(p \exp t X), f(p)) \leq C \varrho(p \exp t X, p)^\alpha = C \varrho(\exp t X, e)^\alpha = C |t X|^\alpha$$

for all  $t \in I$ . Write  $f(q)$  as  $\exp(F(q))$ . Then

$$\begin{aligned} \varrho(f(p \exp t X), f(p)) &= \varrho(\exp(-F(p)) \exp(F(p \exp t X)), e) \\ &= \varrho(\exp \text{BCH}(-F(p), F(p \exp t X)), e) \\ &\simeq \left( \sum_{k=1}^R |\pi_k \text{BCH}(-F(p), F(p \exp t X))|^{2R!/k} \right)^{1/2R!}. \end{aligned}$$

From the hypothesis, it follows that

$$|\pi_l \text{BCH}(-F(p), F(p \exp t X))| \leq C |t|^{l\alpha} \quad \forall t \in I. \tag{6.1}$$

Write  $F_1(q)$  for  $\pi_1 F(q) + \dots + \pi_{l-1} F(q)$ ,  $F_2(q)$  for  $\pi_l F(q)$ , and  $F_3(q)$  for  $\pi_{l+1} F(q) + \dots + \pi_R F(q)$ . As  $\mathfrak{g}_2$  is graded,  $\pi_l \text{BCH}(-F(p), F(p \exp t X))$  is equal to

$$F_2(p \exp t X) - F_2(p) + \pi_l \text{BCH}(-F_1(p), F_1(p \exp t X)), \tag{6.2}$$

and  $\pi_l \text{BCH}(-F_1(p), F_1(p \exp t X))$  is a finite weighted sum of projections of commutators; assuming that  $l \geq 4$ , a typical term is

$$\pi_l [F_1(p), [F_1(p \exp t X), [F_1(p), F_1(p \exp t X)]]].$$

By hypothesis,  $F_1 \in C^{k,\beta}(\Omega_1, \mathfrak{g}_2)$ , so  $F_1(p \exp t X) = F_1(p) + t \tilde{X} F_1(p) + o(t)$ . Substituting this into our typical commutator, we obtain

$$t \pi_l [F_1(p), [F_1(p), [F_1(p), \tilde{X} F_1(p)]]] + o(t).$$

In general, we obtain  $t$  multiplied by a commutator involving  $F_1(p)$  (perhaps more than once) and  $\tilde{X}F_1(p)$  (once only), and higher-order terms. From (6.1), (6.2), and our discussion of  $\pi_l \text{BCH}(-F_1(p), F_1(p \exp tX))$ , we deduce that  $\pi_l \text{BCH}(-F_1(p), F_1(p \exp tX))$  is equal to

$$t \frac{d}{dt} \pi_l \text{BCH}(-F_1(p), F_1(p \exp tX)) \Big|_{t=0} + o(t),$$

and hence,

$$\left| F_2(p \exp tX) - F_2(p) + t \frac{d}{dt} \pi_l \text{BCH}(-F_1(p), F_1(p \exp tX)) \Big|_{t=0} \right| = o(t).$$

From [20, Chapter II, Theorem 1.7],

$$\frac{d}{dt} \text{BCH}(-A, A + tB) \Big|_{t=0} = \frac{\exp(-\text{ad}A) - I}{\text{ad}A} B,$$

whence

$$\begin{aligned} \tilde{X}F_2(p) &= -\frac{d}{dt} \pi_l \text{BCH}(-F_1(p), F_1(p \exp tX)) \Big|_{t=0} \\ &= -\sum_{n=1}^{\infty} \frac{(\text{ad}F_1(p))^{n-1} X F_1(p)}{n!}. \end{aligned}$$

The latter, in turn, is a weighted sum of commutators such as (for instance)  $[F_1(p), [F_1(p), [F_1(p), \tilde{X}F_1(p)]]]$ , so that  $\tilde{X}F_2 \in C^{k-1, \beta}(\Omega_1)$ . Since  $X$  is an arbitrary element of  $V_1$ , we conclude that  $F_2 \in C^{k, \beta}(\Omega_1)$ .  $\square$

## 7. Proof of Theorem 1.1 and applications

We assume throughout this section that  $\Omega_1$  and  $\Omega_2$  are open subsets of Carnot groups  $G_1$  and  $G_2$ .

### *Proof of Theorem 1.1*

Take a  $P1$ -quasiconformal map  $f: \Omega_1 \rightarrow \Omega_2$ . Then  $X_i \cdot \log f$  is smooth when  $1 \leq i \leq m_1$ , by Corollary 5.8, and  $f$  is locally Lipschitz, by Theorem 5.5. Apply Theorem 6.1.  $\square$

### *Definition 7.1*

The diffeomorphism  $f: \Omega_1 \rightarrow \Omega_2$  is said to be *conformal* if it is Pansu differentiable and  $df$ , restricted to  $V_1$ , is a similarity everywhere.

COROLLARY 7.2

For a map  $f : \Omega_1 \rightarrow \Omega_2$ , the following are equivalent:

- (i)  $f$  is 1-quasiconformal;
- (ii)  $f$  is P1-quasiconformal;
- (iii)  $f$  is smooth and conformal.

*Proof*

Lemma 5.3 shows that (i) implies (ii). Conversely, suppose that  $f$  is P1-quasiconformal. By Lemma 5.4, so is  $f^{-1}$ . By Theorem 1.1,  $f$  and  $f^{-1}$  are Pansu differentiable everywhere, and so  $Df$  is invertible, by the chain rule. Apply Lemma 3.3. The equivalence of (ii) and (iii) follows from Theorem 1.1 and (2.1). □

Pansu [28, Proposition 13.1] gives examples of two-step Carnot groups whose Carnot automorphisms are all dilations. We show that all quasiconformal maps of domains in these groups are compositions of translations and dilations. Pansu was clearly aware of this result (see, e.g., [29] and [28, Théorème 4]).

LEMMA 7.3

Suppose that  $\Omega_1$  and  $\Omega_2$  are connected open subsets of a Carnot group  $G$  of step at most 2. Suppose that  $f : \Omega_1 \rightarrow \Omega_2$  is quasiconformal, and suppose that  $Df$  is a dilation almost everywhere in  $\Omega_1$ . Then  $f$  is a translation composed with a dilation.

*Proof*

By hypothesis,  $f$  is P1-quasiconformal, and hence, smooth. Thus  $Df$  is a dilation everywhere. Write  $Df(p)$  as  $\delta_{\alpha(p)}$ . We show first that  $\alpha$  is constant.

Take linearly independent  $X$  and  $Y$  in  $V_1$ . Then  $X$  and  $Y$  generate a two- or three-dimensional subalgebra  $\mathfrak{n}(X, Y)$  of  $\mathfrak{g}$ , depending on whether or not  $[X, Y] = 0$ . Write  $N(X, Y)$  for  $\exp(\mathfrak{n}(X, Y))$ . The vector fields  $\tilde{Z}$  corresponding to  $Z$  in  $\mathfrak{n}(X, Y)$  give rise to an integrable distribution on  $G$ , and  $G$  is foliated by the cosets of  $N(X, Y)$ . The differential of  $f$  maps  $(\tilde{X})_p$  and  $(\tilde{Y})_p$  to multiples of  $(\tilde{X})_{f(p)}$  and  $(\tilde{Y})_{f(p)}$ , so  $f$  respects the foliation and maps connected components of leaves to connected components of leaves.

Fix  $p$  in  $\Omega_1$ , and restrict  $f$  to the leaf through  $p$ . By composing  $f$  with translations, we may suppose that  $p = e$  and  $f(p) = e$ . Thus, at least locally,  $f$  is a map of  $N(X, Y)$  into itself, whose Pansu derivative is a dilation at each point. Suppose that  $N(X, Y) = \mathbb{R}^2$ . We write the restricted map  $f$  in coordinates, and then

$$\begin{pmatrix} \frac{\partial f_1}{\partial x_1} & \frac{\partial f_1}{\partial x_2} \\ \frac{\partial f_2}{\partial x_1} & \frac{\partial f_2}{\partial x_2} \end{pmatrix} = \begin{pmatrix} \alpha & 0 \\ 0 & \alpha \end{pmatrix}.$$

It follows that  $\partial\alpha/\partial x_1 = \partial^2 f_2/\partial x_1 \partial x_2 = 0$ , and similarly,  $\partial\alpha/\partial x_2 = 0$ . We conclude that  $\alpha$  is locally constant on the cosets of  $N(X, Y)$  in this case.

If  $n(X, Y)$  is three-dimensional, then the restriction of  $f$  to  $N(X, Y)$  is a 1-quasiconformal map between domains of the Heisenberg group; hence by [22, Theorem 8], it is the restriction of the action of an element of  $SU(1, 2)$ . These actions are described in [22, Section E], and those whose Pansu differential is a dilation at every point are made of dilations and translations. Thus  $\alpha$  is locally constant on the cosets of  $N(X, Y)$  in this case too.

It follows that  $\tilde{X}\alpha = 0$  on  $\Omega_1$ . But  $X$  was an arbitrary element of  $V_1 \setminus \{0\}$ , so  $\nabla_H \alpha = 0$  and  $\alpha$  is constant.

By composing with  $\delta_\alpha^{-1}$ , we may assume that  $Df = \delta_1$  everywhere in  $\Omega_1$ . By again considering the leaves of the foliation associated to  $N(X, Y)$ , we see that  $f$  acts as a translation on each leaf. This means that if  $q \in \Omega_1$  and  $f(q_1) = q_2$ , then  $f(q_1 \exp(tX)) = q_2 \exp(tX)$  for all sufficiently small  $t$ .

Suppose that  $X_1, \dots, X_N \in V_1$ , and suppose that  $f(q_1) = q_2$ . By iterating the preceding argument, it may be seen that

$$f(q_1 \exp(t_1 X_1) \cdots \exp(t_N X_N)) = q_2 \exp(t_1 X_1) \cdots \exp(t_N X_N)$$

for all sufficiently small  $t_1, \dots, t_N$ . From [12, Lemma 5.1], it follows that  $f(q_1 x) = q_2 x$  for all  $x$  in some neighborhood of the identity in  $G_1$ .

Let  $\Omega_0$  be  $\{p \in \Omega_1 \mid f(p) = q_2 q_1^{-1} p\}$ . Then  $\Omega_0$  is clearly nonempty and closed and, by the argument above, is also open, so it is all  $\Omega_1$ . This shows that  $f$  is a translation on  $\Omega_1$ .  $\square$

#### COROLLARY 7.4

*Suppose that  $G$  is a step-two Carnot group whose Carnot automorphisms are all dilations. Then the 1-quasiconformal maps between domains of  $G$  are all translations composed with dilations.*

Before we state our final result, we recall that the so-called  $H$ -type groups (see [8], [21]) are important examples of Carnot groups.

#### COROLLARY 7.5

*Suppose that  $G$  is an  $H$ -type group whose Lie algebra has center of dimension larger than 2. The 1-quasiconformal maps between open subsets of  $G$  are smooth and form a finite-dimensional space.*

#### *Proof*

This follows from the work of Reimann [30], who established the corresponding infinitesimal result. Our main theorem allows us to obtain information about 1-quasiconformal maps by the standard method.  $\square$

For many  $H$ -type groups, all 1-quasiconformal maps are composed of translations and dilations. However, for the Iwasawa  $N$ -groups of real rank-one simple Lie groups, the group of 1-quasiconformal maps is like the Möbius group and contains inversions as well.

*Acknowledgments.* The authors thank J. Heinonen and H. M. Reimann for their interest in this work and for many useful discussions.

## References

- [1] Z. M. BALOGH, I. HOLOPAINEN, and J. T. TYSON, *Singular solutions, homogeneous norms, and quasiconformal mappings in Carnot groups*, *Math. Ann.* **324** (2002), 159–186. [MR 1931762](#) [457, 470](#)
- [2] L. CAPOGNA, *Regularity for quasi-linear equations in the Heisenberg group*, *Comm. Pure Appl. Math.* **50** (1997), 867–889. [MR 1459590](#) [456](#)
- [3] ———, *Regularity for quasilinear equations and 1-quasiconformal maps in Carnot groups*, *Math. Ann.* **313** (1999), 263–295. [MR 1679786](#) [457, 460](#)
- [4] É. CARTAN, *Sur la géométrie pseudo-conforme des hypersurfaces de l'espace de deux variables complexes, II*, *Ann. Scuola Norm. Sup. Pisa* (2) **1** (1932), 333–354. [456](#)
- [5] S.-S. CHERN and J. K. MOSER, *Real hypersurfaces in complex manifolds*, *Acta Math.* **133** (1974), 219–271. [MR 0425155](#) [456](#)
- [6] T. COULHON, I. HOLOPAINEN, and L. SALOFF-COSTE, *Harnack inequality and hyperbolicity for subelliptic  $p$ -Laplacians with applications to Picard type theorems*, *Geom. Funct. Anal.* **11** (2001), 1139–1191; *Erratum*, *Geom. Funct. Anal.* **12** (2001), 217. [MR 1878317](#); [MR 1904564](#) [457](#)
- [7] M. COWLING, F. DE MARI, A. KORÁNYI, and H. M. REIMANN, “Contact and conformal maps on Iwasawa  $N$  groups” in *Harmonic Analysis on Complex Homogeneous Domains and Lie Groups (Rome, 2001)*, *Atti Accad. Naz. Lincei Cl. Sci. Fis. Mat. Natur. Rend. Lincei* (9) *Mat. Appl.* **13**, Acad. Naz. Lincei, Rome, 2002, 219–232. [MR 1984102](#) [457](#)
- [8] M. COWLING, A. H. DOOLEY, A. KORÁNYI, and F. RICCI,  *$H$ -type groups and Iwasawa decompositions*, *Adv. Math.* **87** (1991), 1–41. [MR 1102963](#) [456, 476](#)
- [9] M. COWLING and H. M. REIMANN, “Quasiconformal mappings on Carnot groups: Three examples” in *Harmonic Analysis at Mount Holyoke (South Hadley, Mass., 2001)*, *Contemp. Math.* **320**, Amer. Math. Soc., Providence, 2003, 111–118. [MR 1979935](#) [457](#)
- [10] N. S. DAIRBEKOV, *The morphism property for mappings with bounded distortion on the Heisenberg group*, *Siberian Math. J.* **40** (1999), 682–694. [MR 1721676](#) [457, 471](#)
- [11] H. FEDERER, *Geometric Measure Theory*, *Grundlehren Math. Wiss.* **153**, Springer, New York, 1969. [MR 0257325](#) [469](#)
- [12] G. B. FOLLAND, *Subelliptic estimates and function spaces on nilpotent Lie groups*, *Ark. Mat.* **13** (1975), 161–207. [MR 0494315](#) [460, 476](#)



- [13] G. B. FOLLAND and E. M. STEIN, *Hardy Spaces on Homogeneous Groups*. Math. Notes **28**, Princeton Univ. Press, Princeton, 1982. MR 0657581 458, 460
- [14] F. W. GEHRING, *Rings and quasiconformal mappings in space*, Trans. Amer. Math. Soc. **103** (1962), 353–393. MR 0139735 455, 457
- [15] J. HEINONEN, *A capacity estimate on Carnot groups*, Bull. Sci. Math. **119** (1995), 475–484. MR 1354248
- [16] ———, “Calculus on Carnot groups” in *Fall School in Analysis (Jyväskylä, Finland, 1994)*, Report **68**, Univ. Jyväskylä, Jyväskylä, Finland, 1995, 1–31. MR 1351042
- [17] J. HEINONEN and I. HOLOPAINEN, *Quasiregular maps on Carnot groups*, J. Geom. Anal. **7** (1997), 109–148. MR 1630785 464, 466, 471
- [18] J. HEINONEN and P. KOSKELA, *Definitions of quasiconformality*, Invent. Math. **120** (1995), 61–79. MR 1323982 455, 457, 468
- [19] ———, *Quasiconformal maps in metric spaces with controlled geometry*, Acta Math. **181** (1998), 1–61. MR 1654771 468
- [20] S. HELGASON, *Differential Geometry, Lie Groups, and Symmetric Spaces*, Pure Appl. Math. **80**, Academic Press, New York, 1978. MR 0514561 461, 474
- [21] A. KAPLAN, *Fundamental solutions for a class of hypoelliptic PDE generated by composition of quadratic forms*, Trans. Amer. Math. Soc. **258** (1980), 147–153. MR 0554324 456, 476
- [22] A. KORÁNYI and H. M. REIMANN, *Foundations for the theory of quasiconformal mappings on the Heisenberg group*, Adv. Math. **111** (1995), 1–87. MR 1317384 456, 476
- [23] ———, *Quasiconformal mappings on the Heisenberg group*, Invent. Math. **80** (1985), 309–338. MR 0788413 456
- [24] G. A. MARGULIS and G. D. MOSTOW, *The differential of a quasi-conformal mapping of a Carnot-Carathéodory space*, Geom. Funct. Anal. **5** (1995), 402–433. MR 1334873 455
- [25] G. D. MOSTOW, *Strong Rigidity of Locally Symmetric Spaces*, Ann. of Math. Stud. **78**, Princeton Univ. Press, Princeton, 1973. MR 0385004 456
- [26] A. NAGEL, E. M. STEIN, and S. WAINGER, *Balls and metrics defined by vector fields, I. Basic properties*, Acta Math. **155** (1985), 103–147. MR 0793239 459
- [27] R. NEVANLINNA, “On differentiable mappings” in *Analytic Functions*, Princeton Math. Ser. **24**, Princeton Univ. Press, Princeton, 1960, 3–9. MR 0116280 455
- [28] P. PANSU, *Métriques de Carnot-Carathéodory et quasiisométries des espaces symétriques de rang un*, Ann. of Math. (2) **129** (1989), 1–60. MR 0979599 456, 457, 468, 470, 475
- [29] ———, “Quasiconformal mappings and manifolds of negative curvature” in *Curvature and Topology of Riemannian Manifolds (Katata, Japan, 1985)*, Lecture Notes in Math. **1201**, Springer, Berlin, 1986, 212–229. MR 0859587 475
- [30] H. M. REIMANN, *Rigidity of H-type groups*, Math. Z. **237** (2001), 697–725. MR 1854087 456, 457, 476
- [31] H. M. REIMANN and F. RICCI, “The complexified Heisenberg group” in *Proceedings on Analysis and Geometry (Novosibirsk Akademgorodok, Russia, 1999)*, Izdat.

- Ross. Akad. Nauk Sib. Otd. Inst. Mat., Novosibirsk, Russia, 2000, 465–480. [MR 1847532](#) [456](#)
- [32] JU. G. REŠETNJAK [YU. G. RESHETNYAK], *Liouville's conformal mapping theorem under minimal regularity hypotheses* (in Russian), *Sibirsk. Mat. Ž.* **8** (1967), 835–840; English translation in *Sib. Math. J.* **8** (1967), 631–634. [MR 0218544](#) [455](#)
- [33] E. M. STEIN, *Harmonic Analysis: Real-Variable Methods, Orthogonality, and Oscillatory Integrals*, Princeton Math. Ser. **43**, Princeton Univ. Press, Princeton, 1993. [MR 1232192](#)
- [34] N. TANAKA, *On the pseudo-conformal geometry of hypersurfaces of the space of  $n$  complex variables*. *J. Math. Soc. Japan* **14** (1962), 397–429. [MR 0145555](#) [456](#)
- [35] P. TANG, *Regularity and extremality of quasiconformal homeomorphisms on CR 3-manifolds*, *Ann. Acad. Sci. Fenn. Math.* **21** (1996), 289–308. [MR 1404088](#) [456](#)
- [36] J. VÄISÄLÄ, *Lectures on  $n$ -Dimensional Quasiconformal Mappings*, Lecture Notes in Math. **229**, Springer, Berlin, 1971. [MR 0454009](#) [455](#)
- [37] ———, *Quasiconformal maps of cylindrical domains*, *Acta Math.* **162** (1989), 201–225. [MR 0989396](#) [468](#)
- [38] S. K. VODOP'YANOV, *Monotone functions and quasiconformal mappings on Carnot groups*, *Siberian Math. J.* **37** (1996), 1113–1136. [MR 1440383](#) [457](#)

### Capogna

Department of Mathematics, University of Arkansas, Fayetteville, Arkansas 72701, USA;  
[lcapogna@uark.edu](mailto:lcapogna@uark.edu)

### Cowling

School of Mathematics and Statistics, University of New South Wales, Sydney New South Wales 2052, Australia; [m.cowling@unsw.edu.au](mailto:m.cowling@unsw.edu.au)