



Proceeding Paper

How Does Mycorrhiza Interact with Different Levels of Fertilization on *Prosopis alba*? †

María Gracia Senilliani ^{1,*} , Analía del Valle Guzmán ¹, Adriana Teresita Gomez ², Ana Carolina Santacruz-García ³ , Monica Sagadin ⁴, Mauricio Ewens ⁵, Cristian Coria ¹, Joaquín Gómez ¹, Rodrigo Frias ¹ and Mónica Azucena Nazareno ³

¹ Instituto de Silvicultura y Manejo de Bosques INSIMA, Facultad de Ciencias Forestales, Universidad Nacional de Santiago del Estero, Santiago del Estero C.P. 4200, Argentina

² Estación Experimental Agropecuaria Santiago del Estero, Instituto Nacional de Tecnología Agropecuaria INTA, Santiago del Estero C.P. 4200, Argentina

³ Consejo Nacional de Investigaciones Científicas y Técnicas CONICET and Instituto de Ciencias Químicas, Facultad de Agronomía y Agroindustrias, Universidad Nacional de Santiago del Estero, Santiago del Estero C.P. 4200, Argentina

⁴ Instituto de Fisiología y Recursos Genéticos Vegetales (IFRGV), Centro de Investigaciones Agropecuarias (CIAP), Instituto Nacional de Tecnología Agropecuarias (INTA), Unidad de Estudios Agropecuarios (UDEA) INTA-CONICET, Córdoba C.P. 5000, Argentina

⁵ Estación experimental Fernández, Universidad Católica de Santiago del Estero (UCSE), Santiago del Estero C.P. 4200, Argentina

* Correspondence: senilliani@gmail.com

† Presented at the 3rd International Electronic Conference on Forests—Exploring New Discoveries and New Directions in Forests, 15–31 October 2022. Available online: <https://iecf2022.sciforum.net/>.



Citation: Senilliani, M.G.; Guzmán, A.d.V.; Gomez, A.T.; Santacruz-García, A.C.; Sagadin, M.; Ewens, M.; Coria, C.; Gómez, J.; Frias, R.; Nazareno, M.A. How Does Mycorrhiza Interact with Different Levels of Fertilization on *Prosopis alba*? *Environ. Sci. Proc.* **2022**, *22*, 31. <https://doi.org/10.3390/IECF2022-13088>

Academic Editor: Rodolfo Picchio

Published: 21 October 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Abstract: This study assessed how the interactions between chemical fertilization levels and arbuscular mycorrhizal fungus species (AMF) affect the seedling's morphology and biochemical traits in *Prosopis alba*. Subsequently, the seedlings were inoculated or not with a mixture of native AMF from two origins of contrasting sites in the Chaco Region. Preliminarily, we observed a positive mycorrhizal response to the AMF in interaction with chemical fertilization (at 60% fertilizer concentration). These results were reflected mainly over some morphological parameters than over biochemical parameters. The lack of a clear answer is probably since the benefits of the symbiosis on the host plant could be observed in the establishment phase in the field.

Keywords: arbuscular mycorrhizal fungus; biofertilizers; biochemical parameters; mycorrhizal response; morphological parameters; quality seedlings

1. Introduction

The production and application of microbial fertilizers are gaining global increase owing to the negative impacts of overuse or improper usage of chemical-based fertilizer and the increased awareness about the association between rhizosphere microorganisms and plants [1–4]. AMF have even shown a significant potential to be used as a biofertilizer to improve the initial growth and establishment of tree seedlings in restoration projects [5]. Inoculated plants could show higher growth rates, lower resource needs, and may be more tolerant to transplanting stress, critical features in restoring degraded areas [6]. In this sense, the study species *Prosopis alba* has considerable potential as native species in restoring areas in the Chaco Semi-arid Region [7]. AMF are soil-borne fungi that could significantly improve plant nutrient uptake and resistance to several abiotic stress factors [5,8]. AMF hyphae exclusively colonize the root cortex and form highly branched structures inside the cells, i.e., arbuscules, which are considered the functional site of nutrient exchange [9]. Therefore, we aimed to evaluate the effect of the interactions between the levels of chemical

fertilization and the inoculum of arbuscular mycorrhizal fungi (AMF) on the production of *P. alba* during the nursery stage.

2. Materials and Methods

2.1. Plant Material and Experimental Conditions

The experiment was carried out at the Experimental Station “Fernández” (Agreement Catholic University of Sgo del Estero-Province of Santiago del Estero) in Santiago del Estero, Argentina (−27°56'0 S, 65°52.50' W). An experiment was conducted from 28 October 2021 to 28 January 2022. Seedlings were produced in trays of individual cells in a nursery with 50% of shading under natural light conditions for 45 days. After that, the plants were exposed to full sun in the acclimation phase until completing 90 days. The levels of chemical fertilization applied according to concentration were: F0%, F30%, F60% and F100%, varying from no fertilizer application to maximum concentration. The foliar fertilizer was applied by spraying once a week in the first hours of the day. In the nursery phase, “YOGUEN N° 3” fertilizer (NPK 25-14-8) was used in a proportion of 5 g/L (F100). In the acclimation phase, “HAKAPHOS Base” fertilizer (NPK 7-12-40) was used in a proportion of 2 g/L (F100). A control situation without application of mycorrhizae (SM) was compared with two inoculum of arbuscular mycorrhizal fungi (M1 and M2) associated with two different rainfall patterns (explained in Section 2.2).

2.2. Mycorrhizal: Isolation, Multiplication and Application of AMF Inocula

Mixed inocula of native AMF were selected from *P. alba* stands located in the Argentine Chaco Region, with two different rainfall regimes: Padre Lozano (PL = M1) in the Western Chaco Domain, Salta province, with 650 mm annual precipitation, and Colonia Benítez (CB = M2) in the Eastern Chaco Domain, Chaco province, with annual precipitation of 1300 mm [10]. Six soil samples per tree were collected at a depth of 20 cm. The AMF inoculum is composed of mycorrhizal roots, spores and hyphae of alfalfa (*Medicago sativa*) and sorghum (*Sorghum sp.*) plants. As trap plants, this inoculum was initially used with “Algarrobo blanco” (*Prosopis alba*). M1: *Claroideogloium claroideum*, *Claroideogloium etunicatum*; *Diversispora spurca*, *Funneliformismosseae* and *Rhizophagus intraradices*; M2: *Claroideogloium claroideum*, *Claroideogloium etunicatum*, *Funneliformis constrictum*, *Funneliformis mosseae*, *Rhizophagus clarus* [11]. Inoculation was performed at sowing by applying 20 g of AMF inoculum (consisting of a mixture of substrate, spores, mycelium and fragments of mycorrhizal roots of trap plants) in the planting hole per container for inoculum (M), and a control treatment was not inoculated (NI = SM) [12].

2.3. Morphological Characteristics

Morphological traits were measured to evaluate the effects of the treatments on plants. Ten plants per treatment were selected and registered: (i) stem neck diameter (SND), (ii) shoot height (SH) and (iii) slenderness index (SI) according to standard protocols. The slenderness index was determined considering the diameter and shoot height ratio. For biomass evaluation, three plants were randomly selected and each plant was divided into shoots and roots. Root-balls were carefully washed to remove substrate L and weighed immediately. The material was dried to constant weight in an oven at 72 °C (±5 °C) for 48 h to determine dry weight. Data were processed and registered the aerial dry weight (ADW) and root dry weight (RDW).

2.4. Mycorrhizal Response

The Mycorrhizal Response (MR) was calculated for the diameter (SND) according to the relationship described by Cavagnaro et al. (2003) [13]:

$$MR = [(M - \text{media NM}) / \text{media NM}] \times 100]$$

M corresponds to the diameter parameter of the inoculated plants, and the mean NM corresponds to the non-inoculated plants.

2.5. Biochemical Characteristics

For this assay, chlorophylls and carotenoids contents were evaluated according to methodology followed by Santacruz-Garcia et al. (2022) [14]. Leaf samples from three plants per treatment were collected for this assay.

2.6. Experimental Design and Statistical Analysis

The experiment was performed using a factorial design consisting of two factors, the AMF inoculum with three levels (two different AMF inoculation strains and control) and the chemical fertilizer with four levels (three different concentrations and a control). The treatments were distributed entirely randomized. For assessments of biochemical and morphological responses to the treatment application, data were analyzed by an ANOVA using a factorial model. The variable MR was evaluated for the test non-parametric Kruskal–Wallis. The statistical software used was Infostat/2017 (with an $\alpha = 0.05$).

3. Results and Discussion

3.1. Evaluation of the Interaction Effect of AMF with Chemical Fertilizer

3.1.1. Morphological Response

The growth of SND showed an interaction between factors ($p < 0.0002$). However, SH was not significantly affected by fertilizer and mycorrhiza factors or the interaction between factors. The SND was significantly lower in SMF60 treatment (value 3.07 mm) than in all other seedlings. However, M1F60 and M1F30 showed the best performance of SND (3.46 mm), stimulating plant growth compared to M1F0 and M1F100 (3.1 to 3.2 mm). No significant differences were detected between M2 and SM, except SMF100 with SND similar to M1F60 or M1F30. It could be said that M1 inoculum in determined concentrations had a positive effect. No significant differences between the slenderness index and ADW were detected among treatments. RDW showed no significant effects between interaction or fertilizer. However, the mycorrhiza factor showed significant differences in RDW ($p < 0.02$). This M2 treatment showed the lowest values (0.73 g) compared to SM (1.13 g) (Table 1). The inoculation with AMF has not always resulted in an enhancement for the plants. Thus, diverse authors reported a growth depression attributed to different reasons [15,16]. Often this behavior is attributed to AMF parasitism, where carbon demand from the fungus exceeds the benefits of increased nutrient uptake [6,17]. The application of the inoculum showed a more pronounced effect on the growth of diameter than the development of roots.

Table 1. Mean and standard deviation for the morphological variables: slenderness index (SI), root dry weight (RDW, g), aerial dry weight (ADW, g). Stars indicate the significance level. Significance levels: * < 0.05 ; < 0.1 . Different letters indicate significant differences, according to Kruskal–Wallis test with $\alpha: 0.05$.

Treatments	SM	M1	M2
Morphological variables			
SI	9.4 ± 0.3 A	9.1 ± 0.3 A	8.9 ± 0.3 A
RDW *	1.13 ± 0.1 A	0.84 ± 0.1 AB	0.73 ± 0.1 B
ADW	1.71 ± 0.1 A	1.71 ± 0.1 A	1.75 ± 0.1 A

SM: without mycorrhizal treatment, M1: Padre Lozano inoculum, M2: Colonia Benitez inoculum.

3.1.2. Mycorrhizal Response

The mycorrhizal response (MR) was evaluated. Results suggest that there were differences between inoculum (AMF). For example, this response on stem neck diameter was significantly higher ($p < 0.0001$) in the treatment inoculated with M1 and 60% fertilizer concentration (M1F60) followed by M1F30, M2F60 and M2F30. We observed a positive MR in this concentration (60%). However, a negative response at 0% and 100% fertilizer concentrations for both inoculum were detected (Figure 1). This coincides with Malusa et al.

(2007) [15], which considered that low-to-moderate substrate fertility favors AMF formation and increases the supply of mineral nutrients, particularly from minimally soluble sources.




Figure 1. Mycorrhizal response of stem neck diameter for AMF and level chemical fertilizer. Different letters indicate significant differences according to the Kruskal–Wallis test.

3.1.3. Biochemical Response

Although no positive effect was found in the interaction of AMF and chemical fertilizer on photosynthetic pigments, the biochemical response was significantly affected by themycorrhizal factor. The SM treatment showed significantly higher contents of photosynthetic pigments chlorophylls ($p < 0.01$), carotenoids and xanthophylls ($p < 0.0002$) (Table 2). These results could be related to higher physiological stability of the seedlings of the SM treatment compared to other treatments. Photosynthetic pigments could be considered an indicator of the stability of plant regulatory functions [18]. Although previous studies have reported that the effect of AM fungi improved chlorophyll content in leaves under drought or salt stress conditions [19], the mycorrhizas did not induce a clear biochemical response in this study.

Table 2. Mean and standard deviation for thebiochemical variables: total contents of chlorophylls ($\mu\text{g}/\text{mL}$), carotenoids and xanthophylls ($\mu\text{g}/\text{gMF}$). Stars indicate the significance level. Significance levels: *** < 0.001 ; ** < 0.01 ; Different letters indicate significant differences, according to Kruskal–Wallis test with $\alpha: 0.05$.

Treatments	SM	M1	M2
Biochemical variables			
Carotenoids and xanthophylls ***	228.67 \pm 6.8 A	188.28 \pm 6.8 B	186.04 \pm 6.8 B
Chlorophylls **	1258.85 \pm 51.2 A	1065.32 \pm 51.2 B	1054.34 \pm 51.2 B

SM: without mycorrhizal treatment, M1: Padre Lozano inoculum, M2: Colonia Benitez inoculum.

4. Conclusions

Our results suggest that reducing the doses of chemical fertilizers by incorporating biofertilizers in the production of *Prosopis alba* nursery plants is possible.

Although the AMF did not influence specific morphological and biochemical parameters as expected concerning the plants without mycorrhizae, maybe it is necessary

to advance in isolating species of native AMF compatible with the site and the *Prosopis alba* species. For this reason, deepen studies of the ecological complexity of mycorrhizal systems.

Author Contributions: M.G.S., A.d.V.G., A.T.G., M.S. and A.C.S.-G. were responsible for conceptualization, methodology, sample collection, analyses and writing. M.E. assisted with conceptualization, methodology and resources. C.C., J.G. and R.F. participated in sample dating. M.A.N. contributed to the conceptualization, methodology, and writing—review and editing. All authors have read and agreed to the published version of the manuscript.

Funding: The authors are grateful to the Universidad Nacional de Santiago del Estero (UNSE), Instituto Nacional de Tecnología Agropecuaria (INTA), Instituto de Fisiología y Recursos Genéticos Vegetales (IFRGV), Centro de Investigaciones Agropecuarias (CIAP), Instituto Nacional de Tecnología Agropecuarias (INTA), Unidad de Estudios Agropecuarios (UDEA) INTA-CONICET, Estación experimental Fernández, Universidad Católica de Santiago del Estero (UCSE), Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET) and the Ministry of Science, Technology and Productive Innovation. In addition, A.C.S.-G. acknowledges for her fellowship granted by CONICET. It is confirm that meaning was retained.

Institutional Review Board Statement: This study was waived for ethical review and approval because it did not involve human or animal research.

Informed Consent Statement: Not applicable.

Data Availability Statement: The data are not publicly available because the data will be used in the future studies.

Acknowledgments: To the projects “Development of a sustainable forestry of implanted forests of high productivity (PE 016)” and “Development and application of technologies for the control of erosion and soil degradation (PE 052)” of INTA and to the projects “Initial forestry of native species in implanted forests in irrigation zones of Santiago del Estero” and “Biostimulants evaluation to improve the productivity of plant species of regional interest under stress conditions” of the CICyT-UNSE. All individuals included in this section have consented to the acknowledgement.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Babalola, O.O. Beneficial Bacteria of Agricultural Importance. *Biotechnol. Lett.* **2010**, *32*, 1559–1570. [[CrossRef](#)] [[PubMed](#)]
2. Babalola, O.O.; Sanni, A.I.; Odhiambo, G.D.; Torto, B. Plant Growth-Promoting Rhizobacteria Do Not Pose Any Deleterious Effect on Cowpea and Detectable Amounts of Ethylene Are Produced. *World J. Microbiol. Biotechnol.* **2007**, *23*, 747–752. [[CrossRef](#)]
3. Hou, M.P.; Oluranti, B.O. Evaluation of Plant Growth Promoting Potential of Four Rhizobacterial Species for Indigenous System. *J. Cent. South Univ.* **2013**, *20*, 164–171. [[CrossRef](#)]
4. Alori, E.; Dare, M.; Babaloba, O. Sustainable Agriculture Reviews. In *Sustainable Agriculture Reviews*; Springer: Berlin/Heidelberg, Germany, 2012; Volume 11, pp. 281–307. ISBN 978-94-007-5448-5.
5. H.M.S.P.Madawala Arbuscular Mycorrhizal Fungi as Biofertilizers: Current Trends, Challenges, and Future Prospects. In *Biofertilizers Advances in Bio-Inoculant*; Amitava, R.; Vijay, S.; Manoj, P.; Sing, H.B.; Sing, A.K., Eds.; Elsevier: Amsterdam, The Netherlands, 2021; Volume 1, pp. 83–93. ISBN 9780128216675.
6. Vandresen, J.; Nishidate, F.R.; Torezan, J.M.D.; Zangara, W. Inoculação de Fungos Micorrízicos Arbusculares e Adubação Na Formação e Pós-Transplante de Mudanças de Cinco Espécies Arbóreas Nativas Do Sul Do Brasil. *Acta Bot. Bras.* **2007**, *21*, 753–765. [[CrossRef](#)]
7. Senilliani, M.G.; Brassiolo, M.; Bruno, C. Dominant Height Curves and Site Index in *Prosopis Alba* Plantations. *AgriScientia* **2021**, *38*, 13–25. [[CrossRef](#)]
8. Sun, Z.; Song, J.; Xin, X.; Xie, X.; Zhao, B. Arbuscular Mycorrhizal Fungal 14-3-3 Proteins Are Involved in Arbuscule Formation and Responses to Abiotic Stresses during AM Symbiosis. *Front. Microbiol.* **2018**, *9*, 91. [[CrossRef](#)] [[PubMed](#)]
9. Balestrini, R.; Lumini, E.; Borriello, R.; Bianciotto, V. Plant Soil Biota Interactions. In *Soil Microbiology, Ecology and Biochemistry*; Academic Press: London, UK, 2015; pp. 311–329. ISBN 9780124159556.
10. Cabrera, A.L. Fitogeografía de La República Argentina. *Bol. Soc. Argent. Bot.* **1971**, *14*, 1–42.
11. Sagadin, M.B.; Monteoliva, M.I.; Luna, C.M.; Cabello, M.N. Diversidad e Infektividad de Hongos Micorrízicos Arbusculares Nativos Provenientes de Algarrobales Del Parque Chaqueño Argentino Con Características Edafoclimáticas Contrastantes. *AgriScientia* **2018**, *35*, 19. [[CrossRef](#)]

12. Salto, C.S.; Sagadin, M.B.; Luna, C.M.; Oberschelp, G.P.J.; Harrand, L.; Cabello, M.N. Interactions between Mineral Fertilization and Arbuscular Mycorrhizal Fungi Improve Nursery Growth and Drought Tolerance of *Prosopis alba* Seedlings. *Agrofor. Syst.* **2020**, *94*, 103–111. [[CrossRef](#)]
13. Cavagnaro, T.R.; Smith, F.A.; Ayling, S.M.; Smith, S. Growth and Phosphorus Nutrition of a Paris-type Arbuscular Mycorrhizal Symbiosis. *New Phytol.* **2003**, *157*, 127–134. [[CrossRef](#)] [[PubMed](#)]
14. Santacruz-García, A.C.; Senilliani, M.G.; Gómez, A.T.; Ewens, M.; Yonny, M.E.; Villalba, G.F.; Nazareno, M.A. Biostimulants as Forest Protection Agents: Do These Products Have an Effect against Abiotic Stress on a Forest Native Species? Aspects to Elucidate Their Action Mechanisms. *For. Ecol. Manag.* **2022**, *522*, 120446. [[CrossRef](#)]
15. Malusa, E.; Sas-Paszt, L.; Popinska, W.; Zurawich, E. The Effect of a Substrate, Containing Arbuscular Mycorrhizal Fungi, and Rhizosphere Microorganisms (Trichoderma, Bacillus, Pseudomonas and Streptomyces), and Foliar Fertilization on Growth Response and Rhizosphere PH of Three Strawberry Cultivars. *Int. J. Fruit Sci.* **2007**, *6*, 25–41. [[CrossRef](#)]
16. Zangaro, W.; Torezan, J.; Domingues, Rostirola, L.; de Souza, P.; Nogueira, M. Influência de Micorrizas, Substratos Orgânicos e Volume de Recipientes Para o Crescimento de *Heliocarpus popayanensis* Kunth. *CERNE* **2015**, *21*, 395–403. [[CrossRef](#)]
17. Veiga, R.S.L.; Jansa, J.; Frossard, E.; Van der Heijden, M.G. Can Arbuscular Mycorrhizal Fungi Reduce the Growth of Agricultural Weeds? *PLoS ONE* **2011**, *6*, e27825. [[CrossRef](#)] [[PubMed](#)]
18. Santacruz-García, A.; Bravo, S.; Del Corro, F.; García, E.; Molina-Terrén, D.; Nazareno, M. How Do Plants Respond Biochemically to Fire? The Role of Photosynthetic Pigments and Secondary Metabolites in the Post-Fire Resprouting Response. *Forests* **2021**, *12*, 56. [[CrossRef](#)]
19. Lin, J.; Wang, Y.; Sun, S.; Mu, C.; Yan, X. Effects of Arbuscular Mycorrhizal Fungi on the Growth, Photosynthesis and Photosynthetic Pigments of *Leymus Chinensis* Seedlings under Salt-Alkali Stress and Nitrogen Deposition. *Sci. Total Environ.* **2017**, *576*, 234–241. [[CrossRef](#)] [[PubMed](#)]