

# Mortality risk attributable to wildfire-related PM<sub>2.5</sub> pollution: a global time series study in 749 locations



Gongbo Chen, Yuming Guo, Xu Yue, Shilu Tong, Antonio Gasparrini, Michelle L Bell, Ben Armstrong, Joel Schwartz, Jouni J K Jaakkola, Antonella Zanobetti, Eric Lavigne, Paulo Hilario Nascimento Saldiva, Haidong Kan, Dominic Royé, Ai Mijovic, Ala Overcenco, Aleš Urban, Alexandra Schneider, Alireza Entezari, Ana Maria Vicedo-Cabrera, Ariana Zeka, Aurelio Tobias, Baltazar Nunes, Barrak Alahmad, Bertil Forsberg, Shih-Chun Pan, Carmen Iñiguez, Caroline Ameling, César De la Cruz Valencia, Christofer Åström, Danny Houthuijs, Do Van Dung, Evangelia Samoli, Fatemeh Mayvaneh, Francesco Sera, Gabriel Carrasco-Escobar, Yadong Lei, Hans Orru, Ho Kim, Iulian-Horia Holobaca, Jan Kyselý, João Paulo Teixeira, Joana Madureira, Klea Katsouyanni, Magali Hurtado-Díaz, Marek Maasikmets, Martina S Ragetti, Masahiro Hashizume, Massimo Stafoggia, Mathilde Pascal, Matteo Scortichini, Micheline de Sousa Zanotti Stagliorio Coelho, Nicolás Valdés Ortega, Niilo R I Rytty, Noah Scovronick, Patricia Matus, Patrick Goodman, Rebecca M Garland, Rosana Abrutzky, Samuel Osorio García, Shilpa Rao, Simona Fratianni, Tran Ngoc Dang, Valentina Colistro, Veronika Huber, Whanhee Lee, Xerxes Seposo, Yasushi Honda, Yue Leon Guo, Tingting Ye, Wenhua Yu, Michael J Abramson, Jonathan M Samet, Shanshan Li

## Summary

**Background** Many regions of the world are now facing more frequent and unprecedentedly large wildfires. However, the association between wildfire-related PM<sub>2.5</sub> and mortality has not been well characterised. We aimed to comprehensively assess the association between short-term exposure to wildfire-related PM<sub>2.5</sub> and mortality across various regions of the world.

**Methods** For this time series study, data on daily counts of deaths for all causes, cardiovascular causes, and respiratory causes were collected from 749 cities in 43 countries and regions during 2000–16. Daily concentrations of wildfire-related PM<sub>2.5</sub> were estimated using the three-dimensional chemical transport model GEOS-Chem at a 0.25° × 0.25° resolution. The association between wildfire-related PM<sub>2.5</sub> exposure and mortality was examined using a quasi-Poisson time series model in each city considering both the current-day and lag effects, and the effect estimates were then pooled using a random-effects meta-analysis. Based on these pooled effect estimates, the population attributable fraction and relative risk (RR) of annual mortality due to acute wildfire-related PM<sub>2.5</sub> exposure was calculated.

**Findings** 65.6 million all-cause deaths, 15.1 million cardiovascular deaths, and 6.8 million respiratory deaths were included in our analyses. The pooled RRs of mortality associated with each 10 µg/m<sup>3</sup> increase in the 3-day moving average (lag 0–2 days) of wildfire-related PM<sub>2.5</sub> exposure were 1.019 (95% CI 1.016–1.022) for all-cause mortality, 1.017 (1.012–1.021) for cardiovascular mortality, and 1.019 (1.013–1.025) for respiratory mortality. Overall, 0.62% (95% CI 0.48–0.75) of all-cause deaths, 0.55% (0.43–0.67) of cardiovascular deaths, and 0.64% (0.50–0.78) of respiratory deaths were annually attributable to the acute impacts of wildfire-related PM<sub>2.5</sub> exposure during the study period.

**Interpretation** Short-term exposure to wildfire-related PM<sub>2.5</sub> was associated with increased risk of mortality. Urgent action is needed to reduce health risks from the increasing wildfires.

**Funding** Australian Research Council, Australian National Health & Medical Research Council.

**Copyright** © 2021 The Author(s). Published by Elsevier Ltd. This is an Open Access article under the CC BY-NC-ND 4.0 license.

## Introduction

Recently, large and unprecedented wildfires have been occurring frequently across the world. During the past 3 years, wildfires have been observed in many locations of the world, including Australia, British Columbia in Canada, the western USA, and the Amazon rainforest.<sup>1</sup> For example, since the start of 2019, wildfires in California have burned more than 3 million acres, resulting in thousands of destroyed homes and businesses.<sup>2</sup> The wildfires in Australia have affected every state and destroyed more than 2000 homes and burned millions of

acres.<sup>3</sup> Wildfires have both direct and indirect effects on health with potentially lasting consequences. Beyond direct injury, mental health can be harmed by the risks fires pose and loss of possessions and housing. The pollution from wildfire smoke can spread as far as 1000 km away and risk of wildfires is projected to keep increasing as climate change worsens.<sup>1</sup>

Wildfire smoke is a complex mixture of particulate matter (PM) and gaseous pollutants.<sup>4</sup> Among the various air pollutants emitted by wildfires, fine particulate matter (PM<sub>2.5</sub>) is of great concern, as particles in this size range

*Lancet Planet Health* 2021; 5: e579–87

Guangdong Provincial Engineering Technology Research Center of Environmental and Health Risk Assessment, Department of Occupational and Environmental Health, School of Public Health, Sun Yat-sen University, Guangzhou, Guangdong, China (G Chen PhD); Department of Epidemiology and Preventive Medicine, School of Public Health and Preventive Medicine, Monash University, Melbourne, VIC, Australia (Prof Y Guo PhD, T Ye MSc, W Yu MPH, Prof M J Abramson PhD, S Li PhD); Jiangsu Key Laboratory of Atmospheric Environment Monitoring and Pollution Control, Collaborative Innovation Center of Atmospheric Environment and Equipment Technology, School of Environmental Science and Engineering, Nanjing University of Information Science & Technology, Nanjing, China (Prof X Yue PhD); Shanghai Children's Medical Center, Shanghai Jiao-Tong University School of Medicine, Shanghai, China (Prof S Tong PhD); School of Public Health, Institute of Environment and Human Health, Anhui Medical University, Hefei, China (Prof S Tong); Center for Global Health, School of Public Health, Nanjing Medical University, Nanjing, China (Prof S Tong); School of Public Health and Social Work, Queensland University of Technology, Brisbane, QLD, Australia (Prof S Tong); Department of

**Public Health Environments and Society** (Prof A Gasparrini PhD, Prof B Armstrong PhD, A Milojevic PhD, F Sera PhD), **Centre for Statistical Methodology** (Prof A Gasparrini), and **Centre on Climate Change & Planetary Health** (Prof A Gasparrini), **London School of Hygiene & Tropical Medicine, London, UK;** **School of Environment, Yale University, New Haven, CT, USA** (Prof M L Bell PhD, W Lee PhD); **Department of Environmental Health, Harvard T H Chan School of Public Health, Harvard University, Boston, MA, USA** (Prof J Schwartz PhD, A Zanobetti PhD, B Alahmad PhD); **Center for Environmental and Respiratory Health Research, University of Oulu, Oulu, Finland** (Prof J J Jaakkola PhD, N R I Rytty PhD); **School of Epidemiology & Public Health, Faculty of Medicine, University of Ottawa, Ottawa, ON, Canada** (Prof E Lavigne PhD); **Air Health Science Division, Health Canada, Ottawa, ON, Canada** (Prof E Lavigne); **Department of Pathology, Faculty of Medicine, University of São Paulo, São Paulo, Brazil** (Prof P H Nascimento Saldiva PhD, M de Sousa Zanotti Stagliorio Coelho PhD); **Department of Environmental Health, School of Public Health, Fudan University, Shanghai, China** (Prof H Kan PhD); **Department of Geography, University of Santiago de Compostela, CIBER of Epidemiology and Public Health (CIBERESP), Spain** (D Royé PhD); **National Agency for Public Health of the Ministry of Health, Chisinau, Moldova** (A Overcenco PhD); **Institute of Atmospheric Physics, Czech Academy of Sciences, Prague, Czech Republic** (A Urban PhD, J Kyselý PhD); **Faculty of Environmental Sciences, Czech University of Life Sciences, Prague, Czech Republic** (A Urban, J Kyselý); **Institute of Epidemiology, Helmholtz Zentrum München–German Research Center for Environmental Health, Neuherberg, Germany** (A Schneider PhD); **Faculty of Geography and Environmental Sciences, Hakim Sabzevari**

## Research in context

### Evidence before this study

Many regions of the world are now facing more frequent and unprecedentedly large wildfires. Wildfire-related air pollution has become a major public health concern, as it can travel widely and cause various adverse health effects. Previous studies have found wildfire-related air pollution to be significantly associated with increased mortality risk. We searched PubMed, Web of Science, Google Scholar, and China National Knowledge Infrastructure using the terms “wildfire”, “bushfire”, “fine particulate matter”, “fine particles”, “PM<sub>2.5</sub>”, “death”, and “mortality” in English and Chinese for studies published up to Dec 25, 2020. We identified several studies exploring the impact of wildfire-related PM<sub>2.5</sub> on mortality. These studies showed that wildfire-related PM<sub>2.5</sub> had adverse effects on all-cause, cardiovascular, and respiratory mortality. However, the existing evidence comes from single-city or single-region studies, and not from studies with global reach.

### Added value of this study

To the best of our knowledge, this is the largest study evaluating associations between acute wildfire-related PM<sub>2.5</sub>

enter into the lungs and reach the alveoli where the small particles can translocate through the alveolar epithelium and enter the circulation.<sup>5,6</sup> Compared with PM<sub>2.5</sub> from urban sources, wildfire-related PM<sub>2.5</sub> tends to be more toxic due to its chemical composition and smaller particle size, and is often accompanied by co-exposure to other harmful environmental factors, particularly high temperatures.<sup>1</sup>

By contrast with numerous studies on total or urban background PM<sub>2.5</sub>, far fewer studies have focused on health effects of wildfire-related PM<sub>2.5</sub> specifically, although some previous studies do suggest harm to public health.<sup>7,8</sup> Wildfire-related PM<sub>2.5</sub> exposure has been found to be associated with adverse health outcomes, such as premature mortality, asthma, and reduced lung function.<sup>9–12</sup> Studies examining the health effects of wildfire smoke in the USA, Canada, Australia, and Europe have found adverse health effects.<sup>9,13–15</sup> However, existing evidence mainly comes from single-city or single-region studies, and not from global studies. One study estimated that 339 000 deaths could have been attributable to global landscape fire smoke annually during 1997–2006,<sup>16</sup> but updated evidence from global-scale studies has not been subsequently reported.

In this study, associations between daily exposure to wildfire-related PM<sub>2.5</sub> and mortality were evaluated using the Multi-City Multi-Country (MCC) Collaborative dataset for 749 cities from 43 countries and regions.

## Methods

### Mortality and socioenvironmental data

Mortality data in this study were obtained from the MCC Collaborative Research Network, an international

and mortality, and the first to do so comprehensively across various regions of the world, using daily death count data between 2000 and 2016 from 749 cities in 43 countries and regions. We found that the pooled relative risks of mortality associated with a 10 µg/m<sup>3</sup> increase in the 3-day moving average of wildfire-related PM<sub>2.5</sub> concentrations were 1.019 (95% CI 1.016–1.022) for all-cause mortality, 1.017 (1.012–1.021) for cardiovascular mortality, and 1.019 (1.013–1.025) for respiratory mortality. Overall, 0.62% (95% CI 0.48–0.75) of all-cause deaths, 0.55% (0.43–0.67) of cardiovascular deaths, and 0.64% (0.50–0.78) of respiratory deaths were attributable to the acute impacts of wildfire-related PM<sub>2.5</sub> exposure during the study period.

### Implications of all the available evidence

This study provides robust epidemiological evidence of the acute effects from wildfire-related PM<sub>2.5</sub> exposure on mortality, based on a large multicountry dataset and standard statistical method. Policy makers and public health professionals should raise awareness of wildfire pollution to prompt public responses and take actions to avoid exposure.

collaboration of research teams established to perform epidemiological studies on associations between environmental stressors and health.<sup>17,18</sup> The current MCC Network covers 750 cities from 43 countries and regions (appendix pp 6–7). Daily counts of all-cause deaths were collected from relevant authorities of each country or region. Mortality data for non-external causes (International Classification of Diseases [ICD] 9th Revision codes 0–799 or 10th Revision [ICD-10] codes A0–R99) were alternatively collected if all-cause mortality data were unavailable. In addition, mortality counts were collected specifically for cardiovascular (ICD-10 codes I00–I99) and respiratory (ICD-10 codes J00–J99) causes. Mortality data for all causes or non-external causes were available for 749 cities during the study period, while cardiovascular mortality data were available for 629 cities in 28 countries and respiratory mortality data for 647 cities in 29 countries. Other location-specific information was also collected: meteorological parameters (daily mean temperature and relative humidity) and gross domestic product (GDP) per capita.

### Estimation of wildfire-related PM<sub>2.5</sub>

From our previous work on global fire air pollution,<sup>19</sup> daily concentrations of wildfire-related PM<sub>2.5</sub> from Jan 1, 2000, to Dec 31, 2016, were estimated at a 0.25° × 0.25° resolution. Briefly, the three-dimensional chemical transport model GEOS-Chem (version 12.0.0) was used to estimate global fire-induced perturbations in PM<sub>2.5</sub>. A biomass burning inventory was adopted from the Global Fire Emissions Database (GFED; version 4.1), which estimated emissions based on satellite retrieval of burn area and active fire information. The GFED detected fires from five sources, including agricultural waste burning; boreal forest fires;




Figure 1: Maximum levels of estimated daily wildfire-related  $PM_{2.5}$  in study locations during 2000–16

tropical forest fires; savanna, grassland, and shrubland fires; and temperate forest fires.<sup>20</sup>

Daily enhancements of  $PM_{2.5}$  concentrations by fires during the study period were estimated as the differences between simulations with and without fire emissions. Daily concentration of wildfire-related  $PM_{2.5}$  was first estimated globally using GEOS-Chem at a spatial resolution of  $2.0^\circ \times 2.5^\circ$ , and then was adjusted and downscaled at a spatial resolution of  $0.25^\circ \times 0.25^\circ$  using ground-level measurements of  $PM_{2.5}$  and other predictors (eg, temperature, precipitation, wind speed, and day of the week). As wildfire-related  $PM_{2.5}$  was not routinely monitored, the GEOS-Chem-derived estimates of all-source  $PM_{2.5}$  were compared with ground-level measurements and their differences were further used to adjust the GEOS-Chem-derived wildfire-related  $PM_{2.5}$ . Results of a ten-fold cross-validation method showed that the adjusted all-source daily  $PM_{2.5}$  concentrations derived from GEOS-Chem explained 86.5% of the variability of ground-level measurements. Details of model validation, adjustment, and downscaling are shown in the appendix (pp 3–5). Based on the raster data on estimation of global wildfire-related  $PM_{2.5}$  at a spatial resolution of  $0.25^\circ \times 0.25^\circ$  (roughly 28 km<sup>2</sup> at the equator), the concentration of the pollutant in each city on each day was assigned as the average of all the cell values that fell at least partly in each city.

### Statistical analysis

To examine the association between exposure to daily wildfire-related  $PM_{2.5}$  and mortality, a two-stage analytical approach was adopted.<sup>21,22</sup> In the first stage, a quasi-Poisson regression was employed to examine the city-specific


association between daily concentration of wildfire-related  $PM_{2.5}$  and death counts. Based on our previous work,<sup>23,24</sup> the single-day effect of wildfire-related  $PM_{2.5}$  exposure on mortality on the current day and its lagged effects up to 7 days (from lag 0 to lag 7 days) were considered in city-specific models. Moving average lag models (eg, lag 0–1 and lag 0–2) were also implemented to examine cumulative effects of wildfire-related  $PM_{2.5}$  exposure.<sup>25</sup> The seasonality and long-term trends were controlled using a natural cubic spline of time with 7 degrees of freedom per year.<sup>21</sup> The moving averages of temperature (for all cities) and relative humidity (applied to 556 out of 749 cities with available humidity data) during lag 0–7 days were controlled using natural cubic splines with 4 degrees of freedom.<sup>26</sup> Additionally, categorical variables for day of the week were included in the model.

In the second stage, the effect estimates from the city-specific models were pooled to derive overall effect estimates at the global and national levels using a random-effects meta-analysis.<sup>27</sup> The pooled  $PM_{2.5}$ –mortality association was shown as relative risk (RR) of death associated with a 10  $\mu\text{g}/\text{m}^3$  increase in wildfire-related  $PM_{2.5}$ . The heterogeneity of effect estimated across cities was tested using the Cochran Q test and  $I^2$  statistic.<sup>28</sup> To check for non-linear associations, the moving average of wildfire-related  $PM_{2.5}$  was fitted using a B-spline function and two knots placed at the 25th and 75th percentiles of mean  $PM_{2.5}$  concentration across all cities.<sup>18</sup> Then concentration–response relationships between wildfire-related  $PM_{2.5}$  exposure and mortality were pooled at the global level.

Our initial analyses showed moderate heterogeneity in effect estimates across cities for all-cause mortality

University, Sabzevar, Khurasan Razavi, Iran (A Entezari PhD, F Mayvaneh PhD); Institute of Social and Preventive Medicine and Oeschger Center for Climate Change Research, University of Bern, Bern, Switzerland (A M Vicedo-Cabrera PhD); Institute of Environment, Health and Societies, Brunel University London, London, UK (A Zeka MSc); Institute of Environmental Assessment and Water Research (IDAEA), Spanish Council for Scientific Research (CSIC), Barcelona, Spain (A Tobias PhD); School of Tropical Medicine and Global Health, Nagasaki University, Nagasaki, Japan (A Tobias, X Seposo PhD); Department of Epidemiology, Instituto Nacional de Saúde Dr Ricardo Jorge, Lisbon, Portugal (B Nunes PhD); Department of Public Health and Clinical Medicine, Umeå University, Umeå, Sweden (Prof B Forsberg PhD, C Åström PhD); National Institute of Environmental Health Science, National Health Research Institutes, Zhunan, Taiwan (S-C Pan PhD, Prof Y L Guo PhD); Department of Statistics and Computational Research, Universitat de València, Valencia, CIBERESP, Spain (C Iñiguez PhD); National Institute for Public Health and the Environment (RIVM), Centre for Sustainability and Environmental Health, Bilthoven, Netherlands (C Ameling BS, D Houthuijs MSc); Department of Environmental Health, National Institute of Public Health, Cuernavaca, Morelos, Mexico (C De la Cruz Valencia MSc, Prof M Hurtado-Díaz PhD); Department of Environmental Health, Faculty of Public Health, University of Medicine and Pharmacy at Ho Chi Minh City, Ho Chi Minh City, Vietnam (D V Dung PhD, T N Dang PhD); Department of Hygiene, Epidemiology and Medical Statistics, National and Kapodistrian University of Athens, Athens, Greece (Prof E Samoli PhD, Prof K Katsouyanni PhD); Department of Statistics, Computer Science and Applications “G Parenti”, University of Florence,

Florence, Italy (F Sera); Institute of Tropical Medicine “Alexander von Humboldt”, Universidad Peruana Cayetano Heredia, Lima, Peru (G Carrasco-Escobar MSc); Climate Change Research Center, Institute of Atmospheric Physics, Chinese Academy of Sciences, Beijing, China (Y Lei MD); Institute of Family Medicine and Public Health, University of Tartu, Tartu, Estonia (H Orru PhD); Department of Public Health Sciences, Graduate School of Public Health, Seoul National University, Seoul, South Korea (H Kim PhD); Faculty of Geography, Babes-Bolyai University, Cluj-Napoca, Romania (I-H Holobaca PhD); Department of Environmental Health, Instituto Nacional de Saúde Dr Ricardo Jorge, Porto, Portugal (J P Teixeira PhD, J Madureira PhD); EPIUnit-Instituto de Saúde Pública, Universidade do Porto, Porto, Portugal (J Madureira); Estonian Environmental Research Centre, Tallinn, Estonia (M Maasikmets PhD); Swiss Tropical and Public Health Institute, Basel, Switzerland (M S Ragetti PhD); Swiss Tropical and Public Health Institute, University of Basel, Basel, Switzerland (M S Ragetti); Department of Global Health Policy, Graduate School of Medicine, The University of Tokyo, Tokyo, Japan (M Hashizume PhD); Department of Epidemiology, Lazio Regional Health Service, Rome, Italy (M Stafoggia PhD, M Scortichini MSc); Santé Publique France, Department of Environmental and occupational Health, French National Public Health Agency, Saint Maurice, France (M Pascal PhD); Department of Public Health, Universidad de los Andes, Santiago, Chile (N Valdés Ortega MSc, P Matus PhD); Gangarosa Department of Environmental Health, Rollins School of Public Health, Emory University, Atlanta, GA, USA (N Scovronick PhD); Technological University Dublin, Dublin, Ireland (Prof P Goodman PhD); Council for Scientific and Industrial Research, Pretoria, South Africa (R M Garland PhD); Department of Geography, Geoinformatics and Meteorology, University of



**Figure 2:** Pooled relative risks of mortality associated with a 10 µg/m<sup>3</sup> increase in wildfire-related PM<sub>2.5</sub> during lag 0–7 days. Estimates show the single-day effects or 3-day moving average effect of wildfire-related PM<sub>2.5</sub> on mortality, with bars representing 95% CIs.

(I<sup>2</sup> 13–50%), and almost no heterogeneity for cardiovascular (I<sup>2</sup>: 0–10%) and respiratory (I<sup>2</sup>: 0–18%) mortality (appendix p 8). Therefore, based on the pooled global-level risk estimates and assuming that the observed relationship was causal, the population attributable fraction (PAF) of annual deaths due to short-term exposure to wildfire-related PM<sub>2.5</sub> was calculated.<sup>24</sup> First, the number of annual deaths attributable to wildfire-related PM<sub>2.5</sub> was calculated for each city using pooled global-level effect estimates. Then, the total number of attributable deaths was divided by the total number of deaths across all cities to derive the pooled PAF at a global level. Additional analyses were done by pooling city-specific results at country, WHO region, and GDP levels. If only one city of a country was included in this study, the results for that city were used to represent its country. These analyses were done separately for all-cause, cardiovascular, and respiratory mortality. The specific formulas used are shown in the appendix (pp 5–6).

To examine the potential confounding effects of PM<sub>2.5</sub> from other sources, the results controlling for

other-source PM<sub>2.5</sub> were compared with those that did not, using data from cities with available ground-measured PM<sub>2.5</sub>. To test whether 7 days were sufficient to capture the lag effects of PM<sub>2.5</sub>, sensitivity analyses were done by extending the maximum lag time from 7 to 10 days. To test the robustness of the results, the degrees of freedom for meteorological variables were changed to 3, 5, and 6, and lag times up to 10 days were considered for these variables. The city-specific models were also checked by only controlling for ambient temperature. All analyses were done using R software (version 4.0.1) and the mvmeta R package.<sup>28</sup>

**Role of the funding source**

The funders of the study had no role in study design, data collection, data analysis, data interpretation, or writing of the report.

**Results**

A summary of study locations, periods, and number of deaths is shown in the appendix (pp 6–7). In total, 65·6 million all-cause deaths, 15·1 million cardiovascular deaths, and 6·8 million respiratory deaths were included in the analyses. Countries and regions contributed a median of 14·0 years (IQR 6·5). The maximum concentrations of estimated daily PM<sub>2.5</sub> induced by wildfires varied substantially by study location (figure 1). The highest daily concentrations of wildfire-related PM<sub>2.5</sub> (>100 µg/m<sup>3</sup>) were mainly estimated for cities in North America and east Asia, such as Saskatoon (Canada), Spokane (USA), Regina (Canada), and Chuncheon and Icheon (South Korea), whereas the lowest concentrations (<10 µg/m<sup>3</sup>) were mainly observed in Europe, such as Rennes and Paris (France), Bern (Switzerland), and Turin (Italy). 665 (89%) of the 749 cities had a mean concentration of estimated daily wildfire-related PM<sub>2.5</sub> of less than 2 µg/m<sup>3</sup>, with IQRs of less than 5 µg/m<sup>3</sup> across all cities (appendix pp 17–18). Additional statistical information of wildfire-related PM<sub>2.5</sub> in study locations are shown in the appendix (pp 9–11).

When considering pooled associations between daily exposure to wildfire-related PM<sub>2.5</sub> and daily mortality during lag 0–7 days, we found that the effects tended to disappear after lag 2 days (figure 2); we thus focused on the effect estimates during lag 0–2 days. Wildfire-related PM<sub>2.5</sub> exposure was significantly associated with all-cause mortality at lag between 0 and 2 days, with the greatest risk at lag 0 days (RR 1·021 [95% CI 1·018–1·024] per 10 µg/m<sup>3</sup> increase), followed by lag 1 day (1·014 [1·012–1·016]) and lag 2 days (1·005 [1·002–1·007]). Similar results were seen for cardiovascular mortality (1·017 [1·014–1·021] at lag 0 days, 1·013 [1·009–1·016] at lag 1 day, and 1·005 [1·001–1·009] at lag 2 days) and respiratory mortality (1·020 [1·015–1·024] at lag 0 days, 1·014 [1·010–1·019] at lag 1 day, and 1·006 [1·001–1·011] at lag 2 days).

The 3-day moving average of wildfire-related PM<sub>2.5</sub> (lag 0–2 days) was significantly associated with the three causes of mortality: RR 1.019 (95% CI 1.016–1.022) for all-cause mortality, 1.017 (1.012–1.021) for cardiovascular mortality, and 1.019 (1.013–1.025) for respiratory mortality. The pooled results for 3-day moving average of wildfire-related PM<sub>2.5</sub> at the country level are shown in table 1 and those pooled by WHO region and by GDP level are presented in the appendix (p 12). The highest unit RRs for all-cause mortality were observed in Europe, particularly in France, Italy, Germany, and Romania. The highest unit RRs for cardiovascular mortality were observed in Europe (including Portugal, Spain, and the Czech Republic) and the highest RRs for respiratory mortality were observed in Europe and Asia (including the Philippines, Sweden, and Kuwait; table 1).

When assessing the pooled concentration–response relationships between mortality and the 3-day moving average of wildfire-related PM<sub>2.5</sub>, RRs initially increased with respect to concentrations for both all-cause and cardiovascular mortality, levelling out at around 20 µg/m<sup>3</sup> (figure 3). For respiratory mortality, the same relationship occurred at lower concentrations, with RRs levelling out at around 15 µg/m<sup>3</sup>; however, a marked increase in RR was observed at concentrations greater than 30 µg/m<sup>3</sup> (figure 3).

Based on the pooled global associations between mortality and the 3-day moving average of wildfire-related PM<sub>2.5</sub>, an estimated 33 510 all-cause deaths (95% CI 26 204–40 763), 6993 cardiovascular deaths (5466–8510), and 3503 respiratory deaths (2739–4259) were attributable to acute wildfire-related PM<sub>2.5</sub> exposure annually on average, corresponding to PAFs of 0.62% (95% CI 0.48–0.75) for all-cause mortality, 0.55% (0.43–0.67) for cardiovascular mortality, and 0.64% (0.50–0.78) for respiratory mortality. PAFs are shown by country or region in table 2, and by WHO region and GDP level in the appendix (p 13), alongside the corresponding attributable numbers of deaths (p 16). The highest PAFs for all-cause mortality due to acute wildfire-related PM<sub>2.5</sub> exposure were observed in Thailand, Guatemala, Mexico, Paraguay, and Peru. WHO regions showing the highest PAFs for all-cause mortality were Central America (1.73%, 1.35–2.10), South-East Asia (1.63%, 95% CI 1.29–1.97), and South Africa (0.99%, 0.78–1.21); these three regions, alongside South America, also showed the highest PAFs for cardiovascular and respiratory mortality, with PAFs greater than 1.00% (appendix p 13).

Sensitivity analyses showed that the pooled results did not change substantially by further controlling for other-source PM<sub>2.5</sub> (appendix pp 19–20). The pooled results using adjusted and unadjusted wildfire-related PM<sub>2.5</sub> were consistent, although greater uncertainties were observed for results using unadjusted data (appendix pp 20–21). Lags of up to 2 days were sufficient to capture the lag effects of PM<sub>2.5</sub>, as no significant associations of wildfire-related PM<sub>2.5</sub> exposure were observed during lags

	All-cause mortality	Cardiovascular mortality	Respiratory mortality
Argentina	1.040 (1.017–1.063)	NA	NA
Australia	1.002 (0.991–1.012)	NA	NA
Brazil	1.011 (0.998–1.024)	NA	NA
Canada	0.992 (0.978–1.007)	0.997 (0.974–1.021)	1.023 (0.994–1.053)
Chile	1.033 (1.010–1.056)	NA	NA
China	1.030 (1.005–1.055)	1.026 (0.982–1.073)	1.006 (0.977–1.036)
Colombia	1.028 (1.011–1.046)	1.023 (0.992–1.055)	1.002 (0.975–1.029)
Costa Rica	1.069 (1.032–1.106)	1.061 (0.995–1.131)	1.000 (0.994–1.006)
Czech Republic	1.073 (1.009–1.142)	1.070 (0.954–1.200)	1.039 (0.951–1.137)
Ecuador	1.010 (0.935–1.091)	1.033 (0.911–1.172)	1.066 (0.886–1.281)
Estonia	0.995 (0.955–1.036)	NA	NA
Finland	1.001 (0.997–1.005)	1.006 (0.999–1.012)	0.974 (0.869–1.070)
France	1.253 (1.079–1.455)	NA	1.022 (0.989–1.056)
Germany	1.126 (1.068–1.188)	NA	NA
Greece	1.009 (1.002–1.016)	1.015 (1.005–1.024)	1.010 (0.991–1.030)
Guatemala	0.996 (0.990–1.002)	NA	NA
Iran	1.014 (0.998–1.030)	1.030 (1.006–1.055)	1.006 (0.987–1.024)
Ireland	0.891 (0.751–1.057)	1.059 (0.713–1.573)	1.033 (0.997–1.070)
Italy	1.139 (1.081–1.199)	NA	NA
Japan	1.027 (1.022–1.032)	1.027 (1.017–1.036)	1.022 (1.005–1.040)
Kuwait	1.017 (0.989–1.045)	1.044 (1.006–1.083)	1.105 (0.973–1.239)
Mexico	1.002 (0.997–1.007)	1.001 (0.991–1.011)	1.072 (0.943–1.218)
Moldova	1.041 (0.916–1.182)	NA	NA
Netherlands	0.990 (0.842–1.164)	NA	NA
Norway	1.016 (0.995–1.036)	1.034 (1.000–1.067)	1.006 (0.987–1.024)
Panama	1.011 (0.918–1.111)	0.989 (0.839–1.161)	1.035 (0.949–1.126)
Paraguay	1.001 (0.999–1.003)	1.000 (0.997–1.003)	1.004 (0.999–1.009)
Peru	0.975 (0.957–0.994)	NA	NA
Philippines	1.008 (0.983–1.034)	1.000 (0.946–1.058)	1.171 (0.963–1.423)
Portugal	1.062 (0.999–1.130)	1.101 (0.967–1.254)	1.009 (0.975–1.043)
Puerto Rico	1.053 (1.003–1.106)	NA	NA
Romania	1.116 (1.075–1.158)	NA	NA
South Africa	1.018 (1.011–1.024)	1.016 (1.005–1.027)	1.016 (1.003–1.029)
South Korea	1.012 (1.000–1.024)	0.992 (0.974–1.010)	1.003 (0.991–1.015)
Spain	1.066 (1.030–1.104)	1.074 (1.026–1.126)	1.015 (0.994–1.038)
Sweden	0.946 (0.868–1.032)	1.007 (0.882–1.149)	1.145 (0.986–1.330)
Switzerland	1.026 (0.827–1.274)	1.059 (0.765–1.464)	0.926 (0.843–1.017)
Taiwan	1.034 (1.013–1.055)	1.009 (0.966–1.053)	1.035 (0.958–1.118)
Thailand	1.016 (1.013–1.020)	1.012 (1.005–1.018)	1.005 (0.996–1.015)
UK	1.023 (0.955–1.095)	1.061 (0.960–1.174)	1.022 (1.007–1.038)
Uruguay	1.019 (1.008–1.029)	NA	NA
USA	1.010 (1.001–1.020)	1.014 (0.998–1.031)	1.023 (1.015–1.030)
Vietnam	1.009 (0.950–1.071)	1.006 (0.953–1.062)	0.990 (0.786–1.246)

Data are RR (95% CI). RRs were associated with per 10 µg/m<sup>3</sup> increase in moving average of wildfire-related PM<sub>2.5</sub> during lag 0–2 days. NA=not available. RR=relative risk.

**Table 1: Relative risks of mortality associated with exposure to wildfire-related PM<sub>2.5</sub> during lag 0–2 days in 43 countries and regions.**

3–10 days (appendix p 22). The results did not change substantially with use of 3, 5, or 6 degrees of freedom and 10-day lag effects for meteorological variables, or with controlling only for temperature in city-specific models (appendix pp 23–27).

Pretoria, Pretoria, South Africa (R M Garland); Unit for Environmental Sciences and Management, North West University, South Africa (R M Garland); Instituto de

Investigaciones Gino Germani, Facultad de Ciencias Sociales, Universidad de Buenos Aires, Buenos Aires, Argentina (R Abrutzky PhD); Department of Environmental Health, University of São Paulo, São Paulo, Brazil (S O Garcia PhD); Norwegian Institute of Public Health, Oslo, Norway (S Rao PhD); Department of Earth Sciences, University of Torino, Turin, Italy (S Fratianni PhD); Department of Quantitative Methods, School of Medicine, University of the Republic, Montevideo, Uruguay (V Colistro PhD); Potsdam Institute for Climate Impact Research, Potsdam, Germany (V Huber PhD); Department of Physical, Chemical and Natural Systems, Universidad Pablo de Olavide, Seville, Spain (V Huber); Faculty of Health and Sport Sciences, University of Tsukuba, Tsukuba, Japan (Prof Y Honda PhD); Environmental and Occupational Medicine, and Institute of Environmental Health and Occupational Health Sciences, National Taiwan University and National Taiwan University Hospital, Taipei, Taiwan (Prof Y L Guo); The Colorado School of Public Health, University of Colorado, Aurora (Prof J M Samet MD)

Correspondence to:

Prof Yuming Guo, School of Public Health and Preventive Medicine, Monash University, Melbourne, VIC 3004, Australia [yuming.guo@monash.edu](mailto:yuming.guo@monash.edu)


or

Dr Shanshan Li, School of Public Health and Preventive Medicine, Monash University, Melbourne, VIC 3004, Australia [shanshan.li@monash.edu](mailto:shanshan.li@monash.edu)

For more on the MCC Collaborative Research Network see <http://mccstudy.lshtm.ac.uk>

See Online for appendix

For more on the GEOS-Chem model see <http://wiki.seas.harvard.edu/geos-chem>



**Figure 3:** The pooled concentration-response relationships between mortality and the 3-day moving average of wildfire-related  $PM_{2.5}$  during lag 0–2 days. Shaded areas represent 95% CIs.

## Discussion

To our knowledge, this is the largest study evaluating associations between acute wildfire-related  $PM_{2.5}$  and mortality, and the first to do so comprehensively across various regions of the world. We found that exposure to wildfire-related  $PM_{2.5}$  was significantly associated with increased all-cause, cardiovascular, and respiratory mortality at a global level, but the associations varied across countries and regions.

The wildfire-related  $PM_{2.5}$ –mortality associations were assessed across various geographical regions and populations during a relatively long study period, based on the largest mortality dataset covering 43 countries and regions worldwide. With the use of a two-stage design, all city-specific associations between wildfire-related  $PM_{2.5}$  and mortality were analysed in the same way, facilitating the comparison of results across different populations and regions.<sup>17,18</sup> The second stage random-effects meta-analysis has been widely used to examine both within-city and between-city variations regarding risk estimates.<sup>29</sup> The PAF was estimated using the pooled effect estimates with the same lag structure for each location, which provides essential information for public health planning and potential interventions.<sup>30</sup>

The results of our study are consistent with those of previous investigations, despite different effect estimates and exposure periods. However, previous studies were mainly restricted to a single study area or country, or a particular fire season. For example, a study in 27 countries in Europe estimated that 1483 premature deaths in 2005 and 1080 in 2008 could be attributable to vegetation fire-related  $PM_{2.5}$ .<sup>15</sup> Another study in Canada found that 54–240 premature deaths were attributable to wildfire-related  $PM_{2.5}$  annually between 2013–15 and 2017–18.<sup>9</sup> Fixed and temporary ground monitors and satellite-based data have alternatively been used to estimate exposure to wildfire-related air pollutants, but these methods provide limited spatiotemporal coverage, low data quality of surface pollution level, and cannot quantify the contribution of fire smoke.<sup>31</sup> The GEOS-Chem

model can address these problems by considering both non-fire and fire emissions. However, uncertainty in emissions data might affect the accuracy of estimation. For example, a study in North America reported that GFED-driven estimates matched well with observations, but showed overestimates and underestimates in some species and regions.<sup>32</sup>

Wildfire-related  $PM_{2.5}$  undergoes long-range transport and continues to contribute to poor air quality even after fire seasons.<sup>33</sup> Therefore, evaluating health effects of wildfires should not be restricted to areas and time periods where and when wildfires occur. The pooled PAF of mortality attributable to acute wildfire-related  $PM_{2.5}$  might seem low in terms of relative increase (<0.7%). This is caused by the special distribution of concentrations of wildfire-related  $PM_{2.5}$  over time. Extremely high concentrations of wildfire-related  $PM_{2.5}$  only occurred during fire seasons, which constituted a very short period relative to the whole study period, while wildfire-related  $PM_{2.5}$  remained at a very low level during the long periods between fire seasons, with nearly 90% of cities having a mean concentration of estimated daily wildfire-related  $PM_{2.5}$  of less than  $2 \mu\text{g}/\text{m}^3$ . However, the overall health impacts of wildfire-related  $PM_{2.5}$  would be generally underestimated by this study. Wildfire-related  $PM_{2.5}$  has both short-term and long-term health effects, but our study only focused on its short-term effects on mortality. More studies are needed in future to systematically examine its long-term effects on various health outcomes.

Our previous work on ambient  $PM_{2.5}$  (mainly urban background  $PM_{2.5}$ ) and daily mortality in 652 cities showed that all-cause mortality increased by 0.44% (95% CI 0.39–0.50), cardiovascular mortality by 0.36% (0.30–0.43), and respiratory mortality by 0.47% (0.35–0.58) with every  $10 \mu\text{g}/\text{m}^3$  increase in  $PM_{2.5}$  at lag 0–1 days.<sup>18</sup> By comparison, we found that wildfire-related  $PM_{2.5}$  exposure had stronger effects on mortality (higher RRs) and a longer lag time than urban  $PM_{2.5}$ . The potential greater toxicity of wildfire  $PM_{2.5}$  could reflect its higher fractions of small particles (eg, sub-micrometre particles and ultrafine particles) and

	All-cause mortality	Cardiovascular mortality	Respiratory mortality
Argentina	0.77% (0.60–0.93)	NA	NA
Australia	0.88% (0.70–1.07)	NA	NA
Brazil	0.70% (0.54–0.85)	NA	NA
Canada	0.33% (0.26–0.41)	0.33% (0.26–0.40)	0.32% (0.25–0.39)
Chile	0.43% (0.33–0.52)	NA	NA
China	0.66% (0.51–0.80)	0.67% (0.52–0.81)	0.65% (0.50–0.79)
Colombia	0.97% (0.76–1.18)	0.97% (0.76–1.18)	0.95% (0.74–1.16)
Costa Rica	0.94% (0.73–1.14)	0.92% (0.72–1.13)	0.92% (0.71–1.12)
Czech Republic	0.14% (0.11–0.17)	0.13% (0.10–0.16)	0.13% (0.10–0.16)
Ecuador	0.98% (0.76–1.19)	0.94% (0.74–1.15)	0.99% (0.77–1.20)
Estonia	0.17% (0.13–0.21)	NA	NA
Finland	0.14% (0.11–0.18)	0.15% (0.12–0.18)	0.13% (0.10–0.16)
France	0.12% (0.09–0.14)	NA	0.11% (0.09–0.14)
Germany	0.13% (0.10–0.16)	NA	NA
Greece	0.33% (0.26–0.40)	0.33% (0.25–0.40)	0.34% (0.26–0.41)
Guatemala	3.04% (2.39–3.68)	NA	NA
Iran	0.34% (0.26–0.41)	0.33% (0.26–0.41)	0.28% (0.22–0.34)
Ireland	0.09% (0.07–0.11)	0.09% (0.07–0.11)	0.08% (0.06–0.10)
Italy	0.30% (0.23–0.36)	NA	NA
Japan	0.63% (0.49–0.76)	0.61% (0.47–0.74)	0.61% (0.47–0.74)
Kuwait	0.37% (0.29–0.46)	0.37% (0.29–0.46)	0.36% (0.28–0.44)
Mexico	1.72% (1.35–2.09)	1.69% (1.32–2.05)	1.77% (1.39–2.15)
Moldova	0.27% (0.21–0.32)	NA	NA
Netherlands	0.13% (0.10–0.15)	NA	NA

(Table 2 continues in next column)

	All-cause mortality	Cardiovascular mortality	Respiratory mortality
(Continued from previous column)			
Norway	0.09% (0.07–0.12)	0.09% (0.07–0.12)	0.09% (0.07–0.11)
Panama	0.37% (0.29–0.45)	0.38% (0.29–0.46)	0.34% (0.26–0.41)
Paraguay	2.09% (1.64–2.54)	2.10% (1.65–2.55)	2.19% (1.72–2.65)
Peru	1.61% (1.26–1.96)	NA	NA
Philippines	0.79% (0.62–0.97)	0.80% (0.63–0.98)	0.78% (0.61–0.95)
Portugal	0.28% (0.22–0.34)	0.26% (0.20–0.31)	0.27% (0.21–0.33)
Puerto Rico	0.27% (0.21–0.34)	NA	NA
Romania	0.35% (0.27–0.42)	NA	NA
South Africa	0.99% (0.78–1.21)	1.00% (0.78–1.21)	1.10% (0.86–1.34)
South Korea	0.53% (0.41–0.64)	0.53% (0.41–0.64)	0.52% (0.41–0.63)
Spain	0.19% (0.15–0.23)	0.19% (0.14–0.23)	0.18% (0.14–0.22)
Sweden	0.10% (0.08–0.12)	0.10% (0.08–0.12)	0.10% (0.07–0.12)
Switzerland	0.15% (0.12–0.18)	0.15% (0.12–0.18)	0.15% (0.12–0.18)
Taiwan	0.58% (0.45–0.70)	0.57% (0.44–0.69)	0.57% (0.45–0.70)
Thailand	2.32% (1.83–2.80)	2.43% (1.92–2.93)	2.44% (1.92–2.94)
UK	0.09% (0.07–0.11)	0.09% (0.07–0.11)	0.08% (0.07–0.10)
Uruguay	0.51% (0.40–0.62)	NA	NA
USA	0.26% (0.20–0.32)	0.26% (0.20–0.31)	0.26% (0.20–0.31)
Vietnam	0.99% (0.78–1.21)	1.04% (0.81–1.27)	0.97% (0.76–1.18)

Data are PAF (95% CI). PAFs were calculated using the pooled global-level risk estimates. The corresponding number of deaths is shown in the appendix (pp 14–15). NA=not available. PAF=population attributable fraction.

**Table 2: PAF of annual mortality due to exposure to wildfire-related PM<sub>2.5</sub> during lag 0–2 days in 43 countries and regions**

more oxidative and proinflammatory components, such as polycyclic aromatic hydrocarbons and aldehydes.<sup>34</sup> Moreover, the joint effects of wildfire-related PM<sub>2.5</sub> and other pollutants, such as oxidant gases, might result in amplified health effects.<sup>1</sup>

Several limitations of this study should be noted. Although our MCC mortality data covered 43 countries and regions, they were not evenly distributed on every

continent. The pooled mortality risk should not be interpreted as providing global results with high representativeness, as the analyses were mainly performed for urban populations. Some country-specific results might not fully represent the health effects for those countries owing to the small number of cities included in this study; in particular, 11 countries only had data for one city (appendix pp 6–7). Moreover, due to missing values or unavailability of data, the mortality data in some locations did not cover the full study period. Fire emissions generate a dynamic mixture of air pollutants that varies over space and time and that cannot be fully captured by the GEOS-Chem model.<sup>35</sup> We did not consider other air

pollutants from wildfires including carbon monoxide, carbon dioxide, or ozone. Moreover, the spatial resolution of estimation is coarse, which might underestimate the spatial variations of exposure and introduce exposure misclassification. The accuracy and spatial resolution of estimated wildfire-related PM<sub>2.5</sub> can be improved in future by including more detailed exposure data (eg, satellite-based data and weather data) with novel models. Finally, we did not analyse the association between wildfire-related PM<sub>2.5</sub> and mortality in susceptible subgroups of the populations (eg, by age or sex) owing to unavailability of individual information. If possible, such stratified analyses should be done in future studies to identify subpopulations vulnerable to wildfire air pollution.

This study provides robust epidemiological evidence for acute effects of wildfire-related PM<sub>2.5</sub> exposure on mortality, based on a large multicountry dataset and standard statistical method. Policy makers and public health professionals should raise awareness of wildfire pollution to guide prompt public responses and take actions to reduce exposure. Effective wildland management policies and practices should be implemented to manage vegetation and mitigate climate change as far as possible.

#### Contributors

YG, AG, MH, and BAr set up the collaborative network. YG and SL conceived, designed, and coordinated the study. YG, SL, and GC developed the statistical methods, took the lead in drafting the manuscript and interpreting the results, and verified the underlying data. YG and XY did the exposure assessment of wildfire-related PM<sub>2.5</sub>. Other authors provided the data on mortality and temperature, and contributed to the interpretation of the results and to the submitted version of the manuscript. All authors had full access to all data and final responsibility to submit this paper for publication.

#### Declaration of interests

We declare no competing interests.

#### Data sharing

Data used in this study were collected by collaborators within the MCC Network under a data sharing agreement and cannot be made available publicly.

#### Acknowledgments

This study was supported by the Australian Research Council (DP210102076) and the Australian National Health and Medical Research Council (APP2000581). YG was supported by a Career Development Fellowship of the Australian National Health and Medical Research Council (APP1163693). SL was supported by an Early Career Fellowship of the Australian National Health and Medical Research Council (APP1109193). GC was supported by the National Natural Science Foundation of China (81903279). AU and JK were supported by the Czech Science Foundation (20-28560S). VH received support from the Spanish Ministry of Economy, Industry and Competitiveness (PCIN-2017-046). XY was supported by the National Key Research and Development Program of China (2019YFA0606802). SR acknowledges funding from EU's Horizon 2020 Project Exhaustion (820655). S-CP and YLG were supported by the Ministry of Science and Technology of Taiwan (MOST 109-2621-M-002-021). AG was supported by the Medical Research Council UK (MR/M022625/1), the Natural Environment Research Council UK (NE/R009384/1), and the EU's Horizon 2020 Project Exhaustion (820655); JM was supported by a fellowship of the Fundação para a Ciência e a Tecnologia (SFRH/BPD/115112/2016). ST was supported by the Science and Technology Commission of Shanghai Municipality (18411951600). NS was supported by the National Institute of Environmental Health Sciences-funded HERCULES Center (P30ES019776).

Editorial note: the *Lancet* Group takes a neutral position with respect to territorial claims in published maps and institutional affiliations.

#### References

- Xu R, Yu P, Abramson MJ, et al. Wildfires, global climate change, and human health. *N Engl J Med* 2020; **383**: 2173–81.
- Kizer KW. Extreme wildfires—a growing population health and planetary problem. *JAMA* 2020; **324**: 1605–06.
- Cousins S. Bushfires expose weaknesses in Australia's health system. *Lancet* 2020; **395**: 175–76.
- Black C, Tesfaigzi Y, Bassein JA, Miller LA. Wildfire smoke exposure and human health: significant gaps in research for a growing public health issue. *Environ Toxicol Pharmacol* 2017; **55**: 186–95.
- Feng S, Gao D, Liao F, Zhou F, Wang X. The health effects of ambient PM<sub>2.5</sub> and potential mechanisms. *Ecotoxicol Environ Saf* 2016; **128**: 67–74.
- Lu F, Xu D, Cheng Y, et al. Systematic review and meta-analysis of the adverse health effects of ambient PM<sub>2.5</sub> and PM<sub>10</sub> pollution in the Chinese population. *Environ Res* 2015; **136**: 196–204.
- Reid CE, Brauer M, Johnston FH, Jerrett M, Balmes JR, Elliott CT. Critical review of health impacts of wildfire smoke exposure. *Environ Health Perspect* 2016; **124**: 1334–43.
- Liu JC, Pereira G, Uhl SA, Bravo MA, Bell ML. A systematic review of the physical health impacts from non-occupational exposure to wildfire smoke. *Environ Res* 2015; **136**: 120–32.
- Matz CJ, Egyed M, Xi G, et al. Health impact analysis of PM<sub>2.5</sub> from wildfire smoke in Canada (2013–2015, 2017–2018). *Sci Total Environ* 2020; **725**: 138506.
- Orr A, Al Migliaccio C, Buford M, Ballou S, Migliaccio CT. Sustained effects on lung function in community members following exposure to hazardous PM<sub>2.5</sub> levels from wildfire smoke. *Toxics* 2020; **8**: 53.
- Liu JC, Wilson A, Mickley LJ, et al. Wildfire-specific fine particulate matter and risk of hospital admissions in urban and rural counties. *Epidemiology* 2017; **28**: 77.
- Arbex MA, Martins LC, de Oliveira RC, et al. Air pollution from biomass burning and asthma hospital admissions in a sugar cane plantation area in Brazil. *J Epidemiol Commun H* 2007; **61**: 395–400.
- Xi Y, Kshirsagar AV, Wade TJ, et al. Mortality in US hemodialysis patients following exposure to wildfire smoke. *J Am Soc Nephrol* 2020; **31**: 1824–35.
- Johnston FH, Borchers-Arriagada N, Morgan GG, et al. Unprecedented health costs of smoke-related PM<sub>2.5</sub> from the 2019–20 Australian megafires. *Nat Sustain* 2021; **4**: 42–47.
- Kollanus V, Prank M, Gens A, et al. Mortality due to vegetation fire-originated PM<sub>2.5</sub> exposure in Europe—assessment for the years 2005 and 2008. *Environ Health Perspect* 2017; **125**: 30–37.
- Johnston FH, Henderson SB, Chen Y, et al. Estimated global mortality attributable to smoke from landscape fires. *Environ Health Perspect* 2012; **120**: 695–701.
- Gasparrini A, Guo Y, Hashizume M, et al. Mortality risk attributable to high and low ambient temperature: a multicountry observational study. *Lancet* 2015; **386**: 369–75.
- Liu C, Chen R, Sera F, et al. Ambient particulate air pollution and daily mortality in 652 cities. *N Engl J Med* 2019; **381**: 705–15.
- Yue X, Unger N. Fire air pollution reduces global terrestrial productivity. *Nat Commun* 2018; **9**: 1–9.
- Oak Ridge National Laboratory Distributed Active Archive Center. Global Fire Emissions Database, Version 4.1 (GFEDv4). Feb 14, 2018. [https://daac.ornl.gov/VEGETATION/guides/fire\\_emissions\\_v4\\_R1.html](https://daac.ornl.gov/VEGETATION/guides/fire_emissions_v4_R1.html) (accessed April 15, 2021).
- Guo Y, Gasparrini A, Armstrong BG, et al. Heat wave and mortality: a multicountry, multicity study. *Environ Health Perspect* 2017; **125**: 087006.
- Gasparrini A, Guo Y, Hashizume M, et al. Temporal variation in heat-mortality associations: a multicountry study. *Environ Health Perspect* 2015; **123**: 1200–07.
- Chen G, Li S, Zhang Y, et al. Effects of ambient PM<sub>2.5</sub> air pollution on daily emergency hospital visits in China: an epidemiological study. *Lancet Planet Health* 2017; **1**: e221–29.
- Chen G, Zhang Y, Zhang W, et al. Attributable risks of emergency hospital visits due to air pollutants in China: a multi-city study. *Environ Pollut* 2017; **228**: 43–49.



- 25 Chen G, Zhang W, Li S, et al. The impact of ambient fine particles on influenza transmission and the modification effects of temperature in China: a multi-city study. *Environ Int* 2017; **98**: 82–88.
- 26 Guo Y, Gasparrini A, Armstrong BG, et al. Temperature variability and mortality: a multi-country study. *Environ Health Perspect* 2016; **124**: 1554–59.
- 27 Shah ASV, Langrish JP, Nair H, et al. Global association of air pollution and heart failure: a systematic review and meta-analysis. *Lancet* 2013; **382**: 1039–48.
- 28 Gasparrini A, Armstrong B, Kenward M. Multivariate meta-analysis for non-linear and other multi-parameter associations. *Stat Med* 2012; **31**: 3821–39.
- 29 Raaschou-Nielsen O, Andersen ZJ, Beelen R, et al. Air pollution and lung cancer incidence in 17 European cohorts: prospective analyses from the European Study of Cohorts for Air Pollution Effects (ESCAPE). *Lancet Oncol* 2013; **14**: 813–22.
- 30 Gasparrini A, Leone M. Attributable risk from distributed lag models. *BMC Med Res Methodol* 2014; **14**: 55.
- 31 Zou Y, O'Neill SM, Larkin NK, et al. Machine learning-based integration of high-resolution wildfire smoke simulations and observations for regional health impact assessment. *Int J Env Res Pub He* 2019; **16**: 2137.
- 32 Carter TS, Heald CL, Jimenez JL, et al. How emissions uncertainty influences the distribution and radiative impacts of smoke from fires in North America. *Atmos Chem Phys* 2020; **20**: 2073–97.
- 33 Cottle P, Strawbridge K, McKendry I. Long-range transport of Siberian wildfire smoke to British Columbia: Lidar observations and air quality impacts. *Atmos Environ* 2014; **90**: 71–77.
- 34 Verma V, Polidori A, Schauer JJ, Shafer MM, Cassee FR, Sioutas C. Physicochemical and toxicological profiles of particulate matter in Los Angeles during the October 2007 southern California wildfires. *Environ Sci Technol* 2009; **43**: 954–60.
- 35 Adetona O, Reinhardt TE, Domitrovich J, et al. Review of the health effects of wildland fire smoke on wildland firefighters and the public. *Inhal Toxicol* 2016; **28**: 95–139.