

RESEARCH

Open Access



Genomics and prevalence of bacterial and archaeal isolates from biogas-producing microbiomes

Irena Maus^{1†}, Andreas Bremges^{1,2,3,4†}, Yvonne Stolze¹, Sarah Hahnke⁵, Katharina G. Cibis⁶, Daniela E. Koeck⁷, Yong S. Kim⁸, Jana Kreubel⁶, Julia Hassa¹, Daniel Wibberg¹, Aaron Weimann³, Sandra Off⁸, Robbin Stantscheff^{6,10}, Vladimir V. Zverlov^{7,9}, Wolfgang H. Schwarz⁷, Helmut König⁶, Wolfgang Liebl⁷, Paul Scherer⁸, Alice C. McHardy³, Alexander Sczyrba^{1,2}, Michael Klocke⁵, Alfred Pühler¹ and Andreas Schlüter^{1*}

Abstract

Background: To elucidate biogas microbial communities and processes, the application of high-throughput DNA analysis approaches is becoming increasingly important. Unfortunately, generated data can only partially be interpreted rudimentary since databases lack reference sequences.

Results: Novel cellulolytic, hydrolytic, and acidogenic/acetogenic *Bacteria* as well as methanogenic *Archaea* originating from different anaerobic digestion communities were analyzed on the genomic level to assess their role in biomass decomposition and biogas production. Some of the analyzed bacterial strains were recently described as new species and even genera, namely *Herbinix hemicellulosilytica* T3/55^T, *Herbinix luporum* SD1D^T, *Clostridium bornimense* M2/40^T, *Proteiniphilum saccharofermentans* M3/6^T, *Fermentimonas caenicola* ING2-E5B^T, and *Petrimonas mucosa* ING2-E5A^T. High-throughput genome sequencing of 22 anaerobic digestion isolates enabled functional genome interpretation, metabolic reconstruction, and prediction of microbial traits regarding their abilities to utilize complex bio-polymers and to perform specific fermentation pathways. To determine the prevalence of the isolates included in this study in different biogas systems, corresponding metagenome fragment mappings were done. *Methanoculleus bourgensis* was found to be abundant in three mesophilic biogas plants studied and slightly less abundant in a thermophilic biogas plant, whereas *Defluviitoga tunisiensis* was only prominent in the thermophilic system. Moreover, several of the analyzed species were clearly detectable in the mesophilic biogas plants, but appeared to be only moderately abundant. Among the species for which genome sequence information was publicly available prior to this study, only the species *Amphibacillus xylanus*, *Clostridium clariflavum*, and *Lactobacillus acidophilus* are of importance for the biogas microbiomes analyzed, but did not reach the level of abundance as determined for *M. bourgensis* and *D. tunisiensis*.

Conclusions: Isolation of key anaerobic digestion microorganisms and their functional interpretation was achieved by application of elaborated cultivation techniques and subsequent genome analyses. New isolates and their genome information extend the repository covering anaerobic digestion community members.

Keywords: Anaerobic digestion, Biomethanation, Genome sequencing, Fragment recruitment, *Defluviitoga tunisiensis*, *Methanoculleus bourgensis*

*Correspondence: aschluet@cebitec.uni-bielefeld.de

[†]Irena Maus and Andreas Bremges contributed equally to this work

¹ Center for Biotechnology (CeBiTec), Bielefeld University, Universitätsstrasse 27, 33615 Bielefeld, Germany

Full list of author information is available at the end of the article

Background

Anaerobic digestion (AD) and biomethanation are commonly applied for the treatment and decomposition of organic material and bio-waste, finally yielding methane (CH₄)-rich biogas. The whole AD process can be divided into four phases: hydrolysis, acidogenesis, acetogenesis, and methanogenesis. Organic polymers are hydrolyzed into sugar molecules, fatty acids, and amino acids by hydrolytic enzymes. These metabolites are further degraded into the intermediate volatile fatty acids (VFA), acetate, alcohols, carbon dioxide (CO₂), and hydrogen (H₂) during acidogenesis and acetogenesis. Finally, CH₄ is produced either from acetate or from H₂ and CO₂. The challenges in each of these steps are reflected within the complexity of the microbial community converting biomass to biogas. Community compositions and dynamics were frequently investigated using different molecular biological methods. Among these, quantitative 'real-time' polymerase chain reaction (qPCR), e.g., [1–5], terminal restriction fragment length polymorphism (TRFLP) [6–8], and the 16S rRNA gene amplicon [9, 10] as well as metagenome sequencing approaches [9, 11–14] applying high-throughput (HT) technologies are the most commonly used methods. In these studies, bacterial members belonging to the classes *Clostridia* and *Bacteroidia* were identified to dominate the biogas microbial communities, followed by *Proteobacteria*, *Bacilli*, *Flavobacteria*, *Spirochaetes*, and *Erysipelotrichi*. Within the domain *Archaea*, members from the orders *Methanomicrobiales*, *Methanosarcinales*, and *Methanobacteriales* were described to be abundant in biogas systems.

However, all recently published metagenome and metatranscriptome studies addressing elucidation of the biogas microbiology reported on a huge fraction of unassignable sequences suggesting that most of the microorganisms in biogas communities are so far unknown [15–18]. This is due to the limiting availability of reference strains and their corresponding genome sequences in public databases. Moreover, reference sequences are often derived from only distantly related strains isolated from different environments. For a better understanding of the microbial trophic networks in AD and any further biotechnological optimization of the biomethanation process, extension of public databases regarding relevant sequence information seems to be an indispensable prerequisite.

Recently, studies on the isolation, sequencing, and physiological characterization of novel microbial strains from various mesophilic and thermophilic biogas reactors were published, e.g., [18–29]. However, only few of these studies addressed the question of whether the described strain played a dominant role within the analyzed microbial community. Accordingly, the objective

of this work was to sequence and analyze a collection of recently described as well as newly isolated bacterial and archaeal strains from different biogas microbial communities to provide insights into their metabolic potential and life-style, and to estimate their prevalence in selected agricultural biogas reactors. In total, 22 different strains originating from meso- and thermophilic anaerobic digesters utilizing renewable primary products and/or organic wastes were analyzed. Based on genome analyses, isolates were functionally classified and assigned to functional roles within the AD process. Moreover, refinement of the metagenome fragment recruitment approach was used for the evaluation of an isolate's prominence in different biogas communities. Overall the aim of this study was the considerable complementation of the reference repository by new genome information regarding AD communities.

Methods

Microbial strains used in this study and isolation of novel strains

In this study, 22 bacterial and archaeal strains were studied from eight meso- and thermophilic, laboratory-scale and agricultural biogas plants (BGPs) utilizing renewable primary products as well as from three further AD sources (detailed information listed in Table 1). The strains *Methanoculleus chikugoensis* L21-II-0 and *Sporanaerobacter* sp. PP17-6a were isolated within this study as follows.

Methanoculleus chikugoensis L21-II-0 Reactor material was diluted fivefold in DSMZ medium 287 [30] containing 20 mM acetate and H₂/CO₂ as the only carbon and energy sources. Initial incubation occurred at 37 °C for 4 weeks without antibiotics. Subsequent cultivation was performed by successive transfer of culture aliquots after incubation periods of 4 weeks into the same medium supplemented with different combinations of the antibiotics tetracycline HCl (15 µg ml⁻¹), vancomycin HCl (50 µg ml⁻¹), ampicillin (100 µg ml⁻¹), and bacitracin (15 µg ml⁻¹) or with penicillin (350 µg ml⁻¹). After a total of 12 cultivation cycles, purity of the culture was confirmed by microscopic inspection and by denaturing gradient gel electrophoresis (DGGE) fingerprint analysis. Strain *M. chikugoensis* L21-II-0 is available from the Leibniz Institute German Collection of Microorganisms and Cell Cultures (DSMZ, Braunschweig, Germany) under the Accession No. DSM 100195. *Sporanaerobacter* sp. PP17-6a: Reactor material was diluted 5 × 10⁶-fold in DSMZ medium 120 [31]. After 4 weeks of incubation at 37 °C, an aliquot of the culture was transferred into the same medium supplemented with penicillin (350 µg ml⁻¹). Transfer and incubation in the same medium were repeated four times. Subsequently,

Table 1 Summary of 22 bacterial and archaeal strains used in this study

Species and strain	Family	Origin		Type of reactor	Fed substrate	T (°C) of reactor	Reference for the isolation strategy or strain origin	Closest related NCBI GenBank entry with a validly published taxonomic affiliation	Similarity of 16S rRNA gene between isolate and GenBank entry (%)	NCBI GenBank entry of closest relative
		Latitude	Longitude							
<i>Bacteria</i>										
<i>Clostridium cellulosi</i> DG5	Clostridiaceae	51.255499	6.396524	Liquid pump/wet fermentation	Maize, pig manure, grass	54	[18] ^b	<i>Clostridium cellulosi</i> AS1.1777	98.8	LN881577
<i>Clostridium</i> sp. N3C		51.255499	6.396524	Liquid pump/wet fermentation	Maize, pig manure, grass	54	[18] ^c	<i>Clostridium putrefaciens</i> DSM 1291 ¹	93.0	NR113324
<i>Clostridium borinimense</i> M2/40 ¹		52.3871	13.0993	Lab-scale UASS/wet fermentation	Maize silage, wheat straw	37	[20]	<i>Clostridium borinimense</i> M2/40 ¹	100	JO388596
<i>Clostridium thermocellum</i> BC1		48.135125	11.581981	Bio-waste compost treatment site close to BGP		60	[18] ^d	<i>Clostridium thermocellum</i> DSM 1237 ¹	99.0	NR074629
<i>Proteiniborus</i> sp. DW1	Clostridiales incertae sedis	49.512893	7.083068	CSTR, wet fermentation	Maize silage, grass, cattle manure	39	[21]	<i>Proteiniborus ethanolyticus</i> GW ¹	96.0	NR044093
<i>Sporanaerobacter</i> sp. PP17-6a		51.255499	6.396524	Lab-scale CSTR/wet fermentation	Maize silage, pig manure, cattle manure	37	This study	<i>Sporanaerobacter acetigenes</i> Lup33	91.0	NR025151
<i>Herbinix hemicellulosilytica</i> T3/55 ¹	Lachnospiraceae	51.255499	6.396524	Liquid pump/wet fermentation	Maize, pig manure, grass	54	[18, 54] ^b	<i>Herbinix hemicellulosilytica</i> T3/55 ¹	100	LN626355
<i>Herbinix luporum</i> SD1D ¹		51.255499	6.396524	Liquid pump/wet fermentation	Maize, pig manure, grass	54	[18, 55] ^b	<i>Herbinix luporum</i> SD1D ¹	100	LN626359
<i>Peptoniphilaceae bacterium</i> str. ING2-D1G	Peptoniphilaceae	51.255499	6.396524	Lab-scale CSTR/wet fermentation	Maize silage, pig manure, cattle manure	37	[22]	<i>Peptoniphilus indolicus</i> DSM 20464 ¹	90.6	AY153431
<i>Propionispora</i> sp. 2/2-37	Veillonellaceae	48.3924	11.7569	CSTR, wet fermentation	Maize silage, grass	38	[18] ^e	<i>Propionispora hippiei</i> KS ¹	95.0	NR036875
<i>Bacillus thermoamylovorans</i> 1A1	Bacillaceae	48.3924	11.7569	CSTR, wet fermentation	Maize silage, pig manure	52	[18] ^f	<i>Bacillus thermoamylovorans</i> DKP ¹	99.0	NR029151

Table 1 continued

Species and strain	Family	Origin		Type of reactor	Fed substrate	T (°C) of reactor	Reference for the isolation strategy or strain origin	Closest related NCBI GenBank entry with a validly published taxonomic affiliation	Similarity of 16S rRNA gene between isolate and GenBank entry (%)	NCBI GenBank entry of closest relative
		Latitude	Longitude							
<i>Proteiniphilum saccharofermentans</i> M3/6 ^T	Porphyromonadaceae	52.3871	13.0993	Lab-scale UASS/wet fermentation	Maize silage, wheat straw	37	[26]	<i>Proteiniphilum saccharofermentans</i> M3/6 ^T	100	KP233809
<i>Fermitimonas caenicola</i> ING2-E5B ^T		51.255499	6.396524	Lab-scale CSTR/wet fermentation	Maize silage, pig manure, cattle manure	37		<i>Fermitimonas caenicola</i> ING2-E5B ^T	100	KP233810
<i>Petrimonas mucosa</i> ING2-E5A ^T		51.255499	6.396524	Lab-scale CSTR/wet fermentation	Maize silage, pig manure, cattle manure	37		<i>Petrimonas mucosa</i> ING2-E5A ^T	100	KP233808
<i>Defluviitoga tunisiensis</i> L3	Petrogocaceae	51.255499	6.396524	Liquid pump/wet fermentation	Maize, pig manure, grass	54	[27]	<i>Defluviitoga tunisiensis</i> Sulflac1 ^T	99.9	NR122085
Archaea										
<i>Methanobacterium formicicum</i> MF ^T	Methanobacteriaceae	DSMZ ^a				37	[50]	<i>Methanobacterium formicicum</i> MF ^T	100	NR115168
<i>Methanobacterium formicicum</i> Mb9		49.878359	6.481390	CSTR, wet fermentation	Maize silage, grass, cattle manure	40	[21]	<i>Methanobacterium formicicum</i> MF ^T	100	NR115168
<i>Methanobacterium</i> sp. Mb1		49.512893	7.083068	CSTR, wet fermentation	Maize silage, grass, cattle manure	39		<i>Methanobacterium formicicum</i> MF ^T	98.0	NR115168
<i>Methanobacterium congolense</i> Buetzberg		53.736687	10.083949	CSTR, dry fermentation	Household garbage	37	[18] ^g	<i>Methanobacterium congolense</i> C ^T	99.0	NR028175
<i>Methanothermobacter wolfei</i> SV6		51.255499	6.396524	Liquid pump/wet fermentation	Maize, pig manure, grass	54	[18] ^h	<i>Methanothermobacter wolfei</i> VKM B-1829 ^T	100	NR040964.1

Table 1 continued

Species and strain	Family	Origin		Type of reactor	Fed substrate	T (°C) of reactor	Reference for the isolation strategy or strain origin	Closest related NCBI GenBank entry with a validly published taxonomic affiliation	Similarity of 16S rRNA gene between isolate and GenBank entry (%)	NCBI GenBank entry of closest relative
		Latitude	Longitude							
<i>Methanococcus bourgenis</i> MS2 ¹	<i>Methanomicrobacteriaceae</i>	DSMZ				37	[49]	<i>Methanococcus bourgenis</i> MS2 ¹	100	NR042786
<i>Methanococcus chikugoensis</i> L21-II-0		51.255499	6.396524	Lab-scale CSTR/wet fermentation	Maize silage, pig manure, cattle manure	37	This study	<i>Methanococcus chikugoensis</i> MG62 ¹	99.0	NR028152

CSTR, continuously stirred tank reactor; UASS, upflow anaerobic solid-state reactor

^a DSMZ, Leibniz Institute DSMZ-German Collection of Microorganisms and Cell Cultures, Braunschweig, Germany

^b Isolation strategy number four described in more detail by [18]

^c Isolation strategy number eight (a) published in [18]

^d Isolation strategy number five published in [18]

^e Isolation strategy number seven published in [18]

^f Isolation strategy number two published in [18]

^g Isolation strategy number ten published in [18]

^h Isolation strategy number eleven published in [18]

cultivation occurred by successive transfer of culture aliquots after incubation periods of 4 weeks into fresh medium supplemented with different combinations of antibiotics as mentioned above for isolation of the strain L21-II-0. After 14 cultivation cycles, isolation of the bacterial strain was performed by plating of the culture material on BBL™ Columbia Agar Base medium (Th. Geyer, Germany) supplemented with 5% laked horse blood (Oxoid, Germany). For purification, single colonies were picked and re-streaked, and incubation occurred at 37 °C.

Phylogenetic classification of the analyzed bacterial and archaeal strains

To determine the phylogenetic relationship between the different strains and closely related type strains, a phylogenetic tree was constructed. For this, the 16S rRNA gene sequences retrieved from the genome sequences of the analyzed strains were aligned using the SINA alignment service v.1.2.11, which is provided online [32]. Subsequently, the SINA alignment and the All-Species Living Tree LTPs123 [33] from the SILVA ribosomal RNA project [34], only consisting of the 16S rRNA gene sequences of validly described type strains, were loaded into the ARB program [35]. Finally, the SINA alignment was placed into the existing LTP tree using ARB's parsimony method. Only type strains closely related to the corresponding isolate analyzed within this study are shown in the tree, whereas the remaining type strains were hidden manually applying "remove species from the tree" function implemented in ARB.

Genomic DNA extraction, sequencing, and bioinformatic analyses of biogas community members

Whole genome sequences of 13 strains, which were used in this study, were published previously (references given in Table 2). Genome sequencing of the following strains was performed within this study: *Proteiniborus* sp. DW1, *Clostridium* sp. N3C (DSM 100067), *Sporanaerobacter* sp. PP17-6a, *Proteiniphilum saccharofermentans* M3/6^T, *Petrimonas mucosa* ING2-E5A^T, *Methanobacterium formicicum* Mb9, *Methanobacterium congolense* Buetzberg, [36] *Methanothermobacter wolfeii* SIV6, and *M. chikugoensis* L21-II-0. In the case of *Clostridium* sp. N3C, *Sporanaerobacter* sp. PP17-6a, and *P. saccharofermentans* M3/6^T, genomic DNA was extracted applying the innuPREP Bacteria DNA Kit (Analytik Jena, Germany). Genomic DNA of *P. mucosa* ING2-E5A^T and *M. chikugoensis* L21-II-0 was extracted as described previously [37]. Genomic DNA of the strain *Proteiniborus* sp. DW1 was obtained applying the protocol published previously [19] and genomic DNA from *M. congolense* Buetzberg was extracted from 10 × 10 ml of a liquid culture using

the Gene Matrix stool DNA purification kit (Roboklon, Germany). DNA of strain *M. wolfeii* SIV6 was obtained applying the FastDNA Spin Kit for Soil (MP Biomedicals).

For bacterial strains mentioned above, 4 µg of purified chromosomal DNA was used to construct an 8-k mate-pair sequencing library (Nextera Mate Pair Sample Preparation Kit, Illumina Inc., Eindhoven, Netherlands) and sequenced applying the mate-pair protocol on an Illumina MiSeq system. Sequencing libraries of the archaeal strains *M. chikugoensis* L21-II-0 and *M. wolfeii* SIV6 were made from 2 µg of chromosomal DNA using the TruSeq DNA PCR-Free Library Preparation Kit (Illumina Inc., Eindhoven, Netherlands) and sequenced applying the paired-end protocol on an Illumina MiSeq system.

The obtained sequences were de novo assembled using the GS de novo Assembler Software (version 2.8, Roche). An in silico gap closure approach was performed [38], which resulted in a draft genome sequence or in a circular chromosome. Gene prediction and annotation of the genomes were performed within the GenDB 2.0 annotation system [39]. Manual metabolic pathway reconstruction was carried out by means of the KEGG pathway mapping implemented in GenDB that compares gene sequences with the corresponding gene product sequences of the NCBI database, with pairwise protein sequence identity being at least 30%. To predict genes encoding carbohydrate-active enzymes, the carbohydrate-active enzyme database (CAZy) annotation web-server dbCAN [40] was used.

Prevalence of the investigated strains within microbial communities of four different agricultural biogas plants applying the metagenome fragment recruitment approach

To evaluate the prevalence of the 22 analyzed strains within the microbial communities of the four different BGPs described previously [41], the corresponding metagenome sequences available for these BGPs (metagenome Accession Nos. at the NCBI database: SRA357208-09, SRA357211, SRA357213-14, SRA357221-23) were mapped on the genome sequences of these isolates with FR-HIT (v0.7; [42]) to sensitively recruit also metagenomic reads with lower sequence identity (global alignment down to 75% nucleotide sequence identity; Additional file 1).

As a baseline to compare against, four known and abundant metagenome-assembled genomes (MAGs) published previously [41] were included (the fifth genome bin 206_Thermotogae matching *Defluviitoga tunisiensis* L3 was excluded, because it is contained in the isolate collection; Table 1).

Furthermore, Mash (v1.1; [43]) was used to quickly identify potentially abundant and publicly available genome sequences in RefSeq (as of June 14, 2016; [44]).

Table 2 Genome features of 22 bacterial and archaeal strains used in this study

Species and strain	Assembly status		Genome size (bp)	GC content (%)	No. of genes	No. of <i>rrm</i> operons	No. of rRNA genes	No. of protein coding genes	EBI accession no.	References
	Genome structure	No. of contigs								
<i>Bacteria</i>										
<i>Clostridium cellulosi</i> DG5	CCC	n.a.	2,229,578	44.15	2088	6	59	2017	ERP006074	[53]
<i>Clostridium</i> sp. N3C	Draft genome	109	3,037,440	32.43	2880	3	66	2880	FMJL01000001–FMJL01000109	This study
<i>Clostridium bornimense</i> M2/40 ^T	CCC	n.a.	2,917,864	29.78	2694	8	56	2613	HG917868	[37]
<i>Clostridium thermocellum</i> BC1	Chromid		699,161	28.09	680	0	0	680	HG917869	
<i>Proteiniborus</i> sp. DW1 ^a	Draft genome	139	3,454,918	39.10	3094	4	52	3095	CBQ0010000001–CBQ0010000139	[61]
<i>Sporanaerobacter</i> sp. PP17-6a	Draft genome	62	3,121,392	32.44	2795	3	40	1793	FMDO01000001–FMDO01000062	This study
<i>Herbinix hemi-cellulosilytica</i> T3/55 ^T	Draft genome	53	3,296,672	33.45	3148	1	46	3148	FMIF01000001–FMIF01000053	This study
<i>Herbinix luporum</i> SD1D ¹	CCC	n.a.	3,037,031	36.69	2681	4	35	1726	CVTD0200000001–CVTD0200000035	[24]
<i>Peptoniphilaceae</i> bacterium str. ING2-D1G	CCC	n.a.	1,601,846	34.85	1541	4	53	1517	LN879430	[78]
<i>Propionispora</i> sp. 2/2–37	Draft genome	43	4,122,013	45.58	3690	1	76	2685	CYSP01000001–CYSP01000043	[29]
<i>Bacillus thermoamylovorans</i> 1A1	Draft genome	106	3,708,331	37.28	3472	10	59	2957	CCRF01000001–CCRF01000106	[79]
<i>Proteomphilum saccharofermentans</i> M3/6 ^T	CCC	n.a.	4,414,963	43.63	3450	3	48	3447	LT605205	This study
<i>Fermentimonas caenicala</i> ING2-ESB ^T	CCC	n.a.	2,808,926	37.30	2455	2	44	2405	LN515532	[25]
<i>Petrimonas mucosa</i> ING2-E5A ¹	CCC	n.a.	3,362,317	48.24	2693	2	46	2693	ERS1319466	This study

Table 2 continued

Species and strain	Assembly status		Genome size (bp)	GC content (%)	No. of genes	No. of <i>rrn</i> operons	No. of tRNA genes	No. of protein coding genes	EBI accession no.	References
	Genome structure	No. of contigs								
<i>DeFluviitoga tunisiensis</i> L3	CCC	n.a.	2,053,097	31.38	1881	3	47	1815	LN824141	[23]
Archaea										
<i>Methanobacterium formicicum</i> MF ^T	CCC	n.a.	2,478,074	41.23	2409	2	44	2100	LN515531	[80]
<i>Methanobacterium formicum</i> Mb9	CCC	n.a.	2,494,510	41.14	2416	2	43	2126	ERS549551	This study
<i>Methanobacterium</i> sp. Mb1	CCC	n.a.	2,029,766	39.74	2021	2	41	1689	HG425166	[19]
<i>Methanobacterium congolense</i> Buetzberg	CCC	n.a.	2,459,553	38.48	2351	3	41	2351	LT607756	This study
	Plasmid		18,118	36.05	24	0	0	24	LT607757	
<i>Methanothermobacter wolfei</i> SIV6	CCC	n.a.	1,686,891	48.89	1793	2	36	1444	ERS1319767	This study
<i>Methanoculleus bourgensis</i> MS2 ^T	CCC	n.a.	2,789,773	60.64	2586	1	45	2586	HE964772	[81]
<i>Methanoculleus chikugoensis</i> L21-II-0	Draft genome	70	2,649,997	61.83	2671	1	45	2671	FMID01000001–FMID01000070	This study

CCC, circularly closed chromosome; n.a., not applicable

^a The strain *Proteiniborus* sp. DW1 was cultivated together with *Methanobacterium* sp. Mb1; the DW1 genome sequence was recovered from sequencing of a mixed culture consisting of strains DW1 and Mb1

The meaning of abundance in this context refers exclusively to the number of metagenome sequences mapped to the genome sequence. For a sketch size of 1,000,000 and a k-mer size of 21, pairwise distances between the metagenomic read sets and all 5061 genomes in RefSeq (plus, as a control, the 22 strains from this study) were calculated. Requiring a minimum of 20 k-mer hits not only confirmed the potential relevance of the selected 22 strains, but additionally identified 46 publicly available strains from RefSeq for further analyses.

All metagenome sequences available for the four BGPs were mapped on the genome sequences of these isolates, the four MAGs, and the 46 reference strains with Kallisto [45] (v0.43.1). For each genome, the GPM (genomes per million) values were calculated using the TPM (transcripts per million) values reported by Kallisto (see Additional file 3).

Results and discussion

Selection of a set of microbial isolates from different biogas-producing communities

Limited availability of genome sequence information in public databases for AD community members generally constrains the interpretation of metagenomic and metatranscriptomic data of such communities leading to large amounts of non-classifiable metagenome sequences from AD habitats [15–18, 46, 47]. Accordingly, parallel application of both traditional culturomics [48] as well as molecular analysis combined with HT sequencing techniques is necessary for detailed studies of complex microbial biogas consortia. Applying 16 different isolation strategies, bacterial and archaeal isolates were obtained from different mesophilic and thermophilic production- and laboratory-scale BGPs (Table 1). Furthermore, two archaeal members, namely *M. bourgensis* MS2^T [49] and *M. formicicum* MF^T [50], were obtained from the DSMZ and included in this study as the reference strains for methanogenic *Archaea* since they were also isolated from AD communities. German BGPs sampled for this study differed in utilized substrates ranging from maize silage, grass, and wheat straw to cattle and/or pig manure. Moreover, one digester analyzed was fed with organic residues and waste material as substrate. Additionally, a bio-waste compost treatment site close to the city of Munich (Germany) was sampled to isolate cellulolytic bacteria. Besides different renewable biomass sources utilized for the AD process, the biogas reactors differed regarding digester design, fermentation technology, and the applied temperature regime ranging from 37 to 54 °C.

This study comprises the analysis of 15 bacterial strains classified as belonging to the phyla *Firmicutes*, *Thermotogae*, and *Bacteroidetes* and seven archaeal isolates of the phylum *Euryarchaeota*. Details on all isolates of this

study, their taxonomy, their origin, and the respective isolation strategy applied are provided in Table 1.

Phylogenetic classification of the microbial isolates selected from different biogas communities


To determine the taxonomic position of the strains analyzed, their 16S rRNA gene sequences were compared to the corresponding sequences from closely related type strains deposited in the SILVA database (Fig. 1). The calculated phylogenetic tree comprises four main groups representing the phyla *Bacteroidetes*, *Firmicutes*, *Thermotogae*, and *Euryarchaeota*. Among the *Bacteroidetes* members, the strains *P. saccharofermentans* M3/6^T, *P. mucosa* ING2-E5A^T, and *Fermentimonas caenicola* ING2-E5B^T were recently described as novel species and were suggested to participate in hydrolysis and acidogenesis of the AD process [26].

Most of the bacterial strains analyzed were allocated to the phylum *Firmicutes*, and within this taxon to the classes *Clostridia*, *Bacilli*, *Tissierellia*, and *Negativicutes*. A diverse group of isolates belong to the class *Clostridia*. They are related to characterized species such as *Clostridium cellulosi* (also denominated as '*Ruminiclostridium cellulosi*'), *Clostridium thermocellum* (also denominated as '*Ruminiclostridium thermocellum*' [51], *Clostridium cellulovorans*, and *Clostridium bornimense*. The latter one was recently described as novel species [20]. All mentioned species represent lignocellulosic biomass degraders [20, 52, 53]. Two other *Clostridia* isolates, namely T3/55^T and SD1D^T, were recently assigned to the species *Herbinix hemicellulosilytica* [54] and *Herbinix luporum* [55], respectively, of the new genus *Herbinix*. Both strains are distantly related to the type strain *Mobilitalea sibirica* P3M-3^T [56] and were described to be involved in thermophilic degradation of lignocellulosic biomass.

The isolates 1A1, ING2-D1G, and 2/2-37 are closely related to the species *Bacillus thermoamylovorans* (class *Bacilli*), *Peptoniphilus indolicus* (class *Tissierellia*), and *Propionispora hippie* (class *Negativicutes*), respectively. The corresponding reference strains were described to perform hydrolytic and acidogenic functions in the AD process [57–59].

Another isolate from a thermophilic BGP was classified as *D. tunisiensis* (phylum *Thermotogae*, class *Thermotogae*) representing an isolated branch of the bacterial part of the tree (Fig. 1). The strain *D. tunisiensis* L3 was described to be adapted to high temperatures and able to utilize different complex carbohydrates to produce ethanol, acetate, H₂, and CO₂ [27, 28]. The latter three metabolites represent substrates for methanogenic *Archaea*.

The strains *Sporanaerobacter* sp. PP17-6a and *Pep-toniphilaceae* bacterium str. ING2-D1G are only distantly related to known bacterial species of the family



Clostridiales incertae sedis and *Peptoniphilaceae* (90–91% identity), respectively, suggesting that they represent new species.

The fourth group of the phylogenetic tree represents methanogenic *Archaea* classified as members of the classes *Methanomicrobia* and *Methanobacteria* (both belonging to the phylum *Euryarchaeota*). Members of these classes were described to perform hydrogenotrophic methanogenesis utilizing CO₂ and H₂ as substrates for CH₄ synthesis [18, 21].

Genome sequence analyses of the whole set of microbial isolates selected

To gain insights into the functional potential of all strains listed in Table 1, their genomes were completely sequenced by application of HT sequencing technologies. Genome sequence information provides the basis for metabolic reconstruction and assignment of functional roles within the AD process, thus enabling biotechnological exploitation of genome features involved in fermentation processes utilizing renewable primary products.

Out of 22 genome sequences, nine, namely those of *Proteiniborus* sp. DW1, *Clostridium* sp. N3C, *Sporan-aerobacter* sp. PP17-6a, *P. saccharofermentans* M3/6^T, *P.*


mucosa ING2-E5A^T, *M. formicicum* Mb9, *M. congolense* Buetzberg, *M. wolfeii* SIV6, and *M. chikugoensis* L21-II-0, were newly established in this study. Genome sequences of the remaining 13 strains were published previously mainly in the form of Genome Announcements (for references, refer to Table 2). The genome sequences of the microorganisms analyzed were established on an Illumina MiSeq system. In silico and PCR-based gap closure strategies resulted in 13 finished and nine draft genome sequences. General genome features, e.g., genome structure, assembly status, size, GC content, and numbers of predicted genes, are summarized in Table 2. Established genomes range in size from 1.6 to 4.4 Mb and feature GC contents from 28.09 to 61.83%. Moreover, *C. bornimense* M2/40^T, in addition to the chromosome, harbors a 699,161-bp chromid (secondary replicon) in its genome containing 680 coding sequences [37]. The methanogen *M. congolense* Buetzberg also harbors an accessory genetic element, namely a plasmid featuring a size of 18,118 bp. Genome annotation applying the GenDB 2.0 platform enabled functional interpretation of genes and reconstruction of metabolic pathways involved in the AD process. Genome analyses provided insights into the lifestyle and functional roles of bacterial and archaeal strains.

Screening of the subset of bacterial genomes to identify genes encoding carbohydrate-active enzymes potentially involved in biomass degradation

To elucidate genes encoding carbohydrate-active enzymes, functional genome annotation applying the HMM-based carbohydrate-active enzyme annotation database dbCAN [40] was performed (Fig. 2). Between 71 and 358 genes encoding enzymes or modules with predicted activity on carbohydrates were identified in each of the bacterial strains analyzed. Among them are dockerin-containing glycoside hydrolases (GH), representing putative cellulosomal enzymes, corresponding cohesin-containing scaffoldins, enzymes acting on large carbohydrate molecules, and carbohydrate-binding motifs involved in sugar binding. The obtained results separate the analyzed strains into two groups: group I strains were predicted to degrade cellulose and hemicellulose, whereas group II strains represent secondary fermentative bacteria relying on metabolites (mainly mono-, di-, and oligosaccharides) produced by group I members (as obvious presence of cellulolytic genes). The *Clostridiaceae* strains DG5, T3/55^T, SD1D^T, M2/40^T, and BC1 harbor a more diverse repertoire of genes involved in the degradation of complex polysaccharides such as cellulose (GH5, GH8, GH9, GH48), xylan (GH10, GH11), and cellobiose- or cellodextrin-phosphorylase genes (GH94). Furthermore, genes for cohesin-containing putative scaffoldins and the corresponding dockerin-containing glycoside hydrolases with a potential for cellulosome formation were also identified in the genomes of these strains. Previous studies reported on the importance of the phylum *Firmicutes*

for hydrolysis of cellulosic material in biogas digesters [12, 60]. In particular, *Clostridiaceae* and *Ruminococcaceae* members are involved in this first step of biomass digestion [11, 18]. *Clostridiaceae* strains *Proteiniborus* sp. DW1 and *Clostridium* sp. N3C were predicted to represent non-cellulolytic isolates (Fig. 2), whereas the cellulolytic strain *C. thermocellum* BC1 [61] is known to be a very efficient cellulose degrader since it encodes cellulosome components and is able to degrade hemicelluloses and pectins [60]. In contrast to the cellulolytic *Clostridiaceae*, the *Porphyromonadaceae* members, namely *P. saccharofermentans* M3/6^T, *P. mucosa* ING2-E5A^T, and *F. caenicola* ING2-E5B^T, encode enzymes predicted to degrade pectins and a variety of hemicelluloses (GH16, GH26, GH28, GH30, GH53, GH74). These strains do not seem to be able to hydrolyze arabinoxylan (lack of GH10, GH11) and crystalline cellulose (lack of GH48). Likewise, *D. tunisiensis* L3 (*Petrotogaceae* family) also possesses a large set of genes predicted to facilitate cleavage of a variety of sugars including cellobiose, arabinosides (GH27), chitin (GH18), pullulan and starch (GH13), and lichenan (GH16) [28].

Another strain supposed to represent a secondary fermentative bacterium, namely *B. thermoamylovorans* 1A1 (*Bacillaceae* family), may contribute to oligosaccharide degradation with genes for GH1, GH2, GH3, or GH43 enzymes. In addition, genes required for growth on cellobiose are present in its genome. Considering the fact that strain 1A1 originally was isolated from a co-culture also containing *C. thermocellum* [61], it is assumed that *B. thermoamylovorans* 1A1 further metabolizes cellobiose produced by cellulolytic *Clostridia*.



Members of the genus *Propionispora* (*Veillonellaceae*) previously were identified in AD communities [62] and predicted to utilize mostly sugars and sugar alcohols, e.g., glucose, fructose, xylitol, or mannitol for growth [59]. The strain *Propionispora* sp. 2/2–37 analyzed in this study additionally harbors genes encoding enzymes participating in cellobiose, starch, and chitin degradation as determined by means of the CAZy analysis.

In contrast, the results obtained for *Peptoniphilaceae* bacterium str. ING2-D1G showed that this bacterium does not encode enzymes involved in the degradation of complex carbohydrates. However, the strain ING2-D1G encodes all enzymes needed to utilize amino acids and monomeric carbohydrates as a carbon source [22]. Its function in the anaerobic digestion process can be hypothesized to be associated with acidogenesis, which was supported by reconstruction of corresponding metabolic pathways.

Prediction of fermentation pathways based on sequence information for the subset of bacterial genomes

Bacteria involved in AD perform a number of different fermentation pathways to recycle reduction equivalents that are produced in the course of metabolite utilization. To determine the fermentation type and the functional role of a given isolate within the biogas process, enzymes encoded in its genome were assigned to selected fermentation pathways represented in the KEGG database (Table 3, Additional file 2 and Fig. 3). Pathways leading to propionate, ethanol, formate, butyrate, acetate, and lactate synthesis were considered in this approach.

Certain bacteria are able to convert sugars, acids, alcohols, or amino acids to propionic acid under anaerobic conditions utilizing the methylmalonyl-CoA or the acrylyl-CoA pathways of the propanoate metabolism [27]. Among the analyzed bacteria, the strains *Propionispora* sp. 2/2–37, *P. saccharofermentans* M3/6^T, *P. mucosa* ING2-E5A^T, and *F. caenicola* ING2-E5B^T encode all enzymes of the methylmalonyl-CoA pathway for the production of propionic acid from pyruvate. Only the strain *Proteiniborus* sp. DW1 was predicted to utilize lactate for propionic acid production via the acrylyl-CoA pathway. Since the enrichment of propionic acid was described as an indicator for process imbalance [27, 63], data on the physiology of propionic acid-producing bacteria can be valuable for the optimization of the biogas plants.

Butyric acid-forming bacteria in biogas systems have been insufficiently characterized so far [27]. Genes encoding enzymes required for butyric acid formation via the butanoate pathway were found in the genomes of the strains *Propionispora* sp. PP16-6a, *Peptoniphilaceae* bacterium str. ING2-D1G, *C. bornimense* M2/40^T, *P. saccharofermentans* M3/6^T, *Clostridium* sp. N3C, *P. mucosa* ING2-E5A^T, *F. caenicola* ING2-E5B^T, and *B.*

thermoamylovorans 1A1. Butanoate production was recently described for the strains *H. luporum* SD1D^T [55] and *H. hemicellulosilytica* T3/55^T [54]. However, the genomes of these bacteria only encode the last two enzymes of the butanoate pathway, namely the phosphate butyryl transferase Ptb and butyrate kinase Buk, predicted to be responsible for butanoate synthesis in these strains.

During acidogenesis, volatile organic compounds such as ethanol, acetate, and formate are produced in the course of the AD process. The latter two metabolites are substrates for methanogenic *Archaea*. Analysis of pathways involved in ethanol, acetate, and formate synthesis, i.e., the mixed-acid fermentation, revealed that all analyzed bacteria harbor genes encoding enzymes of this pathway (see Additional file 2). With the exception of the *Peptoniphilaceae* bacterium str. ING2-D1G, in all other isolates the necessary genes to produce ethanol from pyruvate were identified. Moreover, genes encoding enzymes participating in formate production were found in the *C. cellulosi* DG5, *C. bornimense* M2/40^T, *D. tunisiensis* L3, *C. thermocellum* BC1, and *B. thermoamylovorans* 1A1 genomes. Furthermore, all analyzed bacteria were predicted to be able to produce acetate from acetyl-CoA. Genes encoding the enzymes phosphate acetyltransferase Pta (EC: 2.3.1.8) and acetate kinase Ack (EC: 2.7.2.1), converting acetyl-CoA to acetyl phosphate and subsequently to acetate, were found. In addition, genes encoding the enzymes pyruvate decarboxylase Pdc (EC: 4.1.1.1) and alcohol dehydrogenase Adh (EC: 1.1.1.1), converting pyruvate to acetaldehyde and finally to ethanol, were found in all genomes with the exception of the strain *Peptoniphilaceae* bacterium str. ING2-D1G, which does not possess an *adh* gene. Surprisingly, in the case of the strains *P. mucosa* ING2-E5A^T, *F. caenicola* ING2-E5B^T, and *P. saccharofermentans* M3/6^T, no ethanol production was observed in growth experiments [26]. Possibly, the growth conditions tested might not be favorable to support ethanol synthesis.

Many bacterial species produce 2,3-butanediol under anaerobic conditions from glucose, with *Klebsiella oxytoca* and *Bacillus licheniformis* described as efficient 2,3-butanediol producers [64]. Among the bacteria analyzed, only *Propionispora* sp. 2/2–37 harbors a full set of genes encoding all necessary enzymes (refer to Additional file 2).

Lactic acid was found to be the main fermentation product from household waste digestion [65]. Members of the genera *Bacillus*, *Lactobacillus*, *Leuconostoc*, *Pediococcus*, and *Streptococcus* were previously described to produce lactic acid from several types of sugars [12, 47, 66]. To determine whether the analyzed bacteria have the potential to produce lactic acid, the genomes were

Table 3 continued

Pathway analyzed	Predicted product after fermentation		Peptoniphilaceae bacterium str. ING2-DTG		Propionispora sp. 2/2-37		Bacillus thermoamylovorans 1A1		Proteophilum saccharofermentans M3/6 ^T		Fermentimonas caenicola ING2-E5B ^T		Petrimonas mucosa ING2-E5A ^T		Defluviitoga tunisiensis L3	
	GP	EP ^a	GP	EP ^a	GP	EP ^a	GP	EP ^a	GP	EP ^e	GP	EP ^e	GP	EP ^e	GP	EP ^f
Propionic acid fermentation ⁹	-	ND	-	D	-	D	-	ND	-	D	-	D	-	D	-	ND
Acrylyl-CoA pathway	-	-	+	+	+	+	+	+	+	+	+	+	+	+	-	-
Methylmalonyl-CoA pathway	-	-	+	+	+	+	+	D	+	ND	+	ND	+	ND	+	+
Ethanol fermentation	-	-	+	+	+	+	+	+	+	+	+	+	+	+	+	+
Formic acid fermentation	-	-	+	+	+	+	+	+	+	+	+	+	+	+	+	+
2,3-Butanediol fermentation	-	-	+	ND	+	ND	+	ND	-	-	-	-	-	-	-	-
Formic acid	-	-	+	+	+	+	+	+	+	+	+	+	+	+	+	-
CO ₂ and H ₂	-	-	-	-	-	-	-	-	-	-	-	-	-	-	+	+
Ethanol	-	-	+	D	+	D	+	D	+	+	+	+	+	+	-	-
Mixed-acid fermentation	+	D	+	+	+	+	+	+	+	+	+	+	+	+	+	D
Acetate	+	ND	+	ND	+	ND	+	ND	+	+	+	+	+	+	+	ND
Lactate	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
Succinate	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Butyrate	+	D	+	D	+	D	+	D	+	+	+	+	+	+	+	D
Acetate	+	D	+	D	+	D	+	D	+	+	+	+	+	+	+	D
Lactic acid fermentation	+	ND	+	ND	+	ND	+	ND	+	+	+	+	+	+	+	+
Homolactic acid fermentation	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
Heterolactic acid fermentation	+	D	+	D	+	D	+	D	+	+	+	+	+	+	+	-
Acetate	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	D
Ethanol	+	ND	+	+	+	+	+	+	+	+	+	+	+	+	+	ND

Genomic loci encoding enzymatic functions participating to the corresponding fermentation type for each bacterial strain analyzed are listed in Additional file 2

+; synthesis of the corresponding fermentation end-product is predicted; -, pathway incomplete or misses key enzymes, the synthesis of the corresponding fermentation end-product is doubtful; EP, experimental proof; D, the corresponding fermentation product has been experimentally detected; GP, genes predicted applying metabolic reconstruction within the GenDB 2.0 system [39]; NA, not analyzed; NC, not confirmed; ND, fermentation product has been experimentally not detected

^a Unpublished data

^b Data published in [20]

^c Data published in [54]

^d Data published in [55]

^e Data published in [26]

^f Data published in [27]

⁹ Pathways for propionic acid synthesis via succinate decarboxylation or amino acid degradation were not included

screened for encoded enzymes involved in homolactic and heterolactic acid fermentation. With the exception of the strain *Sporanaerobacter* sp. PP17-6a, all other bacterial genomes were predicted to perform homolactic acid fermentation. They harbor all genes encoding necessary enzymes including the gene for lactate dehydrogenase Ldh (EC: 1.1.1.27) converting pyruvate to lactic acid. Furthermore, some genetic determinants of the heterolactic acid fermentation pathway were identified. However, none of the strains encodes a full set of the genes needed. Hence, the question which strains are responsible for lactic acid production remains unsolved.

Prediction of methanogenesis pathways based on sequence information for the subset of archaeal genomes

The formation of CH₄, the last step in the AD of biomass, is performed by methanogenic *Archaea* (Fig. 3). Based on their genetic repertoire, methanogens are able to perform either the hydrogenotrophic, acetoclastic, or methylotrophic pathway utilizing CO₂ and H₂, acetate, or methylamine and methanol, respectively, for CH₄ production [67]. To predict the pathway by which the analyzed *Archaea* produce CH₄, genes involved in the different methanogenesis pathways mentioned above




Fig. 3 Overview of the four phases of the conversion of biomass into biogas and allocation of the analyzed microbial strains to the different conversion steps. Functional roles of the organisms were determined considering relevant KEGG pathways, namely the propionic acid, ethanol, formic acid, butyric acid, and lactic acid fermentation

Table 4 Predicted genome features and traits of archaeal strains included in this study

Strain name	Features predicted						
	<i>Methanobacterium formicicum</i> MF ^T	<i>Methanobacterium formicicum</i> Mb9	<i>Methanobacterium</i> sp. Mb1	<i>Methanobacterium congolense</i> Buetzberg	<i>Methanothermobacter wolfei</i> SIV6	<i>Methanoculleus bourgensis</i> MS2 ^T	<i>Methanoculleus chikugoensis</i> L21-II-0
Methanogenesis-related hydrogenase genes encoded in the genome	<i>eha, ehb, frh, mvh, hdr</i>	<i>eha, ehb, frh, mvh, hdr</i>	<i>eha, ehb, frh, mvh, hdr</i>	<i>eha, ehb, frh, mvh, hdr</i>	<i>eha, ehb, frh, mvh, hdr</i>	<i>ech, frh, mvh, hdr</i>	<i>ech, frh, mvh, hdr</i>
Substrates used for methanogenesis	H ₂ /CO ₂ , F	H ₂ /CO ₂ , F	H ₂ /CO ₂ , F	H ₂ /CO ₂ , F	H ₂ /CO ₂ , F	H ₂ /CO ₂ , F	H ₂ /CO ₂ , F
Predicted metabolites required for growth	Acetate, cysteine ^a , vitamin B ^a	Acetate	Acetate	Acetate, lactate	Acetate	Acetate, lactate ^b	Acetate, lactate ^b

F, formate; H₂, hydrogen; CO₂, carbon dioxide

^a Utilization of cysteine and vitamin B by the strain MF^T was described previously [50]

^b No growth or methane production was detected on lactate for *Methanoculleus* species described previously [49, 82]

were examined interpreting functional KEGG assignments calculated within GenDB (Table 4).

All *Archaea* analyzed encode a full set of genes involved in CH₄ production from CO₂ and H₂. This result was as expected, as members of the families *Methanobacteriaceae* and *Methanomicrobiaceae* are known to solely perform hydrogenotrophic methanogenesis [68]. Additionally, genes for the formate dehydrogenase complex FdhA-B and a formate transporter FdhC for growth on formate as an alternative methanogenic substrate were identified in all seven analyzed genomes. For acetyl-CoA production from acetate, all seven genomes encode the acetyl-CoA synthetase Acs. Interestingly, methanogens from the genus *Methanoculleus*, namely the strains MS2^T and L21-II-0, also harbor a lactate dehydrogenase gene involved in conversion of lactate to pyruvate or vice versa. However, no growth or CH₄ production from lactate has been described for the *Methanoculleus* species so far.

For activation of H₂ during methanogenesis, all seven *Archaea* analyzed encode the cytoplasmic coenzyme F₄₂₀-reducing [NiFe]-hydrogenases FrhA-D, the cytoplasmic [NiFe]-hydrogenase MvhADG, and the heterodisulfide reductase HdrABC in their genomes. The latter two enzyme complexes interact with the cytoplasmic [NiFe]-hydrogenase MvhADG, which was also identified in all investigated methanogens, for the coupled H₂-driven reduction of ferredoxin and heterodisulfide CoM-S-S-CoB [69]. Furthermore, methanogens of the family *Methanobacteriaceae* encode the membrane-bound energy-converting [NiFe]-hydrogenases EhaA-T and EhbA-Q [70], whereas the *Methanomicrobiaceae* strains encode the energy-converting [NiFe]-hydrogenase EchA-F in their genomes. Members of the order *Methanomicrobiales* were described to exhibit a high affinity for H₂ (ca. 0.1 μM resp. 15 Pa H₂ pressure [71]), possibly providing an advantage over certain *Methanobacteriales* under conditions of low H₂ partial pressure.

Prevalence of bacterial and archaeal isolates in different microbial biogas communities analyzed by metagenome fragment mappings

To determine the prevalence or rather the abundance of the bacterial and archaeal isolates analyzed in this study in communities of production-scale BGPs, metagenome fragment mappings were done using deeply sequenced metagenomes from three mesophilic (BGP1-3) and one thermophilic (BGP4) agricultural BGPs which were published recently [41]. Configurations and process parameters corresponding to these BGPs are documented in the publication cited above. To identify metagenome sequence reads of the BGPs that match the genome sequences of the biogas isolates, these

were mapped to the genomes applying Kallisto. Reads assigned to certain genomes were summed up and normalized according to dataset and genome sizes analogous to TPM (transcripts per million, [72]) values in RNASeq studies, to allow for quantitative comparisons.


Metagenome fragment mapping results were distinguished into the following groups: (I) abundant fully covered genomes, (II) less abundant but fully covered genomes, (III) rare but fully covered genomes, and (IV) rare, partially covered genomes (examples for each group are shown in Additional file 1).

Only three genomes, namely those of *Methanoculleus bourgensis* MS2^T, *D. tunisiensis* L3, and *Clostridium* sp. N3C, fall into group I. *M. bourgensis* is abundant in all mesophilic BGPs studied and slightly less abundant in the thermophilic BGP, whereas *D. tunisiensis* and *Clostridium* sp. N3C are prominent in the thermophilic BGP (Fig. 4, Additional file 3).

Several of the analyzed strains were clearly detectable in the mesophilic BGPs but appeared to be only moderately abundant (group II). The strains *H. luporum* SD1D^T, *M. chikugoensis* L21-II-0, *Sporanaerobacter* sp. PP17-6a, and *M. wolfeii* SIV6 fall into this category. They are supposed to perform functions that are also taken by other community members. In other words, the corresponding microbial guilds are composed of several species featuring similar functionalities. Specific adaptation of species within a guild may refer to slight fluctuations in environmental conditions with one or the other species being more competitive under a particular condition.

The strains *C. bornimense* M2/40^T, *F. caenicola* ING-E5B^T, *H. hemicellulosilytica* T3/55^T, and *C. thermocellum* BC1 seem to be rare in most of the analyzed BGPs (group III), whereas the isolates *Proteiniborus* sp. DW1, *Peptoniphilaceae* bacterium str. ING-D1G, *P. mucosa* ING-E5A^T, *Methanobacterium* sp. Mb1, *P. saccharofermentans* M3/6^T, *B. thermoamylovorans* 1A1, *Propionispora* sp. 2/2-37, *M. formicicum* MF^T, *M. formicicum* Mb9, *M. congolense* Buetzberg, and *C. cellulosi* DG5 seem to be, if at all, of minor importance in most BGPs (group IV).

Furthermore, the non-cultivable fractions of the biogas microbiomes residing in BGPs 1 to 4 were studied by Stolze et al. [41], applying metagenome assembly combined with a binning method. This approach enabled the identification of novel and uncharacterized species represented by MAGs, namely 206_Thermotogae, 175_Fusobacteria, 138_Spirochaetes, 244_Cloacimonetes, and 120_Cloacimonetes. To determine the prevalence of these MAGs in the biogas microbiomes analyzed, fragment recruitments were performed. The obtained results showed that the species represented by the bin 175_Fusobacteria is abundant in the mesophilic BGP3, whereas both *Cloacimonetes* MAGs were abundant in BGP2 and



BGP3. Furthermore, all three MAGs represent fully covered genomes and therefore fall into the groups I and II in the case of 175_ *Fusobacteria* and both *Cloacimonetes* MAG, respectively. The bin 138_ *Spirochaetes* is detectable in the mesophilic BGP3 but appeared to be only moderately abundant (group III). The MAG 206_ *Thermotogae* is very similar to *D. tunisiensis* L3 showing an ANI (average nucleotide identity) value of 99.25%, indicating that these two members belong to the same species [73]. Fragment recruitments for such closely related microorganisms lead to random distribution of the corresponding metagenome sequences to both genome sequences resulting in underestimation of the abundances of both strains. Hence, the

206_ *Thermotogae* MAG was not further considered for fragment recruitments.

Among the publicly available reference species, only the genomes of *M. bourgensis* MAB1 [74] originating from a laboratory-scale biogas reactor and *Amphibacillus xylanus* NBRC 15112 [75], isolated from compost of manure with grass and rice straw, were almost completely covered with metagenome sequences featuring high matching accuracy. The bacterial species *A. xylanus* NBRC 15112 was found to be highly abundant within the BGP1 microbiome, whereas the hydrogenotrophic methanogen *M. bourgensis* MAB1 was dominant in the mesophilic digesters 2 and 3 (Fig. 4). The genomes of both strains fall into group I regarding their fragment

recruitment profiles. Among the microorganisms of group II, the species *C. clariflavum* involved in hydrolysis of cellulose and hemicellulose [76] and *Streptococcus suis* BM407, a human pathogen [77], were found to be nearly fully covered but less abundant.

Based on these findings, metagenome fragment mappings clearly showed that the culturomics approach led to isolation and characterization of dominant and therefore important members of the biogas microbiome. However, since it is assumed that many biogas community members cannot be cultured by currently available cultivation techniques, further prevalent key microorganisms remain to be discovered.

Conclusions

Application of high-throughput and *-omics* technologies such as metagenomics, metatranscriptomics, metaproteomics, and genomics for the analysis of biogas microbial communities is becoming increasingly important. However, currently, the interpretation of generated data is limited due to the restricted availability of the corresponding and appropriate reference genome sequences connected with functional and metabolic information in public databases.

In this study, whole genome sequence information for 22 bacterial and archaeal strains was analyzed with respect to their metabolic functions in AD communities. For 15 bacterial strains, their participation in hydrolysis and/or acidogenesis/acetogenesis of plant biomass decomposition was predicted and partially verified by *in vivo* characterization of pure cultures. *Clostridium cellulosi* DG5, *H. hemicellulosilytica* T3/55^T, *H. luporum* SD1D^T, and *C. thermocellum* BC1 represent cellulose degraders, while the nine remaining bacteria presumably play a role in acidogenesis and/or acetogenesis. The seven analyzed methanogenic *Archaea* were predicted to produce CH₄ via the hydrogenotrophic pathway, representing the final phase of the AD chain.

Among the microorganisms analyzed in this study, only two species, namely *M. bourgensis* and *D. tunisiensis*, were identified to play a dominant role within biogas microbial communities. *Defluviitoga tunisiensis* was proposed as a marker organism for the thermophilic biogas processes. This species is very versatile in the utilization of different sugars that can be converted to metabolites serving as substrates for methanogenesis. *Methanoculleus bourgensis* has frequently been found to dominate methanogenic sub-communities residing in production-scale BGP and is assumed to be well adapted to high-osmolarity conditions and ammonia/ammonium concentrations prevailing when manure is used as a substrate for biogas production. Furthermore, the fragment recruitment analysis of MAGs published by Stolze et al. [41] could also show that in addition to the classical cultivation and

isolation strategy, the metagenome assembly and binning approach may also enable the identification and characterization of previously unknown but abundant species featuring important functional potential in the context of the anaerobic digestion process.

It appeared that among the publicly available genomes only those of the species *A. xylanus*, *C. clariflavum*, and *C. thermocellum* were found to be well represented within biogas microbiomes, but do not reach the level of abundance as observed for *M. bourgensis* and *D. tunisiensis*. Surprisingly, among 5061 complete genome sequences archived in the public database NCBI, only those mentioned above seem to be of pronounced importance for agricultural biogas systems. Accordingly, the applied culturomics approach led to the isolation of further key AD species, thus providing genome sequence information for novel biogas community members. In the future, the non-cultivable fraction of AD communities should also be accessed by single-cell genomics to uncover genome sequence information of further, so far unknown biogas community members.

Additional files

Additional file 1. Fragment recruitment of metagenome sequences derived from four biogas-producing microbiomes to the genome sequences of the exemplarily chosen strains *Amphibacillus xylanus* NBRC 15112^T, *Clostridium* sp. N3C, *Fermentimonas caenicola* ING2-E5B^T, *Methanobacterium formicicum* MF^T and *Methanoculleus bourgensis* MAB1. The x-axis: microbial genome analyzed, y-axis: percent identities of mapped metagenome reads.

Additional file 2. Genomic loci encoding enzymatic functions participating in the propionic acid, ethanol, formic acid, butyric acid and lactic acid fermentation for each strain analyzed.

Additional file 3. List of the 72 most abundant bacterial and archaeal strains within the biogas microbial communities analyzed, their GPM (genomes per million) values and further coverage statistics.

Abbreviations

AD: anaerobic digestion; BGP: biogas plant; CCC: circularly closed chromosome; CAZymes: carbohydrate-active enzymes; CSTR: continuous stirred tank reactor; DSMZ: Leibniz Institute German Collection of Microorganisms and Cell Cultures; GH: glycosyl hydrolase; GPM: genomes per million; HT: high-throughput; KEGG: Kyoto Encyclopedia of Genes and Genomes; qPCR: quantitative 'real-time' polymerase chain reaction; TPM: transcripts per million; TRFLP: terminal restriction fragment length polymorphism; UASS: upflow anaerobic solid-state reactor; VFA: volatile fatty acids.

Authors' contributions

IM performed the phylogenetic classification, genome assembly, and annotation of microbial isolates, participated in the prediction of bacterial fermentation pathways based on genome sequence information, coordinated drafting, and drafted the corresponding parts of the manuscript. AB carried out the fragment recruitment analyses for the 5061 publicly available genomes plus the 22 strains from this study, contributed to the "Results and discussion" section, and revised the manuscript. YS participated in the prediction of bacterial fermentation pathways based on genome sequence information and revised the manuscript. SH contributed to isolation and characterization of acidogenic bacterial strains and additional methanogenic *Archaea* and drafted the corresponding parts of the manuscript. KGC isolated and characterized acidogenic

bacterial strains and revised the manuscript. DEK isolated and characterized cellulolytic, hydrolytic, and acidogenic bacterial strains, participated in the analyses of bacterial genes encoding carbohydrate-active enzymes and revised the manuscript. YSK and JK contributed to the isolation of methanogenic *Archaea* and revised the manuscript. JH contributed to the phylogenetic classification of the analyzed bacterial and archaeal isolates and drafted the corresponding part of the manuscript. DW participated in the genome assembly and annotation of microbial isolates, submitted the 22 genome sequences to the EBI database, and revised the manuscript. AW participated in bioinformatic data analysis and revised the manuscript. SO participated in the isolation and characterization of methanogenic archaeal strains and contributed to the results and discussion part of the manuscript on archaeal isolates. RS participated in the isolation and characterization of methanogenic archaeal strains and contributed to the revision of the manuscript. VVZ and WHS contributed to the design of the study and in the analyses of bacterial genes encoding carbohydrate-active enzymes. HK and WL contributed to the discussion section and revised the manuscript. PS participated in the analysis of the hydrogenase genes in methanogenic archaeal isolates and revised the manuscript. ACM participated in bioinformatic data analysis and revised the manuscript. AScz participated in bioinformatic data analysis and discussion of bioinformatics results. MK participated in the design of this study, contributed to the “Results and discussion” section, and revised the manuscript. AP and AS conceived the study, participated in manuscript coordination, drafted the fragment recruitment section, supervised all biological analyses, and revised the manuscript. All authors read and approved the final manuscript.

Author details

¹ Center for Biotechnology (CeBiTec), Bielefeld University, Universitätsstrasse 27, 33615 Bielefeld, Germany. ² Faculty of Technology, Bielefeld University, Universitätsstrasse 25, 33615 Bielefeld, Germany. ³ Computational Biology of Infection Research, Helmholtz Centre for Infection Research, Inhoffenstraße 7, 38124 Brunswick, Germany. ⁴ German Center for Infection Research (DZIF), partner site Hannover-Braunschweig, Inhoffenstraße 7, 38124 Brunswick, Germany. ⁵ Department Bioengineering, Leibniz Institute for Agricultural Engineering and Bioeconomy (ATB), Max-Eyth-Allee 100, 14469 Potsdam, Germany. ⁶ Johannes Gutenberg-University, Institute of Microbiology and Wine Research, Johann-Joachim Becherweg 15, 55128 Mainz, Germany. ⁷ Department of Microbiology, Technische Universität München, Emil-Ramann-Str. 4, 85354 Freising-Weißenstephan, Germany. ⁸ Faculty Life Sciences/Research Center ‘Biomass Utilization Hamburg’, University of Applied Sciences Hamburg (HAW), Ulmenliet 20, 21033 Hamburg-Bergedorf, Germany. ⁹ Institute of Molecular Genetics, Russian Academy of Science, Kurchatov Sq. 2, Moscow 123182, Russia. ¹⁰ Institut für Forensische Genetik GmbH, Im Derdel 8, 48168 Münster, Germany.

Acknowledgements

The authors acknowledge the German Federal Ministry of Education and Research (BMBF) for support of the research project BIOGAS-MARKER, Grant Number 03SF0440C, and the German Federal Ministry of Food and Agriculture for support of the research project Biogas-Messprogramm III (BMP-III), Grant Number 22404015. The authors thank the Fachagentur für Nachwachsende Rohstoffe (FNR) and Projektträger Jülich (PTJ) for their highly valuable support in project management.

Competing interests

The authors declare that they have no competing interests.

Availability of supporting data

The metagenome datasets supporting the conclusions of this article are available at the NCBI database in the short read archive (SRA): <http://www.ncbi.nlm.nih.gov/sra/?term=SRA357211>. <http://www.ncbi.nlm.nih.gov/sra/?term=SRA357213>. <http://www.ncbi.nlm.nih.gov/sra/?term=SRA357208>. <http://www.ncbi.nlm.nih.gov/sra/?term=SRA357209>. <http://www.ncbi.nlm.nih.gov/sra/?term=SRA357214>. <http://www.ncbi.nlm.nih.gov/sra/?term=SRA357221>. <http://www.ncbi.nlm.nih.gov/sra/?term=SRA357222>. <http://www.ncbi.nlm.nih.gov/sra/?term=SRA357223>.

Consent for publication

Not applicable.

Ethical approval and consent to participate

Not applicable.

Funding

This work was part of the joint project BIOGAS-CORE supported by the German Federal Ministry of Food and Agriculture (BMEL), Grant Nos. 22006712, 22006812, 22007012, 22017111. IM, AB, and DW were supported by a fellowship from the CLIB Graduate Cluster Industrial Biotechnology. The work of AP was supported by the BMBF-funded project ‘Bielefeld-Gießen Center for Microbial Bioinformatics—BiGi (Grant Number 031A533)’ within the German Network for Bioinformatics Infrastructure (de.NBI).

Publisher’s Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Received: 13 December 2016 Accepted: 1 November 2017

Published online: 13 November 2017

References

- Klocke M, Nettmann E, Bergmann I, Mundt K, Souidi K, Mumme J, et al. Characterization of the methanogenic *Archaea* within two-phase biogas reactor systems operated with plant biomass. *Syst Appl Microbiol*. 2008;31:190–205.
- Blume F, Bergmann I, Nettmann E, Schelle H, Rehde G, Mundt K, et al. Methanogenic population dynamics during semi-continuous biogas fermentation and acidification by overloading. *J Appl Microbiol*. 2010;109:441–50.
- Nettmann E, Bergmann I, Pramschüfer S, Mundt K, Plogsties V, Herrmann C, et al. Polyphasic analyses of methanogenic archaeal communities in agricultural biogas plants. *Appl Environ Microbiol*. 2010;76:2540–8.
- Westerholm M, Dolfig J, Sherry A, Gray ND, Head IM, Schnürer A. Quantification of syntrophic acetate-oxidizing microbial communities in biogas processes. *Environ Microbiol Rep*. 2011;3:500–5.
- Heeg K, Pohl M, Sontag M, Mumme J, Klocke M, Nettmann E. Microbial communities involved in biogas production from wheat straw as sole substrate within a two-phase solid-state anaerobic digestion. *Syst Appl Microbiol*. 2014;37:590–600.
- Rademacher A, Nolte C, Schönberg M, Klocke M. Temperature increases from 55 to 75 °C in a two-phase biogas reactor result in fundamental alterations within the bacterial and archaeal community structure. *Appl Microbiol Biotechnol*. 2012;96:565–76.
- Klang J, Theuerl S, Szewzyk U, Huth R, Tölle R, Klocke M. Dynamic variation of the microbial community structure during the long-time monofermentation of maize and sugar beet silage: microbial dynamics during anaerobic digestion. *Microb Biotechnol*. 2015;8:764–75.
- Theuerl S, Kohrs F, Benndorf D, Maus I, Wibberg D, Schlüter A, et al. Community shifts in a well-operating agricultural biogas plant: how process variations are handled by the microbiome. *Appl Microbiol Biotechnol*. 2015;99:7791–803.
- Stolze Y, Zakrzewski M, Maus I, Eikmeyer F, Jaenicke S, Rottmann N, et al. Comparative metagenomics of biogas-producing microbial communities from production-scale biogas plants operating under wet or dry fermentation conditions. *Biotechnol Biofuels*. 2015;8:14.
- Goux X, Calusinska M, Lemaigre S, Klocke M, Udelhoven T, Benizri E, et al. Microbial community dynamics in replicate anaerobic digesters exposed sequentially to increasing organic loading rate, acidosis, and process recovery. *Biotechnol Biofuels*. 2015;8:122.
- Jaenicke S, Ander C, Bekel T, Bisdorf R, Dröge M, Gartemann KH, et al. Comparative and joint analysis of two metagenomic datasets from a biogas fermenter obtained by 454-pyrosequencing. *PLoS ONE*. 2011;6:e14519.
- Wirth R, Kovács E, Maróti G, Bagi Z, Rákhely G, Kovács KL. Characterization of a biogas-producing microbial community by short-read next generation DNA sequencing. *Biotechnol Biofuels*. 2012;5:41.
- Hanreich A, Schimpf U, Zakrzewski M, Schlüter A, Benndorf D, Heyer R, et al. Metagenome and metaproteome analyses of microbial

- communities in mesophilic biogas-producing anaerobic batch fermentations indicate concerted plant carbohydrate degradation. *Syst Appl Microbiol.* 2013;36:330–8.
14. Treu L, Kougias PG, Campanaro S, Bassani I, Angelidaki I. Deeper insight into the structure of the anaerobic digestion microbial community; the biogas microbiome database is expanded with 157 new genomes. *Bioresour Technol.* 2016;216:260–6.
 15. Bremges A, Maus I, Belmann P, Eikmeyer F, Winkler A, Albersmeier A, et al. Deeply sequenced metagenome and metatranscriptome of a biogas-producing microbial community from an agricultural production-scale biogas plant. *GigaScience.* 2015;4:33.
 16. Ortseifen V, Stolze Y, Maus I, Szczyrba A, Bremges A, Albaump SP, et al. An integrated metagenome and -proteome analysis of the microbial community residing in a biogas production plant. *J Biotechnol.* 2016;231:268–79.
 17. Campanaro S, Treu L, Kougias PG, De Francisci D, Valle G, Angelidaki I. Metagenomic analysis and functional characterization of the biogas microbiome using high throughput shotgun sequencing and a novel binning strategy. *Biotechnol Biofuels.* 2016;9:26.
 18. Maus I, Koeck DE, Cibis KG, Hahnke S, Kim YS, Langer T, et al. Unraveling the microbiome of a thermophilic biogas plant by metagenome and metatranscriptome analysis complemented by characterization of bacterial and archaeal isolates. *Biotechnol Biofuels.* 2016;9:171.
 19. Maus I, Wibberg D, Stantscheff R, Cibis K, Eikmeyer FG, König H, et al. Complete genome sequence of the hydrogenotrophic *Archaeon Methanobacterium* sp. Mb1 isolated from a production-scale biogas plant. *J Biotechnol.* 2013;168:734–6.
 20. Hahnke S, Striesow J, Elvert M, Mollar XP, Klocke M. *Clostridium bornimense* sp. nov., isolated from a mesophilic, two-phase, laboratory-scale biogas reactor. *Int J Syst Evol Microbiol.* 2014;64:2792–7.
 21. Stantscheff R, Kuever J, Rabenstein A, Seyfarth K, Dröge S, König H. Isolation and differentiation of methanogenic *Archaea* from mesophilic corn-fed on-farm biogas plants with special emphasis on the genus *Methanobacterium*. *Appl Microbiol Biotechnol.* 2014;98:5719–35.
 22. Tomazetto G, Hahnke S, Langer T, Wibberg D, Blom J, Maus I, et al. The completely annotated genome and comparative genomics of the *Peptoniphilaceae* bacterium str. ING2-D1G, a novel acidogenic bacterium isolated from mesophilic biogas reactor. *J Biotechnol.* 2017; 257:178–86. <https://doi.org/10.1016/j.jbiotec.2017.05.027>.
 23. Maus I, Cibis KG, Wibberg D, Winkler A, Stolze Y, König H, et al. Complete genome sequence of the strain *Defluviitoga tunisiensis* L3, isolated from a thermophilic, production-scale biogas plant. *J Biotechnol.* 2015;203:17–8.
 24. Koeck DE, Maus I, Wibberg D, Winkler A, Zverlov VV, Liebl W, et al. Draft genome sequence of *Herbinix hemicellulosilytica* T3/55T, a new thermophilic cellulose degrading bacterium isolated from a thermophilic biogas reactor. *J Biotechnol.* 2015;214:59–60.
 25. Hahnke S, Maus I, Wibberg D, Tomazetto G, Pühler A, Klocke M, et al. Complete genome sequence of the novel *Porphyromonadaceae* bacterium strain ING2-E5B isolated from a mesophilic lab-scale biogas reactor. *J Biotechnol.* 2015;193:34–6.
 26. Hahnke S, Klocke M, Langer T, Koeck DE. Description of *Proteiniphilum saccharofermentans* sp. nov., *Petrimonas mucosa* sp. nov. and *Fermentimonas caenicola* gen. nov., sp. nov., isolated from mesophilic laboratory-scale biogas reactors, and emended description of the genus *Proteiniphilum*. *Int J Syst Evol Microbiol.* 2016;66:1466–75.
 27. Cibis KG, Gneipel A, König H. Isolation of acetic, propionic and butyric acid-forming bacteria from biogas plants. *J Biotechnol.* 2016;220:51–63.
 28. Maus I, Cibis KG, Bremges A, Stolze Y, Wibberg D, Tomazetto G, et al. Genomic characterization of *Defluviitoga tunisiensis* L3, a key hydrolytic bacterium in a thermophilic biogas plant and its abundance as determined by metagenome fragment recruitment. *J Biotechnol.* 2016;232:50–60.
 29. Koeck DE, Maus I, Wibberg D, Winkler A, Zverlov VV, Liebl W, et al. Draft genome sequence of *Propionispora* sp. strain 2/2-37, a new xylan-degrading bacterium isolated from a mesophilic biogas reactor. *Genome Announc.* 2016. <https://doi.org/10.1128/genomeA.00609-16>.
 30. DSMZ medium 287. http://www.dsmz.de/microorganisms/medium/pdf/DSMZ_Medium287.pdf. Accessed 4 May 2016.
 31. DSMZ medium 120. http://www.dsmz.de/microorganisms/medium/pdf/DSMZ_Medium120.pdf. Accessed 4 May 2016.
 32. SINA alignment service. <https://www.arb-silva.de/aligner/>. Accessed 12 Sept 2016.
 33. Munoz R, Yarza P, Ludwig W, Euzéby J, Amann R, Schleifer KH, et al. Release LTPs104 of the all-species living tree. *Syst Appl Microbiol.* 2011;34:169–70.
 34. Quast C, Pruesse E, Yilmaz P, Gerken J, Schweier T, Yarza P, et al. The SILVA ribosomal RNA gene database project: improved data processing and web-based tools. *Nucleic Acids Res.* 2013;41:D590–6.
 35. Ludwig W. ARB: a software environment for sequence data. *Nucleic Acids Res.* 2004;32:1363–71.
 36. Tejerizo GT, Kim YS, Maus I, Wibberg D, Winkler A, Off S, et al. (2017) Genome sequence of *Methanobacterium congolense* strain Buetzberg, a hydrogenotrophic, methanogenic archaeon, isolated from a mesophilic industrial-scale biogas plant utilizing bio-waste. *J Biotechnol.* 2017;247:1–5
 37. Tomazetto G, Hahnke S, Koeck DE, Wibberg D, Maus I, Pühler A, et al. Complete genome analysis of *Clostridium bornimense* strain M2/40T: a new acidogenic *Clostridium* species isolated from a mesophilic two-phase laboratory-scale biogas reactor. *J Biotechnol.* 2015;232:38–49.
 38. Wibberg D, Blom J, Jaenicke S, Kollin F, Rupp O, Scharf B, et al. Complete genome sequencing of *Agrobacterium* sp. H13-3, the former *Rhizobium lupini* H13-3, reveals a tripartite genome consisting of a circular and a linear chromosome and an accessory plasmid but lacking a tumor-inducing Ti-plasmid. *J Biotechnol.* 2011;155:50–62.
 39. Meyer F, Goesmann A, McHardy AC, Bartels D, Bekel T, Clausen J, et al. GenDB—an open source genome annotation system for prokaryote genomes. *Nucleic Acids Res.* 2003;31:2187–95.
 40. Yin Y, Mao X, Yang J, Chen X, Mao F, Xu Y. dbCAN: a web resource for automated carbohydrate-active enzyme annotation. *Nucleic Acids Res.* 2012;40:W445–51.
 41. Stolze Y, Bremges A, Rummig M, Henke C, Maus I, Pühler A, et al. Identification and genome reconstruction of abundant distinct taxa in microbiomes from one thermophilic and three mesophilic production-scale biogas plants. *Biotechnol Biofuels.* 2016;9:156.
 42. Niu B, Zhu Z, Fu L, Wu S, Li W. FR-HIT, a very fast program to recruit metagenomic reads to homologous reference genomes. *Bioinformatics.* 2011;27:1704–5.
 43. Ondov BD, Treangen TJ, Melsted P, Mallonee AB, Bergman NH, Koren S, et al. Mash: fast genome and metagenome distance estimation using MinHash. *Genome Biol.* 2016;17:132.
 44. Tatusova T, Ciuffo S, Fedorov B, O'Neill K, Tolstoy I. RefSeq microbial genomes database: new representation and annotation strategy. *Nucleic Acids Res.* 2014;42:D553–9.
 45. Bray NL, Pimentel H, Melsted P, Pachter L. Near-optimal probabilistic RNA-seq quantification. *Nat Biotechnol.* 2016;5:525–7.
 46. Zakrzewski M, Goesmann A, Jaenicke S, Jünemann S, Eikmeyer F, Szczepanowski R, et al. Profiling of the metabolically active community from a production-scale biogas plant by means of high-throughput metatranscriptome sequencing. *J Biotechnol.* 2012;158:248–58.
 47. Eikmeyer FG, Rademacher A, Hanreich A, Hennig M, Jaenicke S, Maus I, et al. Detailed analysis of metagenome datasets obtained from biogas-producing microbial communities residing in biogas reactors does not indicate the presence of putative pathogenic microorganisms. *Biotechnol Biofuels.* 2013;6:49.
 48. Lagier JC, Hugon P, Khelaifa S, Fournier PE, La Scola B, Raoult D. The rebirth of culture in microbiology through the example of culturomics to study human gut microbiota. *Clin Microbiol Rev.* 2015;28:237–64.
 49. Ollivier BM, Mah RA, Garcia JL, Boone DR. Isolation and characterization of *Methanogenium bourgense* sp. nov. *Int J Syst Evol Microbiol.* 1986;36:297–301.
 50. Bryant MP, Boone DR. Isolation and characterization of *Methanobacterium formicum* MF. *Int J Syst Evol Microbiol.* 1987;37:171.
 51. Yutin N, Galperin MY. A genomic update on clostridial phylogeny: Gram-negative spore formers and other misplaced clostridia. *Environ Microbiol.* 2013;15:2631–41.
 52. Izquierdo JA, Pattathil S, Guseva A, Hahn MG, Lynd LR. Comparative analysis of the ability of *Clostridium clariflavum* strains and *Clostridium thermocellum* to utilize hemicellulose and untreated plant material. *Biotechnol Biofuels.* 2014;7:136.
 53. Koeck DE, Wibberg D, Maus I, Winkler A, Albersmeier A, Zverlov VV, et al. Complete genome sequence of the cellulolytic thermophile

- Ruminoclostridium cellulosi* wild-type strain DG5 isolated from a thermophilic biogas plant. *J Biotechnol.* 2014;188:136–7.
54. Koeck DE, Zverlov VV, Schwarz WH, Wanner G, Liebl W, Schwarz WH. *Herbinix hemicellulosilytica* gen. nov., sp. nov., a thermophilic cellulose-degrading bacterium isolated from a thermophilic biogas reactor. *Int J Syst Evol Microbiol.* 2015;65:2365–71.
 55. Koeck DE, Hahnke S, Zverlov VV. *Herbinix luporum* sp. nov., a thermophilic cellulose-degrading bacterium isolated from a thermophilic biogas reactor. *Int J Syst Evol Microbiol.* 2016;66:4132–7.
 56. Podosokorskaya OA, Bonch-Osmolovskaya EA, Beskorovanyy AV, Toshchakov SV, Kologanova TV, Kublanov IV. *Mobilitalea sibirica* gen. nov., sp. nov., a halotolerant polysaccharide-degrading bacterium. *Int J Syst Evol Microbiol.* 2014;64:2657–61.
 57. Combet-Blanc Y, Ollivier B, Streicher C, Patel BKC, Dwivedi PP, Pot B, et al. *Bacillus thermoamylovorans* sp. nov., a moderately thermophilic and amyolytic bacterium. *Int J Syst Evol Microbiol.* 1995;45:9–16.
 58. Ezaki T, Li N, Shu S, Zhao L, Kawamura Y, Li ZY. Proposal of the genera *Anaerococcus* gen. nov., *Peptoniphilus* gen. nov. and *Gallicola* gen. nov. for members of the genus *Peptostreptococcus*. *Int J Syst Evol Microbiol.* 2001;51:1521–8.
 59. Abou-Zeid DM. *Propionispora hippei* sp. nov., a novel Gram-negative, spore-forming anaerobe that produces propionic acid. *Int J Syst Evol Microbiol.* 2004;54:951–4.
 60. Koeck DE, Zverlov VV, Liebl W, Schwarz WH. Comparative genotyping of *Clostridium thermocellum* strains isolated from biogas plants: genetic markers and characterization of cellulolytic potential. *Syst Appl Microbiol.* 2014;37:311–9.
 61. Koeck DE, Wibberg D, Koellmeier T, Blom J, Jaenicke S, Winkler A, et al. Draft genome sequence of the cellulolytic *Clostridium thermocellum* wild-type strain BC1 playing a role in cellulosic biomass degradation. *J Biotechnol.* 2013;168:62–3.
 62. Zhang F, Yang JH, Dai K, Chen Y, Li QR, Gao FM, et al. Characterization of microbial compositions in a thermophilic chemostat of mixed culture fermentation. *Appl Microbiol Biotechnol.* 2016;100:1511–21.
 63. Ahlert S, Zimmermann R, Ebling J, König H. Analysis of propionate-degrading consortia from agricultural biogas plants. *Microbiologyopen.* 2016. <https://doi.org/10.1002/mbo3.386>.
 64. Nilegaonkar S, Bhosale SB, Kshirsagar DC, Kapadi AH. Production of 2,3-butanediol from glucose by *Bacillus licheniformis*. *World J Microbiol Biotechnol.* 1992;8:378–81.
 65. Zhang B, Cai W, He P. Influence of lactic acid on the two-phase anaerobic digestion of kitchen wastes. *J Environ Sci.* 2007;19:244–9.
 66. Ohara H, Uchida K, Yahata M, Kondo H. NAD-specific 6-phosphogluconate dehydrogenase in lactic acid bacteria. *Biosci Biotechnol Biochem.* 1996;60:692–3.
 67. Ferry JG. How to make a living by exhaling methane. *Annu Rev Microbiol.* 2010;64:453–73.
 68. Boone DR, Castenholz RW. *Bergey's manual of systematic bacteriology: the Archaea and the deeply branching and phototrophic Bacteria.* New York: Springer Press; 2001.
 69. Kaster AK, Moll J, Parey K, Thauer RK. Coupling of ferredoxin and heterodisulfide reduction via electron bifurcation in hydrogenotrophic methanogenic *Archaea*. *Proc Natl Acad Sci.* 2011;108:2981–6.
 70. Tersteegen A, Hedderich R. *Methanobacterium thermoautotrophicum* encodes two multisubunit membrane-bound [NiFe] hydrogenases. Transcription of the operons and sequence analysis of the deduced proteins. *Eur J Biochem.* 1999;264:930–43.
 71. Lee MJ, Zinder SH. Hydrogen partial pressures in a thermophilic acetate-oxidizing methanogenic coculture. *Appl Environ Microbiol.* 1998;54:1457–61.
 72. Li B, Ruotti V, Stewart RM, Thomson JA, Dewey CN. RNA-Seq gene expression estimation with read mapping uncertainty. *Bioinformatics.* 2010;4:493–500.
 73. Konstantinidis KT, Tiedje JM. Genomic insights that advance the species definition for prokaryotes. *Proc Natl Acad Sci USA.* 2005;102:2567–72.
 74. Manzoor S, Schnürer A, Bongcam-Rudloff E, Müller B. Complete genome sequence of *Methanoculleus bourgensis* strain MAB1, the syntrophic partner of mesophilic acetate-oxidising bacteria (SAOB). *Stand Genom Sci.* 2016;11:80.
 75. Nimura Y, Koh E, Yanadiga F, Suzuki KI, Komagata K, Kozaki M. *Amphibacillus xylanus* gen. nov., sp. nov., a facultatively anaerobic spore-forming xylan-digesting bacterium which lacks cytochrome, quinone, and catalase. *Int J Syst Evol Microbiol.* 1990;40:297–301.
 76. Shiratori H, Sasaya K, Ohiwa H, Ikeno H, Ayame S, Kataoka N, et al. *Clostridium clariflavum* sp. nov. and *Clostridium caenicola* sp. nov., moderately thermophilic, cellulose-/cellobiose-digesting bacteria isolated from methanogenic sludge. *Int J Syst Evol Microbiol.* 2009;59:1764–70.
 77. Holden MT, Hauser H, Sanders M, Ngo TH, Cherevach I, Cronin A, et al. Rapid evolution of virulence and drug resistance in the emerging zoonotic pathogen *Streptococcus suis*. *PLoS ONE.* 2009;4:e6072.
 78. Koeck DE, Maus I, Wibberg D, Winkler A, Zverlov VV, Liebl W, et al. Complete genome sequence of *Herbinix luporum* SD1D, a new cellulose degrading bacterium isolated from a thermophilic biogas reactor. *Genome Announc.* 2016. <https://doi.org/10.1128/genomeA.00687-16>.
 79. Koeck DE, Wibberg D, Maus I, Winkler A, Albersmeier A, Zverlov VV, et al. First draft genome sequence of the amyolytic *Bacillus thermoamylovorans* wild-type strain 1A1 isolated from a thermophilic biogas plant. *J Biotechnol.* 2014;192:154–5.
 80. Maus I, Stantscheff R, Wibberg D, Stolze Y, Winkler A, Pühler A, et al. Complete genome sequence of the methanogenic neotype strain *Methanobacterium formicicum* MFT. *J Biotechnol.* 2014;192:40–1.
 81. Maus I, Wibberg D, Stantscheff R, Stolze Y, Blom J, Eikmeyer FG, et al. Insights into the annotated genome sequence of *Methanoculleus bourgensis* MS2T, related to dominant methanogens in biogas-producing plants. *J Biotechnol.* 2015;201:43–53.
 82. Zellner G, Messner P, Winter J, Stackebrandt E. *Methanoculleus palmolei* sp. nov., an irregularly coccoid methanogen from an anaerobic digester treating wastewater of a palm oil plant in north-Sumatra, Indonesia. *Int J Syst Bacteriol.* 1998;4:1111–7.

Submit your next manuscript to BioMed Central and we will help you at every step:

- We accept pre-submission inquiries
- Our selector tool helps you to find the most relevant journal
- We provide round the clock customer support
- Convenient online submission
- Thorough peer review
- Inclusion in PubMed and all major indexing services
- Maximum visibility for your research

Submit your manuscript at
www.biomedcentral.com/submit

