

UvA-DARE (Digital Academic Repository)

The transient IGR J17361-4441 in the globular cluster NGC 6388

an unusual very-faint transient X-ray binary or an outburst from an intermediate-mass black hole

Wijnands, R.; Yang, Y.J.; Degenaar, N.; Altamirano, D.; Tudose, V.; Heinke, C.O.

Publication date 2011 Document Version Final published version Published in The astronomer's telegram License Unspecified

Link to publication

Citation for published version (APA):

Wijnands, R., Yang, Y. J., Degenaár, N., Altamirano, D., Tudose, V., & Heinke, C. O. (2011). The transient IGR J17361-4441 in the globular cluster NGC 6388: an unusual very-faint transient X-ray binary or an outburst from an intermediate-mass black hole. *The astronomer's telegram*. https://www.astronomerstelegram.org/?read=3595

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please Ask the Library: https://uba.uva.nl/en/contact, or a letter to: Library of the University of Amsterdam, Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

UvA-DARE is a service provided by the library of the University of Amsterdam (https://dare.uva.nl)

ATel #3595: The transient IGR J17361-4441 in the globular cluster NGC 6388: an unusual very-faint transient X-ray binary or an outburst from an intermediate-mass blac...



Thanks to Patrons, The Astronomer's Telegram is free to read, free to publish and always will be. Thank you.

[Previous | Next | ADS]

The transient IGR J17361-4441 in the globular cluster NGC 6388: an unusual very-faint transient X-ray binary or an outburst from an intermediate-mass black hole

ATel #3595; R. Wijnands, Y. J. Yang, N. Degenaar, D. Altamirano (University of Amsterdam), V. Tudose (ASTRON), C. O. Heinke (University of Alberta) on 26 Aug 2011; 10:49 UT

Credential Certification: Rudy Wijnands (rudy@space.mit.edu)

Subjects: Radio, X-ray, Request for Observations, Black Hole, Globular Cluster, Neutron Star, Transient

Referred to by ATel #: 3627

Tweet

Gibaud et al. (2011, ATel **#3565**) reported the INTEGRAL discovery of a new transient, IGR J17361-4441, located in the Galactic globular cluster NGC 6388 (Ferrigno et al. 2011, ATel **#3566**). Swift/XRT has monitored this source from 16 to 25 August on a nearly daily basis (see ATel **#3566** for the first observation). The 0.5-10 keV unabsorbed flux in those observations was roughly constant, typically (4-6)e-11 erg/s/cm^2, corresponding to a 0.5-10 keV X-ray luminosity of (6-9)e35 erg/s (for a distance of 11.5 kpc), making this source a very-faint X-ray transient (VFXT; see Wijnands et al. 2006, A&A, 449, 1117, for a discussion about VFXTs). The spectrum is very hard, with a photon index of 0.6-1.0 (see also ATel **#3566**).

Considering its luminosity and location in a globular cluster, the system is most probably a neutron-star low-mass X-ray binary (NS-LMXB), although the hard X-ray spectrum is atypical for LMXB transients, which normally show indices of 1.6-2.2 in this luminosity range. However, there exist a small sub-group of NS-LMXBs with similarly hard spectra: accreting X-ray pulsars with high magnetic field (>1e12 Gauss) neutron stars (e.g., Her X-1, GRO J1744-28; see Becker et al. 1977, ApJ, 214, 879; Giles et al. 1996, ApJ, 469, L25). Sensitive X-ray observations (e.g., Chandra, XMM-Newton) could detect pulsations with periods above ~0.1 second as observed for this class. This would be the first such transient found in a globular cluster.

Alternatively, the source could be an LMXB with an ordinary, low magnetic field (<1e10 Gauss) neutron star. In that case, long observations might detect type-I X-ray bursts and high-sensitivity, high-time resolution observations (e.g., with XMM-Newton) might detect

Related	
3958	IGR J17361-4441: ~200 days of monitoring with Swift/XRT
	Chandra localisation of IGR J17361-4441 in the globular cluster NGC 6388
3617	ATCA radio non-detection of IGR J17361-4441
3595	The transient IGR J17361- 4441 in the globular cluster NGC 6388: an unusual very- faint transient X-ray binary or an outburst from an intermediate-mass black hole
3566	Swift/XRT localization of IGR J17361-4441 in the globular cluster NGC 6388
3565	A new hard X-ray transient discovered by INTEGRAL: IGR J17361-4441

ATel #3595: The transient IGR J17361-4441 in the globular cluster NGC 6388: an unusual very-faint transient X-ray binary or an outburst from an intermediate-mass blac...

pulsations > 100 Hz. This source probably has a rather low time-averaged accretion rate and it such systems often harbor accreting millisecond X-ray pulsars (e.g. Wijnands 2008, AIPC, 1010, 382). Detecting either fast pulsations or type-I X-ray bursts from the system will prove it is an ordinary NS-LMXB, although its enigmatic X-ray spectrum still needs to be explained.

Thirdly, NGC 6388 may harbor an intermediate mass black hole (of (6-20)E3 solar masses, Lanzoni et al. 2007, ApJ, 670, 1065, Lutzgendorf et al. 2011, A&A, 533, 36). This outburst could be an accretion outburst from an intermediate mass black hole, as the X-ray position is currently consistent with the cluster center. A scheduled Chandra observation of this source will determine its position, and may rule out this hypothesis. Alternatively, sensitive radio observations might detect elevated radio emission at a level of a few tens of mJy (using the results of Merloni et al. 2003, MNRAS, 345, 1057; Maccarone 2004, MNRAS, 351, 1049, and the observed X-ray luminosity) which would also provide support for this hypothesis.

Since this VFXT in NGC 6388 is very unusual in its behavior, we strongly urge follow-up observations of this source at all wavelengths to gain more insight in its nature.

[Telegram Index]

R. E. Rutledge, Editor-in-Chief Derek Fox, Editor rrutledge@astronomerstelegram.org dfox@astronomerstelegram.org