

POSTILLA

Published from 1950 to 2004, the short papers of the *Postilla* series reported on original research by the Yale Peabody Museum of Natural History's curators, staff, and research associates, and their colleagues, in the natural science disciplines represented by the collections of the Museum's curatorial divisions.

The *Postilla* series, which ceased publication with Number 232 (2004), was incorporated into the journal *Bulletin of the Peabody Museum of Natural History*, available from BioOne Complete at <https://bioone.org/>.

Yale Peabody Museum scholarly publications are archived through EliScholar, a digital platform for scholarly publishing provided by Yale University Library at <https://elischolar.library.yale.edu/>.



This work is licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License.
<https://creativecommons.org/licenses/by-nc-sa/4.0/>

Yale PEABODY MUSEUM OF NATURAL HISTORY

P.O. Box 208118 | New Haven CT 06520-8118 USA | peabody.yale.edu

POSTILLA
PEABODY MUSEUM
YALE UNIVERSITY


NUMBER 168

15 APRIL 1976

**A NEW SPECIES OF *APTERODON* (MAMMALIA,
CREODONTA) FROM THE UPPER EOCENE
QASR EL-SAGHA FORMATION OF EGYPT**

ELWYN L. SIMONS

PHILIP D. GINGERICH





POSTILLA

Published by the Peabody Museum of Natural History, Yale University

Postilla includes results of original research on systematic, evolutionary, morphological, and ecological biology, including paleontology. Syntheses and other theoretical papers based on research are also welcomed. *Postilla* is intended primarily for papers by the staff of the Peabody Museum or on research using material in this Museum.

Editor: John H. Ostrom

Managing Editor: Zelda Edelson

Postilla is published at frequent but irregular intervals. Manuscripts, orders for publications, and all correspondence concerning publications should be directed to:

Publications Office

Peabody Museum of Natural History

New Haven, Conn. 06520, USA

Lists of the publications of the Museum are available from the above office. These include *Postilla*, *Bulletin*, *Discovery*, and special publications. *Postilla* and the *Bulletin* are available in exchange for relevant publications of other scientific institutions anywhere in the world.

Inquiries regarding back numbers of the discontinued journal, *Bulletin of the Bingham Oceanographic Collection*, should be directed to:

Kraus Reprint Co.

Route 100

Millwood, New York 10546

**A NEW SPECIES OF *APTERODON* (MAMMALIA, CREODONTA)
FROM THE UPPER EOCENE QASR EL-SAGHA FORMATION OF EGYPT**

Elwyn L. Simons

Peabody Museum of Natural History and
Department of Geology and Geophysics
Yale University
New Haven, Connecticut 06520

Philip D. Gingerich

Museum of Paleontology
The University of Michigan
Ann Arbor, Michigan 48109

Received 6 January 1975

ABSTRACT

The mandible of a medium-sized carnivore discovered by a recent Yale paleontological expedition to the Fayum Province of Egypt represents a new species of the creodont *Apterodon*. The type and only specimen of this new species, *Apterodon saghensis*, comes from the Upper Eocene Qasr el-Sagha Formation and thus represents the earliest terrestrial carnivore yet described from the African continent. Unfortunately the specimen contributes little to our understanding of the origin and relationships of *Apterodon*.

POSTILLA 168: 9 p.

15 April 1976

INTRODUCTION

From 1961 through 1967 a series of paleontological expeditions to the Fayum Province of Egypt was undertaken by Yale University field parties, and these expeditions have added considerably to our knowledge of the Paleogene mammalian faunas of Africa. The Oligocene faunas of the Jebel el-Qatrani Formation proved to be very rich and they have consequently received the most attention. Recent publications on these faunas include description of a new species of the primate *Parapithecus* (Simons, 1974), two new genera and species of carnivores (Simons and Gingerich, 1974), and a new genus and species of hyrax (Meyer, 1973). Detailed analysis of the sediments of the Jebel el-Qatrani Formation indicates that they were deposited on a low, featureless deltaic plain with gallery forests along the rivers and open savannah between (Bowen and Vondra, 1974).

Some attention was devoted to the underlying Qasr el-Sagha Formation by the Yale expedition. Study of the marine invertebrate fauna from this formation indicates that it is probably of late Bartonian age [ca. 40 million years (m.y.) B.P.] and correlative with the Mokattam and Maadi Formations of the Cairo region (Said, 1962, p. 102-103). Of particular interest is a fragmentary mammalian mandible found by one of us in an interbedded claystone, siltstone, and quartz sandstone facies of the Qasr el-Sagha Formation, which detailed sedimentary and stratigraphic analysis has indicated represents a prograding delta front (Vondra, 1974).

The new specimen, a very corroded mandibular ramus preserving the crowns of three teeth and parts of two others, is important as the first good record of a wholly terrestrial mammal from the Qasr el-Sagha Formation and the earliest confirmed record of a terrestrial carnivore from the African continent.

ABBREVIATIONS

CGM	Cairo Geological Museum, Cairo, Egypt
MNHN	Muséum National d'Histoire Naturelle, Paris, France
UCMP	Museum of Paleontology, University of California, Berkeley, California
UM	Museum of Paleontology, The University of Michigan, Ann Arbor, Michigan
YPM	Peabody Museum of Natural History, Yale University, New Haven, Connecticut

SYSTEMATICS

CLASS MAMMALIA
ORDER CREODONTA
FAMILY HYAENODONTIDAE
GENUS *Apterodon* Fischer, 1880

Apterodon Fischer, 1880, p. 288. Type species: *Apterodon gaudryi* Fischer.
Dasyurodon Andreae, 1887, p. 126. Type species: *Dasyurodon flonheimensis*
Andreae.

Pterodon (in part), Andrews, 1904, p. 211.

INCLUDED SPECIES. *Apterodon gaudryi* Fischer, 1880; *A. flonheimensis* (Andreae, 1887); *A. macrognathus* (Andrews, 1904); *A. altidens* Schlosser, 1910; *A. saghensis* new species.

DISTRIBUTION. The type and only specimen of *Apterodon gaudryi* comes from the French Quercy Phosphorites of early or middle Oligocene age (ca. 32-37 m.y.) and *A. flonheimensis*, a possible synonym of *A. gaudryi*, comes from the middle Oligocene (ca. 32-34 m.y.) of Flonheim in the German Mainz Basin. *Apterodon macrognathus* and *A. altidens* both come from the lower fossil wood zone of the Jebel el-Qatrani Formation in Egypt (early to middle Oligocene, ca. 32-37 m.y.). *Apterodon saghensis*, described below, comes from the Egyptian Qasr el-Sagha Formation and extends the range of the genus into the late Eocene (ca. 40 m.y.) These approximate ages are based on Romer (1966, p. 335) and Berggren (1972, p. 198).

***Apterodon saghensis*, new species**

Figures 1, 2, 3C

?*Apterodon* sp. nov., Simons, 1968, p. 15.

TYPE. CGM 40006, a left mandible with the root of the lower canine, the crowns of P₂₋₄, and the damaged crown of M₁. Sharp casts of the type specimen are deposited at Yale (YPM 30241) and the University of Michigan (UM 63389).

HYPODIGM. Type specimen only.

TYPE LOCALITY. The type specimen was collected in 1963 by E. L. Simons from a locality in the Fayum depression of Egypt some twenty feet above the

base of the deltaic sandstone facies of the Qasr el-Sagha Formation and approximately one-half mile north or northwest of the Qasr el-Sagha temple, near the road up the escarpment leading to the overlying lower fossil wood zone of the Jebel el-Qatrani Formation.

ETYMOLOGY. Named with reference to the specimen's provenance, from the Qasr el-Sagha Formation near the temple Qasr el-Sagha.

DIAGNOSIS. *Apterodon saghensis* differs from the species previously described in being significantly smaller (see Table 1), and in having a higher, straighter, and less robust fourth lower premolar (see Fig. 3).

TABLE 1. Tooth measurements of the type specimen of *Apterodon saghensis* and a comparison with the Fayum species *A. macrognathus* and *A. altidens* from the lower fossil wood zone. All measurements in millimeters. Mandibular depth measured beneath M_1 .

	<i>Apterodon saghensis</i> CGM 40006	<i>Apterodon macrognathus</i> UCMP 62218	<i>Apterodon altidens</i> YPM 18160
P_2 L	8.5*	14.8	9.8*
W	4.2	8.3	5.7
P_3 L	10.7	15.8	—
W	4.7	8.5	—
P_4 L	12.0	18.7	15.1
W	5.9	8.9	7.1
M_1 L	9.5	13.4*	12.7
W	5.5	7.0*	6.4
Mandibular Depth	18.1	39.0	26.6

*estimated

DESCRIPTION. The type and only specimen of *Apterodon saghensis* is a left mandible fragment six centimeters in length. The bone is highly corroded, but a portion of the symphyseal surface remains on the medial side of the mandible. The front of the jaw is missing, and no trace remains of the incisors or their alveoli.

The crown of the lower canine is also missing, but its root is preserved and measures 9.7 mm by 7.1 mm in cross section. The mandible is damaged above and lateral to the canine root and the first premolar is missing, though it was undoubtedly retained in this species.

The crowns of three premolars, P_{2-4} , are well preserved. P_2 has a single blunt protoconid cusp, with cristids extending anteriorly and posteriorly from




FIG. 1. Lateral view of type specimen of *Apterodon saghensis*, CGM 40006, showing the root of the lower canine, and P_2 - M_1 . Note the badly corroded mandibular bone.

the protoconid. A very weak cingulid is present on the buccal side of the crown, and a slightly more prominent cristid is developed at the anterolingual corner. The posterior portion is damaged slightly, but the crown length was approximately 8.5 mm. The width of the crown of P_2 is 4.2 mm. P_3 is slightly larger, but is otherwise very similar in morphology to P_2 . P_3 measures 10.7 mm in length, and 4.7 mm in width. A portion of the lingual cingulid is missing, but this shelf broadens posteriorly making a very shallowly basined heel medial to the posterior cristid from the protoconid. P_2 and P_3 both lack any accessory cusps.

The lower fourth premolar is very well preserved, apart from an antero-posterior split in the enamel of the protoconid predating mineralization of the specimen. The protoconid is the dominant cusp on the tooth, and it is relatively high and straight (see Figure 3). A cristid runs down the anterior surface of the protoconid, connecting with a small but distinct anterior basal cusp or paraconid. The posterior cristid of the protoconid joins the posterior basal cusp (which is apparently serially homologous with the hypoconid of a typical molar) in a distinct notch toward the back of the crown. A strong lingual cingulid connects the anterior and posterior basal cusps, and a very weak cingulid is developed on the buccal margin of the tooth. The crown of P_4 measures 12.0 mm in length and 5.9 mm in width. Just as on P_{2-3} , the enamel on this crown appears to be very slightly crenulated.

The protoconid of M_1 appears possibly to have had a straighter posterior margin than is typical in *Apterodon*, which suggests that perhaps a small




FIG. 2. Type specimen of *Apterodon saghensis* in lateral (A) and occlusal (B) view, showing the root of the lower canine (C), lower premolars (P_2-4), and the first lower molar (M_1).

metaconid was retained, but the crown of this tooth is too damaged to offer any positive evidence of this. A distinct constriction separating the trigonid of the tooth from the talonid is apparent, and a buccal cingulid is weakly developed. The tooth is too poorly preserved to show any other details. M_1 measures approximately 9.5 mm in length and 5.5 mm in width. The mandible is broken behind M_1 , and no trace of M_2 or M_3 remains.

DISCUSSION

Species of *Apterodon* are the most common carnivores found in the Oligocene Jebel el-Qatrani Formation of the Fayum, and the specimen described here indicates that species of this genus were probably common in the Late Eocene as well. The abundance of *Apterodon* in Egyptian deposits and its




FIG. 3. Comparison of the lower fourth premolar of four species of *Apterodon*. *A*, *Apterodon altidens* from the Fayum lower fossil wood zone, Jebel el-Qatrani Formation (YPM 18160). *B*, *Apterodon macrognathus* from the Fayum lower fossil wood zone, Jebel el-Qatrani Formation (YPM 18165, reversed). *C*, *Apterodon saghensis* from the Fayum, Qasr el-Sagha Formation (CGM 40006). *D*, *Apterodon gaudryi* from the French Quercy Phosphorites (from a cast of the type, MNHN).

rarity elsewhere (the two European species are known from a single specimen each) suggests that *Apterodon* probably evolved in isolation in Africa, with the exception of a limited migration around the Tethyan Sea resulting in a minor invasion of Europe in the early Oligocene.

Unfortunately, only one other carnivore specimen possibly equal in age to *Apterodon saghensis* is known from Africa, a fragmentary specimen of a "hyaenodont" mentioned by Savage (1969) from Dor el Talha in southern Libya. Neither specimen contributes significantly to understanding the origin

and relationships of *Apterodon*. In his original description of *Apterodon*, Fischer (1880) recognized its similarities to *Pterodon*. Andrews (1904) at first placed *Apterodon macrognathus* in *Pterodon*. In describing the first skulls of the genus, Osborn (1909) retained *Apterodon* in the Hyaenodontidae. However, the long-supported relationship of *Apterodon* to *Pterodon* and the Hyaenodontidae was questioned by Van Valen (1966, p. 85), who proposed transferring *Apterodon* to the condylarth family Mesonychidae, a suggestion followed by Romer (1966, p. 385). Szalay (1967) disputed many of the points raised by Van Valen. Szalay concluded that as a result of his discussion "the undoubted hyaenodontid affinity of *Apterodon* is confirmed." In the absence of an adequate Eocene fossil record, we follow Lange (1967) in regarding any definitive phylogenetic or systematic placement of *Apterodon* as premature. It seems preferable to retain provisionally for *Apterodon* the place it has long occupied among the creodont Hyaenodontidae; however, if and when true mesonychids are discovered in Africa the question might be reopened.

ACKNOWLEDGMENTS

We thank Mr. D. Alfar of the Cairo Geological Museum, Dr. R. Hamilton of the British Museum (Natural History), Dr. M. C. McKenna, the American Museum of Natural History, and Dr. D. E. Savage, University of California, Berkeley, for access to comparative material in their care. Mr. Grant Meyer of the Yale Peabody Museum assisted in preparing the manuscript. We also thank Mrs. Krystyna Butterfield, Mr. Karoly Kutasi, and Mrs. Gladys Newton of the University of Michigan Museum of Paleontology for, respectively, their drawing, photography, and typing skills.

LITERATURE CITED

- Andreae, Achilles. 1887. Ein neues Raubtier aus dem mitteloligocänen Meeressand des Mainzer Beckens. Ber. Senckenberg. naturforsch. Ges., Frankfurt a. M. 1887: 125–133.
- Andrews, Charles W. 1904. Further notes on the mammals of the Eocene of Egypt. Geol. Mag. 1: 211–215.
- Berggren, William A. 1972. A Cenozoic time-scale—some implications for regional geology and paleobiogeography. Lethaia 4: 195–215.
- Bowen, Bruce E. and Carl F. Vondra. 1974. Paleoenvironmental interpretations of the Oligocene Gabal el Qatrani Formation, Fayum depression, Egypt. Ann. Geol. Surv. Egypt 4: 115–138.
- Fischer, Paul. 1880. Note sur un nouveau genre de mammifère fossile (*Apterodon gaudryi*) des Phosphorites de Quercy. Bull. Soc. Géol. France 8: 288–290.
- Lange, Brigitte. 1967. Créodontes des phosphorites du Quercy *Apterodon gaudryi*. Ann. Paléontologie 53: 79–90.

- Meyer, Grant E. 1973. A new Oligocene hyrax from the Jebel el Qatrani Formation, Fayum, Egypt. Postilla (Peabody Mus. Natur. Hist., Yale Univ.) no. 163: 1-11.
- Osborn, Henry F. 1909. New carnivorous mammals from the Fayum Oligocene, Egypt. Bull. Amer. Mus. Natur. Hist. 26: 415-424.
- Romer, Alfred S. 1966. Vertebrate paleontology. 3rd ed. Univ. Chicago Press, Chicago. 468 p.
- Said, Rushdi. 1962. The geology of Egypt. Amer. Elsevier Pub. Co., New York. 377 p.
- Savage, Robert J. G. 1969. Early Tertiary mammal locality in southern Libya. Proc. Geol. Soc. London. 1969: 167-171.
- Schlosser, Max. 1910. Über einige fossile Säugetiere aus dem Oligocän von Ägypten. Zool. Anz. 35: 500-508.
- Simons, Elwyn L. 1968. African Oligocene mammals: Introduction, history of study, and faunal succession. Bull. Peabody Mus. Natur. Hist. (Yale Univ.) 28: 1-21.
- 1974. *Parapithecus grangeri* (Parapithecidae, Old World Higher Primates): New species from the Oligocene of Egypt and the initial differentiation of Cercopithecoidea. Postilla (Peabody Mus. Natur. Hist., Yale Univ.) no. 166: 1-12.
- Simons, Elwyn L. and Philip D. Gingerich. 1974. New carnivorous mammals from the Oligocene of Egypt. Ann. Geol. Surv. Egypt 4: 157-166.
- Szalay, Frederick S. 1967. The affinities of *Apterodon* (Mammalia, Deltatheridia, Hyaenodontidae). Amer. Mus. Natur. Hist. Novitates, no. 2293: 1-17.
- Van Valen, Leigh. 1966. Deltatheridia, a new order of mammals. Bull. Amer. Mus. Natur. Hist. 132: 1-126.
- Vondra, Carl F. 1974. Upper Eocene transitional and near-shore marine Qasr el Sagha Formation, Fayum depression, Egypt. Ann. Geol. Surv. Egypt 4: 79-94.

INFORMATION FOR AUTHORS

REVIEW The Publications Committee of the Peabody Museum of Natural History reviews and approves manuscripts for publication. Papers will be published in approximately the order in which they are accepted; delays may result if manuscript or illustrations are not in proper form. To facilitate review, the original and one carbon or xerox copy of the typescript and figures should be submitted. The author should keep a copy.

STYLE Authors of biological papers should follow the *CBE Style Manual* Third Edition (Amer. Inst. Biol. Sci.). Authors of paleontological manuscripts may choose to follow the *Suggestions to Authors of the Reports of the U.S. Geological Survey*, Fifth Edition (U.S. Govt. Printing Office).

FORM Maximum size is 80 printed pages including illustrations (= about 100 manuscript pages including illustrations). Manuscripts must be typewritten, with wide margins, on one side of good quality 8½ x 11" paper. *Double space everything. Do not underline anything except genera and species.* The editors reserve the right to adjust style and form for conformity.

TITLE Should be precise and short. Title should include pertinent key words which will facilitate computerized listings. Names of new taxa are not to be given in the title.

ABSTRACT The paper must begin with an abstract. Authors must submit completed BioAbstract forms; these can be obtained from the *Postilla* editors in advance of submission of the manuscripts.

NOMENCLATURE Follow the International Codes of Zoological and Botanical Nomenclature.

ILLUSTRATIONS Must be planned for reduction to 4¼ x 7" (to allow for running head and two-line caption). If illustration must go sideways on page, reduction should be to 4 x 7¼". All illustrations should be called "Figures" and numbered in arabic, with letters for parts within one page. It is the author's responsibility to see that illustrations are properly lettered and mounted. Captions should be typed double-spaced on a separate page.

FOOTNOTES Should not be used, with rare exceptions. If unavoidable, type double-spaced on a separate page.

TABLES Should be numbered in arabic. Each must be typed on a separate page. Horizontal rules should be drawn lightly in pencil; vertical rules must not be used. Tables are expensive to set and correct; cost may be lowered and errors prevented if author submits tables typed with electric typewriter for photographic reproduction.

REFERENCES The style manuals mentioned above must be followed for form and for abbreviations of periodicals. Double space.

AUTHOR'S COPIES Each author receives 50 free copies of his *Postilla*. Additional copies may be ordered at cost by author when he returns galley proof. All copies have covers.

PROOF Author receives galley proof and manuscript for checking printer's errors, but extensive revision cannot be made on the galley proof. Corrected galley proof and manuscript must be returned to editors within seven days.

COPYRIGHT Any issue of *Postilla* will be copyrighted by Peabody Museum of Natural History only if its author specifically requests it.