POSTILLA

Published from 1950 to 2004, the short papers of the *Postilla* series reported on original research by the Yale Peabody Museum of Natural History's curators, staff, and research associates, and their colleagues, in the natural science disciplines represented by the collections of the Museum's curatorial divisions.

The *Postilla* series, which ceased publication with Number 232 (2004), was incorporated into the journal *Bulletin of the Peabody Museum of Natural History*, available from BioOne Complete at https://bioone.org/.

Yale Peabody Museum scholarly publications are archived through EliScholar, a digital platform for scholarly publishing provided by Yale University Library at https://elischolar.library.yale.edu/.



This work is licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License. https://creativecommons.org/licenses/by-nc-sa/4.0/

Yale peabody museum of natural history

Yale Peabody Museum of Natural History
New Haven, CT, 06511

Postilla Number 188 28 March 1983

A New Species of *Isectolophus* (Mammalia, Tapiroidea) from the Middle Eocene of Wyoming

Robert Milton Schoch

(Received 12 October 1982)

Abstract

A partial right dentary bearing M_3 from the middle Eocene (Bridger) of Wyoming represents a new species of *Isectolophus* here designated *I. radinskyi*. *I. radinskyi* differs from all other *Isectolophus* in its smaller size; it is intermediate in size between *Homogalax protapirinus* and its supposed descendant *Isectolophus latidens*.

Key Words

Isectolophus, Tapiroidea, Bridgerian, Eocene, fossil mammal.

Introduction

Among the previously uncatalogued middle Eocene fossil mammal remains from the Green River (Bridger) Basin, Wyoming, housed in the Yale Peabody Museum, New Haven (YPM), is a partial dentary bearing M₃ of a tapiroid which represents a new species of *Isectolophus*.

Family Isectolophidae Peterson, 1919 Genus *Isectolophus* Scott and Osborn, 1887

Isectolophus radinskyi Schoch, new species (Fig. 1, Table 1)

Holotype

YPM 40262, partial right dentary with posterior root of P_3 , roots of P_4 – M_2 and complete M_3 .

Horizon and Locality of the Type

Middle Eocene (Bridgerian) aged strata of the Bridger Formation, Green River (Bridger) Basin, Wyoming. Collected by L. LaMothe and J. W. Chew at Lone Springs, Wyoming, July 1874.

Hypodigm

Known only from the holotype.

Etymology

Named for Leonard B. Radinsky in honor of his many contributions toward a better understanding of tapiroid evolution.

Systematic Paleontology Class Mammalia Linnaeus, 1758 Order Perissodactyla Owen, 1848 Suborder Ceratomorpha Wood, 1937 Superfamily Tapiroidea Burnett, 1830 (Gill, 1872)

[©] Copyright 1983 by the Yale Peabody Museum of Natural History. All rights reserved. No part of this publication, except brief quotations for scholarly purposes, may be reproduced without the written permission of the Director, Yale Peabody Museum of Natural History.




Fig. 1Isectolophus radinskyi Schoch, new species, YPM 40262 (holotype), right dentary with M₃: A, occlusal view, stereo pair; B, labial view; C, lingual view. Scales are in mm.

Diagnosis

Smallest known species of *Isectolophus:* length of M₁₋₃ approximately 35.2 mm; dentary slightly shallower than that of *Isectolophus latidens.*

Description

YPM 40262 consists of a partial right dentary which bears a fragment of the posterior root of P_3 , the roots of P_4 – M_2 and the complete M_3 . P_3 – M_3 are all double-rooted and increase in size posteriorly. The preserved portion of the dentary is comparable in morphology to that of *Isectolophus latidens*, although somewhat shallower and more gracile (depth of ramus under $M_{1-2} = 21.0$ mm).

 M_3 of Isectolophus radinskyi is virtually identical in morphology to the M_3 of other species of Isectolophus, although much smaller in size. The transverse protolophid and hypolophid are parallel to one another, and very slightly notched in their centers (such that the protoconid and metaconid, hypoconid and entoconid remain relatively distinct) and trend slightly anterolabially. The protolophid is very slightly higher than

the hypolophid. The metaconid is very slightly higher than the protoconid; there is no metastylid. The paralophid is represented by a low anterolabial and anterior ridge. The metalophid is low and extends from the apex of the hypoconid to the posterolingual base of the protoconid. The trigonid and talonid (excluding the hypoconulid lobe) are subequal in length and width. The hypoconulid forms a large, broad, posterior heel bounded labially and posteriorly by a ridge which is highest labially (about half the height of that of the hypoconid) and decreases in height posterolingually. This ridge is minutely cuspidate and bears a shallow, posterior notch positioned slightly lingual to the anteroposterior midline of the tooth. M₂ bears a low, minutely cuspidate cingulid on its anterior and labial sides.

Discussion

Radinsky (1963) rediagnosed, revised and described *Isectolophus*, recognizing two species, the smaller and earlier *I. latidens* (middle Eocene, mean length $M_{1-3}=40.4$ mm) and the somewhat larger and later *I. annectens* (late Eocene, mean length $M_{1-3}=46.8$ mm). In preserved parts YPM 40262

Table 1

Measurements (in mm) of M_3 s of *Isectolophus latidens* and *Isectolophus radinskyi*, new species.

,			
	Length	Width	
lsectolophus latidens			
YPM 12563	17.9	9.0	
YPM 15297	18.7	9.1	
YPM 15298	18.0 ±	_	
YPM 16337	17.5	8.5+	
lsectolophus radinskyi			
YPM 40262	15.2	7.8	

4

is identical to *I. latidens*, except for its smaller size (Table 1). YPM 40262 is within the size range of the early Eocene isectolophid Homogalax protapirinus. However, it differs from Homogalax in lacking a metastylid on M₃. Metastylids are always present on the M_{1-3} of Homogalax (Radinsky 1963). Among other known Eocene North American tapiroids, YPM 40262 is within the size range of the early Eocene helaletid Heptodon calciculus and the middle Eocene Helaletes nanus. However, YPM 40262 differs from these species, and all Helaletidae, in bearing a relatively unreduced metalophid and a long and broad (large) hypoconulid on M₂. In helaletids the M₁₋₃ metalophids are greatly reduced and the M₃ hypoconulid is short and narrow or lost (Radinsky 1963). Thus, based on the morphology of M₃, YPM 40262 is referable only to the Isectolophidae, and of the two genera composing this family it is closest morphologically to other species of Isectolophus. Based on the lengths of M₁₋₃ in the known species of *Isectolophus* (Radinsky 1963), I. latidens is approximately 14 percent smaller than I. annectens and YPM 40262 is approximately 13 percent smaller than I. latidens. On this basis I coin the name Isectolophus radinskyi, new species, for YPM 40262, the smallest known species of Isectolophus.

Radinsky (1963, p. 74) postulated that *Homogalax protapirinus* gave rise to *Isectolophus latidens*. The discovery of *Isectolophus radinskyi* supports this hypothesis. *Isectolophus radinskyi* is intermediate between these two forms; it is within the size range of *Homogalax protapirinus*, but in nonmetric morphology is identical to *Isectolophus latidens*.

Acknowledgments

I thank Professor John H. Ostrom (YPM) for allowing me to study specimens in his care. I thank Mary Ann Turner (YPM) for help in locating specimens and locality information. The photographs were taken by William K. Sacco (YPM). Part of the costs of publication were supported by the O. C. Marsh Publication Fund, Yale Peabody Museum

Literature Cited

Radinsky, L. B. 1963. Origin and early evolution of North American Tapiroidea. Bull. Peabody Mus. Nat. Hist. (Yale Univ.) 17:1–115.

The Author

Robert Milton Schoch. Department of Geology and Geophysics and Yale Peabody Museum of Natural History, Yale University, 170 Whitney Avenue, P.O. Box 6666, New Haven, CT 06511.