

Source apportionment, identification and characterization, and emission inventory of ambient particulate matter in 22 Eastern Mediterranean Region countries

Faridi, Sasan; Yousefian, Fatemeh; Roostaei, Vahid; Harrison, Roy M.; Azimi, Faramarz; Niazi, Sadegh; Naddafi, Kazem; Momeniha, Fatemeh; Malkawi, Mazen; Moh'd Safi, Heba Adel; Rad, Mona Khaleghy; Hassanvand, Mohammad Sadegh

DOI:

[10.1016/j.envpol.2022.119889](https://doi.org/10.1016/j.envpol.2022.119889)

License:

Creative Commons: Attribution-NonCommercial-NoDerivs (CC BY-NC-ND)

Document Version

Peer reviewed version

Citation for published version (Harvard):

Faridi, S, Yousefian, F, Roostaei, V, Harrison, RM, Azimi, F, Niazi, S, Naddafi, K, Momeniha, F, Malkawi, M, Moh'd Safi, HA, Rad, MK & Hassanvand, MS 2022, 'Source apportionment, identification and characterization, and emission inventory of ambient particulate matter in 22 Eastern Mediterranean Region countries: a systematic review and recommendations for good practice', *Environmental Pollution*, vol. 310, 119889. <https://doi.org/10.1016/j.envpol.2022.119889>

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

1 **Source Apportionment, Identification and Characterization, and Emission**
2 **Inventory of Ambient Particulate Matter in 22 Eastern Mediterranean Region**
3 **Countries: A Systematic Review and Recommendations for Good Practice***

4 Sasan Faridi ^{a, b}, Fatemeh Yousefian ^c, Vahid Roostaei ^b, Roy M. Harrison ^{d, e}, Faramarz Azimi ^f, Sadegh Niazi ^g,
5 Kazem Naddafi ^{a, b}, Fatemeh Momeniha ^h, Mazen Malkawi ⁱ, Heba Adel Moh'd Safi ⁱ, Mona Khaleghy Rad ⁱ,
6 Mohammad Sadegh Hassanvand ^{a, b*}

7 ^a Center for Air Pollution Research (CAPR), Institute for Environmental Research (IER), Tehran University of
8 Medical Sciences, Tehran, Iran

9 ^b Department of Environmental Health Engineering, School of Public Health, Tehran University of Medical Sciences,
10 Tehran, Iran

11 ^c Department of Environmental Health Engineering, Faculty of Health, Kashan University of Medical Sciences,
12 Kashan, Iran

13 ^d School of Geography Earth and Environmental Science, University of Birmingham, Birmingham, UK

14 ^e Department of Environmental Sciences, Faculty of Meteorology, Environment and Arid Land Agriculture, King
15 Abdulaziz University, Jeddah, Saudi Arabia

16 ^f Environmental Health Research Center, Lorestan University of Medical Sciences, Khorramabad, Iran

17 ^g International Laboratory for Air Quality and Health, School of Earth and Atmospheric Sciences, Science and
18 Engineering Faculty, Queensland University of Technology (QUT), Brisbane, Queensland, Australia

19 ^h Center for Solid Waste Research, Institute for Environmental Research, Tehran University of Medical Sciences,
20 Tehran, Iran

21 ⁱ Environmental Health Exposures Centre for Environmental Health Action (CEHA), World Health Organization
22 (WHO), Jordan

23

24 * **Corresponding Author:** MS. Hassanvand, PhD, Center for Air Pollution Research (CAPR), Institute for
25 Environmental Research (IER) and Department of Environmental Health Engineering, School of Public Health,
26 Tehran University of Medical Sciences, Phone: +98 88978395, Fax: +98 88978397, 8th Floor, No. 1547, North Kargar
27 Avenue, Tehran, Iran (hassanvand@tums.ac.ir).

28 ★ This study is a part of a consultancy to provide a background assessment of air quality (AQ) management system
29 for the Eastern Mediterranean countries in order to improve their capabilities to use the updated World Health
30 Organization Air Quality Guidelines in compiling national AQ standards.

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49 **Abstract**

50 Little is known about the main sources of ambient particulate matter (PM) in the 22 Eastern Mediterranean
51 Region (EMR) countries. We designed this study to systematically review all published and unpublished
52 source apportionment (SA), identification and characterization studies as well as emission inventories in
53 the EMR. Of 440 articles identified, 82 (11 emission inventory ones) met our inclusion criteria for final
54 analyses. Of 22 EMR countries, Iran with 30 articles had the highest number of studies on source specific
55 PM followed by Pakistan (n = 15 articles) and Saudi Arabia (n = 8 papers). By contrast, there were no
56 studies in Afghanistan, Bahrain, Djibouti, Libya, Somalia, Sudan, Syria, Tunisia, United Arab Emirates and
57 Yemen. Approximately 72% of studies (51) were published within a span of 2015-2021. 48 studies
58 identified the sources of PM_{2.5} and its constituents. Positive matrix factorization (PMF), principal
59 component analysis (PCA) and chemical mass balance (CMB) were the most common approaches to
60 identify the source contributions of ambient PM. Both secondary aerosols and dust, with 12-51% and 8-
61 80% (33% and 30% for all EMR countries, on average) had the greatest contributions in ambient PM_{2.5}.
62 The remaining sources for ambient PM_{2.5}, including mixed sources (traffic, industry and residential (TIR)),
63 traffic, industries, biomass burning, and sea salt were in the range of approximately 4-69%, 4-49%, 1-53%,
64 7-25% and 3-29%, respectively. For PM₁₀, the most dominant source was dust with 7-95% (49% for all
65 EMR countries, on average). The limited number of SA studies in the EMR countries (one study per
66 approximately 9.6 million people) in comparison to Europe and North America (1 study per 4.3 and 2.1
67 million people respectively) can be augmented by future studies that will provide a better understanding of
68 emission sources in the urban environment.

69 **Keywords:** Source apportionment; Particulate matter; PM_{2.5}; PM₁₀; Eastern Mediterranean Region

70

71

72

73

74

75 **1. Introduction**

76 Long- and short-term exposures to ambient PM and its constituents are recognized as one of the most
77 pressing topics in modern-day public health (Brook et al. 2017; Faridi et al. 2017; Lelieveld et al. 2015).
78 Ambient PM is globally pervasive and is emitted from a number of sources, such as vehicular emissions,
79 industrial units, power generation, agricultural burning, wildfires, resuspended dust and dust storm events
80 (Goudie 2014; Hadley et al. 2018a; Hadley et al. 2018b; Querol et al. 2019). Source apportionment (SA)
81 studies have become increasingly conducted to determine the origins of ambient PM, particularly PM_{2.5} and
82 its constituents (Hopke et al. 2020; Karagulian et al. 2015; Münzel et al. 2021). Identification and
83 quantitative apportionment of ambient PM and its components (e.g., carbon species, sulfates, nitrates, metal
84 (loid)s, and unknown constituents) to their sources is an important step in providing information that guides
85 the development of air quality management strategies (Hopke et al. 2020; Karagulian et al. 2015; Münzel
86 et al. 2021; Taghvaei et al. 2018a). Additionally, source specific PM concentrations are also being applied
87 in health impact studies to be able to focus attention on those sources most likely to be responsible for the
88 observed adverse health effects (Hopke et al. 2020; Karagulian et al. 2015; Münzel et al. 2021). Based on
89 the undeniable importance of SA studies in designing mitigation strategies to improve ambient air quality
90 and its beneficial health consequences, the World Health Organization (WHO) released its initial
91 compilation of SA studies in 2015. This initial database was constructed by Karagulian and colleagues and
92 has been published by the Atmospheric Environment journal in 2015 (Hopke et al. 2020; Karagulian et al.
93 2015; Münzel et al. 2021). Moreover, the other review on SA for both PM₁₀ and PM_{2.5} at global scale has
94 been more recently conducted by Hopke et al. in 2020 (Hopke et al. 2020). In regional studies, Singh et al.
95 (2016) have quantitatively investigated the source contributions of ambient PM_{2.5} and PM₁₀ in South Asia
96 (Almeida et al. 2020; Singh et al. 2017). As a well-known multivariate receptor model, PMF has been
97 widely applied for SA of PM mass concentrations and its composition around the world (Soleimanian et al.
98 2019; Taghvaei et al. 2018a; Taghvaei et al. 2018b). Other receptor models, including CMB and PCA, and
99 also emission inventory approaches have been extensively used for identifying sources of PM (Arhami et
100 al. 2017; Esmaeilirad et al. 2020; Shahbazi et al. 2016a). These models/approaches can link the ambient

101 PM mass concentration and its constituents to a number of sources such as traffic related emissions,
102 secondary aerosols, industrial emissions, ship emissions, soil, road dust, biomass burning, sea salt, to name
103 but a few (Taghvaei et al. 2018b). Consequently, the findings provided by the aforementioned
104 models/approaches can be beneficial for authorities and policy-makers in each country or region in order
105 to mitigate the leading contributor of ambient air pollution (PM_{2.5}) and its associated health impacts
106 (Esmailirad et al. 2020; Taghvaei et al. 2018b).

107 The WHO reported that nearly 95% of the world's population lived in areas with ambient PM_{2.5}
108 concentrations exceeding the guideline (10 µg m⁻³) recommended by the 2005 WHO health-based air-
109 quality guidelines (Barn et al. 2018; Barzeghar et al. 2020; Brook et al. 2017; Rajagopalan et al. 2018).
110 However, the updated WHO Air Quality Guideline that was recently published (September 2021) is much
111 stricter (e.g., annual average guideline level for PM_{2.5} is 5 µg m⁻³ which is halved compared to previous
112 value). Though ambient PM_{2.5} levels are declining in many developed nations, the concentrations in the
113 EMR countries remain unchanged or continue to increase because successful implementation of air
114 pollution mitigation policies for the reduction of PM_{2.5} exposures may be near impossible in these nations
115 over a short- and even long-time periods (Barn et al. 2018; Bennett et al. 2019; Brook et al. 2017; Faridi et
116 al. 2020; Hadley et al. 2018b). As a result, the majority of the PM_{2.5}-related health and economic impacts
117 will occur in these countries (Barzeghar et al. 2020; Faridi et al. 2020). The underlying reasons for high
118 level of ambient PM_{2.5} concentrations across the EMR countries may be associated with dust storm events,
119 unsustainable development, continuing urbanization and industrialization, increasing mobile sources and
120 associated emissions alongside ineffective ambient air quality standards and ambient air pollution
121 abatement policies at national and subnational levels (Amini et al. 2017; Amini et al. 2019; Danaei et al.
122 2019; Shahsavani et al. 2020; Shamsipour et al. 2019). Based on the aforementioned issues, we designed
123 this study to systematically review all published and unpublished studies regarding the SA, identification
124 and characterization of ambient PM as well as emission inventories for it in the 22 countries in the EMR
125 and identify the most important sources for more effective air pollution control measures in this region.

126 **2. Methods**

127 **2.1. Search strategy, Study inclusion criteria, Article selection and Data extraction**

128 To identify the conducted studies (published and unpublished) concerning the SA, identification and
129 characterization of PM in 22 EMR countries (**Figure S1**) as well as emission inventories, several
130 approaches were used as follows: firstly, we conducted a systematic search of the articles based on the
131 Preferred Reporting and Items for Systematic Review and Meta-Analysis (PRISMA) criteria (Faridi et al.
132 2021; Faridi et al. 2022), as shown in the **Figure S2a** and **b**. The search was performed on December 23,
133 2021. To access the relevant studies, we queried three English language databases, including Scopus,
134 PubMed, and Web of Science Core Collection using the search keywords mentioned in **Table S1** and **S2**
135 (Supplemental file). We had two inclusion criteria as follows: **1)** study methods: any types of SA (e.g.,
136 PMF, CMB, and PCA) and emission inventory approaches, and **2)** language: English. To identify and select
137 the relevant papers, two authors (S.F (Sasan Faridi) and F.Y (Fatemeh Yousefian)) screened all articles,
138 independently and in duplicate. We selected eligible articles based on the title and abstract, if they fulfilled
139 both aforementioned inclusion criteria. Then, if the title and abstract of studies did not provide sufficient
140 detail for a decision, two authors reviewed the full text of articles independently. To justify the exclusion
141 of any article, a more rigorous second round of screening of all selected articles has been made by S.F. and
142 F.Y, and any conflict and discrepancies between the preceding reviewers on the studies was resolved by
143 M.S.H (Mohammad Sadegh Hassanvand) through verbal discussion and consensus. As another approach,
144 to increase the sensitivity and gather the relevant records that may be unpublished, we have tried to identify
145 them through health experts at the World Health Organization/ Regional office of the Eastern
146 Mediterranean Regional Office/Regional Centre for Environmental Health Action (WHO/EMRO/CEHA).
147 S.F and F.Y independently extracted the detailed information on the characteristics of studies, including
148 study ID, country/city of the studies, the used approaches to identify sources, air pollutants and identified
149 sources and their contributions (**Table 1** (for the studies of SA, identification and characterization of PM)
150 and **Table S3** (for emission inventories)).


151

152 3. Results and Discussion

153 3.1. Overview of the included SA, identification and characterization studies

154 **Table 1** provides a summary of SA, identification and characterization studies for PM in the EMR
155 countries. As shown in the **Table 1**, detailed information on study ID, time interval of sampling, the used
156 method for identifying SA of PM, the country and city of study and the apportioned air pollutants have
157 been mentioned. Based on the information supplied in **Figure S2** and **Table S1**, of 333 articles identified,
158 71 met our inclusion criteria for final analyses. Amongst all 22 countries in the EMR countries, Iran with
159 30 articles had the highest number of studies on source specific PM followed by Pakistan (n = 15 articles)
160 and Saudi Arabia (n = 8 papers), as shown in **Table 1**. Note that there were no studies in Afghanistan,
161 Bahrain, Djibouti, Libya, Somalia, Sudan, Syria, Tunisia, United Arab Emirates and Yemen.
162 Approximately 72% of all studies (51) were published within a span of 2015-2021 (**Figure S3**). **Figure S4**
163 shows the pie chart of the number of papers with a given from the EMR countries with the listed particle
164 metric. Almost 68% of published papers (48) have been performed to identify the sources of PM_{2.5} and its
165 constituents, including polycyclic aromatic hydrocarbons (PAHs), carbonaceous aerosols (like total organic
166 carbon (OC), elemental carbon (EC) and black carbon (BC)) and heavy metals (**Figure 1**). In 12, 4 and 10
167 studies, the authors of published papers have identified the source contributions of PM₁₀, PM_{10-2.5} and TSP.
168 Interestingly, one study in Lebanon has determined the source contribution of PM₁ (**Figure S4**). Moreover,
169 To specify the sources of ambient PM using different models and approaches mentioned above, the included
170 studies have analyzed various chemical constituents (metals, water-soluble ion, PAHs, OC, EC, and BC),
171 as presented in **Figure 1**. 34 studies have analyzed simultaneously two and more than two chemical
172 constituents (metals + PAHs (two studies), (metals + BC (four studies), (metals + water soluble ions (six
173 studies), water soluble ions + OC + EC (one study), metals + water soluble ions + BC (2 studies), metals +
174 water soluble ions + OC + EC (10 studies) and metals + water soluble ions + PAHs + OC + EC (nine
175 studies)) for specifying ambient PM air pollution sources. However, one group of chemical constituents
176 (PAHs (19 studies) or metals (15 studies) or water-soluble ions (one study) or OC (one study)) has been
177 analyzed specifying ambient PM air pollution sources. As presented in **Table 1**, the number of samples


178 used to conduct models or approaches for specifying ambient PM sources in the countries of the EMR
 179 countries ranged from 12 to 1802 samples (153 for all studies, on average), though several studies have not
 180 reported the number of samples used. **Figure 2** gives the data on the number of studies with the application
 181 of different approaches. Given the used approaches, PMF, PCA and CMB were the most common
 182 approaches to identify the source contributions of PM because they were used in 39, 15 and 4 studies
 183 respectively.



184 **Figure 1.** Analyzed various chemical constituents (the number and percentage of studies) to conduct SA
 185 models/ approaches (Considering simultaneously two and more than two chemical constituents for
 186 specifying ambient PM sources).

187

188



189

190 **Figure 2.** The number of studies with the application of different approaches for SA, identification and
 191 characterization of ambient PM in the EMR countries.

192

193

194

195

196

197

198

199

200

Table 1. Summary of SA, identification and characterization studies for PM air pollution in the EMR countries.

Study ID	Time interval	Site type of sampling	Method	Country/city	Air pollutants	# of samples
(Taghvaei et al. 2018a)	May 2012 to June 2013	Urban (residential)	PMF	Iran/ Tehran	PM _{2.5} -bound PAHs	74
(Altwayjiri et al. 2021; Taghvaei et al. 2018b)	May 2012 to June 2013	Urban (residential)	PMF	Iran/ Tehran	PM _{2.5}	78
(Soleimani et al. 2019)	May 2012 to June 2013	Urban (residential)	PMF	Iran/ Tehran	PM _{10-2.5}	60
(Esmailirad et al. 2020)	February 2014 to February 2015 and from late-January 2017 to mid-April 2017	Residential	PMF	Iran/ Tehran	PM _{2.5}	116
(Ali-Taleshi et al. 2021a; Ali-Taleshi et al. 2021b; Ali-Taleshi et al. 2021c)	March 2018 to February 2019	Urban	PMF	Iran/ Tehran	PM _{2.5} -bound PAHs	135
(Maleki et al. 2021)	March 2018 to March 2019	Urban (low, moderate and high traffic)	UNMIX model	Iran/ Tehran	PM _{2.5} -bound PAHs	60
(Shahsavani et al. 2017)	Every 3 days during the spring of 2015	Urban and Sub-urban	PCA	Iran/ Shiraz	PM ₁₀ -bound PAHs	60
(Kermani et al. 2021b)	March 2018 to March 2019	Urban (residential)	PCA	Iran/ Karaj	PM _{2.5} -bound heavy metals	68
(Kermani et al. 2021a)	March 2018 to March 2019	Industrial area, High traffic areas; Middle and low traffic areas,	UNMIX model	Iran/ Tehran	PM _{2.5} -bound PAHs and heavy metals	108
(Gholampour et al. 2015)	January 2013 to September 2013	Sub-urban (Lake region)	PCA	Iran/ Urmia	TSP and PM ₁₀	48
(Akhbarizadeh et al. 2021)	During December 2016 and September 2017	Urban	PMF	Iran/ Bushehr	PM _{2.5} -bound PAHs	46
(Eivazzadeh et al. 2021)	April 2017 to March 2018	Urban and Sub-urban	PCA	Iran/ Tabriz	TSP	12
(Gholampour et al. 2016)	September 2012 to June 2013	Urban and Sub-urban (industrial)	PCA	Iran/ Tabriz	TSP and PM ₁₀	78

Study ID	Time interval	Site type of sampling	Method	Country/city	Air pollutants	# of samples
(Soleimani et al. 2018)	from March 2014 to March 2015	Urban and Industrial	PCA	Iran/ Isfahan	PM _{2.5} -bound heavy metals	Not reported
(Sowlat et al. 2013)	April 2010 to March 2011	Urban	PMF	Iran/ Ahvaz	PM ₁₀	72
(Sowlat et al. 2012)	April 2010 to March 2011	Urban	PMF	Iran/ Ahvaz	TSP	72
(Ashrafi et al. 2018)	Not reported	Urban (residential)	PMF, PMF-CMB, SPECIATE database-CMB	Iran/ Ahvaz	TSP	74
(Alidadi et al. 2021)	September 2018 to August 2019	City center-traffic, Near industrial zone, Far from urban traffic, Semi-traffic, Background	UNMIX model	Iran/ Ahvaz	PM ₁₀	100
(Najmeddin and Keshavarzi 2019)	February 8 to March 9, 2017	Urban (residential), Highway and Industrial	PMF	Iran/ Ahvaz	PM ₁₀	30
(Kholdebarin et al. 2015)	9 months (from 30/07/2012 up to 21/04/2013)	Urban (residential), Traffic zone and Industrial	CMB	Iran/ Tehran	PM ₁₀	32
(Halek et al. 2010)	April 2004 to March 2005	Urban	Diagnostic ratios and PCA	Iran/ Tehran	PAHs (particulate and gaseous phases)	Not reported
(Azimi-Yancheshmeh et al. 2021)	Feb. 2018 to Jan. 2019	Urban (residential), Highway and Industrial	PMF	Iran/ Karaj	PM _{2.5} -bound PAHs	130
(Moeinaddini et al. 2014)	October 2011 to March 2012	Traffic zone (Between two highways)	PMF	Iran/ Tehran	Respirable particles-bound PAHs (PM ₄ -bound PAHs), Respirable particles-bound n-alkanes (PM ₄ -bound n-alkanes)	155
(Soleimani et al. 2021)	December 2017 to September 2018	Urban and Industrial	PMF	Iran/ Isfahan	PM _{2.5} -bound PAHs	200
(Abbasi et al. 2020)	August 6th to 31st, 2017 and January 28th to February 6th, 2018	Urban and Industrial	PMF	Iran/ Asaluyeh	PM ₁₀	48

Study ID	Time interval	Site type of sampling	Method	Country/city	Air pollutants	# of samples
(Arhami et al. 2018)	February 2014 to February 2015	Residential	CMB	Iran/ Tehran	PM _{2.5} and Carbonaceous compounds (organic and elemental)	51
(Arhami et al. 2017)	February 2014 to February 2015	Residential	PCA-CMB	Iran/ Tehran	PM _{2.5}	51
(Altuwajjiri et al. 2022)	December 2019– March 2020 and May 2020–August 2020	Urban (residential)	PCA-MLR	Saudi Arabia/ Riyadh	Oxidative potential of PM ₁₀	Not reported
(Nayebare et al. 2018)	February 26, 2014 – January 27, 2015	Urban (residential) and Semi-industrial	PMF	Saudi Arabia/ Makkah	PM _{2.5}	40
(Khodeir et al. 2012)	Between June and September, 2011	Urban (residential and new residential) and Sub-urban	SPlus Factor Analysis model with Varimax rotations	Saudi Arabia/ Jeddah	PM _{2.5} and PM ₁₀	84 for PM _{2.5} and 80 for PM ₁₀
(Nayebare et al. 2016)	May 6 th – June 17 th 2013	Urban (residential) and Semi-industrial	PMF	Saudi Arabia/ Rabigh	PM _{2.5}	40
(Bian et al. 2016)	September 2011 to September 2012	Urban (residential) and Industrial	PMF	Saudi Arabia/ Riyadh city	PM ₁₀	167
(Cusack et al. 2020)	September 2015 and December 2017	Semi-urban and Industrial	PMF	Saudi Arabia/ Jeddah	TSP	238
(Alghamdi et al. 2015)	Between 23 February 2013 and 23 April 2013	Urban (residential and background) and Industrial	PMF	Saudi Arabia/ Jeddah	PAHs (particulate and gaseous phases)	180
(Shaltout et al. 2017)	December 2012 to January 2014	Urban	PMF	Saudi Arabia/ Makkah	PM _{2.5}	Not reported
(Lurie et al. 2019)	August 2008 through August 2009	Urban (residential and commercial) and Industrial	PMF	Pakistan/ Karachi	PM _{2.5}	377
(Harrison et al. 1997)	A period of 12 months (10/92 to 10/93)	Urban	PCA-MLR	Pakistan/ Lahore	TSP and PM ₁₀	62
(Mehmood et al. 2020)	Four seasons in 2017	Urban	PCA-MLR	Pakistan/ Islamabad	PM _{2.5} -bound PAHs and PM ₁₀ -bound PAHs	160
(Lodhi et al. 2009)	November 2005 to March 2006	Urban (residential)	PMF	Pakistan/ Lahore	PM _{2.5}	310

Study ID	Time interval	Site type of sampling	Method	Country/city	Air pollutants	# of samples
(Shaheen et al. 2005)	From October 2002 to May 2003	Urban	PCA	Pakistan/ Islamabad	TSP	134
(Shah et al. 2012)	September 2008– March 2009	Urban	PCA	Pakistan/ Islamabad	TSP	153
(Mansha et al. 2012)	January 2006 to January 2008	Urban (residential)	PMF	Pakistan/ Karachi	PM _{2.5}	402
(Ishtiaq et al. 2021)	Summer in 2014	Urban	PCA-MLR	Pakistan (Karachi, Lahore, Faisalabad, Rawalpindi, Multan, Quetta, Peshawar and Gilgit)	PAHs (particulate (PM _{2.5}) and gaseous phases)	Not reported
(Alvi et al. 2020)	September 2015 to December 2016	Urban	PCA	Pakistan/ Faisalabad	PM ₁₀	117
(Raja et al. 2010)	November 2005 to January 2006	Urban	PMF	Pakistan/ Lahore	PM _{2.5}	129
(Ahmad et al. 2020)	July 2018	Urban	PCA	Pakistan/ Lahore	PM _{2.5}	30
(Ahmad et al. 2021)	February 2019	Urban	PCA-MLR	Pakistan (Lahore and Peshawar)	PM _{2.5}	60
(Shahid et al. 2018)	March – April 2009	Urban	PMF	Pakistan/ Karachi	PM ₁₀	Not reported
(Hamid et al. 2018)	From 15 th February to 22 nd April 2014	Urban, Traffic and Industrial	PCA-MLR	Pakistan (Islamabad and Rawalpindi)	PAH	25
(Khanum et al. 2021)	February 2014 and March 2014	Urban	PCA	Pakistan/ Lahore	PM ₁₀	30
(Javed et al. 2019)	May to December 2015	Urban (residential and commercial) and Sub-urban	PMF	Qatar/ Doha	PM _{2.5} and PM _{10-2.5}	278
(Javed and Guo 2021)	May to December 2015	Urban (residential and commercial)	PMF	Qatar/ Doha	PM _{2.5} and PM _{10-2.5}	108 for each of air pollutants
(Heo et al. 2017)	January to December 2007	Urban (residential)	PMF	Middle Eastern cities in Jordan (Amman, Aqaba, and Rachma), and Palestine (East Jerusalem, and Hebron)	PM _{2.5}	580

Study ID	Time interval	Site type of sampling	Method	Country/city	Air pollutants	# of samples
(Heo et al. 2017)	January to December 2007	Urban (residential)	PMF	Middle Eastern cities in Jordan (Amman, Aqaba, and Rachma), and Palestine (East Jerusalem, and Hebron)	PM _{2.5}	580
(von Schneidemesser et al. 2010)	January to December 2007	Urban (residential and commercial)	PMF	Middle Eastern cities in Jordan (Amman and Rachma) and Palestine (East Jerusalem and Hebron)	PM _{2.5} -carbonaceous aerosols	120
(von Schneidemesser et al. 2010)	January to December 2007	Urban (residential and commercial)	PMF	Middle Eastern cities in Jordan (Amman and Rachma) and Palestine (East Jerusalem and Hebron)	PM _{2.5} -carbonaceous aerosols	120
(Hamad et al. 2015)	From September 2012 through September 2013	Urban (commercial)	CMB	Iraq/ Baghdad	PM _{2.5} -carbonaceous aerosols	53
(Alahmad et al. 2021)	October 2017 to October 2019	Urban (residential)	PMF	Kuwait (Kuwait city and Ali Sabah Al-Salem)	PM _{2.5}	1186
(Al-Dabbous and Kumar 2015)	31 days during the summer months of May and June 2013	Urban	PMF	Kuwait/ Fahaheel	Concentration of particles with 5 – 1000 nm	Not reported
(Waked et al. 2015)	July 2011	Sub-urban	Chemical-transport model (CTM)	Lebanon/ Beirut	PM _{2.5} -Total organic carbon (OC)	20
(Jaafar et al. 2021)	November 2016 and November 2017	Urban (background)	PMF	Lebanon/ Beirut	PM _{2.5} -bound PAHs	55
(Saliba et al. 2010)	Between 2003 and 2007	Urban (residential and commercial)	Mettler-Toledo microgram balance (model UMX2)	Lebanon/ Beirut	PM _{2.5} and PM ₁₀	42 for PM _{2.5} and 41 for PM ₁₀
(Alolayan et al. 2013)	Between February 2004 and October 2005	Urban (residential)	PMF	Kuwait/ Kuwait city	PM _{2.5}	1802
(Shaltout et al. 2020)	October 2010 to May 2011	Urban (residential) and Industrial	PCA	Egypt/ Greater Cairo	PM _{2.5}	32
(Boman et al. 2013)	September 2010 and May 2011	Urban (residential)	PCA	Egypt/ Central Cairo	PM _{2.5}	54



Study ID	Time interval	Site type of sampling	Method	Country/city	Air pollutants	# of samples
(Khairy and Lohmann 2013)	(July–August, 2010) and (December–January, 2011)	Urban (residential), Traffic and Industrial	FA/MLR, PMF and UNMIX	Egypt/ Alexandria	PAHs (gaseous + particulate phases)	150
(Abu-Allaban et al. 2002; Abu-Allaban et al. 2007)	21 February–3 March and 29 October–27 November 1999	Urban (residential and background), Traffic, Industrial	CMB	Egypt/ Cairo	PM _{2.5} and PM ₁₀	73
(Tahri et al. 2013)	February 2007 to February 2008	Traffic	PMF	Morocco/ Kenitra city	PM _{2.5} and PM _{10-2.5}	63 for each of air pollutants
(Abdul-Wahab 2004)	Not Reported	Urban (residential) and Industrial	PCA	Oman/ Sohar	TSP-heavy metals	31

3.2. Source contributions to ambient PM_{2.5} and PM₁₀ in the EMR countries based on the SA, identification and characterization studies


Figure 3 and **4** reveal the relative concentrations of sources to PM_{2.5} and PM₁₀ mass concentrations in the included studies. Included studies reported seven major sources for ambient PM_{2.5} and PM₁₀ including dust, traffic, industries, sea salt, biomass burning, and secondary aerosols. Since, some of them stated no specific sources for ambient PM_{2.5} and PM₁₀, categorized into an unspecified category. As shown in **Figure 3**, both secondary aerosols and dust, with 12-51% (33% for all EMR countries, on average) and 8-80% (30% for all EMR countries, on average) made the greatest contributions to ambient PM_{2.5} air pollution. The remaining sources for ambient PM_{2.5} mass concentrations, including mixed sources (traffic, industry and residential(TIR)), traffic, industries, biomass burning, and sea salt were in the range of approximately 4-69%, 4-49%, 1-53%, 7-25% and 3-29%, respectively. Of the above-mentioned sources, dust had the greatest contribution to ambient PM_{2.5} in Egypt (68%) and Kuwait (49%), while the figures for Saudi Arabia, Iran and Qatar were lower at 24%, 20% and 12%, respectively. In the studies conducted in Palestine, Jordan, Pakistan and Qatar, approximately 42%, 39%, 32% and 30% of ambient PM_{2.5} mass concentrations were attributed to secondary aerosols, respectively. For the study performed in Morocco, nearly 53% of ambient PM_{2.5} mass concentration was attributed to industries, whereas the figure for Jordan was lower at 3%. Traffic was the most dominant source for ambient PM_{2.5} mass concentrations in Iran (41%), while it had the lowest contribution in PM_{2.5} sources for Egypt (9%) and Jordan (10%). Biomass burning had nearly the same contribution in ambient PM_{2.5} (13%) in the studies conducted in both Jordan and Palestine.

Regarding PM₁₀ (**Figure 4**), the most dominant source of ambient PM₁₀ mass concentrations was dust with 7-95% (49% for all EMR countries, on average), whereas the figure for secondary aerosols in these countries had the lowest contributions and ranged from 7-25% (18% on average). The contribution of mixed sources (traffic, industry and residential), sea salt, traffic, industry, and biomass burning were in the range of approximately 18-94%, 9-60%, 4-72%, <1-31% and 9-25%, respectively. In the studies performed in Iran, the most dominant source for ambient PM₁₀ mass concentrations were sea salt and dust (26 % and

25%, on average). Given studies in Lebanon and Pakistan, the contribution of traffic for ambient PM₁₀ mass concentrations (54% and 34%), was higher than that of in Iran and Saudi Arabia (21% and 11%). Comparing all EMR regions, industries made the highest contributions to PM₁₀ mass concentrations (26.0%) in Pakistan, while these were the lowest for Iran (10.1%).




Saudi Arabia




- Traffic (%)
- Mixed sources (TIR) (%)
- Industries (%)
- Dust (%)
- Others (%)

Egypt




- Traffic (%)
- Mixed sources (TIR) (%)
- Industries (%)
- Dust (%)
- Others (%)

Qatar




- Traffic (%)
- Mixed sources (TIR) (%)
- Secondary aerosols (%)
- Dust (%)
- Sea salt (%)
- Others (%)

Kuwait




- Traffic (%)
- Mixed sources (TIR) (%)
- Dust (%)
- Others (%)

Jordan




- Traffic (%)
- Mixed sources (TIR) (%)
- Secondary aerosols (%)
- Industries (%)
- Biomass burning (%)
- Dust (%)
- Sea salt (%)

Palestine




- Traffic (%)
- Mixed sources (TIR) (%)
- Secondary aerosols (%)
- Industries (%)
- Biomass burning (%)
- Dust (%)
- Sea salt (%)

Morocco



- Traffic (%)
- Industries (%)
- Dust (%)

Lebanon



- Secondary aerosols (%)
- Biomass burning (%)
- Dust (%)
- Others (%)





Figure 3. Relative contributions of sources to PM_{2.5} mass concentrations in the included studies and the average of relative contributions of each source in the EMR countries.







Figure 4. Relative contributions of sources to PM_{10} mass concentrations in the included studies and the average of relative contributions of each source in the EMR countries.

3.3. Comparison to previous SA studies

We have surveyed three systematic reviews regarding SA of ambient PM air pollution previously conducted at a global scale (Hopke et al. 2020; Hopke et al. 2022; Karagulian et al. 2015). Moreover, we have explored a more recently paper authored by McDuffie and colleagues on source sector and fuel contributions to ambient PM_{2.5} and attributable mortality across multiple spatial scales (McDuffie et al. 2021). As shown in **Table S4**, two studies (out of three) have surveyed the contributions of sources of PM_{2.5} and PM₁₀ air pollution, and another study has explored SA of particle number concentrations. As reported, in the findings of our study above, we have reviewed and reported not only the included studies in these reviews but also additional studies in detail. Figure S6 gives information on the relative source of PM_{2.5} mass in 2017 based on the study of (McDuffie et al. 2021). As shown in **Figure S5** and **Table S5**, the study of (McDuffie et al. 2021) has specified 20 sources (AFCID (anthropogenic, fugitive, combustion, and industrial) dust, windblown dust, agriculture, energy coal, energy noncoal, industry coal, industry noncoal, nonroad transport, road transport, and others: Residential Coal Combustion, Residential Biofuel Combustion, Residential Other Combustion, Commercial Combustion, Other Combustion, Waste, International Shipping, Agricultural Waste Burning, Other Open Fire, Solvent, and Remaining Sources) for ambient PM_{2.5} air pollution in all countries of the EMR. Of these sources, windblown dust with 33-78% (56% for all EMR countries, on average) had the greatest contribution in ambient PM_{2.5} air pollution of all EMR countries, followed by energy (noncoal) with 1-23% (about 10% for all EMR countries, on average) and road transport with 2-11% (6% for all EMR countries, on average). The lowest contributions were reported for the sources of nonroad transport (< 1% for all EMR countries, on average) and industry (coal) (< 1% for all EMR countries, on average). Though this study has surveyed and reported the contribution of sources to PM_{2.5} mass concentrations for all EMR countries, there were considerable differences with the findings of our study. Detailed information on the contribution of each of the sources to ambient PM_{2.5} air pollution in all EMR countries based on the study of (McDuffie et al. 2021) is presented in **Table S5**. It should be noted that the McDuffie et al (2021) study uses a global scale dispersion model based upon an emissions inventory, and not receptor modelling based upon atmospheric measurement data and receptor modelling.

3.4. Overview of the included emission inventory studies and their findings

Table S3 shows the detailed information concerning the emission inventory studies for ambient air pollution in the EMR countries. As shown in **Table S3**, 11 studies have been conducted on the emission inventory of ambient air pollutants in 6 countries (out of 22) of the EMR. Of these studies, 6 have been conducted in Iran (five studies for Tehran and one for Mashhad city), while the remainder were conducted in Afghanistan (Kabul), Kuwait (Jahara), Pakistan (Khyber Pakhtunkhwa and Balochistan regions), Morocco (Casablanca) and Lebanon (Beirut and its 26 sub-regions). **Figure S6** reveals just relative contributions (%) of each sector to PM emissions in the EMR countries. Given Iran, the studies of (Heger and Sarraf 2018; Hosseini and Shahbazi 2016; Shahbazi et al. 2016a; Shahbazi et al. 2016b; Shahbazi et al. 2019) have reported that the mobile sources contributed on average 70% of ambient PM air pollution in Tehran, while the figures for other sources such as energy conversion, industry, household or commercial sectors and terminals were significantly lower, at 20%, 7%, 2% and 1%, respectively. Based on the study of (Shahbazi et al. 2021), nearly half of TSP, PM₁₀ and PM_{2.5} emissions in Mashhad was contributed by mobile sources. The second in terms of contribution was energy conversion with 32-37% of TSP, PM₁₀ and PM_{2.5} emissions. With regard to Lebanon (Beirut), the largest contributor to ambient PM₁₀ and PM_{2.5} air pollution was industry, accounting for 62% and 59%, followed by domestic sector and mobile sources (Waked et al. 2012). For Afghanistan (Kabul), the highest contribution to ambient PM₁₀ emissions was attributed mobile and dust (52%), followed by domestic sector (31%) and industry (12%) (Karim 2020)

4. Perspectives and Recommendations

Ambient air pollution is a complex mixture of particulate and gaseous primary and secondary pollutants that varies in levels and composition according to time and place (Rajagopalan and Landrigan 2021). Among all air pollutants, PM air pollution has been recognized as the most notable environmental cause of respiratory-, cardiovascular-, and cancer-related deaths worldwide (Faridi et al. 2021; McDuffie et al. 2021; Rajagopalan et al. 2018; Southerland et al. 2022; Tian et al. 2019). In the recent Global Burden of Disease (GBD) study, it was indicated that exposure to PM_{2.5} was globally responsible for 4.14 million (95% Confidence Interval (CI): 3.45-4.80) deaths in 2019 (GBD 2019). One of the major challenges to improve

this sobering situation is to understand the origin of the ambient air pollution to make sure whether air quality plans are targeting/have targeted the major sources to ensure effective results (Hopke et al. 2020; Thunis et al. 2019). Overall, air quality plans include the following objectives: **1)** identify and quantify the sources that contribute most to air pollution levels; **2)** inform on the efficiency of mitigation strategies; **3)** identify the most appropriate measures to be applied to each of these sources and/or **4)** evaluate scenarios for future emissions to assess the effectiveness of mitigation measures to control air quality levels (Hopke et al. 2020; Thunis et al. 2019). SA studies aim at understanding the origin of ambient air pollution and are generally used for the two first inter-related tasks (Hopke et al. 2020; Thunis et al. 2019). As a result, accurately designing and conducting SA studies can affect the objectives 2, 3 and 4. Though full details are reviewed elsewhere (Hopke et al. 2020; Hopke and Jaffe 2020; Thunis et al. 2019), hereunder we introduce several recommendations for good practice in the future studies to make reported SA more useful to researchers and even policymakers.

- As mentioned above, more than 25 studies have apportioned the sources of ambient PM air pollution in the EMR countries using PCA and PCA-MLR (20 studies), SPlus Factor Analysis model with Varimax rotations (2 studies), and UNMIX (3 studies). These outdated data analysis methods have been replaced with more quantitative data analytical tools (like PMF and CMB) that provide much greater and more appropriate information on sources of variation in the data (Hopke et al. 2020; Mousavi et al. 2018; Paatero et al. 2014; Sowlat 2019). Unlike PCA, PMF is a weighted method which is able to give most weight to those chemical components with the least measurement uncertainty, hence giving more reliable results. Consequently, the more recently published studies could reanalyze their results using PMF or possibly CMB. Use of CMB depends upon availability of local chemical profiles of sources which may not be readily available. We recommend that the researchers of future studies use these approaches and tools (PMF and CMB) rather than outdated ones to specify ambient PM air pollution sources.
- As mentioned in **Table 1**, 5, 8 and 23 studies have specified the sources of ambient PM (PM_{2.5}, PM₁₀ and PM_{10-2.5}) and its constituents based on < 50, 51-75 and > 100 samples using PMF.

Additionally, 5, 6 and 11 studies have specified the sources of ambient PM_{2.5} air pollution based on < 50, 51-75 and > 100 samples using PMF. This tool is often used on speciated ambient PM_{2.5} datasets with 10-20 chemical species > 100 samples (EPA 2015). As a general rule, the multivariate statistical methods (PMF and PCA) are unreliable unless the number of samples exceeds the number of analytical variables by at least threefold, and that criterion does not appear to have been met in all cases.

- As reported in **Figure 1**, to specifying ambient PM air pollution sources in the EMR countries, approximately half of studies (36) have analyzed just one group of chemical constituents (PAHs (19 studies) or metals (15 studies) or water-soluble ions (one study) or OC (one study)). To specify sources of air pollution precisely, it is necessary to characterize multiple chemical components of PM_{2.5} and PM₁₀. When a study apportions the sources of air pollution just based on a given chemical component of PM (like PAHs or other components) the results differ compared to a study that applied the model based on a wide range of components (like PAHs + heavy metals + water soluble ions). The receptor model will provide more precise findings in order to allow adoption of better air pollution control abatement strategies, if the study can characterize a wide range of components of PM (PAHs, heavy metals, OC, EC, BC and other components). Consequently, we recommend that the researchers consider more than one generic component of PM to characterize ambient air pollution sources.
- It needs to be recognized that some of these studies are based upon short-term sampling campaigns (see **Table 1** for sampling dates), and are hence unlikely to be representative of long-term average conditions. They should hence be regarded with caution, as seasonal emission and climate factors, and even short-term weather events influence particulate matter concentrations and SA. They should therefore be regarded as indicative of, rather than defining the SA for a location. Air quality data are also site-specific, and this needs to be taken into account in data interpretation and use. There are obvious differences between roadside and urban background sites, coastal and inland sites, and those with predominantly bare soil relative to those with good ground cover. The design

of sampling programmes should aim to use sites representative of a wider area, and not just the immediate locality.

- Amongst all 22 countries in the EMR, Iran with 30 articles had the highest number of studies on source specific PM air pollution followed by Pakistan (n = 15 articles) and Saudi Arabia (n = 8 papers). Note that there were no studies in Afghanistan, Bahrain, Djibouti, Libya, Somalia, Sudan, Syria, Tunisia, United Arab Emirates and Yemen. When we are comparing the number of conducted studies in the EMR countries with their population (**Table S6**), it should be highlighted that the number of studies is almost negligible (one study per approximately 9.6 million people (71 studies per 680 million people in the EMR countries. However, this rate for Europe, North America, South America, Western Asia, South-eastern Asia, Eastern Asia (without China) and China was one study per 4.3, 2.1, 39.7, 26.1, 46.8, 9.5 and 5.7 million people respectively. Thus, the currently limited numbers of SA studies in EMR countries in comparison to Europe and North America can be augmented by additional future studies that will provide a better understanding of emission sources in the urban environment.
- Note that all included studies have been designed and conducted by researchers in the universities. In fact, the included studies have not been requested by organizations responsible for reducing ambient air pollution, or these studies have not been part of an integrated program for specifying ambient PM air pollution sources in order to design air quality plans or explore the efficiency of mitigation strategies. With regard to these issues, to accurately identify the origin of ambient air pollution constitutes as an essential step in air quality management to subsequently identify measures to control air pollution, and a much greater number of studies as part of an integrated program is needed in these countries.
- Also noteworthy is the fact that recent developments indicates that source apportionment done simultaneously with more than one receptor model (PMF + CMB) could provide more robust results (Contini et al. 2016). As a result, we suggest that the researchers consider simultaneously two receptor models to improve the robustness of their findings.

- Finally, one of the most important issues for SA is sampling instruments and analytical procedures (type of samplers employed, type of filters used and the process of its preparation before and after sampling, and analytical procedures) that are used to produce the input data for SA models and tools. Prior to conduct of the study, we recommend that the researchers read the relevant protocols and papers in detail. Full details regarding requirements for the quality of data used for SA by the more modern and statistically appropriate tools and methods like PMF and CMB are presented elsewhere (Belis et al. 2020; Hopke et al. 2020; Karagulian et al. 2015; Thunis et al. 2019).

Acknowledgments

This study was financially supported by Environmental Health Exposures Centre for Environmental Health Action (CEHA), World Health Organization (WHO), Jordan (grant number: 202778822).

References

- Abbasi S, Keshavarzi B, Moore F, Hopke PK, Kelly FJ, Dominguez AO. 2020. Elemental and magnetic analyses, source identification, and oxidative potential of airborne, passive, and street dust particles in asaluyeh county, iran. *Science of The Total Environment* 707:136132.
- Abdul-Wahab SA. 2004. Source characterization of atmospheric heavy metals in industrial/residential areas: A case study in oman. *Journal of the Air & Waste Management Association* 54:425-431.
- Abu-Allaban M, Gertler AW, Lowenthal DH. 2002. A preliminary apportionment of the sources of ambient pm₁₀, pm_{2.5}, and vocs in cairo. *Atmospheric Environment* 36:5549-5557.
- Abu-Allaban M, Lowenthal D, Gertler A, Labib M. 2007. Sources of pm₁₀ and pm_{2.5} in cairo's ambient air. *Environmental Monitoring and Assessment* 133:417-425.
- Ahmad M, Cheng S, Yu Q, Qin W, Zhang Y, Chen J. 2020. Chemical and source characterization of pm_{2.5} in summertime in severely polluted lahore, pakistan. *Atmospheric Research* 234:104715.
- Ahmad M, Yu Q, Chen J, Cheng S, Qin W, Zhang Y. 2021. Chemical characteristics, oxidative potential, and sources of pm_{2.5} in wintertime in lahore and peshawar, pakistan. *J Environ Sci* 102:148-158.
- Akhbarizadeh R, Dobaradaran S, Torkmahalleh MA, Saeedi R, Aibaghi R, Ghasemi FF. 2021. Suspended fine particulate matter (pm_{2.5}), microplastics (mps), and polycyclic aromatic hydrocarbons (pahs) in air: Their possible relationships and health implications. *Environmental Research* 192:110339.
- Al-Dabbous AN, Kumar P. 2015. Source apportionment of airborne nanoparticles in a middle eastern city using positive matrix factorization. *Environmental Science: Processes & Impacts* 17:802-812.
- Alahmad B, Al-Hemoud A, Kang CM, Almarri F, Kommula V, Wolfson JM, et al. 2021. A two-year assessment of particulate air pollution and sources in kuwait. *Environmental Pollution* 282.
- Alghamdi MA, Alam MS, Yin J, Stark C, Jang E, Harrison RM, et al. 2015. Receptor modelling study of polycyclic aromatic hydrocarbons in jeddah, saudi arabia. *Science of the Total Environment* 506:401-408.
- Ali-Taleshi MS, Bakhtiari AR, Moeinaddini M, Squizzato S, Feiznia S, Cesari D. 2021a. Single-site source apportionment modeling of pm_{2.5}-bound pahs in the tehran metropolitan area, iran: Implications for source-specific multi-pathway cancer risk assessment. *Urban Climate* 39:100928.
- Ali-Taleshi MS, Moeinaddini M, Bakhtiari AR, Feiznia S, Squizzato S, Bourliva A. 2021b. A one-year monitoring of spatiotemporal variations of pm_{2.5}-bound pahs in tehran, iran: Source apportionment,

local and regional sources origins and source-specific cancer risk assessment. *Environmental Pollution* 274:115883.

Ali-Taleshi MS, Squizzato S, Bakhtiari AR, Moeinaddini M, Masiol M. 2021c. Using a hybrid approach to apportion potential source locations contributing to excess cancer risk of pm_{2.5} during heating and non-heating periods in a megacity in the middle east. *Environmental research* 201:111617.

Alidadi R, Mansouri N, Hemmasi A, Mirzahosseini S. 2021. Source apportionment of heavy metal with unmix in ambient air of ahvaz city, southwest of iran. *International Journal of Environmental Science and Technology*:1-8.

Almeida S, Manousakas M, Diapouli E, Kertesz Z, Samek L, Hristova E, et al. 2020. Ambient particulate matter source apportionment using receptor modelling in european and central asia urban areas. *Environmental Pollution* 266:115199.

Alolayan MA, Brown KW, Evans JS, Bouhamra WS, Koutrakis P. 2013. Source apportionment of fine particles in kuwait city. *Science of the Total Environment* 448:14-25.

Altuwayjiri A, Taghvaei S, Mousavi A, Sowlat MH, Hassanvand MS, Kashani H, et al. 2021. Association of systemic inflammation and coagulation biomarkers with source-specific pm_{2.5} mass concentrations among young and elderly subjects in central tehran. *Journal of the Air & Waste Management Association* 71:191-208.

Altuwayjiri A, Pirhadi M, Kalafy M, Alharbi B, Sioutas C. 2022. Impact of different sources on the oxidative potential of ambient particulate matter pm₁₀ in riyadh, saudi arabia: A focus on dust emissions. *Science of The Total Environment* 806:150590.

Alvi MU, Mahmud T, Kistler M, Kasper-Giebl A, Shahid I, Alam K, et al. 2020. Elemental composition of particulate matter in south-asian megacity (faisalabad-pakistan): Seasonal behaviors, source apportionment and health risk assessment. *Revista de Chimie (Rev Chim)* 71:288-301.

Amini H, Hosseini V, Schindler C, Hassankhany H, Yunesian M, Henderson SB, et al. 2017. Spatiotemporal description of btex volatile organic compounds in a middle eastern megacity: Tehran study of exposure prediction for environmental health research (tehran sepehr). *Environmental pollution* 226:219-229.

Amini H, Nhung NTT, Schindler C, Yunesian M, Hosseini V, Shamsipour M, et al. 2019. Short-term associations between daily mortality and ambient particulate matter, nitrogen dioxide, and the air quality index in a middle eastern megacity. *Environmental Pollution* 254:113121.

Arhami M, Hosseini V, Shahne MZ, Bigdeli M, Lai A, Schauer JJ. 2017. Seasonal trends, chemical speciation and source apportionment of fine pm in tehran. *Atmospheric environment* 153:70-82.

Arhami M, Shahne MZ, Hosseini V, Haghghat NR, Lai AM, Schauer JJ. 2018. Seasonal trends in the composition and sources of pm_{2.5} and carbonaceous aerosol in tehran, iran. *Environmental pollution* 239:69-81.

Ashrafi K, Fallah R, Hadei M, Yarahmadi M, Shahsavani A. 2018. Source apportionment of total suspended particles (tsp) by positive matrix factorization (pmf) and chemical mass balance (cmb) modeling in ahvaz, iran. *Archives of environmental contamination and toxicology* 75:278-294.

Azimi-Yancheshmeh R, Moeinaddini M, Feiznia S, Riyahi-Bakhtiari A, Savabieasfahani M, van Hullebusch ED, et al. 2021. Seasonal and spatial variations in atmospheric pm_{2.5} in karaj city, iran: Sources, distributions, and health risks. *Sustainable Cities and Society* 72:103020.

Barn P, Gombojav E, Ochir C, Boldbaatar B, Beejin B, Naidan G, et al. 2018. The effect of portable hepa filter air cleaner use during pregnancy on fetal growth: The ugaar randomized controlled trial. *Environment international* 121:981-989.

Barzeghar V, Sarbakhsh P, Hassanvand MS, Faridi S, Gholampour A. 2020. Long-term trend of ambient air pm₁₀, pm_{2.5}, and o₃ and their health effects in tabriz city, iran, during 2006–2017. *Sustainable Cities and Society* 54:101988.

Belis CA, Pernigotti D, Pirovano G, Favez O, Jaffrezo J, Kuenen J, et al. 2020. Evaluation of receptor and chemical transport models for pm₁₀ source apportionment. *Atmospheric environment: X* 5:100053.

Bennett JE, Tamura-Wicks H, Parks RM, Burnett RT, Pope III CA, Bechle MJ, et al. 2019. Particulate matter air pollution and national and county life expectancy loss in the USA: A spatiotemporal analysis. *PLoS medicine* 16:e1002856.

Bian Q, Alharbi B, Collett Jr J, Kreidenweis S, Pasha MJ. 2016. Measurements and source apportionment of particle-associated polycyclic aromatic hydrocarbons in ambient air in riyadh, saudi arabia. *Atmospheric Environment* 137:186-198.

Boman J, Shaltout AA, Abozied AM, Hassan SK. 2013. On the elemental composition of pm_{2.5} in central cairo, egypt. *X-Ray Spectrometry* 42:276-283.

Brook RD, Newby DE, Rajagopalan S. 2017. The global threat of outdoor ambient air pollution to cardiovascular health: Time for intervention. *JAMA cardiology* 2:353-354.

Contini D, Cesari D, Conte M, Donato A. 2016. Application of pmf and cmb receptor models for the evaluation of the contribution of a large coal-fired power plant to pm₁₀ concentrations. *Science of the Total Environment* 560:131-140.

Cusack M, Arrieta J, Duarte CM. 2020. Source apportionment and elemental composition of atmospheric total suspended particulates (tsp) over the red sea coast of saudi arabia. *Earth Systems and Environment* 4:777-788.

Danaei G, Farzadfar F, Kelishadi R, Rashidian A, Rouhani OM, Ahmadnia S, et al. 2019. Iran in transition. *The Lancet* 393:1984-2005.

Eivazzadeh M, Hassanvand MS, Faridi S, Gholampour A. 2021. Source apportionment and deposition of dustfall-bound trace elements around tabriz, iran. *Environmental Science and Pollution Research*:1-13.

EPA. 2015. *Pmf_5.0_user_guide*.

Esmaeilirad S, Lai A, Abbaszade G, Schnelle-Kreis J, Zimmermann R, Uzu G, et al. 2020. Source apportionment of fine particulate matter in a middle eastern metropolis, tehran-iran, using pmf with organic and inorganic markers. *Science of The Total Environment* 705:135330.

Faridi S, Naddafi K, Kashani H, Nabizadeh R, Alimohammadi M, Momeniha F, et al. 2017. Bioaerosol exposure and circulating biomarkers in a panel of elderly subjects and healthy young adults. *Science of the Total Environment* 593:380-389.

Faridi S, Nodehi RN, Sadeghian S, Tajdini M, Hoseini M, Yunesian M, et al. 2020. Can respirator face masks in a developing country reduce exposure to ambient particulate matter? *Journal of Exposure Science & Environmental Epidemiology*:1-12.

Faridi S, Yousefian F, Janjani H, Niazi S, Azimi F, Naddafi K, et al. 2021. The effect of covid-19 pandemic on human mobility and ambient air quality around the world: A systematic review. *Urban Climate*:100888.

Faridi S, Brook RD, Yousefian F, Hassanvand MS, Nodehi RN, Shamsipour M, et al. 2022. Effects of respirators to reduce fine particulate matter exposures on blood pressure and heart rate variability: A systematic review and meta-analysis. *Environmental Pollution*:119109.

GBD. 2019. Health data.

Gholampour A, Nabizadeh R, Hassanvand MS, Taghipour H, Nazmara S, Mahvi AH. 2015. Characterization of saline dust emission resulted from urmia lake drying. *Journal of Environmental Health Science and Engineering* 13:1-11.

Gholampour A, Nabizadeh R, Hassanvand MS, Taghipour H, Rafee M, Alizadeh Z, et al. 2016. Characterization and source identification of trace elements in airborne particulates at urban and suburban atmospheres of tabriz, iran. *Environmental Science and Pollution Research* 23:1703-1713.

Goudie AS. 2014. Desert dust and human health disorders. *Environment international* 63:101-113.

Hadley MB, Baumgartner J, Vedanthan R. 2018a. Developing a clinical approach to air pollution and cardiovascular health. *Circulation* 137:725-742.

Hadley MB, Vedanthan R, Fuster V. 2018b. Air pollution and cardiovascular disease: A window of opportunity. *Nature reviews Cardiology* 15:193.

Halek F, Kianpour-Rad M, Kavousi A. 2010. Characterization and source apportionment of polycyclic aromatic hydrocarbons in the ambient air (tehran, iran). *Environmental Chemistry Letters* 8:39-44.

Hamad SH, Schauer JJ, Heo J, Kadhim AK. 2015. Source apportionment of pm_{2.5} carbonaceous aerosol in baghdad, iraq. *Atmospheric Research* 156:80-90.

Hamid N, Syed JH, Junaid M, Mahmood A, Li J, Zhang G, et al. 2018. Elucidating the urban levels, sources and health risks of polycyclic aromatic hydrocarbons (pahs) in pakistan: Implications for changing energy demand. *Science of the Total Environment* 619-620:165-175.

Harrison RM, Smith D, Piou C, Castro L. 1997. Comparative receptor modelling study of airborne particulate pollutants in birmingham (united kingdom), coimbra (portugal) and lahore (pakistan). *Atmospheric Environment* 31:3309-3321.

Heger M, Sarraf M. 2018. Air pollution in tehran: Health costs, sources, and policies:World Bank.

Heo J, Wu B, Abdeen Z, Qasrawi R, Sarnat JA, Sharf G, et al. 2017. Source apportionments of ambient fine particulate matter in israeli, jordanian, and palestinian cities. *Environmental Pollution* 225:1-11.

Hopke PK, Dai Q, Li L, Feng Y. 2020. Global review of recent source apportionments for airborne particulate matter. *Science of The Total Environment* 740:140091.

Hopke PK, Jaffe DA. 2020. Letter to the editor: Ending the use of obsolete data analysis methods. *Aerosol and Air Quality Research* 20:688-689.

Hopke PK, Feng Y, Dai Q. 2022. Source apportionment of particle number concentrations: A global review. *Science of The Total Environment*:153104.

Hosseini V, Shahbazi H. 2016. Urban air pollution in iran. *Iranian Studies* 49:1029-1046.

Ishtiaq J, Syed JH, Jadoon WA, Hamid N, Iqbal Chaudhry MJ, Shahnawaz M, et al. 2021. Atmospheric polycyclic aromatic hydrocarbons (pahs) at urban settings in pakistan: Spatial variations, sources and health risks. *Chemosphere* 274.

Jaafar W, Zaherddine V, Hussein F, Saliba NA, Hayeck N. 2021. Poor regulation implications in a low and middle income country based on pah source apportionment and cancer risk assessment. *Environmental Science-Processes & Impacts* 23:1986-1996.

Javed W, Iakovides M, Garaga R, Stephanou EG, Kota SH, Ying Q, et al. 2019. Source apportionment of organic pollutants in fine and coarse atmospheric particles in doha, qatar. *Journal of the Air & Waste Management Association* 69:1277-1292.

Javed W, Guo B. 2021. Chemical characterization and source apportionment of fine and coarse atmospheric particulate matter in doha, qatar. *Atmospheric Pollution Research* 12:122-136.

Karagulian F, Belis CA, Dora CFC, Prüss-Ustün AM, Bonjour S, Adair-Rohani H, et al. 2015. Contributions to cities' ambient particulate matter (pm): A systematic review of local source contributions at global level. *Atmospheric environment* 120:475-483.

Karim MM, Nukala, V.,. 2020. *Emission_inventory_and_air_quality_dispersion_modeling_of_kabul_city*.

Kermani M, Asadgol Z, Gholami M, Jafari AJ, Shahsavani A, Goodarzi B, et al. 2021a. Occurrence, spatial distribution, seasonal variations, potential sources, and inhalation-based health risk assessment of organic/inorganic pollutants in ambient air of tehran. *Environmental Geochemistry and Health* 43:1983-2006.

Kermani M, Jafari AJ, Gholami M, Arfaeinia H, Shahsavani A, Fanaei F. 2021b. Characterization, possible sources and health risk assessment of pm_{2.5}-bound heavy metals in the most industrial city of iran. *Journal of Environmental Health Science and Engineering*:1-13.

Khairy MA, Lohmann R. 2013. Source apportionment and risk assessment of polycyclic aromatic hydrocarbons in the atmospheric environment of alexandria, egypt. *Chemosphere* 91:895-903.

Khanum F, Chaudhry MN, Skouteris G, Saroj D, Kumar P. 2021. Chemical composition and source characterization of pm₁₀ in urban areas of lahore, pakistan. *Indoor Built Environ* 30:924-937.

Khodeir M, Shamy M, Alghamdi M, Zhong M, Sun H, Costa M, et al. 2012. Source apportionment and elemental composition of pm_{2.5} and pm₁₀ in jeddah city, saudi arabia. *Atmospheric pollution research* 3:331-340.

Kholdebarin A, Biati A, Moattar F, Shariat SM. 2015. Outdoor pm₁₀ source apportionment in metropolitan cities—a case study. *Environmental monitoring and assessment* 187:1-12.

Lelieveld J, Evans JS, Fnais M, Giannadaki D, Pozzer A. 2015. The contribution of outdoor air pollution sources to premature mortality on a global scale. *Nature* 525:367-371.

Lodhi A, Ghauri B, Khan MR, Rahman S, Shafique S. 2009. Particulate matter (pm_{2.5}) concentration and source apportionment in lahore. *Journal of the Brazilian Chemical Society* 20:1811-1820.

Lurie K, Nayebare SR, Fatmi Z, Carpenter DO, Siddique A, Malashock D, et al. 2019. Pm_{2.5} in a megacity of asia (karachi): Source apportionment and health effects. *Atmospheric Environment* 202:223-233.

Maleki R, Asadgol Z, Kermani M, Jafari AJ, Arfaeinia H, Ghodsi S, et al. 2021. Concentration, sources, and inhalation-based risk assessment of pm_{2.5}-bound pahs and trace elements in ambient air of areas with low and high traffic density in tehran. *Arabian Journal of Geosciences* 14:1-19.

Mansha M, Ghauri B, Rahman S, Amman A. 2012. Characterization and source apportionment of ambient air particulate matter (pm_{2.5}) in karachi. *Science of the total environment* 425:176-183.

McDuffie EE, Martin RV, Spadaro JV, Burnett R, Smith SJ, O'Rourke P, et al. 2021. Source sector and fuel contributions to ambient pm_{2.5} and attributable mortality across multiple spatial scales. *Nature Communications* 12:1-12.

Mehmood T, Zhu T, Ahmad I, Li X. 2020. Ambient pm_{2.5} and pm₁₀ bound pahs in islamabad, pakistan: Concentration, source and health risk assessment. *Chemosphere* 257:127187.

Moeinaddini M, Sari AE, Chan AY-C, Taghavi SM, Hawker D, Connell D. 2014. Source apportionment of pahs and n-alkanes in respirable particles in tehran, iran by wind sector and vertical profile. *Environmental science and pollution research* 21:7757-7772.

Mousavi A, Sowlat MH, Hasheminassab S, Polidori A, Sioutas C. 2018. Spatio-temporal trends and source apportionment of fossil fuel and biomass burning black carbon (bc) in the los angeles basin. *Science of the Total Environment* 640:1231-1240.

Münzel T, Sørensen M, Lelieveld J, Hahad O, Al-Kindi S, Nieuwenhuijsen M, et al. 2021. Heart healthy cities: Genetics loads the gun but the environment pulls the trigger. *European Heart Journal*.

Najmeddin A, Keshavarzi B. 2019. Health risk assessment and source apportionment of polycyclic aromatic hydrocarbons associated with pm₁₀ and road deposited dust in ahvaz metropolis of iran. *Environmental geochemistry and health* 41:1267-1290.

Nayebare SR, Aburizaiza OS, Khwaja HA, Siddique A, Hussain MM, Zeb J, et al. 2016. Chemical characterization and source apportionment of pm_{2.5} in rabigh, saudi arabia. *Aerosol and Air Quality Research* 16:3114-3129.

Nayebare SR, Aburizaiza OS, Siddique A, Carpenter DO, Hussain MM, Zeb J, et al. 2018. Ambient air quality in the holy city of makkah: A source apportionment with elemental enrichment factors (efs) and factor analysis (pmf). *Environmental Pollution* 243:1791-1801.

Paatero P, Eberly S, Brown S, Norris G. 2014. Methods for estimating uncertainty in factor analytic solutions. *Atmospheric Measurement Techniques* 7:781-797.

Querol X, Tobías A, Pérez N, Karanasiou A, Amato F, Stafoggia M, et al. 2019. Monitoring the impact of desert dust outbreaks for air quality for health studies. *Environment international* 130:104867.

Raja S, Biswas KF, Husain L, Hopke PK. 2010. Source apportionment of the atmospheric aerosol in lahore, pakistan. *Water, air, and soil pollution* 208:43-57.

Rajagopalan S, Al-Kindi SG, Brook RD. 2018. Air pollution and cardiovascular disease: Jacc state-of-the-art review. *Journal of the American College of Cardiology* 72:2054-2070.

Rajagopalan S, Landrigan PJ. 2021. Pollution and the heart. *New England Journal of Medicine* 385:1881-1892.

Saliba N, El Jam F, El Tayar G, Obeid W, Roumie M. 2010. Origin and variability of particulate matter (pm₁₀ and pm_{2.5}) mass concentrations over an eastern mediterranean city. *Atmospheric Research* 97:106-114.

Shah MH, Shaheen N, Nazir R. 2012. Assessment of the trace elements level in urban atmospheric particulate matter and source apportionment in islamabad, pakistan. *Atmospheric pollution research* 3:39-45.

Shahbazi H, Reyhanian M, Hosseini V, Afshin H. 2016a. The relative contributions of mobile sources to air pollutant emissions in tehran, iran: An emission inventory approach. *Emission control science and technology* 2:44-56.

Shahbazi H, Taghvaei S, Hosseini V, Afshin H. 2016b. A gis based emission inventory development for tehran. *Urban Climate* 17:216-229.

Shahbazi H, Hassani A, Hosseini V. 2019. Evaluation of tehran clean air action plan using emission inventory approach. *Urban Climate* 27:446-456.

Shahbazi H, Abolmaali AM, Alizadeh H, Salavati H, Zokaei H, Zandavi R, et al. 2021. Development of high-resolution emission inventory to study the relative contribution of a local power plant to criteria air pollutants and greenhouse gases. *Urban Climate* 38:100897.

Shaheen N, Shah MH, Jaffar M. 2005. A study of airborne selected metals and particle size distribution in relation to climatic variables and their source identification. *Water, air, and soil pollution* 164:275-294.

Shahid I, Alvi MU, Shahid MZ, Alam K, Chishtie F. 2018. Source apportionment of pm₁₀ at an urban site of a south asian mega city. *Aerosol and Air Quality Research* 18:2498-2509.

Shahsavani A, Tobías A, Querol X, Stafoggia M, Abdolshahnejad M, Mayvaneh F, et al. 2020. Short-term effects of particulate matter during desert and non-desert dust days on mortality in iran. *Environment international* 134:105299.

Shahsavani S, Hoseini M, Dehghani M, Fararouei M. 2017. Characterisation and potential source identification of polycyclic aromatic hydrocarbons in atmospheric particles (pm₁₀) from urban and suburban residential areas in shiraz, iran. *Chemosphere* 183:557-564.

Shaltout AA, Boman J, Alsulimane ME. 2017. Identification of elemental composition of pm_{2.5} collected in makkah, saudi arabia, using edxrf. *X-Ray Spectrometry* 46:151-163.

Shaltout AA, Boman J, Hassan SK, Abozied AM, Al-Ashkar EA, Abd-Elkader OH, et al. 2020. Elemental composition of pm_{2.5} aerosol in a residential–industrial area of a mediterranean megacity. *Archives of environmental contamination and toxicology* 78:68-78.

Shamsipour M, Hassanvand MS, Gohari K, Yunesian M, Fotouhi A, Naddafi K, et al. 2019. National and sub-national exposure to ambient fine particulate matter (pm_{2.5}) and its attributable burden of disease in iran from 1990 to 2016. *Environmental Pollution* 255:113173.

Singh N, Murari V, Kumar M, Barman S, Banerjee T. 2017. Fine particulates over south asia: Review and meta-analysis of pm_{2.5} source apportionment through receptor model. *Environmental Pollution* 223:121-136.

Soleimani M, Amini N, Sadeghian B, Wang D, Fang L. 2018. Heavy metals and their source identification in particulate matter (pm_{2.5}) in isfahan city, iran. *Journal of environmental sciences* 72:166-175.

Soleimani M, Ebrahimi Z, Mirghaffari N, Moradi H, Amini N, Pulsen KG, et al. 2021. Seasonal trend and source identification of polycyclic aromatic hydrocarbons associated with fine particulate matters (pm_{2.5}) in isfahan city, iran using diagnostic ratio and pmf model.

Soleimani E, Taghvaei S, Mousavi A, Sowlat MH, Hassanvand MS, Yunesian M, et al. 2019. Sources and temporal variations of coarse particulate matter (pm) in central tehran, iran. *Atmosphere* 10:291.

Southerland VA, Brauer M, Mohegh A, Hammer MS, van Donkelaar A, Martin RV, et al. 2022. Global urban temporal trends in fine particulate matter (pm_{2.5}) and attributable health burdens: Estimates from global datasets. *The Lancet Planetary Health*.

Sowlat MH, Naddafi K, Yunesian M, Jackson PL, Shahsavani A. 2012. Source apportionment of total suspended particulates in an arid area in southwestern iran using positive matrix factorization. *Bulletin of environmental contamination and toxicology* 88:735-740.

Sowlat MH, Naddafi K, Yunesian M, Jackson PL, Lotfi S, Shahsavani A. 2013. Pm10 source apportionment in ahvaz, iran, using positive matrix factorization. *Clean—soil, air, water* 41:1143-1151.

Sowlat MH. 2019. Physico-chemical characteristics and sources of ambient pm mass and number concentrations and their associated toxicity, and development of novel techniques for high time-resolution measurement of pm-bound metals for application in source apportionment studies:University of Southern California.

Taghvae S, Sowlat MH, Hassanvand MS, Yunesian M, Naddafi K, Sioutas C. 2018a. Source-specific lung cancer risk assessment of ambient pm2. 5-bound polycyclic aromatic hydrocarbons (pahs) in central tehran. *Environment international* 120:321-332.

Taghvae S, Sowlat MH, Mousavi A, Hassanvand MS, Yunesian M, Naddafi K, et al. 2018b. Source apportionment of ambient pm2. 5 in two locations in central tehran using the positive matrix factorization (pmf) model. *Science of The Total Environment* 628:672-686.

Tahri M, Bounakhla M, Zghaid M, Benchrif A, Zahry F, Noack Y, et al. 2013. Txf characterization and source identification by positive matrix factorization of airborne particulate matter sampled in kenitra city (morocco). *X-Ray Spectrometry* 42:284-289.

Thunis P, Clappier A, Tarrasón L, Cuvelier C, Monteiro A, Pisoni E, et al. 2019. Source apportionment to support air quality planning: Strengths and weaknesses of existing approaches. *Environment international* 130:104825.

Tian Y, Liu H, Wu Y, Si Y, Li M, Wu Y, et al. 2019. Ambient particulate matter pollution and adult hospital admissions for pneumonia in urban china: A national time series analysis for 2014 through 2017. *PLoS medicine* 16:e1003010.

von Schneidemesser E, Zhou J, Stone EA, Schauer JJ, Qasrawi R, Abdeen Z, et al. 2010. Seasonal and spatial trends in the sources of fine particle organic carbon in israel, jordan, and palestine. *Atmospheric Environment* 44:3669-3678.

Waked A, Afif C, Seigneur C. 2012. An atmospheric emission inventory of anthropogenic and biogenic sources for lebanon. *Atmospheric environment* 50:88-96.

Waked A, Afif C, Seigneur C. 2015. Assessment of source contributions to air pollution in beirut, lebanon: A comparison of source-based and tracer-based modeling approaches. *Air Qual Atmos Health* 8:495-505.

Supplemental file

Source Apportionment, Identification and Characterization, and Emission Inventory of Ambient Particulate Matter in 22 Eastern Mediterranean Region Countries: A Systematic Review and Recommendations for Good Practice[★]

Sasan Faridi ^{a, b}, Fatemeh Yousefian ^c, Vahid Roostaei ^b, Roy M. Harrison ^{d, e}, Faramarz Azimi ^f, Sadegh Niazi ^g, Kazem Naddafi ^{a, b}, Fatemeh Momeniha ^h, Mazen Malkawi ⁱ, Heba Adel Moh'd Safi ⁱ, Mona Khaleghy Rad ⁱ, Mohammad Sadegh Hassanvand ^{a, b*}

^a Center for Air Pollution Research (CAPR), Institute for Environmental Research (IER), Tehran University of Medical Sciences, Tehran, Iran

^b Department of Environmental Health Engineering, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran

^c Department of Environmental Health Engineering, Faculty of Health, Kashan University of Medical Sciences, Kashan, Iran

^d School of Geography Earth and Environmental Science, University of Birmingham, Birmingham, UK

^e Department of Environmental Sciences, Faculty of Meteorology, Environment and Arid Land Agriculture, King Abdulaziz University, Jeddah, Saudi Arabia

^f Environmental Health Research Center, Lorestan University of Medical Sciences, Khorramabad, Iran

^g International Laboratory for Air Quality and Health, School of Earth and Atmospheric Sciences, Science and Engineering Faculty, Queensland University of Technology (QUT), Brisbane, Queensland, Australia

^h Center for Solid Waste Research, Institute for Environmental Research, Tehran University of Medical Sciences, Tehran, Iran

ⁱ Environmental Health Exposures Centre for Environmental Health Action (CEHA), World Health Organization (WHO), Jordan

*** Corresponding Author:** MS. Hassanvand, PhD, Center for Air Pollution Research (CAPR), Institute for Environmental Research (IER) and Department of Environmental Health Engineering, School of Public Health, Tehran University of Medical Sciences, Phone: +98 88978395, Fax: +98 88978397, 8th Floor, No. 1547, North Kargar Avenue, Tehran, Iran (hassanvand@tums.ac.ir).

★ This study is a part of a consultancy to provide a background assessment of air quality (AQ) management system for the Eastern Mediterranean countries in order to improve their capabilities to use the updated World Health Organization Air Quality Guidelines in compiling national AQ standards.

Table S1. Full search strategy (components and the results of search) in PubMed, Scopus, and Web of Science for the source apportionment, characterization and identification studies.

Components		Keywords
Countries		Afghanistan Bahrain Djibouti Egypt Islamic Republic of Iran (Iran) Iraq Jordan Kuwait Lebanon Libya Morocco Occupied Palestinian territory (Palestine) Oman Pakistan Qatar Saudi Arabia Somali Sudan Syrian Arab Republic United Arab Emirates Tunisia Yemen
Air Quality		Air Pollution Air Pollutant Particulate Matter PM ₁₀ PM _{2.5}
Approaches		Source Apportionment Source Identification Source characterization
Databases	Results	
Scopus: 196	((TITLE-ABS-KEY (<i>afghanistan</i>) OR TITLE-ABS-KEY (<i>bahrain</i>) OR TITLE-ABS-KEY (<i>diibouti</i>) OR TITLE-ABS-KEY (<i>egypt</i>) OR TITLE-ABS-KEY (" <i>Islamic Republic of Iran</i> ") OR TITLE-ABS-KEY (<i>iran</i>) OR TITLE-ABS-KEY (<i>iraq</i>) OR TITLE-ABS-KEY (<i>jordan</i>) OR TITLE-ABS-KEY (<i>kuwait</i>) OR TITLE-ABS-KEY (<i>lebanon</i>) OR TITLE-ABS-KEY (<i>libya</i>) OR TITLE-ABS-KEY (<i>morocco</i>) OR TITLE-ABS-KEY (" <i>Occupied Palestinian territory</i> ") OR TITLE-ABS-KEY (<i>palestinian</i>) OR TITLE-ABS-KEY (<i>oman</i>) OR TITLE-ABS-KEY (<i>pakistan</i>) OR TITLE-ABS-KEY (<i>qatar</i>) OR TITLE-ABS-KEY (" <i>Saudi Arabia</i> ") OR TITLE-ABS-KEY (<i>somali</i>) OR TITLE-ABS-KEY (<i>sudan</i>) OR TITLE-ABS-KEY (" <i>Syrian Arab Republic</i> ") OR TITLE-ABS-KEY (" <i>United Arab Emirates</i> ") OR TITLE-ABS-KEY (<i>tunisia</i>) OR TITLE-ABS-KEY (<i>yemen</i>))) AND ((TITLE-ABS-KEY (" <i>air pollution</i> ") OR TITLE-ABS-KEY (" <i>air pollutant*</i> ") OR TITLE-ABS-KEY (" <i>particulate matter</i> ") OR TITLE-ABS-KEY (<i>pm2.5</i>) OR TITLE-ABS-KEY (<i>pm10</i>))) AND ((TITLE-ABS-	

	KEY ("Source identification") OR TITLE-ABS-KEY ("Source characterization") OR TITLE-ABS-KEY ("Source apportionment")) AND (LIMIT-TO (DOCTYPE , "ar") OR LIMIT-TO (DOCTYPE , "re")) AND (LIMIT-TO (LANGUAGE , "English"))
PubMed: 50	((("air pollution"[MeSH Terms] OR ("air pollution"[Title/Abstract] OR "air pollutant"[Title/Abstract] OR "particulate matter"[Title/Abstract] OR "PM2.5"[Title/Abstract] OR "PM10"[Title/Abstract]))) AND (((("Iran"[Title/Abstract] OR "Islamic Republic of Iran"[Title/Abstract] OR "Afghanistan"[Title/Abstract] OR "Bahrain"[Title/Abstract] OR "Djibouti"[Title/Abstract] OR "Egypt"[Title/Abstract] OR "Iraq"[Title/Abstract] OR "Jordan"[Title/Abstract] OR "Kuwait"[Title/Abstract] OR ("Lebanon"[Title/Abstract] OR "Libya"[Title/Abstract] OR "Morocco"[Title/Abstract] OR "Occupied Palestinian territory"[Title/Abstract] OR "Palestinian"[Title/Abstract] OR "Oman"[Title/Abstract] OR "Pakistan"[Title/Abstract] OR "Qatar"[Title/Abstract] OR ("Saudi Arabia"[Title/Abstract] OR "Somali"[Title/Abstract] OR "Sudan"[Title/Abstract] OR "Syrian Arab Republic"[Title/Abstract] OR "United Arab Emirates"[Title/Abstract] OR "Tunisia"[Title/Abstract] OR "Yemen"[Title/Abstract]))) AND (((("Source Apportionment"[Title/Abstract] OR ("Source Identification"[Title/Abstract] OR ("Source characterization"[Title/Abstract]))
WOS: 87	((((TI=(Afghanistan) OR AB=(Afghanistan) OR TI=(Bahrain) OR AB=(Bahrain) OR TI=(Djibouti) OR AB=(Djibouti) OR TI=(Egypt) OR AB=(Egypt) OR TI=("Islamic Republic of Iran") OR AB=("Islamic Republic of Iran") OR TI=(Iran) OR AB=(Iran) OR TI=(Iraq) OR AB=(Iraq) OR TI=(Jordan) OR AB=(Jordan) OR TI=(Kuwait) OR AB=(Kuwait) OR TI=(Lebanon) OR AB=(Lebanon) OR TI=(Libya) OR AB=(Libya) OR TI=(Morocco) OR AB=(Morocco) OR TI=("Occupied Palestinian territory") OR AB=("Occupied Palestinian territory") OR TI=(Palestine) OR AB=(Palestine) OR TI=(Oman) OR AB=(Oman) OR TI=(Pakistan) OR AB=(Pakistan) OR TI=(Qatar) OR AB=(Qatar) OR TI=("Saudi Arabia") OR AB=("Saudi Arabia") OR TI=(Somali) OR AB=(Somali) OR TI=(Sudan) OR AB=(Sudan) OR TI=("Syrian Arab Republic") OR AB=("Syrian Arab Republic") OR TI=("United Arab Emirates") OR AB=("United Arab Emirates") OR TI=(Tunisia) OR AB=(Tunisia) OR TI=(Yemen) OR AB=(Yemen)))) AND (TI=("air pollution") OR AB=("air pollution") OR TI=("Air pollutant") OR AB=("Air pollutant") OR TI=("particulate matter") OR AB=("particulate matter") OR TI=(PM2.5) OR AB=(PM2.5) OR TI=(PM10) OR AB=(PM10) AND) TI=("Source Apportionment") OR AB=("Source Apportionment") OR (TI="Source Identification") OR AB=("Source Identification") OR (TI="Source Characterization") OR AB=("Source Characterization") (AND LANGUAGE: (English)

Table S2. Full search strategy (components and the results of search) in PubMed, Scopus, and Web of Science for the emission inventory studies.

Components		Keywords
Countries		Afghanistan Bahrain Djibouti Egypt Islamic Republic of Iran (Iran) Iraq Jordan Kuwait Lebanon Libya Morocco Occupied Palestinian territory (Palestine) Oman Pakistan Qatar Saudi Arabia Somali Sudan Syrian Arab Republic United Arab Emirates Tunisia Yemen
Air Quality		Air Pollution Air Pollutant Particulate Matter PM ₁₀ PM _{2.5}
Approach		Emission Inventory
Databases	Results	
Scopus: 82	(TITLE-ABS-KEY ("Emission Inventory")) AND ((TITLE-ABS-KEY ("air pollution") OR TITLE-ABS-KEY ("air pollutant") OR TITLE-ABS-KEY ("air pollutatns"))) AND ((TITLE-ABS-KEY (bahrain) OR TITLE-ABS-KEY (afghanistan) OR TITLE-ABS-KEY (djibouti) OR TITLE-ABS-KEY (egypt) OR TITLE-ABS-KEY ("Islamic Republic of Iran") OR TITLE-ABS-KEY (iran) OR TITLE-ABS-KEY (iraq) OR TITLE-ABS-KEY (jordan) OR TITLE-ABS-KEY (kuwait) OR TITLE-ABS-KEY (lebanon) OR TITLE-ABS-KEY (libya) OR TITLE-ABS-KEY (morocco) OR TITLE-ABS-KEY ("Occupied Palestinian territory") OR TITLE-ABS-KEY (palestinian) OR TITLE-ABS-KEY (oman) OR TITLE-ABS-KEY (pakistan) OR TITLE-ABS-KEY (qatar) OR TITLE-ABS-KEY ("Saudi Arabia") OR TITLE-ABS-KEY (somali) OR TITLE-ABS-KEY (sudan) OR TITLE-ABS-KEY ("Syrian Arab Republic") OR TITLE-ABS-KEY ("United Arab Emirates") OR TITLE-ABS-KEY (tunisia) OR TITLE-ABS-KEY (yemen))) AND (LIMIT-TO (LANGUAGE , "English")))	

<p>PubMed: 8</p>	<p>((("air pollution"[Title/Abstract] OR "air pollutant"[Title/Abstract] OR "air pollutants"[Title/Abstract]) AND "Iran"[Title/Abstract]) OR "i r iran"[Affiliation] OR "i r iran"[Title/Abstract] OR "i r iran"[Title/Abstract] OR "IR Iran"[Title/Abstract] OR "IR Iran"[Title/Abstract] OR "i r iran"[Title/Abstract] OR "Islamic Republic of Iran"[Affiliation] OR "Islamic Republic of Iran"[Title/Abstract] OR "iran islamic republic of"[Title/Abstract] OR "iran islamic republic of"[Title/Abstract] OR "iran islamic republic"[Title/Abstract] OR "Iranian"[Title/Abstract] OR "Iranians"[Title/Abstract] OR "persia*"[Title/Abstract] OR "Afghanistan"[Title/Abstract] OR "Bahrain"[Title/Abstract] OR "Djibouti"[Title/Abstract] OR "Egypt"[Title/Abstract] OR "Iraq"[Title/Abstract] OR "Jordan"[Title/Abstract] OR "Kuwait"[Title/Abstract] OR "Lebanon"[Title/Abstract] OR "Libya"[Title/Abstract] OR "Morocco"[Title/Abstract] OR "Occupied Palestinian territory"[Title/Abstract] OR "Palestinian"[Title/Abstract] OR "Oman"[Title/Abstract] OR "Pakistan"[Title/Abstract] OR "Qatar"[Title/Abstract] OR "Saudi Arabia"[Title/Abstract] OR "Somali"[Title/Abstract] OR "Sudan"[Title/Abstract] OR "Syrian Arab Republic"[Title/Abstract] OR "United Arab Emirates"[Title/Abstract] OR "Tunisia"[Title/Abstract] OR "Yemen"[Title/Abstract]) AND "Emission Inventory"[Title/Abstract]</p>
<p>WOS: 17</p>	<p>((TI=("Emission Inventory") OR AB=("Emission Inventory")) AND ((TI=(Afghanistan) OR AB=(Afghanistan) OR TI=(Bahrain) OR AB=(Bahrain) OR TI=(Djibouti) OR AB=(Djibouti) OR TI=(Egypt) OR AB=(Egypt) OR TI=("Islamic Republic of Iran") OR AB=("Islamic Republic of Iran") OR TI=(Iran) OR AB=(Iran) OR TI=(Iraq) OR AB=(Iraq) OR TI=(Jordan) OR AB=(Jordan) OR TI=(Kuwait) OR AB=(Kuwait) OR TI=(Lebanon) OR AB=(Lebanon) OR TI=(Libya) OR AB=(Libya) OR TI=(Morocco) OR AB=(Morocco) OR TI=("Occupied Palestinian territory") OR AB=("Occupied Palestinian territory") OR TI=(Palestine) OR AB=(Palestine) OR TI=(Oman) OR AB=(Oman) OR TI=(Pakistan) OR AB=(Pakistan) OR TI=(Qatar) OR AB=(Qatar) OR TI=("Saudi Arabia") OR AB=("Saudi Arabia") OR TI=(Somali) OR AB=(Somali) OR TI=(Sudan) OR AB=(Sudan) OR TI=("Syrian Arab Republic") OR AB=("Syrian Arab Republic") OR TI=("United Arab Emirates") OR AB=("United Arab Emirates") OR TI=(Tunisia) OR AB=(Tunisia) OR TI=(Yemen) OR AB=(Yemen)))) AND (((TI=("Air pollution") OR AB=("Air pollution") OR TI=("Air pollutant") OR AB=("Air pollutant") OR TI=("Air pollutants") OR AB=("Air pollutants")))) AND LANGUAGE: (English) Indexes=SCI-EXPANDED, SSCI Timespan=All years</p>

Table S3. Detailed information concerning the emission inventory studies for ambient air pollution in the EMR countries.

Study ID	Country/City	Applied method	Air pollutants	The relative contribution (%) of sources
(Hosseini and Shahbazi, 2016; Shahbazi et al., 2019; Shahbazi et al., 2016a; Shahbazi et al., 2016b)	Iran/ Tehran	The International Vehicle Emissions (IVE) model	PM	Mobile: 70% (Passenger Cars: 2%; Taxi: 1%; Minibus: 7%; Tehran Municipal Bus: 24%; Private Sector Bus: 30; Truck: 24%; Motorcycle: 12%); Industry: 7%; Household or commercial emissions: 2%; Energy conversion: 20%; Terminals: 1%
			CO	Mobile: 98% (Passenger Cars: 51%; Taxi: 21%; Pickup: 11%; Minibus: 1%; Tehran Municipal Bus: 1%; Motorcycle: 15%); Others: 2%
			SO _x	Mobile: 6% (Passenger Cars: 29%; Taxi: 3%; Pickup: 3%; Minibus: 12%; Tehran Municipal Bus: 16%; Private Sector Bus: 15%; Truck: 21%; Motorcycle: 1%); Industry: 22%; Energy conversion: 68%; Terminals: 4%
			NO _x	Mobile: 46% (Passenger Cars: 37%; Taxi: 11%; Pickup: 6%; Minibus: 7%; Tehran Municipal Bus: 11%; Private Sector Bus: 10%; Truck: 13%; Motorcycle: 5%); Household or commercial emissions: 23%; Energy conversion: 25%; Others: 2%
			VOCs	Mobile: 87% (Passenger Cars: 44%; Taxi: 12%; Pickup: 9%; Minibus: 1%; Tehran Municipal Bus: 1%; Private Sector Bus: 1%; Truck: 1%; Motorcycle: 31%); Gas Station: 10.5%; Others: 2.5%
(Shahbazi et al., 2021)	Iran/ Mashhad	The International Vehicle Emissions (IVE) model	TSP	Mobile: 46%; Industry: 20%; Household and Commercial emissions: 1%; Energy Conversion: 32%; Terminals: 1%
			PM ₁₀	Mobile: 45%; Industry: 18%; Household and Commercial emissions: 1%; Energy Conversion: 35%; Terminals: 1%
			PM _{2.5}	Mobile: 45%; Industry: 15%; Household and Commercial emissions: 1%; Energy Conversion: 37%; Terminals: 2%
			NO _x	Mobile: 24%; Industry: 11%; Household and Commercial emissions: 8%; Energy Conversion: 56%; Terminals: 1%
			SO _x	Mobile: 1%; Industry: 2%; Household and Commercial emissions: 0.2%; Energy Conversion: 97%; Terminals: 1%; Gas Station: 0.1%
			CO	Mobile: 96%; Industry: 2%; Household and Commercial emissions: 1%; Energy Conversion: 97%; Gas Station: 0.4%
			VOCs	Mobile: 82%; Industry: 1%; Household and Commercial emissions: 2%; Energy Conversion: 1%; Terminals: 1%; Gas Station: 13%
(Heger and Sarraf, 2018)	Iran/ Tehran	The International Vehicle Emissions (IVE) model	PM	Mobile: 70%; Industry: 7%; Household and Commercial emissions: 2%; Energy Conversion: 20%; Gas Station: 1%
Md. Masud Karim and Venkata Nukala ^a	Afghanistan/ Kabul	USEPA emission factors and USEPA regulatory dispersion model AERMOD	PM ₁₀	Vehicles and Dust: 52%; Fuel combustion for space heating: 31%; Brick kilns: 12%
			NO ₂	Portable generators: 19%; Residential Heating: 8%
			SO ₂	Fuel combustion for space heating: 15%; Portable generators: 8%
			CO	Fuel combustion for space heating: 39%; Gasoline Vehicles: 36%
(Ettouney et al., 2009)	Kuwait/ Jahara	Industrial source complex short term model (ISCST)	SO ₂	Mobile: 4%; Oil field: 34%; Power plants: 61%; Oil Refineries: 1%
			CO	Mobile: 92%; Oil field: 8%;
			NO _x	Mobile: 27.5%; Oil field: 21%; Power plants: 0.9%; Oil Refineries: 50.6%
(Khan and Ahmad, 2018)	Pakistan/ Khyber Pakhtunkhwa and Balochistan regions	The greenhouse gas and air pollution interactions and synergies (GAINS) model	SO ₂	Industry: 7%; Power and heating sector: 63%; Domestic sector: 13%
(Khatami et al., 1998)	Morocco/ Casablanca	Not reported	SO ₂	Industry: 94%; Transport: 3.1%; Ternary ^b : 1.7%; Ocean: 1.2%
			NO _x	Industry: 74.3%; Transport: 23.7%; Ternary: 2%
			CO	Industry: 62.7%; Transport: 37%; Ternary: 0.3%

Study ID	Country/City	Applied method	Air pollutants	The relative contribution (%) of sources
(Waked et al., 2012)	Lebanon/ Beirut and its 26 sub-regions	Emission factors and activities from EMEP/EEA air pollutant emission inventory guidebook ^c used in this work	CO	Mobile: 93%; Industry: 3%; Forest: 1%
			NO _x	Mobile: 52%; Industry: 23%; Domestic sector: 12%; Forest: 1%
			SO ₂	Mobile: 11%; Industry: 73%; Forest: 1%
			PM ₁₀	Mobile: 11%; Industry: 62%; Domestic sector: 15%; Forest: 4%
			PM _{2.5}	Mobile: 11%; Industry: 59%; Domestic sector: 20%; Forest: 6%

^a<https://www.researchgate.net/publication/341879689> Emission Inventory and Air Quality Dispersion Modeling of Kabul City

^b Emissions from the public and private administrations and services (hotels, hospitals, public offices, banks, trade offices, etc.)

^c<https://www.eea.europa.eu/publications/emep-eea-emission-inventory-guidebook-2009>

Table S4. Summary of information on the previously conducted systematic review studies at global scale in comparison to our study.

Study ID	Title	Summary on the approach used to identify studies	EMR countries (cities, and # of studies) reviewed in the studies	EMR countries (cities, and # of studies) reviewed in our systematic review study	Comments
(Karagulian et al., 2015)	Contributions to cities' ambient particulate matter (PM): A systematic review of local source contributions at global level	Systematic Scopus and Google searches were performed to retrieve city studies of source apportionment for particulate matter. The research covered the period between 1990 and 2014 (year of publication).	Pakistan (Karachi and Lahore, and 3 studies), Iran (Ahvaz, and 1 study), Saudi Arabia (Jeddah, and 1 study), Egypt (Cairo, and 1 study), Kuwait (Kuwait, and 1 study)	Iran (Tehran, Shiraz, Karaj, Urmia, Bushehr, Tabriz, Isfahan, Ahvaz, and Asaluyeh, and 30 studies), Saudi Arabia (Riyadh, Makkah, Jeddah, and Rabigh, and 8 studies), Pakistan (Karachi, Lahore, Islamabad, Faisalabad, Rawalpindi, Multan, Quetta, Peshawar and Gilgit, and 15 studies), Egypt (Cairo and Alexandria, and 5 studies), Kuwait (Kuwait city, Ali Sabah Al-Salem, Fahaheel, and Kuwait, and 3 studies), Lebanon (Beirut, and 3 studies), Qatar (Doha, and 2 studies), Jordan (Amman, Aqaba, and Rachma, 2 studies), and Palestine (East Jerusalem, and Hebron, and 2 studies), Iraq (Baghdad, and 1 study) and Tunisia (Tunis, and 1 study)	As stated the findings of our study on the name of countries and cities under investigation as well as the number of studies in comparison to previous works reported in the two previous columns, we have reviewed and reported not only the included studies in these reviews but also additional studies in detail.
(Hopke et al., 2020)	Global review of recent source apportionments for airborne particulate matter	Searches were performed using the Web of Science, SCOPUS, and Google Scholar using the relevant keywords to identify the published papers since 2014 through December 2019	Iran (Tehran, Ahvaz and Asaluyeh County, and 7 studies), Pakistan (Karachi and Lahore, and 3 studies), Saudi Arabia (Makkah and Rabigh, and 3 studies), Jordan (Amman, Aqaba and Rachma, and 1 study), Palestine (East Jerusalem and Hebron, and 1 study), Qatar (Doha, and 1 study), Iraq (Baghdad, and 1 study) and Tunisia (Tunis, and 1 study)		
(Hopke et al., 2022)	Source apportionment of particle number concentrations: A global review	Searches were performed for papers published prior to August 1, 2021 using the Web of Science, SCOPUS, and Google Scholar with the relevant keywords.	Kuwait (Fahaheel, and 1 study)		

Table S5. Detailed information on the contribution of each of sources to ambient PM_{2.5} air pollution in all EMR countries based on the study of (McDuffie et al., 2021).


Country Name	AFCID Dust (%)	Windblown Dust (%)	Agriculture (%)	Energy Coal (%)	Energy NonCoal (%)	Industry Coal (%)	Industry NonCoal (%)	NonRoad Transport (%)	Road Transport (%)	Others *	Residential Coal Combustion (%)	Residential Biofuel Combustion (%)	Residential Other Combustion (%)	Commercial Combustion (%)	Other Combustion (%)	Waste (%)	International Shipping (%)	Agricultural Waste Burning (%)	Other Open Fire (%)	Remaining Sources (%)	Solvent (%)
Afghanistan	5.1	51.8	5.2	6.6	5	2.2	2.7	0.1	6.2	15.1	0	7.3	0.2	0.2	0.1	3.9	0.1	0	0.5	2.8	-0.2
Bahrain	1.3	55.7	2.9	1.1	22.9	0.4	5.7	0	6.9	3.1	0.1	0.4	0.1	0.1	0.1	1.1	0.1	0	0	1.1	-0.1
Egypt	11.8	42	5.6	5.4	6.3	0.6	4.7	0.5	5.8	17.3	0.2	3.5	0.3	0.3	0.5	5.8	1.9	0.3	0.1	4.7	-0.2
Iran	3.8	52.9	8.1	1.7	9.8	0.4	5.2	0.1	11	7	0.1	1	0.4	0.4	0.6	2.6	0.3	0.2	0.5	0.9	0
Iraq	4.3	56.6	5.1	3.4	11.1	0.6	3.7	0.1	9	6.1	0.2	0.7	0.3	0.3	0.3	3.8	0.3	0.2	0.2		-0.1
Jordan	4.6	44	8.9	8.9	9.4	1	4.4	0.3	6.8	11.7	0.3	1.5	0.2	0.5	0.3	3.4	0.9	0.2	0.1	4.6	-0.2
Kuwait	1.6	66.2	3.5	1.5	15.2	0.3	3.6	0	6.1	2	0.1	0.4	0.1	0.2	0.1	2.2	0.2	0.1	0.1		-0.1
Lebanon	4.9	39.3	8.7	9.2	11.6	1.4	4.2	0.3	6	14.4	0.5	1.8	0.2	0.6	0.4	3.9	0.9	0.3	0.1	5.7	-0.1
Libya	1.9	77.4	1.5	2.1	1.2	0.2	1.1	0.2	3.3	11.1	0.2	1.6	0.7	0.2	0.2	0.6	0.9	0.3	0.5	6.1	-0.1
Morocco	3.3	66.3	3.4	1.8	1.6	0.1	2.7	0.3	5.4	15.1	0	2.6	0.1	1.4	0.5	2	3.3	0.1	0.6	4.5	-0.1
Palestine	5.6	38.9	9.8	9.7	9.4	1	4.5	0.3	7.1	13.7	0.3	1.7	0.2	0.5	0.3	3.9	1.2	0.3	0.1	5.3	-0.2
Oman	1.3	73.5	1.7	0.9	8.5	0.8	4.4	0	3.6	5.3	0.1	1.4	0.1	0.1	0.1	0.7	0.9	0	0.1	2	-0.1
Qatar	1.3	58.8	2.8	1	20.9	0.4	5.4	0	6.4	3	0.1	0.4	0.1	0.1	0.1	1	0.1	0	0	1.1	-0.1
Saudi Arabia	1.6	63.7	2.7	1.6	13.8	0.4	6.8	0.1	5.1	4.2	0.1	0.9	0.1	0.1	0.1	1.3	0.5	0.1	0.1	1.1	-0.1
Sudan	2.1	78	0.7	1.1	4.4	0.2	2.4	0.1	2.4	8.6	0.1	2.3	0.1	0.5	0.1	1.1	0.4	0.2	1.3	2.8	0
Syria	6.9	38.9	8.9	9.1	8.3	1.7	4.2	0.3	6.7	15	0.7	1.7	0.4	0.9	0.6	4.5	0.8	0.5	0.2	4.9	-0.2
Tunisia	16.4	33	3.8	2.2	3.4	0.3	4.1	0.5	9.6	26.7	0.2	5.3	1	0.5	1	2.8	4.9	0.4	1.1	9.7	-0.2
United Arab Emirates	1.4	65	2.8	0.7	14.6	1	5.9	0	5.3	3.3	0	0.7	0.1	0.1	0.1	0.8	0.5	0	0.1	1	-0.1
Yemen	2.1	63.2	2.2	1.5	11.7	0.8	5	0.1	3.9	9.5	0.1	2	0.1	0.2	0.1	2.2	1.1	0.1	0.2	3.6	-0.1

* The sum of the next columns.

1 **Table S6.** The number of conducted studies per population in the different continents or regions of the
 2 world.


Continent or region	Population (million)	Number of conducted studies	Ratio (study per population)	Reference
EMR countries	680	71	1 study per 9.6 million	Present study
Europe	748	174	1 study per 4.3 million	(Hopke et al., 2020)
North America	373	175	1 study per 2.1 million	
South America	437	11	1 study per 39.7 million	
Western Asia	313	12	1 study per 26.1 million	
South-eastern Asia	655	14	1 study per 46.8 million	
Eastern Asia (without China)	255	27	1 study per 9.5 million	
China	1.4 billion	246	1 study per 5.7 million	

3
 4
 5




7
 8

Figure S1. Spatial distribution of 22 EMR countries.




9
10
11



12
13
14
15
16
17


Figure S2. Flowchart illustrating the stages of the literature search on the SA, identification and characterization of PM studies (A) as well as emission inventories (B) in 22 EMR countries.



18


19 **Figure S3.** Chronology of source apportionment, identification and characterization studies over the EMR
 20 countries.

21



22


23 **Figure S4.** Pie chart of the number of papers from the EMR countries with the listed particle metric.



2

25 **Figure S5.** Relative (fractional) source of PM_{2.5} mass in 2017 based on the study of (McDuffie et al.,
 26 2021).

27



28

29 **Figure S6.** Relative contributions (%) of each sector to PM emissions in the EMR countries based on the
 30 conducted emission inventory studies.

31

32

33 References

- 34 Ettouney, R.S., Abdul-Wahab, S., Elkilani, A.S., 2009. Emissions inventory, ISCST, and neural network
35 modelling of air pollution in Kuwait. *International Journal of Environmental Studies* 66, 193-206.
- 36 Heger, M., Sarraf, M., 2018. Air pollution in Tehran: Health costs, sources, and policies. World Bank.
- 37 Hopke, P.K., Dai, Q., Li, L., Feng, Y., 2020. Global review of recent source apportionments for airborne
38 particulate matter. *Science of the Total Environment* 740, 140091.
- 39 Hopke, P.K., Feng, Y., Dai, Q., 2022. Source apportionment of particle number concentrations: A global
40 review. *Science of the Total Environment*, 153104.
- 41 Hosseini, V., Shahbazi, H., 2016. Urban air pollution in Iran. *Iranian Studies* 49, 1029-1046.
- 42 Karagulian, F., Belis, C.A., Dora, C.F.C., Prüss-Ustün, A.M., Bonjour, S., Adair-Rohani, H., Amann, M., 2015.
43 Contributions to cities' ambient particulate matter (PM): A systematic review of local source contributions
44 at global level. *Atmospheric environment* 120, 475-483.
- 45 Khan, A., Ahmad, S.S., 2018. Application of GAINS model for assessing selected air pollutants in Khyber
46 Pakhtunkhwa and Balochistan, Pakistan. *Arabian Journal of Geosciences* 11, 1-10.
- 47 Khatami, A., Ponche, J.-L., Jabry, E., Mirabel, P., 1998. The air quality management of the region of Great
48 Casablanca (Morocco). Part 1: Atmospheric emission inventory for the year 1992. *Science of the Total
49 Environment* 209, 201-216.
- 50 McDuffie, E.E., Martin, R.V., Spadaro, J.V., Burnett, R., Smith, S.J., O'Rourke, P., Hammer, M.S., van
51 Donkelaar, A., Bindle, L., Shah, V., 2021. Source sector and fuel contributions to ambient PM_{2.5} and
52 attributable mortality across multiple spatial scales. *Nature Communications* 12, 1-12.
- 53 Shahbazi, H., Abolmaali, A.M., Alizadeh, H., Salavati, H., Zokaei, H., Zandavi, R., Torbatian, S., Yazgi, D.,
54 Hosseini, V., 2021. Development of high-resolution emission inventory to study the relative contribution
55 of a local power plant to criteria air pollutants and Greenhouse gases. *Urban Climate* 38, 100897.
- 56 Shahbazi, H., Hassani, A., Hosseini, V., 2019. Evaluation of Tehran clean air action plan using emission
57 inventory approach. *Urban Climate* 27, 446-456.
- 58 Shahbazi, H., Reyhanian, M., Hosseini, V., Afshin, H., 2016a. The relative contributions of mobile sources
59 to air pollutant emissions in Tehran, Iran: an emission inventory approach. *Emission control science and
60 technology* 2, 44-56.
- 61 Shahbazi, H., Taghvaei, S., Hosseini, V., Afshin, H., 2016b. A GIS based emission inventory development
62 for Tehran. *Urban Climate* 17, 216-229.
- 63 Waked, A., Afif, C., Seigneur, C., 2012. An atmospheric emission inventory of anthropogenic and biogenic
64 sources for Lebanon. *Atmospheric environment* 50, 88-96.

65

66