

Spartina alterniflora has the highest methane emissions in a St. Lawrence estuary salt marsh

Comer-warner, Sophie A; Ullah, Sami; Ampuero Reyes, Wendy; Krause, Stefan; Chmura, Gail L

DOI:

[10.1088/2752-664X/ac706a](https://doi.org/10.1088/2752-664X/ac706a)

License:

Creative Commons: Attribution (CC BY)

Document Version

Publisher's PDF, also known as Version of record

Citation for published version (Harvard):

Comer-warner, SA, Ullah, S, Ampuero Reyes, W, Krause, S & Chmura, GL 2022, 'Spartina alterniflora has the highest methane emissions in a St. Lawrence estuary salt marsh', *Environmental Research: Ecology*, vol. 1, no. 1, 011003. <https://doi.org/10.1088/2752-664X/ac706a>

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

LETTER • OPEN ACCESS

Spartina alterniflora has the highest methane emissions in a St. Lawrence estuary salt marsh

To cite this article: Sophie A Comer-Warner *et al* 2022 *Environ. Res.: Ecology* 1 011003

View the [article online](#) for updates and enhancements.

You may also like

- [Modeling photosynthesis of *Spartina alterniflora* \(smooth cordgrass\) impacted by the Deepwater Horizon oil spill using Bayesian inference](#)
Wei Wu, Patrick D Biber, Mark S Peterson et al.
- [The weathering of oil after the Deepwater Horizon oil spill: insights from the chemical composition of the oil from the sea surface, salt marshes and sediments](#)
Zhanfei Liu, Jiqing Liu, Qingzhi Zhu et al.
- [Climate and anthropogenic controls on blue carbon sequestration in Hudson River tidal marsh, Piermont, New York](#)
D Peteet, J Nichols, D Pederson et al.

ENVIRONMENTAL RESEARCH ECOLOGY



LETTER

Spartina alterniflora has the highest methane emissions in a St. Lawrence estuary salt marsh

OPEN ACCESS

RECEIVED
13 January 2022

REVISED
9 May 2022

ACCEPTED FOR PUBLICATION
17 May 2022

PUBLISHED
30 August 2022

Original content from
this work may be used
under the terms of the
[Creative Commons
Attribution 4.0 licence](#).

Any further distribution
of this work must
maintain attribution to
the author(s) and the title
of the work, journal
citation and DOI.



Sophie A Comer-Warner^{1,2,*} , Sami Ullah^{2,3} , Wendy Ampuero Reyes¹ , Stefan Krause^{2,3} 
and Gail L Chmura¹ 

¹ Department of Geography, McGill University, Montreal, Quebec H3A 0B9, Canada

² School of Geography, Earth and Environmental Sciences, University of Birmingham, Edgbaston, B15 2TT Birmingham, United Kingdom

³ Birmingham Institute of Forest Research, University of Birmingham, Edgbaston, B15 2TT Birmingham, United Kingdom

* Author to whom any correspondence should be addressed.

E-mail: s.comer-warner@bham.ac.uk

Keywords: salt marsh, elevation zone, vegetation, blue carbon, methane, invasive species, sea level rise

Supplementary material for this article is available [online](#)

Abstract

Salt marshes have the ability to store large amounts of ‘blue carbon’, potentially mitigating some of the effects of climate change. Salt marsh carbon storage may be partially offset by emissions of CH₄, a highly potent greenhouse gas. Sea level rise and invasive vegetation may cause shifts between different elevation and vegetation zones in salt marsh ecosystems. Elevation zones have distinct soil properties, plant traits and rhizosphere characteristics, which affect CH₄ fluxes. We investigated differences in CH₄ emissions between four elevation zones (mudflat, *Spartina alterniflora*, *Spartina patens* and invasive *Phragmites australis*) typical of salt marshes in the northern Northwest Atlantic. CH₄ emissions were significantly higher from the *S. alterniflora* zone ($17.7 \pm 9.7 \text{ mg C m}^{-2} \text{ h}^{-1}$) compared to the other three zones, where emissions were negligible ($<0.3 \text{ mg C m}^{-2} \text{ h}^{-1}$). These emissions were high for salt marshes and were similar to those typically found in oligohaline marshes with lower salinities. CH₄ fluxes were significantly correlated with soil properties (salinity, water table depth, bulk density and temperature), plant traits (rhizome volume and biomass, root volume and dead biomass volume all at 0–15 cm) and CO₂ fluxes. The relationships between CH₄ emissions, and rhizome and root volume suggest that the aerenchyma tissues in these plants may be a major transport mechanism of CH₄ from anoxic soils to the atmosphere. This may have major implications for the mitigation potential carbon sink from salt marshes globally, especially as *S. alterniflora* is widespread. This study shows CH₄ fluxes can vary over orders of magnitude from different vegetation in the same system, therefore, specific emissions factors may need to be used in future climate models and for more accurate carbon budgeting depending on vegetation type.

1. Introduction

Salt marshes have high rates of carbon sequestration, which have the potential to reduce some of the impacts of climate change (Chmura *et al* 2003, McLeod *et al* 2011). The strength of the carbon sink resulting from carbon sequestration may be offset by emissions of the potent greenhouse gas (GHG), CH₄, which has a sustained-flux global warming potential 45 times higher than that of CO₂ over 100 years (Neubauer and Megonigal 2015). Salt marshes are often assumed to have negligible CH₄ fluxes (Chmura *et al* 2003, McLeod *et al* 2011), however, some studies have observed CH₄ fluxes from these environments (e.g. Chmura *et al* 2011, Roughan *et al* 2018, Yang *et al* 2021). Despite CH₄ emissions potentially switching salt marshes from net C sinks to net C sources (Bridgman *et al* 2006), understanding of the effect of how global change will affect CH₄ emissions from these systems remains inadequately quantified (McLeod *et al* 2011).

Methane fluxes from salt marshes vary between elevation zones due to changes in soil properties, plant species composition, plant traits and rhizosphere characteristics (Burke *et al* 2002, Philippot *et al* 2009, Noyce and Megonigal 2021, Rosentreter *et al* 2021). The soil properties affecting CH₄ emissions include temperature, bulk density, redox conditions, hydrological regime and substrate supply (Boeckx *et al* 1997, Zhao *et al* 2020, Noyce and Megonigal 2021, Rosentreter *et al* 2021). Salinity (as a proxy for sulphate supply from tidal floodwaters) is a major control (Mitsch and Gosselink 2007, Poffenbarger *et al* 2011) and sulphate-reducing bacteria in the soil outcompete methanogenic archaea (Villa 2020). However, alternative methanogenic pathways have been observed recently, which enable significant CH₄ emissions even at high salinities (Kelley *et al* 2015, Yuan *et al* 2019, Villa 2020). CH₄ emissions may vary between vegetation zones due to changes in both production versus consumption and transport from the soil to the atmosphere (Villa *et al* 2020, Noyce and Megonigal 2021).

A key mechanism of CH₄ transport is through pore spaces within the aerenchyma system of rhizomes, roots and stems, which can account for 80%–90% of total GHG emissions from some wetlands (Schutz *et al* 1991). Aerenchyma provide a direct channel between anoxic soils and the atmosphere, allowing CH₄ to be released without oxidation, therefore, increasing emissions (Brix *et al* 1992, Verville *et al* 1998). The transport mechanism and capacity can vary greatly depending on plant species and composition and so is expected to vary with marsh elevation zone (Laanbroek 2010, Villa 2020). Stomatal conductance may also be responsible for some plant-mediated transport, however, conflicting results of both higher CH₄ emissions and no effect of stomatal conductance have been observed (e.g. Whiting and Chanton 1996, Garnet *et al* 2005).

Salt marshes are under threat from changes in many environmental factors including sea level rise and invasive vegetation (Kirwan and Megonigal 2013, Martin and Moseman-Valtierra 2015). Environmental change effects on salt marsh vegetation include landward shifts in vegetation zones, transitions from native to invasive vegetation and vegetation dieback (e.g. Kirwan and Megonigal 2013, Martin and Moseman-Valtierra 2015). This may result in changes in the size or species composition of different elevation zones and conversion of vegetated salt marshes to tidal flats. The effect of shifting vegetation zones due to sea level rise on GHG fluxes and the understanding of the dominant controls on GHG fluxes remain critical knowledge gaps (Moseman-Valtierra *et al* 2016, Villa 2020).

In this study we examine whether fluxes of CH₄ vary with different elevation zones of a northern salt marsh, particularly considering differences in vegetation as a key driver of CH₄ fluxes. We considered all four of the major elevation zones present in the region: mudflat, *Spartina alterniflora*, *Spartina patens* and invasive *Phragmites australis*, therefore, incorporating marsh zones that may shift with both sea level rise and invasion of non-native species. We measured soil and plant properties to determine the environmental factors driving high CH₄ emissions and controlling differences observed between zones.

2. Materials and methods


2.1. Study site

This study was conducted in a salt marsh on the St. Lawrence estuary near La Pocatière, Quebec, Canada (47°22'24.7" N 70°03'26.3" W, figure 1(a)). The St. Lawrence estuary is mesohaline at this location (Gauthier 1980) with soil porewater salinities measured in this study ranging between 13 and 23 PSU. The average tidal range at this location is ~4.3 m. The climate is cool and wet with an average annual temperature and precipitation of 4.5 °C and 933 mm, respectively (Environment Canada, 2020). The salt marsh is approximately 160 m wide, with extensive mudflat, and is bordered on the landward side by a dyke. The salt marsh deposits are 1.8–2.0 m deep (van Ardenne 2016).

The zonation of salt marsh vegetation found here reflects that of the St. Lawrence estuary and is common across salt marshes of Eastern Canada and the Northeast U.S. (figure 1(b)). An extensive mudflat, exposed at low tides, is bordered by *S. alterniflora*, which is replaced at higher elevations by *S. patens*. Here, the *S. patens* zone also contains *Bolboschoenus maritimus*, *Plantago maritima*, *Triglochin maritima*, and *Salicornia europaea* in minor abundance. *Solidago sempervirens* and *Spartina cynosuroides* are found as elevation continues to increase, followed by a transition into terrestrial species along the dyke (Gu *et al* 2020). An invasive genetic strain of *P. australis*, widespread across the US and Canada (Saltonstall 2002), was observed at this study site in 2006 and has replaced the native vegetation in the transition zone from high marsh to terrestrial species, creating tall, dense monospecific stands (Gu *et al* 2020).

2.2. Field sampling

Six sampling locations were identified in each of the four elevation zones: mudflat, *S. alterniflora*, *S. patens*, and *P. australis* (*S. patens*/*P. australis* transition zone with 20% aerial coverage of *P. australis*). The sampling locations were chosen to capture similar elevation and species composition within each zone. The average



elevations of sampling locations in the *S. alterniflora*, *S. patens* and *P. australis* were 1.8, 2.5 and 2.7 m higher than the mudflat, respectively.

2.3. Gas sampling

GHG fluxes (CH_4 and CO_2) were measured using a dark, static chamber method designed to enclose both soil and vegetation (Magenheimer *et al* 1996). Chambers were made of 18 l, 26 cm diameter polycarbonate bottles, which were wrapped with bubble wrap and aluminium foil to insulate and block sunlight. Chambers were fitted with a 12 V, battery-powered computer fan (Sunon, Brea, California, U.S.) to ensure mixing of the air within the chamber and a venting tube was wrapped once around the outside of the chamber to minimise pressure differences (Hutchinson and Mosier 1981, Rochette 2011). The air temperature (HOBO Temperature Data Logger, Onset, Bourne, Massachusetts, U.S.) inside and outside of the chambers was measured during gas flux measurements. The average outside air temperature was 28.2 °C, with temperatures inside the chambers <4 °C below this.

Gas samples were taken on the 23rd of August 2020 from all sites, and gas fluxes presented here should represent the growing season fluxes. Gas samples were taken between 12:30 and 17:40, 5.5–10.5 h after high tide. In the vegetated sites, the chambers were deployed onto PVC collars (10 cm high) inserted 2.5 cm into the salt marsh three days before sampling. Chambers were fitted with a sampling assembly that was inserted into the chamber opening before the chambers were placed into the collar rim. The collar rim was filled with ambient water, taken from a nearby pool, to ensure an air-tight seal without altering porewater characteristics if any water overflowed onto the soil. Care was taken to ensure no vegetation was broken during installation of the chambers. The sampling assembly was long enough to minimise disturbance during sampling and had a 35 ml volume, therefore, during sampling, 35 ml of gas was extracted and discarded before a 25 ml sample representing gas inside the chamber was extracted. Five millilitres was expelled through the stopcock and needle of the syringe to minimise cross-contamination between samples and 20 ml was injected into a pre-evacuated 12 ml exetainer (Labco, U.K.). Gas samples were taken at 0, 20, 40 and 60 min. In the mudflat, chambers were carefully inserted directly into the mudflat to approximately 3 cm. Chambers were allowed to equilibrate with the air for 10 min after installation to minimise the effect of any disturbance during installation on initial GHG concentrations (Hamilton *et al* 2020). After this time, sampling assemblies were gently inserted into the chamber opening and sampled as described for the vegetated zones, except that gas samples were taken at 0, 20 and 40 min. Soil temperature was measured during chamber deployment at 10 cm depth within 10 cm of the collars using a soil thermometer (DeltaTrak 11050, Pleasanton, USA).

Soil and vegetation samples were collected from within the collar locations from the 24 to 26 August 2020 and the 19 and 20 September 2020. Vegetation was clipped at the soil surface and two 0–15 cm soil cores were collected, one for bulk density, total organic carbon (TOC) and total nitrogen (TN) analysis, and one for extractable nutrients, dissolved organic carbon (DOC) and total dissolvable nitrogen (TDN). Soil cores for bulk density/TOC/TN were collected using a 2.5 cm dia. Dutch gouge corer, sliced at 0–15 cm, then wrapped in plastic wrap. Soil cores for extractable nutrients were collected using the same corer, except in the mudflat where the top 3 cm were collected into a plastic bag using a metal spoon. Water table depth (WTD) was

measured as in Yu and Chmura (2009). Porewater was collected to measure salinity at 0–15 cm using a porewater sipper or extracted with a syringe from core holes (Yu and Chmura 2009). Neither of these methods were successful in the *S. alterniflora*, therefore, at these locations sub-samples of the extractable nutrient cores were centrifuged to release porewater. Tidal water was sampled for mudflat salinity measurements and salinity was measured using a portable ATC refractometer. All soil cores were collected with minimal compaction and all samples were transported cool and in the dark to the laboratory and stored at 4 °C.

2.4. Analyses

2.4.1. Bulk density and TOC/TN

Bulk density/TOC/TN cores were dried at 60 °C to constant weight, finely ground and analysed for TOC/TN. The bulk density was determined by dividing the dry soil weight by the core volume. TOC and TN samples were processed and analysed by the Soil Ecology Research Group, McGill University. Samples of ~0.5 g were weighed into a crucible and 1–2 ml of HCl was added to eliminate inorganic C. The samples were then dried at 50 °C for 48 h to remove all HCl. TOC and TN were measured by direct combustion at 900 °C with an Elemental Analyser (ThermoFinnigan Flash EA 1112 CN analyser, Carlo Erba, Milan, Italy). Results were obtained with an accuracy of ±5% for N and ±1% for C, and a limit of detection of 0.05% for both N and C.

2.4.2. Extractable nitrate and ammonium

Nitrate and ammonium samples were processed and analysed by the Soil Ecology Research Group, McGill University. Nitrate was measured as nitrate + nitrite and is referred to as nitrate from here on. A volume of 25 ml of 2 M KCl was added to 5 g of field-moist soil, shaken for 1 h at 200 rpm and then centrifuged for 20 min at 4000 rpm within 48 h of sample collection. The extractant was then filtered (0.45 μm) and analysed following the method of Sims *et al* (1995) on a microplate reader with a limit of detection of 0.1 ppm and an accuracy of ±5%.

2.4.3. Extractable DOC and TDN

To extract DOC and TDN 25 ml of ultrapure water (18.2 MΩ) was added to 5 g of field-moist soil, shaken for 1 h at 200 rpm and then centrifuged for 20 min at 4000 rpm. The extractant was then filtered (0.45 μm) and analysed on a TOC/TDN analyser (TOC VCSn + TMN-1, Shimadzu, Kyoto, Japan), with a 50 mg C l⁻¹ standard resulting in an accuracy and precision of 3.0 and ±4.4 mg l⁻¹, respectively.

2.4.4. GHG fluxes

Gas samples were analysed for CH₄ and CO₂ on a gas chromatograph (GC-14, Shimadzu, Kyoto, Japan) fitted with a flame ionisation detector at 250 °C. CO₂ was methanised to CH₄ for analysis. Standards of CH₄ (5.1 ppm) and CO₂ (5000 ppm) resulted in an accuracy and precision of 6.6 ± 1.5 and 0.4 ppm, and 5324 ± 324 and 78 ppm, respectively, for CH₄ and CO₂. GHG fluxes were calculated from the change in headspace gas concentration over time using a linear regression of the linear portion of the flux, which prevented data from being excluded due to low *r*². Where fluxes were below the minimum detectable concentration difference (MDCD) of the GC, fluxes were set to zero (Sgouridis and Ullah 2017).


2.4.5. Statistical analysis

Due to the relatively small sample size the non-parametric Kruskal–Wallis rank sum was used to test for significant differences (*p* < 0.05) in GHG fluxes between sites using R (R Development Core Team 2011). If significant differences were found, a post-hoc Dunn test with Bonferroni correction was performed to determine which groups were significantly different from each other. Spearman's R correlation, which is a non-parametric correlation analysis able to find linear or non-linear relationships, was performed in R (R Development Core Team 2011) to investigate potential relationships between CH₄ fluxes and environmental variables.

3. Results

3.1. CH₄ fluxes

CH₄ emissions were significant (above the MDCD) from the *S. alterniflora* locations resulting in a net CH₄ flux of 17.7 ± 9.7 mg C m⁻² h⁻¹ (figure 2). Only two locations in the *P. australis* zone and one in each of the *S. patens* and mudflat zones had CH₄ fluxes above the MDCD, which were all small at <0.3 mg C m⁻² h⁻¹. This resulted in significant differences in CH₄ fluxes between zones (*p* < 0.01, Kruskal–Wallis rank sum test).



Fluxes were significantly higher from the *S. alterniflora* zone compared to all three of the other zones (mudflat $p < 0.01$, *S. patens* $p < 0.01$, *P. australis* $p = 0.01$, Dunn test).

3.2. Relationships between GHG fluxes and environmental parameters

Zone-averaged environmental parameters are shown in table 1 and include further data from a companion study at this site, which were collected at the same time as the sampling of this study (Ampuero Reyes and Chmura 2022). Spearman correlation analysis on the whole dataset (table 2, figures 3 and S1 (available online at stacks.iop.org/ERE/1/011003/mmedia)) revealed that CH₄ fluxes were significantly related to multiple parameters with highest correlations found with salinity ($r = 0.72$, adjusted $p < 0.01$), WTD ($r = 0.72$, adjusted $p < 0.01$), CO₂ ($r = 0.64$, adjusted $p < 0.01$), rhizome volume at 0–15 cm ($r = 0.62$, adjusted $p < 0.01$), bulk density ($r = 0.61$, adjusted $p = 0.01$) and temperature ($r = 0.60$, adjusted $p = 0.01$). Rhizome biomass, root volume and dead volume at 0–15 cm were also significantly related to CH₄ with Spearman's r correlation coefficients between 0.48 and 0.51 (adjusted $p < 0.04$, table 2). Spearman correlation analysis on the *S. alterniflora* dataset revealed no significant relationships between CH₄ fluxes and environmental parameters.

Table 1. Environmental variables from four elevation zones at La Pocatière on the St. Lawrence River Estuary, August–September 2020. Included are data from this study and from a comparison study conducted at the same time (Ampuero Reyes and Chmura 2022).

Variable	Zone			
	Mudflat	<i>S. alterniflora</i>	<i>S. patens</i>	<i>P. australis</i>
CH ₄ -C (mg m ⁻² h ⁻¹)	0.1 ± 0.2	17.7 ± 9.7	0.1 ± 0.2	0.3 ± 0.4
CO ₂ -C (mg m ⁻² h ⁻¹)	59.7 ± 49.4	333.9 ± 67.7	156.7 ± 48.3	279.2 ± 100.7
NO ₃ ⁻ (μg N g wet soil ⁻¹)	0.7 ± 0.2	0.6 ± 0.2	0.6 ± 0.1	0.6 ± 0.1
NH ₄ ⁺ (μg N g wet soil ⁻¹)	2.6 ± 0.7	3.4 ± 2.1	3.0 ± 0.7	1.7 ± 0.1
DOC (μg C g wet soil ⁻¹)	32.5 ± 15.9	48.7 ± 11.4	43.5 ± 11.1	35.5 ± 5.0
TDN (μg N g wet soil ⁻¹)	13.3 ± 2.4	8.5 ± 2.5	9.4 ± 5.4	7.3 ± 3.2
Salinity	13.0 ± 0.0	20.3 ± 2.0	14.5 ± 0.8	15.2 ± 0.4
Temp. (°C)	19.6 ± 0.2	19.9 ± 0.0	16.9 ± 0.7	16.8 ± 0.3
Depth of water table (cm)	0.0 ± 0.0	7.0 ± 0.0	1.5 ± 0.7	5.0 ± 0.0
TOC (%)	1.5 ± 0.1	3.0 ± 0.3	3.6 ± 0.4	3.5 ± 3.0
TN (%)	0.1 ± 0.0	0.2 ± 0.0	0.2 ± 0.0	0.2 ± 0.0
OC:N	13.9 ± 1.7	15.7 ± 1.8	16.6 ± 1.1	14.8 ± 0.7
Bulk density (g cm ⁻³)	0.6 ± 0.0	0.8 ± 0.0	0.6 ± 0.1	0.6 ± 0.1
Aboveground biomass (g m ⁻²)	NA	26.1 ± 5.0	28.6 ± 8.7	62.2 ± 12.2
Rhizome volume 0–15 cm (cm ³)	NA	25.9 ± 11.9	11.3 ± 6.2	9.9 ± 1.2
Rhizome biomass 0–15 cm (g m ⁻²)	NA	3.2 ± 1.4	2.3 ± 0.9	2.9 ± 0.5
Root volume 0–15 cm (cm ³)	NA	27.8 ± 13.1	23.1 ± 13.8	7.6 ± 1.4
Root biomass 0–15 cm (g m ⁻²)	NA	2.7 ± 1.5	3.0 ± 1.3	0.9 ± 0.3
Dead volume 0–15 cm (cm ³)	NA	47.5 ± 14.9	29.5 ± 15.5	57.0 ± 28.3
Dead biomass 0–15 cm (g m ⁻²)	NA	3.2 ± 0.9	2.3 ± 1.1	4.1 ± 1.5
Rhizome volume 15–30 cm (cm ³)	NA	10.5 ± 5.5	7.4 ± 2.4	14.3 ± 10.4
Rhizome biomass 15–30 cm (g m ⁻²)	NA	1.2 ± 0.7	0.9 ± 0.3	2.4 ± 1.7
Root volume 15–30 cm (cm ³)	NA	12.2 ± 4.9	22.2 ± 9.3	20.6 ± 15.2
Root biomass 15–30 cm (g m ⁻²)	NA	1.6 ± 0.5	2.2 ± 0.4	3.2 ± 2.8
Dead volume 15–30 cm (cm ³)	NA	23.3 ± 16.2	12.0 ± 4.0	52.3 ± 50.8
Dead biomass 15–30 cm (g m ⁻²)	NA	2.2 ± 1.6	1.3 ± 0.4	3.4 ± 2.2


Table 2. Spearman's correlation (*r*) and adjusted *p*-values for the variables found to be significantly correlated with CH₄ fluxes.

Gas	Variable	Spearman's correlation (<i>r</i>)	Adjusted <i>p</i> (FDR)
CH ₄	CO ₂	0.64	<0.01
CH ₄	Salinity	0.72	<0.01
CH ₄	Temp.	0.60	0.01
CH ₄	Water table depth	0.72	<0.01
CH ₄	BD	0.61	0.01
CH ₄	RZ vol. 0–15 cm	0.62	<0.01
CH ₄	RZ bio. 0–15 cm	0.51	0.03
CH ₄	Root vol. 0–15 cm	0.51	0.03
CH ₄	Dead vol. 0–15 cm	0.48	0.04

4. Discussion

4.1. Comparison of CH₄ fluxes from *S. alterniflora* to other salt marshes

CH₄ fluxes observed from the *S. alterniflora* were much higher than those observed from other Northern salt marshes with *S. patens* and *S. alterniflora* as dominant vegetation (Magenheimer *et al* 1996, Chmura *et al* 2011, 2016, Roughan *et al* 2018). When compared to fluxes from other *S. alterniflora* marshes these results aligned with one study in China at a site with low salinity (Yang *et al* 2021) but were higher than most other fluxes previously observed (e.g. Bartlett *et al* 1985, Roughan *et al* 2018, Tong *et al* 2018). Annual *S. alterniflora* fluxes found here (upscaled for the period that the salt marsh is not frozen) were almost 140 times those expected for polyhaline marshes, with fluxes similar to the highest emitting oligohaline marshes (Poffenbarger *et al* 2011). This is especially surprising given that the study salt marsh is typically frozen from November to April i.e. no expected fluxes during this period and average annual temperatures are much lower than those found in other salt marshes. As gas fluxes were only measured once here, the annual fluxes calculated are estimates that should be refined with more annual data points, however, this upscaling suggests that CH₄ fluxes were much higher than those expected for polyhaline marshes.



4.2. Fluxes between zones

To our knowledge, CH₄ fluxes from all four marsh zones studied here have not been previously investigated, however, differences between marsh zones have been observed. Previously, higher CH₄ emissions from *S. alterniflora* than from bare mudflat, *S. patens* and *P. australis* (Yuan *et al* 2015, Roughan *et al* 2018, Liu *et al* 2019, Yang *et al* 2021), as well as other salt marsh species including, *Cyperus malaccensis* and *Suaeda salsa* have been observed (Zhang *et al* 2010, Yuan *et al* 2015). However, similar emissions between *S. alterniflora* and invasive *P. australis*, and *S. alterniflora* and *S. patens*, as well as higher emissions from invasive *P. australis* than native *S. patens*/*Distichlis spicata* have also been measured (Emery and Fulweiler 2014, Martin and Moseman-Valtierra 2015, Moseman-Valtierra *et al* 2016, Mueller *et al* 2016). Here, emissions of invasive *P. australis* and *S. patens* were similar. Species-specific comparisons are critical, therefore, to estimate correct emission factors for GHG budgets of salt marshes for the use in future climate models and for more accurate carbon budgeting depending on vegetation type.

4.3. Environmental controls

Differences in CH₄ emissions observed between elevation zones can be explained by differences in plant traits (such as root and rhizome biomass and volume) and plant-associated microbial assemblages, which can affect CH₄ production and consumption, and transport of CH₄ from anoxic soils to the atmosphere (Moseman-Valtierra *et al* 2016, Noyce and Megoñigal 2021). Distinct differences between microbial communities of *S. alterniflora* and *P. australis* have previously been found (Ravit *et al* 2003).

Belowground or total biomass may be positively correlated with CH₄ emissions or porewater concentrations (Cheng *et al* 2007, Mozdzer and Megoñigal 2013, Martin and Moseman-Valtierra 2017a, Noyce and Megoñigal 2021) or have no significant relationship (Moseman-Valtierra *et al* 2016). Additionally, lower CH₄ emissions with higher belowground biomass from *S. patens* has been attributed to increased methane oxidation in the rhizosphere (Martin and Moseman-Valtierra 2017a). These conflicting results indicate that the controls of the rhizosphere on CH₄ emissions requires further exploration and stage in the growing season, among other factors, may be affecting these relationships. This may be due to the interplay between root exudates promoting methanogenesis and diffusion of O₂ through roots suppressing methanogenesis (Fritz *et al* 2011, Waldo *et al* 2019), which are two processes among many others, which require further investigation.

Vascular plants promote gaseous exchange between the atmosphere and the subsurface with the dominant CH₄ emissions pathway in wetlands being through aerenchyma (Megoñigal *et al* 1999, Villa 2020). *S. alterniflora*, *S. patens* and *P. australis* all have the ability to create extensive aerenchyma in their rhizome and root systems with the extent of aerenchyma formation dependent on abiotic factors affecting soil oxygenation, such as water logging (e.g. Armstrong and Armstrong 1991, Armstrong *et al* 1999, Maricle and Lee 2002, Granse *et al* 2022). In salt marshes of the Northern Northwest Atlantic, *S. alterniflora* is subject to

the longest hydroperiod of any species, therefore, more extensive aerenchyma are expected compared with the *S. patens* and *P. australis*.

The relationship of CH₄ emissions with rhizome volume, rhizome biomass and root volume at 0–15 cm suggests this could be a key metric in predicting high CH₄ emissions and that CH₄ was transported through aerenchyma. An increase in CH₄ emissions where aerenchymous plants are present (e.g. Ford *et al* 2012) and at sites with higher stem densities (Chmura *et al* 2016) have been observed previously in salt marshes. The relationship with both rhizome metrics but only the root volume is likely due to the rhizome system being particularly important in these gaseous transport mechanisms (Armstrong and Armstrong 1991, Brix *et al* 1992).

Higher plant-mediated transport from *S. alterniflora* than *P. australis* has been observed previously, supporting the higher CH₄ emissions found here from the *S. alterniflora* (Tong *et al* 2012). Furthermore, the removal of *P. australis* aboveground biomass has no effect on CH₄ emissions indicating that plant transport is less important than abiotic factors or microbial production (Martin and Moseman-Valtierra 2017b). CH₄ fluxes from *S. alterniflora* have been found to be significantly correlated with porewater CH₄ concentrations (Zhang and Ding 2011) suggesting that the high emissions from this zone may be due to both high CH₄ production and high gas transport, which was found by Tong *et al* (2012).

An inverse relationship between salinity and CH₄ emissions, with negligible CH₄ emissions above a threshold salinity of 18 is widely assumed (Bartlett *et al* 1987, Poffenbarger *et al* 2011). Here, a positive correlation was observed between salinity and CH₄ fluxes contradicting both the inverse relationship between salinity and CH₄ emissions, and the salinity threshold of 18 for negligible CH₄ emissions (Poffenbarger *et al* 2011). Similar observations of highest CH₄ emissions at study sites with highest salinities and high CH₄ emissions at salinities >18 have been reported previously (e.g. Chmura *et al* 2011, 2016, Martin and Moseman-Valtierra 2017a, Emery and Fulweiler 2017), indicating that the assumption of low or negligible CH₄ emissions at high salinities is not valid across all environmental conditions. This may be due to other confounding environmental factors playing a more dominant role in CH₄ production and subsequent transport, for example, temperature being a more dominant control on CH₄ emissions than salinity (Abdul-Aziz *et al* 2018) or higher stem densities in areas of high salinity increasing transport (Chmura *et al* 2016).

High salinity is assumed to result in low CH₄ production due to substrate-competition between sulphate-reducing bacteria and methanogens (King and Wiebe 1980, Villa 2020). Methanogenesis can occur through three different pathways, acetoclastic, hydrogenotrophic and methylotrophic (Villa 2020). Methylotrophic methanogens preferentially use methylated substrates and are not in direct competition with sulphate-reducers (Oremland *et al* 1982, Villa 2020). This is the primary pathway for CH₄ production in salt marshes (Oremland *et al* 1982), therefore, despite high salinities salt marshes may be sources of CH₄ to the atmosphere. A transition from hydrogenotrophic to methylotrophic methanogenesis has been observed after *S. alterniflora* invasion into a tidal flat, further supporting the hypothesis that methanogens were not competing with sulphate-reducing bacteria (Yuan *et al* 2014, 2019). Additionally, salinity is not always a good indicator of sulphate concentrations as local depletion of sulphate in soil microzones may occur independently of salinity (King and Wiebe 1980, Poffenbarger *et al* 2011).

Higher water table or flooding is generally associated with higher CH₄ emissions although there may be a lag between higher water table and higher CH₄ emissions (Turetsky *et al* 2008, Zhao *et al* 2020, Calabrese *et al* 2021, Knox *et al* 2021). The highest CH₄ emissions observed here from the *S. alterniflora* zone where the water table was deepest below the surface were, therefore, unexpected. However, lower water level has been observed to have both no control on and increased rates of CH₄ fluxes from salt marshes indicating that other factors may be more important drivers in these ecosystems (Cheng *et al* 2007, Abdul-Aziz *et al* 2018).

Decomposition of dead vegetation is a likely source of carbon for methanogenesis explaining the significant relationship between dead volume at 0–15 cm and CH₄ fluxes. Additionally, the significant relationship between CH₄ and rhizome volume and biomass and root volume suggest that root-allocated organic carbon is an important source of fresh substrate for methanogenic bacteria. The lack of a significant relationship between CH₄ fluxes and DOC perhaps indicates that exudation by the dense rhizome of *S. alterniflora* was driving high rates of methanogenesis. Methanogenesis tends to occur in deeper, more anoxic soil layers so that dead vegetation at greater depths likely promotes methanogenesis (Conrad 1996, Villa 2020), however, CH₄ production has been observed up to ten times higher in oxic compared to anoxic soils (Angle *et al* 2017), supporting the relationship with dead volume in the upper soil layer observed here.

High temperature and bulk density may promote increased rates of biogeochemical reactivity and methanogenesis over methanotrophy (Dunfield *et al* 1993, Boeckx *et al* 1997). Higher temperatures are well known to increase reaction rates and have previously been observed to drive increased CH₄ emissions in salt and brackish marshes, as well as other wetlands (e.g. Hirota *et al* 2007, Noyce and Megonigal 2021). Despite this, conflicting relationships have been reported including no effect of temperature on CH₄ emissions, lower

CH₄ emissions at higher temperatures and increased CH₄ emissions with temperature in *P. australis* but not *S. alterniflora* (Ford *et al* 2012, Martin and Moseman-Valtierra 2017a, Liu *et al* 2019). Soils with higher bulk densities tend to facilitate more anoxic conditions, which may lead to methanogenesis and limit CH₄ oxidation (Boeckx *et al* 1997, del Grosso *et al* 2000). CH₄ emissions were also correlated with CO₂ emissions, indicating that the *S. alterniflora* zone was characterised by high rates of both aerobic and anaerobic respiration.

CH₄ transport by plant diffusion alone is slower than active exchange mechanisms (Brix *et al* 1992), which occur in the presence of light. Measuring fluxes in dark chambers excludes the active pumping mechanism of GHGs through plants (e.g. Whiting and Chanton 1996) resulting in potential underestimation of GHG fluxes. The relationships observed here between belowground biomass and CH₄ emissions indicate that diffusive plant processes, at least, are still important, supporting previous suggestions of diffusive transport of CH₄ through the aerenchyma of *S. alterniflora* (Zhang and Ding 2011). Additionally, plant community composition determines the extent to which light may alter rates of CH₄ emissions, with both large and negligible effects observed (Bartlett *et al* 1987, van der Nat and Middelburg 2000, Poffenbarger *et al* 2011).

4.4. Effect of sea level rise and invasive vegetation on CH₄ emissions

The relatively high CH₄ emissions from the *S. alterniflora* zone suggest that the strength of the carbon sink of salt marshes vegetated with *S. alterniflora* is greatly reduced. This has large implications for the blue C value of salt marshes, as *S. alterniflora* distribution is widespread and rapidly invading some coastal ecosystems. The patterns of CH₄ fluxes between elevation zones suggest that conversion of *S. alterniflora* to mudflat from sea level rise will significantly reduce CH₄ emissions, by 99.4%. If *S. patens* converts to *S. alterniflora* due to sea level rise this will lead to a large increase in CH₄ emissions (17 600%). Here, where the transition between *S. patens* and *P. australis* was studied, the effect of the invasive *P. australis* on the strength of the carbon sink appears to be minimal. CH₄ fluxes from within the monospecific *P. australis* stands are required to assess how the complete replacement of *S. patens* with invasive *P. australis* may affect the strength of the carbon sink.

5. Conclusions

Methane emissions were dependent on elevation zone with the highest emissions from the *S. alterniflora* while fluxes from the other elevation zones were negligible. These findings indicate that species-specific emission factors should be considered in carbon budgeting and to refine modelling of the strength of the carbon sink in salt marshes.

Methane emissions were high for salt marshes and were similar to those typically found in oligohaline marshes with lower salinities. Methane fluxes were significantly correlated with soil properties, plant traits and CO₂ fluxes. The control of root and rhizome volume on CH₄ fluxes indicates the importance of aerenchyma in transporting CH₄ directly from anoxic soils or porewater into the atmosphere, without being oxidised.

Highest CH₄ emissions from the zone of highest salinity provides further evidence that high salinity may not be a good predictor of low CH₄ emissions. This may be due to evidence that the methanogenic pathway in salt marshes and particularly in *S. alterniflora* does not directly compete with sulphate reducers. Further investigation is required to determine the microbial production versus emissions of CH₄ in these zones, as well as on constraining the methanogenesis mechanism in major salt marsh zones to allow the effect of salinity to be better constrained.

Data availability statement

The data that support the findings of this study are openly available at the following URL/DOI: <https://doi.org/10.5281/zenodo.6500188> (Comer-Warner *et al* 2022). Data will be available from 03 January 2023.

Acknowledgments

This project has received funding from the European Union's Horizon 2020 Research and Innovation Programme under the Marie Skłodowska-Curie Grant Agreement 838296 to S C W, an NSERC Discovery Grant to G L C and a Natural Environment Research Council Grant No. (NE/T012323/1) to S U. We thank Mike Dalva for his invaluable knowledge and insight on building and deploying static chambers.

ORCID iDs

Sophie A Comer-Warner  <https://orcid.org/0000-0003-1260-3151>

Sami Ullah  <https://orcid.org/0000-0002-9153-8847>

Wendy Ampuero Reyes  <https://orcid.org/0000-0002-1065-4951>

Stefan Krause  <https://orcid.org/0000-0003-2521-2248>

Gail L Chmura  <https://orcid.org/0000-0001-7163-3903>

References

- Abdul-Aziz O I, Ishtiaq K S, Tang J, Moseman-Valtierra S, Kroeger K D, Gonnea M E and Morkeski K 2018 Environmental controls, emergent scaling, and predictions of greenhouse gas (GHG) fluxes in coastal salt marshes *J. Geophys. Res.* **123** 2234–56
- Ampuero Reyes W and Chmura G 2022 Contribution of belowground plant components to salt marsh soil volume *Estuar. Coast. Shelf Sci.* accepted
- Angle J C *et al* 2017 Methanogenesis in oxygenated soils is a substantial fraction of wetland methane emissions *Nat. Commun.* **8** 1567
- Armstrong J, Afreeen-Zobayed F, Blyth S and Armstrong W 1999 *Phragmites australis*: effects of shoot submergence on seedling growth and survival and radial oxygen loss from roots *Aquat. Bot.* **64** 275–89
- Armstrong J and Armstrong W 1991 A convective through-flow of gases in *Phragmites australis* (Cav.) Trin. ex Steud *Aquat. Bot.* **39** 75–88
- Bartlett K B, Bartlett D S, Robert C and Sebacher D I 1987 Methane emissions along a salt marsh salinity gradient *Biogeochemistry* **202** 183–202
- Bartlett K B, Harriss R C and Sebacher D I 1985 Methane flux from coastal salt marshes *J. Geophys. Res.* **90** 5710–20
- Boeckx P, van Cleemput O and Villaralvo I 1997 Methane oxidation in soils with different textures and land use *Nutr. Cycl. Agroecosyst.* **49** 91–95
- Bridgman S D, Megonigal J P, Keller J K, Bliss N B, Trettin C and Service U F 2006 The carbon balance of North American Wetlands *Wetlands* **26** 889–916
- Brix H, Sorrel B K and Orr P T 1992 Internal pressurization and convective gas flow in some emergent freshwater macrophytes *Limnol. Oceanogr.* **37** 1420–33
- Burke D J, Hamerlynck E P and Hahn D 2002 Interactions among plant species and microorganisms in salt marsh sediments *Appl. Environ. Microbiol.* **68** 1157–64
- Calabrese S, Garcia A, Wilmoth J L, Zhang X and Porporato A 2021 Critical inundation level for methane emissions from wetlands *Environ. Res. Lett.* **16** 044038
- Cheng X, Peng R, Chen J, Luo Y, Zhang Q, An S, Chen J and Li B 2007 CH₄ and N₂O emissions from *Spartina alterniflora* and *Phragmites australis* in experimental mesocosms *Chemosphere* **68** 420–7
- Chmura G L, Anisfeld S C, Cahoon D R and Lynch J C 2003 Global carbon sequestration in tidal, saline wetland soils *Glob. Biogeochem. Cycles* **17** 1111–22
- Chmura G L, Kellman L and Guntenspergen G R 2011 The greenhouse gas flux and potential global warming feedbacks of a northern macrotidal and microtidal salt marsh *Environ. Res. Lett.* **6** 044016
- Chmura G L, Kellman L, van Ardenne L and Guntenspergen G R 2016 Greenhouse gas fluxes from salt marshes exposed to chronic nutrient enrichment *PLoS One* **11** e0149937
- Comer-Warner S A, Ullah S, Ampuero Reyes W, Krause S and Chmura G L 2022 Methane fluxes from four elevation zones in a St. Lawrence Estuary salt marsh *Zenodo* (<https://doi.org/10.5281/zenodo.6500188>)
- Conrad R 1996 Soil microorganisms as controllers of atmospheric trace gases (H₂, CO, CH₄, OCS, N₂O, and NO) *Microbiol. Rev.* **60** 609–40
- del Grosso S J *et al* 2000 General CH₄ oxidation model and comparisons of CH₄ oxidation in natural and managed systems *Glob. Biogeochem. Cycles* **14** 999–1019
- Dunfield P, Knowles R, Dumont R and Moore T R 1993 Methane production and consumption in temperate and subarctic peat soils: response to temperature and pH *Soil Biol. Biochem.* **25** 321–6
- Emery H E and Fulweiler R W 2014 *Spartina alterniflora* and invasive *Phragmites australis* stands have similar greenhouse gas emissions in a New England marsh *Aquat. Bot.* **116** 83–92
- Emery H E and Fulweiler R W 2017 Incomplete tidal restoration may lead to persistent high CH₄ emission *Ecosphere* **8** e01968
- Ford H, Garbutt A, Jones L and Jones D L 2012 Methane, carbon dioxide and nitrous oxide fluxes from a temperate salt marsh: grazing management does not alter global warming potential *Estuar. Coast. Shelf Sci.* **113** 182–91
- Fritz C, Pancotto V A, Elzenga J T M, Visser E J W, Grootjans A P, Pol A, Iturraspe R, Roelofs J G M and Smolders A J P 2011 Zero methane emission bogs: extreme rhizosphere oxygenation by cushion plants in Patagonia *New Phytol.* **190** 398–408
- Garnet K N, Megonigal J P, Litchfield C and Taylor G E 2005 Physiological control of leaf methane emission from wetland plants *Aquat. Bot.* **81** 141–55
- Gauthier B 1980 Les limites phytogéographiques du Saint-Laurent *Provancheria* **11** 1–103
- Granse D, Titschack J, Ainouche M, Jensen K and Koop-Jakobsen K 2022 Subsurface aeration of tidal wetland soils: root-system structure and aerenchyma connectivity in *Spartina* (Poaceae) *Sci. Total Environ.* **802** 149771
- Gu J, van Ardenne L B and Chmura G L 2020 Invasive *Phragmites* increases blue carbon stock and soil volume in a St. Lawrence estuary marsh *J. Geophys. Res.* **125** e2019JG005473
- Hamilton D J, Bulmer R H, Schwendenmann L and Lundquist C J 2020 Nitrogen enrichment increases greenhouse gas emissions from emergent intertidal sandflats *Sci. Rep.* **10** 6686
- Hirota M, Senga Y, Seike Y, Nohara S and Kunii H 2007 Fluxes of carbon dioxide, methane and nitrous oxide in two contrastive fringing zones of coastal lagoon, Lake Nakaumi, Japan *Chemosphere* **68** 597–603
- Hutchinson G L and Mosier A R 1981 Improved soil cover method for field measurement of nitrous oxide fluxes *Soil Sci. Soc. Am. J.* **45** 311–6
- Kelley C A, Chanton J P and Bebout B M 2015 Rates and pathways of methanogenesis in hypersaline environments as determined by ¹³C-labeling *Biogeochemistry* **126** 329–41

- King G M and Wiebe W J 1980 Regulation of sulfate concentrations and methanogenesis in salt marsh soils *Estuar. Coast. Shelf Sci.* **10** 215–23
- Kirwan M L and Megonigal J P 2013 Tidal wetland stability in the face of human impacts and sea-level rise *Nature* **504** 53–60
- Knox S H *et al* 2021 Identifying dominant environmental predictors of freshwater wetland methane fluxes across diurnal to seasonal time scales *Glob. Change Biol.* **27** 3582–604
- Laanbroek H J 2010 Methane emission from natural wetlands: interplay between emergent macrophytes and soil microbial processes. A mini-review *Ann. Bot.* **105** 141–53
- Liu L, Wang D, Chen S, Yu Z, Xu Y, Li Y, Ge Z and Chen Z 2019 Methane emissions from estuarine coastal wetlands: implications for global change effect *Soil Sci. Soc. Am. J.* **83** 1368–77
- Magenheimer J F, Moore T R, Chmura G L and Daoust R J 1996 Methane and carbon dioxide flux from a macrotidal salt marsh, Bay of Fundy, New Brunswick *Estuaries* **19** 139–45
- Maricle B R and Lee R W 2002 Aerenchyma development and oxygen transport in the estuarine cordgrasses *Spartina alterniflora* and *S. anglica* *Aquat. Bot.* **74** 109–20
- Martin R M and Moseman-Valtierra S 2015 Greenhouse gas fluxes vary between *Phragmites australis* and native vegetation zones in coastal wetlands along a salinity gradient *Wetlands* **35** 1021–31
- Martin R M and Moseman-Valtierra S 2017a Different short-term responses of greenhouse gas fluxes from salt marsh mesocosms to simulated global change drivers *Hydrobiologia* **802** 71–83
- Martin R M and Moseman-Valtierra S 2017b Plant manipulations and diel cycle measurements test drivers of carbon dioxide and methane fluxes in a *Phragmites australis*-invaded coastal marsh *Aquat. Bot.* **137** 16–23
- McLeod E, Chmura G L, Bouillon S, Salm R, Björk M, Duarte C M, Lovelock C E, Schlesinger W H and Silliman B R 2011 A blueprint for blue carbon: toward an improved understanding of the role of vegetated coastal habitats in sequestering CO₂ *Front. Ecol. Environ.* **9** 552–60
- Megonigal J P, Whalen S C, Tissue D T, Bovard B D, Albert D B and Allen A S 1999 A plant-soil-atmosphere microcosm for tracing radiocarbon from photosynthesis through methanogenesis *Soil Sci. Soc. Am. J.* **63** 665–71
- Mitsch W J and Gosselink J G 2007 *Wetlands* 4th edn (Hoboken, NJ: Wiley) (<https://doi.org/10.1016/j.ecoleng.2008.08.011>)
- Moseman-Valtierra S, Abdul-Aziz O I, Tang J, Ishtiaq K S, Morkeksi K, Mora J and Kroeger K D 2016 Carbon dioxide fluxes reflect plant zonation and belowground biomass in a coastal marsh *Ecosphere* **7** e01560
- Mozdzer T J and Megonigal J P 2013 Increased methane emissions by an introduced *Phragmites australis* lineage under global change *Wetlands* **33** 609–15
- Mueller P, Hager R N, Meschter J E, Mozdzer T J, Langley J A, Jensen K and Megonigal J P 2016 Complex invader-ecosystem interactions and seasonality mediate the impact of non-native *Phragmites* on CH₄ emissions *Biol. Invasions* **18** 2635–47
- Neubauer S C and Megonigal J P 2015 Moving beyond global warming potentials to quantify the climatic role of ecosystems *Ecosystems* **18** 1000–13
- Noyce G and Megonigal J P 2021 Biogeochemical and plant trait mechanisms drive enhanced methane emissions in response to whole-ecosystem warming *Biogeosci. Discuss.* **18** 2449–63
- Oremland R S, Marsh L M and Polcin S 1982 Methane production and simultaneous sulphate reduction in anoxic, salt marsh sediments *Nature* **296** 143–5
- Philippot L, Hallin S, Börjesson G and Baggs E M 2009 Biochemical cycling in the rhizosphere having an impact on global change *Plant Soil* **321** 61–81
- Poffenbarger H J, Needelman B A and Megonigal J P 2011 Salinity influence on methane emissions from tidal marshes *Wetlands* **31** 831–42
- R Development Core Team 2011 *R: A Language and Environment for Statistical Computing* (Vienna: R Foundation for Statistical Computing) (available at: www.R-project.org)
- Ravit B, Ehrenfeh J G and Haggblom M M 2003 A comparison of sediment microbial communities associated with *Phragmites australis* and *Spartina alterniflora* in two brackish wetlands of New Jersey *Estuaries* **26** 465–74
- Rochette P 2011 Towards a standard non-steady-state chamber methodology for measuring soil N₂O emissions *Anim. Feed Sci. Technol.* **166–167** 141–6
- Rosentreter J A, Al-Haj A N, Fulweiler R W and Williamson P 2021 Methane and nitrous oxide emissions complicate coastal blue carbon assessments *Glob. Biogeochem. Cycles* **35** e2020GB006858
- Roughan B L, Kellman L, Smith E and Chmura G L 2018 Nitrous oxide emissions could reduce the blue carbon value of marshes on eutrophic estuaries *Environ. Res. Lett.* **13** 044034
- Saltonstall K 2002 Cryptic invasion by a non-native genotype of the common reed, *Phragmites australis*, into North America *Proc. Natl Acad. Sci.* **99** 2445–9
- Schutz H, Schroder P and Rennenberg H 1991 Role of plants in regulating the methane flux to the atmosphere *Trace Gas Emissions from Plants* ed T Sharkey, E Holland and J Mooney (San Diego, CA: Academic) pp 29–63
- Sgouridis F and Ullah S 2017 Soil greenhouse gas fluxes, environmental controls, and the partitioning of N₂O sources in UK natural and seminatural land use types *J. Geophys. Res.* **122** 2617–33
- Sims G K, Ellsworth T R and Mulvaney R L 1995 Microscale determination of inorganic nitrogen in water and soil extracts *Commun. Soil Sci. Plant Anal.* **26** 303–16
- Tong C, Morris J T, Huang J, Xu H and Wan S 2018 Changes in pore-water chemistry and methane emission following the invasion of *Spartina alterniflora* into an oligohaline marsh *Limnol. Oceanogr.* **63** 384–96
- Tong C, Wang W-Q, Huang J-F, Gauci V, Zhang L-H and Zeng C-S 2012 Invasive alien plants increase CH₄ emissions from a subtropical tidal estuarine wetland *Biogeochemistry* **111** 677–93
- Turetsky M R, Treat C C, Waldrop M P, Waddington J M, Harden J W and McGuire A D 2008 Short-term response of methane fluxes and methanogen activity to water table and soil warming manipulations in an Alaskan peatland *J. Geophys. Res.* **113** G00A10
- van Ardenne L B 2016 Quantifying soil carbon storage and losses in natural and drained salt marshes *Master's Thesis* McGill University, Montreal, QC (available at: http://digitool.library.mcgill.ca:80/R/-&func=dbin-jump-full&object_id=143925&silolibrary=GEN01)
- van der Nat F-J and Middelburg J J 2000 Methane emission from freshwater marshes *Biogeochemistry* **49** 103–21
- Verville J H, Hobbie S E, Chapin F S III and Hooper D U 1998 Response of tundra CH₄ and CO₂ flux to manipulation of temperature and vegetation *Biogeochemistry* **41** 215–35

- Villa J A 2020 Functional representation of biological components in methane-cycling processes in wetlands improves modeling predictions *J. Geophys. Res.* **125** e2020JG005794
- Villa J A *et al* 2020 Methane and nitrous oxide porewater concentrations and surface fluxes of a regulated river *Sci. Total Environ.* **715** 136920
- Waldo N B, Hunt B K, Fadely E C, Moran J J and Neumann R B 2019 Plant root exudates increase methane emissions through direct and indirect pathways *Biogeochemistry* **145** 213–34
- Whiting G J and Chanton J P 1996 Control of the diurnal pattern of methane emission from emergent aquatic macrophytes by gas transport mechanisms *Aquat. Bot.* **54** 237–53
- Yang B, Li X, Lin S, Jiang C, Xue L, Wang J, Liu X, Espenberg M, Pärn J and Mander Ü 2021 Invasive *Spartina alterniflora* changes the Yangtze Estuary salt marsh from CH₄ sink to source *Estuar. Coast. Shelf Sci.* **252** 107258
- Yu O T and Chmura G L 2009 Soil carbon may be maintained under grazing in a St Lawrence Estuary tidal marsh *Environ. Conserv.* **36** 312–20
- Yuan J, Ding W, Liu D, Kang H, Freeman C, Xiang J and Lin Y 2015 Exotic *Spartina alterniflora* invasion alters ecosystem—atmosphere exchange of CH₄ and N₂O and carbon sequestration in a coastal salt marsh in China *Glob. Change Biol.* **21** 1567–80
- Yuan J, Ding W, Liu D, Xiang J and Lin Y 2014 Methane production potential and methanogenic archaea community dynamics along the *Spartina alterniflora* invasion chronosequence in a coastal salt marsh *Appl. Microbiol. Biotechnol.* **98** 1817–29
- Yuan J, Liu D, Ji Y, Xiang J, Lin Y, Wu M and Ding W 2019 *Spartina alterniflora* invasion drastically increases methane production potential by shifting methanogenesis from hydrogenotrophic to methylotrophic pathway in a coastal marsh *J. Ecol.* **107** 2436–50
- Zhang Y and Ding W 2011 Diel methane emissions in stands of *Spartina alterniflora* and *Suaeda salsa* from a coastal salt marsh *Aquat. Bot.* **95** 262–7
- Zhang Y, Ding W, Cai Z, Valerie P and Han F 2010 Response of methane emission to invasion of *Spartina alterniflora* and exogenous N deposition in the coastal salt marsh *Atmos. Environ.* **44** 4588–94
- Zhao M, Han G, Li J, Song W, Qu W, Eller F, Wang J and Jiang C 2020 Responses of soil CO₂ and CH₄ emissions to changing water table level in a coastal wetland *J. Clean. Prod.* **269** 122316