



Contribution to the Salpingidae (Coleoptera, Tenebrionoidea) of Greece including new faunistic records and ecological data on a poorly studied family

George Kakiopoulos¹, Jakovos Demetriou²

1 *Iliodos 60–62 street, 11527 Athens (Ampelokipi), Greece*

2 *Joint Services Health Unit Cyprus, BFC RAF Akrotiri BFPO 57, Akrotiri, Cyprus*

Corresponding author: George Kakiopoulos (strepens@yahoo.com)

Received 24 March 2022 | Accepted 8 June 2022 | Published 31 December 2022

Citation: Kakiopoulos G, Demetriou J (2022) Contribution to the Salpingidae (Coleoptera, Tenebrionoidea) of Greece including new faunistic records and ecological data on a poorly studied family. Travaux du Muséum National d'Histoire Naturelle “Grigore Antipa” 65(2): 39–49. <https://doi.org/10.3897/travaux.65.e84293>

Abstract

Salpingidae are a small family of beetles living mainly under the bark of deciduous forest trees and conifers, feeding on fungi, organic matter and other insects. A total of 23 species are known in the Western Palearctic out of which only six have been recorded in Greece. Material surveys of the first author during the last decades unveil ten species of Salpingidae for the country. Four species, *Salpingus ruficollis* (Linnaeus, 1760), *Sphaeriestes castaneus* (Panzer, 1796), *Sphaeriestes exsanguis* (Abeille de Perrin, 1870) and *Vincenzellus ruficollis* (Panzer, 1794) are reported from Greece for the first time. These records are presented alongside a literature survey regarding Greek Salpingidae, their overall distribution, ecology, host-plants and conservation status in the country.

Keywords

mycetophagous beetles, narrow-waisted bark beetles, Salpingidae, saproxylic beetles, *Salpingus ruficollis*, *Sphaeriestes castaneus*, *Sphaeriestes exsanguis*, *Vincenzellus ruficollis*.

Introduction

The family Salpingidae Leach, 1815 is comprised of approximately 350 species distributed across the world. Twenty-three species inhabit the Western Palearctic belonging to the subfamilies Agleninae Horn, 1878 and Salpinginae Leach, 1815

(Pollock and Löbl 2008). Salpingidae (narrow-waisted bark beetles) are mostly found in forests, where they live mainly under the bark of trees feeding on fungi and other insects, although the ecology of most Salpingidae is rather poorly understood (Barnouin and Zagatti 2017). Amongst these species, *Aglenus brunneus* (Gyllenhal, 1813) is considered as cryptogenic (Denux and Zagatti 2010) [i.e. of unknown origin (Carlton 1996)], probably introduced to Europe (Barnouin and Zagatti 2017) and currently widespread (Pollock and Löbl 2008). This species has been strongly associated with human habitations where it infests cultivated mushrooms, stored products and grain but also residing in decomposing organic matter (e.g. guano and manure) (Denux and Zagatti 2010; Barnouin and Zagatti 2017).

According to Pollock and Löbl (2008), four species of the family Salpingidae have been recorded from Greece. However, past overlooked literature records (von Oertzen 1886) as well as more recent research endeavours (Chehlarov et al. 2016) have raised this number to six species in total. In particular, the following species are known to inhabit Greece: *A. brunneus*, *Cariderus aeneus* (Olivier, 1807), *Lissodema denticollis* (Gyllenhal, 1813), *Salpingus planirostris* (Fabricius, 1787), *Sphaeriestes aeratus* (Mulsant, 1859) and *Sphaeriestes reyi* (Abeille de Perrin, 1874) (von Oertzen 1886; Pollock and Löbl 2008; Chehlarov et al. 2016; Popa et al. 2019).

Materials and methods

Locality data on Greek Salpingidae were extrapolated and corrected from the available scientific literature, citizen-science platforms (iNaturalist 2022) as well as specimens deposited in the private collection of the first author (GK) and Mr Thomas Barnouin (Laboratoire National d'Entomologie Forestière, Office National des Forêts) (TB) (Appendix 1). Data were mapped using QGIS Version 3.18.2 free and open source Geographic Information System (<https://qgis.org/en/site/>) (Fig. 1).

Results

Material examined:

Subfamily Agleninae Horn, 1878

Aglenus brunneus (Gyllenhal, 1813)

Peloponnese, Messinia, Mani, near Stoupa village [36.85°N; 22.26°E], 31.I.2004, alt. 100 m, leg. G. Kakiopoulos, shallow cave soil, occasional dead specimens under stones (2 collected) (GK); Peloponnese, Lakonia, Mt. Taygetos near Trypi village [37.09°N; 22.35°E], 15.IV.2007, alt. 700 m, leg. G. Kakiopoulos, entrance of shallow cave, 1 individual (alive) under stone (GK).

Subfamily Salpinginae Leach, 1815

Cariderus aeneus (Olivier, 1807)

Western Greece, Aitoloakarnania, Mt. Arakynthos [38.47°N; 21.47°E], 17.I.2004, alt. 800 m, leg. G. Kakiopoulos, 1 specimen under the bark of a young dry deciduous *Quercus* sp. (GK); Thessaly, Karditsa, near Portitsa village [39.31°N; 21.81°E], 14.XI.2010, alt. 300 m, leg. G. Kakiopoulos, 1 specimen collected by beating dry branches of deciduous *Quercus* sp. (GK).

Lissodema denticollis (Gyllenhal, 1813)

Central Macedonia, Chalkidiki, Mt. Stratonikon between Olympiada and Neochorion villages [40.54°N; 23.70°E], 28.VI.2007, alt. 800 m, leg. G. Kakiopoulos, 2 specimens collected by beating dead *Fagus sylvatica* L. branches (GK); Western Macedonia, Florina, south bank of lake Cheimaditida [40.61°N; 21.57°E], 25.V.2010, alt. 600 m, leg. G. Kakiopoulos, 1 specimen collected by beating *Populus* sp. branches (GK); Western Greece, Achaea, near Valta village [38.11°N; 22.06°E], 13.VII.2014, alt. 950 m, leg. G. Kakiopoulos, 1 specimen collected by beating *Platanus orientalis* L. branches (GK).

Salpingus ruficollis (Linnaeus, 1760) – New record for Greece

East Macedonia and Thrace, Kavala, Mt Pangaiion [40.91°N; 24.09°E], 29.IX.2002, alt. 1450 m, leg. G. Kakiopoulos, 1 specimen under *Fagus sylvatica* L. bark (GK) (Fig. 2A).

Salpingus planirostris (Fabricius, 1787)

Central Macedonia, Mt. Pieria towards Mt. Titaros [40.09°N; 22.26°E], 28.IV.2002, alt. 1600 m, leg. G. Kakiopoulos, 1 specimen under *Fagus sylvatica* L. bark (GK); Central Macedonia, Chalkidiki, Mt. Stratonikon between Olympiada and Neochorion villages [40.54°N; 23.70°E], 28.VI.2007, alt. 800 m, leg. G. Kakiopoulos, 2 specimens collected by beating dead *Fagus sylvatica* L. branches (species very common at the site) (GK); Thessaly, Volos [39.383523°N; 22.9736°E], 23.X.2021, alt. 130 m, obs. C. Kazilas, ident. G. Kakiopoulos, 1 individual (<https://www.inaturalist.org/observations/102398968>).

Sphaeriestes aeratus (Mulsant, 1859)

Central Greece, Euboea, near Pagondas village [38.67°N; 23.58°E], 12.XI.2004, alt. 450 m, leg. G. Kakiopoulos, 1 specimen collected by beating *Pinus halepensis* Mill. branches (GK).

Sphaeriestes castaneus (Panzer, 1796) – New record for Greece

East Macedonia and Thrace, Xanthi, near Leivaditis village, Chaidou forest [41.30°N; 24.66°E], 09.VII.2007, alt. 1150 m, leg. G. Kakiopoulos, 1 teratological specimen (9-segmented antennae), collected by beating *Pinus sylvestris* L. branches (GK) (Fig. 2B).

Sphaeriestes exsanguis (Abeille de Perrin, 1870) – New record for Greece
 North Aegean, Lesvos island, 6 km NW of Myloi Lampou [39.18°N; 26.35°E],
 19.X.2005, leg. G. Kakiopoulos, 2 specimens collected by beating *Pinus brutia* Ten.
 branches (GK) (Fig. 2C); Dodecanese, Rhodes island, Kritinia castel [36.26°N;
 27.81°E], 01.IV.2009, leg. Pierre Berger, 1 specimen (TB).

Sphaeriestes reyi (Abeille de Perrin, 1874)

Western Greece, Achaea, Aroania village [37.88°N; 22.01°E], 29.IX.1996, alt. 700 m,
 leg. G. Kakiopoulos, 1 specimen under the bark of dead tree (GK); Epirus, Ioannina,
 Mt. Smolikas, Palioselli village area [40.04°N; 20.88°E], 07.X.2003, alt. 1100 m,
 leg. G. Kakiopoulos, 1 specimen under the bark of cut *Pinus nigra* J.F. Arnold tree
 (GK); North Aegean, Lesvos island, Mt. Ordymnos, Antissa area [39.23°N; 25.93°E],
 18.X.2005, alt. 650 m, leg. G. Kakiopoulos, 1 specimen on stone (GK).

Vincenzellus ruficollis (Panzer, 1794) – New record for Greece

Western Macedonia, Mt. Smolikas, Samarina village area [40.09°N; 20.99°E],
 11.V.2002, alt. 1900 m, leg. G. Kakiopoulos, 1 specimen on a fallen *Fagus sylvatica*
 L. tree trunk (GK) (Fig. 2D).

Literature records:

Aglenus brunneus (Gyllenhal, 1813)

Attica, leg. E. von Oertzen (von Oertzen 1886); Crete, leg. E. von Oertzen (von
 Oertzen 1886); Greece (Pollock and Löbl 2008); Thessaly, Larissa, between
 Kefolovryso and Palaiokastro, cave Melissotrypa [39.87°N; 22.05°E], 2013–2018,
 alt. 299 m, in hypogenic cave that contains a sulfidic lake (Popa et al. 2019); Crete,
 Chania, Chordaki, cave Koutrouli [35.55°N; 24.16°E] (Coiffait 1955; Beron 2016;
 Paragamian 2021).

Lissodema denticollis (Gyllenhal, 1813)

Central Macedonia, Chalkidiki Peninsula, Arnea [Arnaia] [40.48°N; 23.59°E],
 15.VII.2008, leg. and ident. L. Fancello, 1 individual under *Fagus* sp., leaf-litter sifting
 (in L. Fancello collection) (Chehlarov et al. 2016).

Cariderus aeneus (Olivier, 1807)

Western Greece, Ileia, leg. U. Brenske (von Oertzen 1886); Greece (Pollock and Löbl
 2008).

Salpingus planirostris (Fabricius, 1787)

Western Greece, Ileia, leg. E. von Oertzen (von Oertzen 1886).

Sphaeriestes aeratus (Mulsant, 1859)

Greece (Pollock and Löbl 2008).

Sphaeriestes reyi (Abeille de Perrin, 1874)

Crete (Iablokoff-Khnzorian 1985); Greece (Kriti = Crete) (Pollock and Löbl 2008).

Discussion

In this article, an additional four species are added to the catalogue of Greek Salpingidae, raising their overall number to ten species. The study of Greek Salpingidae can be largely attributed to opportunistic material surveys with early reports providing no detail regarding precise localities, the habitat and ecology of reported species (von Oertzen 1886; Iablokoff-Khnzorian 1985). Despite the small number of records traced in the scientific literature, the total of 18 additional records, provide important distributional and ecological data for ten species inhabiting Greece (Fig. 1). For example, the recent discovery of *L. denticollis* in Greece (Chehlarov et al. 2016) is supplemented by three records of the species from Northern Greece and Peloponnese collected on beech, poplar and plane trees. Although, *L. denticollis* is known to be polyphagous (Barnouin and Zagatti 2017) *P. orientalis* and *Populus* sp. are recorded for the first time as its hosts.

Throughout the examination of specimens collected by the first author and Mr. T. Barnouin, *Salpingus ruficollis* (Linnaeus, 1760), *Sphaeriestes castaneus* (Panzer, 1796),




Figure 1. Georeferenced records of Greek Salpingidae according to the available literature and examined material. Locality records covering a large geographic area e.g. von Oertzen (1886) and Iablokoff-Khnzorian (1985) are not depicted.

Sphaeriestes exsanguis (Abeille de Perrin, 1870) and *Vincenzellus ruficollis* (Panzer, 1794) are recorded for the first time in Greece (Fig. 2A–D). The lone teratological specimen of *S. castaneus* with 9-segmented antennae is of additional importance as it constitutes an interesting record of teratogenesis in Salpingidae. Such an abnormality initially obstructed the correct identification of the specimen although, all remaining morphological characters were consistent with those of *S. castaneus* (Barnouin and Zagatti 2017). Furthermore, *S. exsanguis*, previously known from France (including Corsica) (Iablokoff-Khnzorian 1985), Portugal (Fuente 1932) and Spain (Lencina et al. 2008) was collected from Lesvos and Rhodes islands, suggesting its extended distribution not only in Western Europe but also towards the Eastern Mediterranean and probably in neighbouring Turkey.

Regarding the cryptogenic *A. brunneus*, it has been said to originate from North America (de Peyerimhoff 1945) although, this statement has been reviewed as rather controversial (Buckland et al. 2009). The species has been discovered in multiple archaeological sites across Northern Europe (Kenward 1975, 1976; Griffin et al. 1988; Kenward and Allison 1994; O'Brien et al. 2005; Buckland et al. 2009) as well as in Egypt, where an accidental introduction by the Romans has been hypothesised (Panagiotakopulu and van der Veen 1997). All the aforementioned records correspond to peri-domestic habitats or human mediated transportations, suggesting *A. brunneus* being highly synanthropic, widespread and probably native to the Old World (Buckland et al. 2009; Denux and Zagatti 2010; Barnouin and Zagatti 2017). In Greece, the species has been collected from caves in Crete (Coiffait 1955; Beron 2016; Paragamian 2021), Thessaly (Popa et al. 2019; Paragamian 2021) and Peloponnese (present article) while, such troglophilic encounters have also been reported from Corsica (Coache 2009) and Iran (Sadeghi et al. 2014). It is impossible to know whether these records represent human-mediated transportations to caves although, the presence of *A. brunneus* in such remote habitats where human interaction is rather minimal, raise further questions regarding the nativity or not of *A. brunneus* in Greece and by extend in the Palearctic.

The remaining species were mostly associated with forest trees of the genera *Fagus*, *Pinus*, *Populus* and *Quercus* according to their ecological preferences (Iablokoff-Khnzorian 1985; Barnouin and Zagatti 2017). *Sphaeriestes aeratus*, a rather rare Palearctic species (Iablokoff-Khnzorian 1985) was collected from branches of *P. halepensis* as opposed to the bark of fir and beech trees, as mentioned in the available literature (Barnouin and Zagatti 2017). Nevertheless, the biology of *S. aeratus* is still considered as poorly known (Barnouin and Zagatti 2017), with our results suggesting its additional dwelling on pines.

Up to date the extinction risk of European Salpingidae has been addressed in the regional Red List of Italian saproxylic beetles, where only *Sphaeriestes bimaculatus* (Gyllenhal, 1810) has been assessed as threatened [vulnerable (VU) under the IUCN criteria] (Carpaneto et al. 2015). In addition, nine Salpingidae are regarded as nearly threatened (NT) in Italy, including the native to Greece *S. ruficollis*, *S. aeratus*, *S. castaneus* and *S. reyi* (Carpaneto et al. 2015). Taking into consideration the scarcity

of published data on Greek Salpingidae, as well as the numerous threats encountered by European saproxylic beetles (Cálix et al. 2018), future research in the country could emphasize on material sampling and assessment of the extinction risk of Greek Salpingidae. Conservation actions minimizing some of the major threats to saproxylic beetles such as logging, tree loss and wood harvesting, urbanisation and tourism development or the increase in fire frequency/intensity (Cálix et al. 2018) could also benefit rarely encountered or potentially threatened species of Salpingidae.

Acknowledgments

We would like to thank Dr Darren Andrew Pollock (Eastern New Mexico University) for his valuable information on Salpingidae as well as Mr Thomas Barnouin (Laboratoire National d'Entomologie Forestière, Office National des Forêts) for his comments and provided record of *S. exsanguis* from Rhodes Island. We are also thankful to Mr Christos Kazilas (Institute of Biology, Leiden University, Sylvius Laboratory) for the provision of observational data regarding *S. planirostris* as well as Dr Apostolos Trichas (Natural History Museum of Crete) and Dr Christos Georgiadis




Figure 2. Dorsal view of *Salpingus ruficollis* (Linnaeus, 1760) (A), *Sphaeriestes castaneus* (Panzer, 1796) (B), *Sphaeriestes exsanguis* (Abeille de Perrin, 1870) (C) and *Vincenzellus ruficollis* (Panzer, 1794) (D), new records for Greece deposited in the private collection of G. Kakiopoulos.

(National and Kapodistrian University of Athens) for their corrections, comments and suggestions upon the manuscript, during the review process.

References

- Barnouin T, Zagatti P (2017) Les Salpingidae de la faune de France (Coleoptera Tenebrionoidea). L'Entomologiste 73(6): 353–386. [in French]
- Beron P (2016) Faune cavernicole de la Grèce. Maison d'édition Est-West, Sofia, 229 pp. [in French]
- Buckland PC, Panagiotakopulu E, Sveinbjarnardóttir GA (2009) A failed invader in the North Atlantic, the case of *Aglenus brunneus* Gyll. (Col., Colydiidae), a blind flightless beetle from Iceland. Biological Invasions 11: 1239–1245. <https://doi.org/10.1007/s10530-008-9339-6>
- Cálix M, Alexander KNA, Nieto A, Dodelin B, Soldati F, Telnov D, Vazquez-Albalate X, Aleksandrowicz O, Audisio P, Istrate P, Jansson N, Legakis A, Liberto A, Makris C, Merkl O, Mugerwa Petterson R, Schlaghamersky J, Bologna MA, Brustel H, Buse J, Novák V, Purchart L (2018) European Red List of Saproxylic Beetles. Brussels, Belgium: IUCN. Available at: <https://portals.iucn.org/library/node/47296>
- Carlton JT (1996) Biological invasions and cryptogenic species. Ecology 77: 1653–1655. <https://doi.org/10.2307/2265767>
- Carpaneto GM, Baviera C, Biscaccianti AB, Brandmayr P, Mazzei A, Mason F, Battistoni A, Teofili C, Rondinini C, Fattorini S, Audisio P (2015) A Red List of Italian Saproxylic Beetles: taxonomic overview, ecological features and conservation issues (Coleoptera). Fragmenta Entomologica 47(2): 53–126. <https://doi.org/10.13133/2284-4880/138>
- Chehlarov E, Guéorguiev B, Hristovski S, Fancello L, Cvetkovska-Gorgievska A, Prelík D (2016) New country records and rare and interesting species of Coleoptera from the Balkan Peninsula. Acta Zoologica Bulgarica 68(3): 331–338.
- Coache A (2009) Premier essai de tamisage dans les grottes de Haute-Corse (Coleoptera). L'Entomologiste 65: 39–41. [in French]
- Coiffait H (1955) Sur quelques Coléoptères cavernicoles de Grèce. Notes Biospéologiques 10(2): 205–209. [in French]
- Denux O, Zagatti P (2010) Coleoptera families other than Cerambycidae, Curculionidae sensu lato, Chrysomelidae sensu lato and Coccinelidae. BioRisk 4(1): 315–406. <https://doi.org/10.3897/biorisk.4.61>
- de la Fuente JM (1932) Catálogo sistemático-geográfico de los Coleópteros observados en la Península Ibérica, Pirineos propiamente dichos y Baleares (Continuación). Boletín de la Sociedad Entomológica de España 15: 75–90. [in Spanish]
- Griffin K, Oakland RH, Jones AKG, Kenward HK, Lie RW, Schia E (1988) Animal bones, moss, plant, insect and parasite remains. In: Schia E (Ed.) De arkeologiske utgravninger i Gamlebyen, Oslo 5. Alveheim and Eide, Ovre Ervik, 115–140.
- Iablokoff-Khnzorian SM (1985) Les Pythidae paléarctiques (Coleoptera). Deutsche Entomologische Zeitschrift 32(1–3): 193–229. <https://doi.org/10.1002/mmnd.19850320129> [in French]

- iNaturalist (2022) iNaturalist. <https://www.inaturalist.org> (accessed 20 March 2022)
- Kenward HK (1975) The biological and archaeological implications of the beetle *Aglenus brunneus* (Gyllenhal) in ancient faunas. *Journal of Archaeological Science* 2(1): 63–69. [https://doi.org/10.1016/0305-4403\(75\)90046-1](https://doi.org/10.1016/0305-4403(75)90046-1)
- Kenward HK (1976) Further archaeological records of *Aglenus brunneus* (Gyll.) in Britain and Ireland, including confirmation of its presence in the roman period. *Journal of Archaeological Science* 3(3): 275–277. [https://doi.org/10.1016/0305-4403\(76\)90062-5](https://doi.org/10.1016/0305-4403(76)90062-5)
- Kenward HK, Allison E (1994) A preliminary view of the insect assemblages from the early Christian rath site at Deer Park Farms, Northern Ireland. In: Rackham J (Ed.) *Environment and economy in Anglo-Saxon England*. Council for British Archaeology Research Report, 89–107.
- Lencina LJ, Baena M, Gallego D, Andújar C (2008) Un nuevo representante y nuevos registros de la familia Salpingidae Leach, 1815 (Coleoptera) en la Península Ibérica. *Boletín Sociedad Entomológica Aragonesa* 43: 421–423. [in Spanish]
- O'Brien C, Selby K, Ruiz Z, Brown A, Dinnin M, Caseldine C, Langdon P, Stuijts I (2005) A sediment-based multi-proxy palaeoecological approach to the environmental archaeology of lake dwellings (crannogs), central Ireland. *Holocene* 15: 707–719. <https://doi.org/10.1191%2F0959683605hl845rp>
- von Oertzen E (1886) Verzeichniss der Coleopteren Griechenlands und Cretas, nebst' einigen Bemerkungen über ihre geographische Verbreitung und die Zeit des Vorkommens einiger Arten betreffenden Sammelberichten. *Berliner Entomologische Zeitschrift* 30: 189–293. https://www.zobodat.at/pdf/Berliner-Ent-Zeitschrift_30_0189-0293.pdf [in German]
- Panagiotakopulu E, van der Veen M (1997) Synanthropic insect faunas from Mons Claudianus, a Roman quarry site in the Eastern Desert, Egypt. *Quaternary Proceedings* 5: 199–206.
- Paragamian K (2021) *Aglenus brunneus* (Gyllenhal, 1813). In: Paragamian K, Poulinakis M, Paragamian S, Nikoloudakis I (Eds) Cave fauna of Greece database - Hellenic Institute of Speleological Research. Species data updated 10.03.2020. <https://database.inspee.gr/fauna/browse/Animalia/Arthropoda/Insecta/Coleoptera/Salpingidae/Aglenus/Aglenus%20brunneus> (Accessed 16.11.2021)
- de Peyerimhoff P (1945) Les genres de Coléoptères importés ou acclimatés dans la fauna euro-méditerranéenne. *Revue française d'Entomologie* 12: 5–11. [in French]
- Pollock D, Löbl I (2008) Salpingidae. In: Löbl I, Smetana A (Eds) *Catalogue of Palaearctic Coleoptera*. Vol. 5. Tenebrionoidea. Stenstrup, Apollo Books, Denmark, 417–420.
- Popa I, Brad T, Vaxevanopoulos M, Giurgenca A, Baba ŞC, Iepure S, Plăiaşu R, Sarbu SM (2019) Rich and diverse subterranean invertebrate communities inhabiting Melissotrypa cave in Central Greece. *Travaux de l'Institut de Spéléologie «Émile Racovitz»* 58: 65–78.
- Sadeghi S, Dashan M, Bakhshi Y (2014) First record of the genus *Aglenus* (Coleoptera: Salpingidae) as a cave dweller from Iran (Tadovan Cave). *Cave and Karst Science* 41(3): 129 – 131.

Appendix 1. Literature records and examined material of Greek Salpingidae.

Species	Coordinates (decimal)	Altitude (m)	Locality	Date (DD/MM/YYYY)	Leg. Ident.	Habitat	Notes
<i>Agelenus brunneus</i>	36.85, 22.26	100	Peloponnese, Messini, Mani, near Stoupa village	31.I.2004	G. Kakiopoulos	shallow cave soil, occasional dead specimens under stones	in GK collection
<i>Agelenus brunneus</i>	37.09, 22.35	700	Peloponnese, Lakonia, Mt Taygetos near Trypi village	15.IV.2007	G. Kakiopoulos	entrance of shallow cave	in GK collection
<i>Agelenus brunneus</i>	N/A	N/A	Attica	N/A	E. von Oertzen	N/A	von Oertzen (1886)
<i>Agelenus brunneus</i>	N/A	N/A	Crete	N/A	E. von Oertzen	N/A	von Oertzen (1886)
<i>Agelenus brunneus</i>	N/A	N/A	N/A	N/A	N/A	N/A	Pollock and Lobl (2008)
<i>Agelenus brunneus</i>	39.87, 22.05	299	Thessaly, Larissa, between Kefalovryso and Palaiokastro, cave Melissotrypa Crete, Chania, Chordaki, cave Koutrouli	2013-2018	N/A	in hypogenic cave that contains a sulfidic lake	Popa et al. (2019)
<i>Agelenus brunneus</i>	35.55, 24.16	N/A	N/A	N/A	N/A	N/A	Coiffait (1955); Beron (2016); Paragamian (2021)
<i>Cariderus aeneus</i>	38.47, 21.47	800	Western Greece, Aitolokarnania, Mt Arakynthos	17.I.2004	G. Kakiopoulos	under the bark of a young dry deciduous <i>Quercus</i> sp.	in GK collection
<i>Cariderus aeneus</i>	39.31, 21.81	300	Thessaly, Karditsa, near Portitsa village	14.XI.2010	G. Kakiopoulos	collected by beating dry branches of deciduous <i>Quercus</i> sp.	in GK collection
<i>Cariderus aeneus</i>	N/A	N/A	Western Greece, Ileia	N/A	U. Brenske; E. von Oertzen	N/A	von Oertzen (1886)
<i>Cariderus aeneus</i>	N/A	N/A	N/A	N/A	N/A	N/A	Pollock and Lobl (2008)
<i>Lissodema denicolle</i>	40.54, 23.71	800	Central Macedonia, Chalkidiki, Mt Stratoniuk between Olympiada and Neochorion villages	28.VI.2007	G. Kakiopoulos	collected by beating dead <i>Fagus sylvatica</i> L. branches	in GK collection
<i>Lissodema denicolle</i>	40.61, 21.57	600	Western Macedonia, Florina, south bank of lake Cheimaditida	25.V.2010	G. Kakiopoulos	collected by beating <i>Populus</i> sp. branches	in GK collection
<i>Lissodema denicolle</i>	38.11, 22.06	950	Western Greece, Achaea, near Valta village	13.VII.2014	G. Kakiopoulos	collected by beating <i>Platanus orientalis</i> L. branches	in GK collection
<i>Lissodema denicolle</i>	40.48, 23.59	N/A	Central Macedonia, Chalkidiki Peninsula, Arria [Arriala]	15.VII.2008	L. Fancello	individual under <i>Fagus</i> sp., leaf-litter siftng	Chehlarov et al. (2016)
<i>Salpingus ruficollis</i>	40.91, 24.09	1450	East Macedonia and Thrace, Kavala, Mt Pangion	29.IX.2002	G. Kakiopoulos	under <i>Fagus sylvatica</i> L. bark	in GK collection; new record for Greece

Appendix 1. (continued)

Species	Coordinates (decimal)	Altitude (m)	Locality	Date (DD/MM/YYYY)	Leg. Ident.	Habitat	Notes
<i>Salpingus planirostris</i>	40.09, 22.26	1600	Central Macedonia, Mt Pieria towards Mt Titatos	28.IV.2002	G. Kakiopoulos	under <i>Fagus sylvatica</i> L. bark	in GK collection
<i>Salpingus planirostris</i>	40.54, 23.71	800	Central Macedonia, Chalkidiki, Mt Stratoniou between Olympiada and Neochorion villages	28.VI.2007	G. Kakiopoulos	collected by beating dead <i>Fagus sylvatica</i> L. branches (very common at the site)	in GK collection
<i>Salpingus planirostris</i>	39.3835, 22.9736	120	Thessaly, Volos	23.X.2021	C. Kazilas; G. Kakiopoulos	observed; online at: https://www.inaturalist.org/observations/102358968	iNaturalist
<i>Salpingus planirostris</i>	N/A	N/A	Western Greece, Ileia	N/A	E. von Oertzen	N/A	von Oertzen (1886)
<i>Sphaeriestes aeratus</i>	38.67, 23.58	450	Central Greece, Euboea, near Pagondas village	12.XL.2004	G. Kakiopoulos	collected by beating <i>Pinus halepensis</i> Mill. branches	in GK collection
<i>Sphaeriestes aeratus</i>	N/A	N/A	N/A	N/A	N/A	N/A	Pollock and Lobl (2008)
<i>Sphaeriestes castaneus</i>	41.31, 24.66	1150	East Macedonia and Thrace, Xanthi, near Leivadiatis village, Chaidou forest	09.VII.2007	G. Kakiopoulos	I teratological specimen, collected by beating <i>Pinus sylvestris</i> L. branches	in GK collection; new record for Greece
<i>Sphaeriestes exsanguis</i>	39.18, 26.35	N/A	North Aegean, Lesvos island, 6 km NW of Mylooi Lampou	19.X.2005	G. Kakiopoulos	collected by beating <i>Pinus brutia</i> Ten. branches	in GK collection; new record for Greece
<i>Sphaeriestes exsanguis</i>	36.26, 27.81	N/A	South Aegean, Dodecanese, Rhodes island, Kritina castle	01.IV.2009	Pierre Berger; Thomas Barnouin	specimen under the bark of dead tree	in TB collection; new record for Greece
<i>Sphaeriestes reyi</i>	37.88, 22.01	700	Western Greece, Achaea, Aroania village	29.IX.1996	G. Kakiopoulos	specimen under the bark of cut <i>Pinus nigra</i> J.F. Arnold tree	in GK collection
<i>Sphaeriestes reyi</i>	40.04, 20.88	1100	Epirus, Ioannina, Mt Smolikas, Palioselli village area	07.X.2003	G. Kakiopoulos	specimen under the bark of cut <i>Pinus nigra</i> J.F. Arnold tree	in GK collection
<i>Sphaeriestes reyi</i>	39.23, 25.93	650	North Aegean, Lesvos island, Mt Ordymnos, Antissa area	18.X.2005	G. Kakiopoulos	specimen on stone	in GK collection
<i>Sphaeriestes reyi</i>	N/A	N/A	Crete	N/A	S.M. Iabolkoff-Khnzorian	N/A	Iabolkoff-Khnzorian 1985
<i>Vincenzellus ruficollis</i>	40.09, 20.99	1900	Western Macedonia, Mt Smolikas, Samarina village area	11.V.2002	G. Kakiopoulos	specimen on a fallen <i>Fagus sylvatica</i> L. tree trunk	in GK collection; new record for Greece
						N/A	Pollock and Lobl (2008)