

This is an electronic reprint of the original article.

This reprint *may differ* from the original in pagination and typographic detail.

Author(s): Mladen Ognjenović, Ivan Seletković, Mia Marušić, Mathieu Jonard, Pasi Rautio, Volkmar Timmermann, Melita Perčec Tadić, Miran Lanščak, Damir Ugarković & Nenad Potočić

Title: The Effect of Environmental Factors on the Nutrition of European Beech (*Fagus sylvatica* L.) Varies with Defoliation

Year: 2023

Version: Published version

Copyright: The Author(s) 2023

Rights: CC BY 4.0

Rights url: <http://creativecommons.org/licenses/by/4.0/>






Please cite the original version:

Ognjenović, M.; Seletković, I.; Marušić, M.; Jonard, M.; Rautio, P.; Timmermann, V.; Tadić, M.P.; Lanščak, M.; Ugarković, D.; Potočić, N. The Effect of Environmental Factors on the Nutrition of European Beech (*Fagus sylvatica* L.) Varies with Defoliation. *Plants* 2023, 12, 168.
<https://doi.org/10.3390/plants12010168>

All material supplied via *Jukuri* is protected by copyright and other intellectual property rights. Duplication or sale, in electronic or print form, of any part of the repository collections is prohibited. Making electronic or print copies of the material is permitted only for your own personal use or for educational purposes. For other purposes, this article may be used in accordance with the publisher's terms. There may be differences between this version and the publisher's version. You are advised to cite the publisher's version.

Article

The Effect of Environmental Factors on the Nutrition of European Beech (*Fagus sylvatica* L.) Varies with Defoliation

Mladen Ognjenović ¹, Ivan Seletković ², Mia Marušić ^{2,*}, Mathieu Jonard ³, Pasi Rautio ⁴, Volkmar Timmermann ⁵, Melita Perčec Tadić ⁶, Miran Lanščak ², Damir Ugarković ⁷ and Nenad Potočić ²

¹ Department of BI, Analytics and Research, Njuškalo Ltd., 10000 Zagreb, Croatia

² Croatian Forest Research Institute, 10450 Jastrebarsko, Croatia

³ Earth and Life Institute, Université Catholique de Louvain, 1348 Louvain-la-Neuve, Belgium

⁴ Natural Resources Institute Finland, 00790 Helsinki, Finland

⁵ Norwegian Institute of Bioeconomy Research, 1433 Ås, Norway

⁶ Meteorological and Hydrological Service, 10000 Zagreb, Croatia

⁷ Faculty of Forestry and Wood Technology, University of Zagreb, 10000 Zagreb, Croatia

* Correspondence: miam@sumins.hr

Abstract: Despite being adapted to a wide range of environmental conditions, the vitality of European beech is expected to be significantly affected by the projected effects of climate change, which we attempted to assess with foliar nutrition and crown defoliation, as two different, yet interlinked vitality indicators. Based on 28 beech plots of the ICP Forests Level I network, we set out to investigate the nutritional status of beech in Croatia, the relation of its defoliation and nutrient status, and the effects of environmental factors on this relation. The results indicate a generally satisfactory nutrition of common beech in Croatia. Links between defoliation and nutrition of beech are not very direct or very prominent; differences were observed only in some years and on limited number of plots. However, the applied multinomial logistic regression models show that environmental factors affect the relationship between defoliation and nutrition, as climate and altitude influence the occurrence of differences in foliar nutrition between defoliation categories.

Keywords: tree vitality; foliar composition; stoichiometry; climate change; ICP Forests



Citation: Ognjenović, M.; Seletković, I.; Marušić, M.; Jonard, M.; Rautio, P.; Timmermann, V.; Tadić, M.P.; Lanščak, M.; Ugarković, D.; Potočić, N. The Effect of Environmental Factors on the Nutrition of European Beech (*Fagus sylvatica* L.) Varies with Defoliation. *Plants* **2023**, *12*, 168. <https://doi.org/10.3390/plants12010168>

Academic Editor: Aneta Helena Baczewska-Dąbrowska

Received: 31 October 2022

Revised: 7 December 2022

Accepted: 23 December 2022

Published: 30 December 2022



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Meteorological and climate conditions affect vegetation directly and indirectly. Direct effects include responses to temperature; indirect effects occur primarily as soil-mediated phenomena, such as the influence of precipitation on soil moisture regimes [1–4]. Probably the greatest threat to existing forest ecosystems in Europe comes from a changing climate [5]. Extreme climatic events such as droughts are thought to be important in initiating changes in forest ecosystems [6]. According to IPCC [7] predictions, extreme climate events such as heat waves and long-lasting summer droughts will occur more frequently. Such climate changes will negatively impact the stability, structure and biodiversity of forest ecosystems throughout Europe [8]. The region of southeast Europe is one of the most vulnerable regions in Europe when it comes to climate change impacts, primarily through intensified severity and duration of droughts and heat waves. As these impacts should be stronger and faster than on the rest of the continent [9,10] this region is an ideal model for studying the future impact of changing climatic conditions.

A prerequisite for healthy growth and balanced metabolism is the maintenance of adequate concentrations and relatively stable ratios of nutrients in plant tissues [11]. Concentrations of nutrients and their relationships in the leaves of forest trees are important indicators of their functioning. It is crucial to account for nutrient limitation when studying the forest response to climate change [12], as nutritional status has a significant and diverse impact on tree vitality [13]. The concentration of nutrients in leaves depends on many

factors: the amount and distribution of precipitation and the length of the growing season, the presence of nutrients in the soil, ion antagonisms, ion mobility and uptake capacity, etc. [14]. It seems that the nutritional status of European beech (*Fagus sylvatica* L.) is deteriorating: Jonard et al. [12] and Talkner et al. [15] both report on the negative trend in beech phosphorus (P) nutrition; furthermore, trends of concentrations of calcium and magnesium in the foliage of beech trees in Europe were also found to be negative [12]. Nutrient contents of tree foliage, which reflect atmospheric and soil-related influences, are an important part of the UNECE ICP Forests monitoring scheme [16]. Beech forests in Croatia have a very wide edaphic amplitude [17], but baseline data on the chemical composition of foliage in beech stands has so far been missing.

Tree vitality is defined as the ability of a tree to assimilate, to survive stress, to react to changing conditions, and to reproduce [18]. As vitality cannot be measured directly, various indicators can be used to describe it [19]. Crown defoliation is a commonly used tree vitality indicator [20–22], which can be obtained cost-effectively and relatively quickly in field surveys [23]. Defoliation is defined as the loss of needles or leaves in the assessable part of the crown compared to a reference tree [24]. Trees with defoliation above 25% are regarded as considerably defoliated and are thus defined as damaged [25]. A recent study showed that there are significant morphological and, in this context even more importantly, physiological differences in beech leaves between healthy and considerably defoliated trees [22]. Although European beech had the lowest mean defoliation (20.9%) of all main tree species in the ICP Forests 2021 survey, the 20-year trend in mean plot defoliation shows that defoliation of beech is constantly on the rise (3.4% in 20 years), and that larger deviations from this trend can be linked to drought events [26]. A good example is year 2004, when the annual mean plot defoliation was higher than the trend as a result of the drought in the preceding year which affected large parts of Europe [27–29]. In Croatia, the trend in mean defoliation of beech is much steeper with an 8.6 % rise in the period of 1996–2017 [30].

European beech is often mentioned in the context of sensitivity to high temperatures and drought [31,32], as beech grows best in areas marked by moderately warm summers and high quantities of precipitation [33]. Beech forms vast forest communities which provide multiple ecosystem goods and services, and is the dominant tree species in many deciduous forests in Europe under temperate climate conditions [34]. The projected effects of climate change, particularly drought, are expected to significantly affect the vitality of beech, which we attempted to assess with foliar nutrition and crown defoliation, as two different, yet closely interlinked vitality indicators. Although considered a nonspecific indicator of vitality, defoliation appears to depend on the nutritional status of trees. However, research on the relationship between tree defoliation and nutritional status has so far been limited, especially for beech: Jonard et al. [35] found that defoliation was associated with lower concentrations of calcium (Ca) and magnesium (Mg) in beech leaves. A study conducted in Switzerland did not find a link between nitrogen (N) concentrations and defoliation [36], but Ferretti et al. [37] report that the proportion of common beech trees with more than 25% defoliation increases with elevated nitrogen to calcium and potassium (K) ratios in the leaves, indicating that crown status depends on the balance of nutrients in leaves. Toigo et al. [38] found that defoliation increased due to drought effects, while the influence of nutritional status was less consistent. The study by Ognjenović et al. [39] on a limited sample of 26 beech trees showed that defoliation and calcium leaf concentrations show similar susceptibility to drought, but with different responses in time. Obviously, besides the overall tree health condition, the concentration of nutrients in leaves depends on many outside factors. In this context, we set out to better characterize the link between defoliation and foliar nutrition and determine the influence of environmental factors on the foliar composition of beech trees of two distinct defoliation categories, on a large sample of beech-dominated ICP Forests Level I plots in Croatia, with the hypotheses that (I) environmental factors have a significant influence over the defoliation/nutrition relationship and (II) trees with higher defoliation have lower concentrations of nutrients in leaves. In the

process, we investigated whether tree nutrition plays a role in the loss of vitality of beech trees (and vice versa); and if the link between tree nutrition and vitality loss varies from year to year due to interannual climate variations.

2. Results

2.1. Foliar Nutrient Concentrations

Mean N foliar concentrations in the study area level were within normal range [40] in both defoliation categories and across sampling years. Repeated Measures ANOVA indicated that there was no difference in N concentrations between LD and HD trees, while there was a difference between individual years ($F(2, 556) = 30.4, p < 0.0001$). In 2018, significantly lower N concentrations were recorded for both defoliation categories compared to 2019 and 2020 (Table 1).

Table 1. Results of Holm pairwise comparisons of foliar nutrient concentrations and sample dry mass within a defoliation category and between sampling years (NS—not significant, * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$).

Defoliation Category	Year 1	Year 2	N	P	K	Ca	Mg	Sample Dry Mass
LD	2018	2019	***	ns	***	**	***	***
LD	2018	2020	***	***	***	***	ns	*
LD	2019	2020	ns	***	ns	***	*	*
HD	2018	2019	***	***	***	ns	***	***
HD	2018	2020	***	***	***	***	***	***
HD	2019	2020	ns	*	**	***	ns	*

Although there is no significant main effect of defoliation category on P concentrations, the pairwise comparisons between LD and HD trees indicated significant differences in 2019, when LD trees had a higher concentration than HD trees, but not in 2018 and 2020 (Figure 1). Significant differences in P concentrations ($F(2, 556) = 74.9, p < 0.001$) were also observed between individual sampling years (Table 1) with a decrease during the study period. However, values of P on the plot level were mostly normal throughout the study period except for the year 2020, when P concentrations were slightly below normal range on 53% of the research plots (Figure S3).

No significant differences were established for potassium (K) between defoliation categories and mean concentration values stayed within the normal range in each year (Figure 1).

Repeated Measures ANOVA showed no significant main effect of defoliation category on Mg concentrations. Nevertheless, pairwise comparisons between LD and HD trees revealed significant differences in 2018, with HD trees having higher Mg concentrations (Figure 1). Mean Mg concentrations were within normal range in each year, although they were lower in 2019 and 2020 compared to 2018 for HD trees.

Similar to other elements, there was no significant main effect of defoliation category on calcium (Ca) concentrations ($F(2, 556) = 3.4, p = 0.06$). Pairwise comparisons, however, showed that HD trees in 2018 and 2019 had significantly higher Ca concentrations than LD trees (Figure 1). For these years, values for both groups fall within normal Ca concentration range for beech. In 2020, both categories had significantly higher concentrations than in previous years, bordering on surplus (luxury) values, but no significant difference was found between them (Figure 1).

Although there was no significant main effect of defoliation category on dry leaf mass, the pairwise comparisons between LD and HD trees indicated significant differences in 2018, when LD trees had a higher dry leaf mass than HD trees (Figure 1). However, there was a statistically significant main effect of the sampling year on dry leaf mass ($F(2, 556) = 56.3, p < 0.001$) and pairwise comparisons showed differences between all years for both defoliation categories (Table 1).




Figure 1. Mean foliar concentrations and sample dry mass (dots) with confidence intervals (vertical bars) for LD (green color) and HD trees (orange color) on all plots within sampling years. Dotted lines represent lower and upper critical value for normal range of foliar N concentrations for common beech according to Mellert and Goettlein [40]. Only significant differences in foliar concentrations between LD and HD categories determined by the Holm–Bonferroni method are indicated (* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$).

On the plot level, mean Ca concentrations were mostly in the normal range during the first two years, while excessive Ca concentrations were observed on all plots located in the alpine region in 2020 (Figure S5), and most of the alpine region plots had insufficient P concentrations (Figure S3). In the same year, excessive K concentrations were found on eleven plots (39%), most of them located in the continental region (Figure S4). N values are rather plot dependent and relatively stable in time (Figure S2), while Mg concentrations show a very broad range of values, independent of the area, defoliation category, or sampling year (Figure S6)

Overall, the share of plots with differences in foliar concentrations between defoliation categories was relatively low for most elements during the entire sampling period (Table 2). The highest share of plots with differences in foliar concentrations between defoliation categories was determined for Ca (21.4%) in 2018, followed by Mg (17.9%) in the same year.

Table 2. Number of plots with significantly higher foliar concentrations in each defoliation category within sampling years determined by the independent samples *t*-test (total number of plots: 28).

		N	P	K	Ca	Mg
2018	HD higher	1	-	-	6	4
	LD higher	1	-	2	-	1
2019	HD higher	-	-	1	-	2
	LD higher	1	4	1	1	1
2020	HD higher	-	2	1	1	1
	LD higher	2	-	-	-	1

2.2. Foliar Nutrient Ratios

N/P ratios were especially high and above the upper limit for both LD and HD trees in 2020 at the study area level, mostly due to very low P values that approached deficiency values (Figure 2). Repeated Measures ANOVA did not show a significant main effect of defoliation category on N/P ratio. However, pairwise comparison between LD and HD trees indicated significant differences in N/P ratios in 2019, with values out of optimal range for HD trees due to low P concentration values (Figure 2). The main effect of the sampling year was significant ($F(2556) = 13.4$, $p < 0.001$) with an increase in N/P ratio during the monitoring period.




Figure 2. Mean foliar ratios (dots) and confidence intervals (vertical bars) for LD (green color) and HD trees (orange color) on all plots within sampling years. Dotted lines represent lower and upper critical value for normal range of foliar ratios for common beech according to Mellert and Goettlein [40]. Only significant differences in foliar ratios between LD and HD categories determined by the Holm–Bonferroni method are indicated (* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$).

No significant differences were found for N/K ratio between defoliation groups or sampling years and values were within normal range in all years.

Repeated Measures ANOVA identified a significant main effect of defoliation categories on N/Ca ratios ($F(2556) = 109.9$, $p < 0.0001$). Pairwise comparisons confirmed differences in N/Ca ratios between defoliation groups in 2018 and 2019 (Figure 2). These differences were dictated primarily by Ca concentrations, which showed higher values in HD trees in 2018 and 2019. Sampling year also had a significant effect ($F(2556) = 13.4$, $p < 0.001$) and pairwise comparison established that values within defoliation categories were lower in 2020 compared to 2018 and 2019 (Table 3).

Due to low Mg concentrations in 2019 and 2020, the N/Mg ratios were mostly in the surplus range in those years (Figure 2). No significant effect of defoliation categories on N/Mg ratios was found. However, pairwise comparisons showed a significant difference between LD and HD trees in 2018, with higher values for LD trees.

Table 3. Results of Holm pairwise comparisons of foliar nutrient ratios within a defoliation category and between sampling years (ns—not significant, * $p < 0.05$, *** $p < 0.001$).

Defoliation Category	Year 1	Year 2	N/P	N/K	N/Ca	N/Mg
LD	2018	2019	ns	ns	ns	ns
LD	2018	2020	***	ns	***	ns
LD	2019	2020	***	ns	***	ns
HD	2018	2019	ns	ns	ns	***
HD	2018	2020	*	*	***	***
HD	2019	2020	ns	ns	***	ns

Mean N/P, N/K and N/Ca ratios on the plot level were mostly in normal range during the entire study (Figures S7–S9). However, in 2020, most of the plots located in the alpine biogeographic region had excessive N/P and insufficient N/Ca ratios.

The highest share of plots with differences between defoliation categories (5 out of 28, or 17.9%) was determined for N/Ca in 2018 and N/P in 2019. The share of plots in which we found differences hardly passed 10% for other ratios and years (Table 4).

Table 4. Number of plots with significantly higher foliar ratios in each defoliation category within sampling years determined by the independent samples *t*-test (total number of plots: 28).

		N/P	N/Ca	N/Mg	N/K
2018	HD higher	-	-	1	2
	LD higher	-	5	2	-
2019	HD higher	5	1	1	2
	LD higher	-	2	1	1
2020	HD higher	1	-	-	-
	LD higher	1	2	1	-

2.3. Effects of Environmental Factors on Differences in Foliar Nutrition

While the fitted MLR (multinomial logistic regression) models for N and P did not indicate a significant influence of any of the tested factors, we found that environmental factors influence the occurrence of differences in foliar concentrations of K, Ca, and Mg between defoliation categories.

Altitude (alt) and mean annual temperature (MAT) are the only environmental factors that influence the occurrence of Differences in Foliar Concentrations (DFC) of K. Keeping all other variables constant, the odds for LD trees having higher K concentration is 0.5% higher for every meter increase in altitude. On the other hand, odds are 35% lower for HD trees to have higher K concentrations when the mean temperature increases.

Trees on higher altitudes are more likely to have higher Ca concentrations in HD trees (Table 5). An opposite effect can be seen with mean annual precipitation (MAP) where, keeping all other variables constant, with an increase in precipitation the odds are 90% higher for LD trees and 2% lower for HD trees to have higher Ca concentrations. With increasing maximum temperatures (Tmax) it is 1.3 times likely that HD trees will have significantly higher Ca concentrations.

Beech trees on higher altitudes are also more likely to have higher Mg concentrations in HD trees, but the effect on LD trees is not significant. Keeping all other variables constant, with a 1 °C increase in mean temperature it is 6.6 times likely that HD trees will have higher Mg concentrations, while at the same time the likelihood of LD having higher concentrations is negligible (Table 5).

Table 5. Results of multinomial logistic regression for Differences in Foliar Concentrations (DFC) of potassium, calcium and magnesium and corresponding model AIC. Odds ratios are reported for each environmental factor while the confidence intervals are given in the brackets (* $p < 0.1$, ** $p < 0.05$, *** $p < 0.01$).

Element	Environmental Factor ¹	DFC	
		HD Higher	LD Higher
K	Alt	1.000 (0.996, 1.003)	1.005 *** (1.002, 1.008)
	MAT	0.644 *** (0.480, 0.809)	0.884 (0.659, 1.108)
	AIC	48.042	48.042
Ca	Tmax	1.298 *** (1.112, 1.485)	0.00002 *** (0.00002, 0.00002)
	sand	0.926 (0.809, 1.043)	0.048 *** (0.048, 0.048)
	MAP	0.988 ** (0.977, 0.999)	1.900 *** (1.900, 1.900)
	Alt	1.005 ** (1.001, 1.009)	0.372 *** (0.372, 0.372)
	AIC	56.908	56.908
Mg	MAT	6.578 *** (6.330, 6.826)	1.671 * (1.149, 2.194)
	MAP	1.002 (0.996, 1.007)	1.006 (0.998, 1.015)
	Alt	1.009 *** (1.005, 1.013)	1.001 (0.992, 1.009)
	sand	0.991 (0.885, 1.097)	1.121 ** (1.020, 1.222)
	pH	2.600 ** (1.708, 3.492)	1.015 (−0.815, 2.846)
	AIC	80.311	80.311

Environmental factors ¹ Alt—altitude, MAT—Mean Annual Temperature, MAP—Mean Annual Precipitation, sand—soil sand fraction, pH—pH value, AIC—Akaike information criterion.

The results of the fitted MLR models indicate that beech on higher altitudes is less likely to have significant differences in N/P ratios and increasing maximum temperature has the same effect (Table 6). A similar effect of maximum temperatures can be seen for differences in N/Ca. Holding all other variables constant, with an increase in precipitation the odds are 0.6% higher that HD trees will have higher N/P ratios and 0.8% higher N/Mg ratios. When mean temperatures and soil pH increase, it is 2.5 and 3.7 times, respectively, more likely that LD trees will have higher N/Mg ratios. Keeping all other variables constant, with a unit increase in PDSI, i.e., less drought, the odds are 83.5% higher that LD trees will have higher N/Ca ratios. The fitted MLR models for N/K did not indicate a significant influence of any of the tested environmental factors on the occurrence probability of differences in foliar nutrient ratios.

Table 6. Results of multinomial logistic regression for Differences in Foliar Ratios (DFR) of potassium, calcium and magnesium. Odds ratios are reported for each environmental factor while the confidence intervals are given in the brackets (* $p < 0.1$, ** $p < 0.05$, *** $p < 0.01$).

	Environmental Factor ¹	DFC	
		HD Higher	LD Higher
N/P	MAP	1.006 *	0.989
		(1.000, 1.012)	(0.967, 1.011)
	Tmax	0.820 ***	0.006 ***
		(0.696, 0.943)	(−0.438, 0.449)
	Alt	0.992 **	0.960 ***
		(0.986, 0.999)	(0.937, 0.983)
	AIC	59.378	59.378
N/Ca	Tmax	0.070 ***	0.402 ***
		(−1.553, 1.693)	(0.029, 0.775)
	PDSI	1.401	1.835 **
		(−0.956, 3.758)	(1.251, 2.419)
	Ca ²⁺	0.534	0.909 **
		(−0.314, 1.381)	(0.829, 0.988)
	MAP	0.996	0.993 ***
		(0.957, 1.036)	(0.989, 0.998)
	silt	0.741	1.158 *
		(−0.353, 1.835)	(0.996, 1.319)
	AIC	72.131	72.131
N/Mg	MAP	0.999	1.008 ***
		(0.989, 1.009)	(1.002, 1.014)
	MAT	1.070	2.502 ***
		(0.648, 1.493)	(2.198, 2.805)
	pH	0.631	3.708 ***
		(−1.419, 2.681)	(2.770, 4.646)
	SPEI	2.613	0.071 **
		(−1.249, 6.474)	(−2.192, 2.333)
	AIC	65.799	65.799

Environmental factors ¹ Alt—altitude, MAT—Mean Annual Temperature, Tmax—Mean annual maximum temperature, MAP—Mean Annual Precipitation, silt—soil silt fraction, SPEI—Standardised Precipitation-Evapotranspiration Index, PDSI—Palmer Drought Severity Index, Ca²⁺—Soil exchangeable calcium, AIC—Akaike information criterion.

3. Discussion

Assessing the nutritional status of trees using the values of element concentrations in needles or leaves is a common diagnostic practice in forestry [40–42]. In Croatia there have been only a few studies into the nutritional status of beech [41,43,44], however these studies were performed on a limited research area. The results obtained in this study indicate a generally satisfactory nutrition of common beech in Croatia. Mean concentrations on the study and plot levels were within the normal range for beech [40]. Similar results were reported by Ognjenović et al. [39] from a long-term monitoring based case study of beech foliar composition on one ICP Forests intensive monitoring plot.

Large deviations from the mean concentration values of longer periods in individual years have been reported for several mineral nutrients [45–47]. In contrast, foliar K concentrations were reported to be relatively stable [39], and this finding is repeated in our study, although concentrations of all investigated elements stayed mostly within limit values [40]. Very similar relationships of N and K to limit values has also been observed in other studies [48,49]. On the other hand, P foliar levels are a reason for concern at the European level [12,15]. While beech P nutrition on a study area level seemed to be sufficient [40], on average 33% of the plots had insufficient P nutrition during the study

period (Figure S3). Additionally, the ratio of nitrogen to phosphorus shows that beech nutrition in 2019 and 2020 was not balanced in this respect, which is partly caused by diminishing P, and partly by increasing N concentrations, respectively.

The fact that we found surplus N/Mg ratios in 2019 and 2020, both due to increasing N and decreasing Mg values, echoes results of other studies that report an unbalanced Mg nutrition on national [48,50] and European level [12].

Excessive Ca concentrations on an extremely high proportion of plots (90%) were determined in a study of beech nutrition in northern Spain, but since no symptoms of unbalanced nutrition have been identified in the field, the authors believe that the deviations of the obtained values may be based on ecological and geological differences between the studied forests and central European forests on the basis of which reference values were obtained [51]. This consideration is consistent with the hypothesis that the optimal leaf composition of each species depends on the specific ecological biogeochemical niche that the species occupies [52]. Taking into account the wide ecological amplitude of common beech, there is a possibility that the optimal range of concentrations of nutrients is adapted to the specific habitat in which a particular stand develops. Thus Sardans and Peñuelas [53] assume that plant species have a certain degree of flexibility in changing stoichiometry in response to changes in environmental conditions such as climate gradients.

Nutritional status and tree crown defoliation, as indicators of vitality, can provide a basis for monitoring the effects of long-term environmental changes on forest ecosystems [22,48]. According to Simon and Wild [54], if the concentration of a certain element remains in the normal range, a decrease in mineral nutrition should be regarded more as a consequence than the cause of damage. If, on the other hand, the concentrations are inadequate, we can suspect nutrition to be the cause of damage. Although we did not find a universal relationship between the defoliation categories and nutritional status of beech trees, subsequent comparisons revealed differences in the element concentrations and ratios for certain years. For instance, we found significantly higher P concentrations in LD trees in year 2019. On the contrary, a study in northern Spain recorded higher P concentrations in more damaged trees [51]. The most pronounced differences in concentrations between defoliation categories were determined for Ca, with HD trees having significantly higher concentrations in 2018 and 2019. Higher concentrations in HD trees were also recorded for Mg in 2018. Conversely, Jonard et al. [35] report that higher defoliation values are associated with lower Ca and Mg concentrations in a liming and P/K fertilization experiment of beech stands in Belgium.

A case study investigating interrelations of various common beech vitality indicators did not find significant links between foliar nutrient concentrations and defoliation [39]. It should be taken into account that (i) the study was performed on a limited area, (ii) element concentrations were mostly within the normal range, and (iii) the range of defoliation values was narrow and seldom exceeded 25%. The lack of association between N concentrations and defoliation values, also recorded in this study, Thimonier et al. [36] attribute to the same kind of limitations. Unlike Ouimet and Moore [55], who observed that an increase in K concentration in needles resulted in a decrease in defoliation in balsam fir stands, we found no significant differences in K concentrations between defoliation categories. Compared to fertilization experiments, the analysis of the nutritional status in natural stands is more demanding due to the high variability of data and the inability to control many factors that may affect the nutritional status. Ewald [56] states heterogeneous environmental conditions as the reason for the lack of significant relationship between nutrient concentrations and spruce defoliation status in a study conducted in Bavaria. On the other hand, with this study we do have a full picture of beech nutritional status in Croatia, including the insight into the significant variations between years and defoliation categories. Significant differences in concentrations only during certain years are not a phenomenon specific to this research alone. High year-to-year variability in nutrient concentrations is characteristic of nutritional studies [15,35,48,57].

The quantification of forest ecosystem response in a climate driven changing environment is fundamental for maintenance, enhancement and restoration of future forest ecosystem goods and services. The results presented in this study are a novel approach in investigating the effects of climate and environmental properties on the defoliation/nutrition relationship. Both tree nutrition and defoliation have been reported to depend on climate properties in various studies. A study by Braun [58] showed that air temperature and precipitation have a considerable impact on the foliar nutrient concentration. Jonard et al. [49] observed a positive relationship between precipitation and foliar Ca concentrations, while insufficient water supply during dry spells have a negative impact on the uptake of calcium, according to Bergmann [59]. Our results indicate that the increase in precipitation is better utilized in low defoliation trees since the odds are 90% higher for LD trees to have higher Ca concentrations with an increase in precipitation. Lukac et al. [60] state that elements with primarily biologically controlled cycles such as nitrogen can show a different reaction to temperature changes than elements whose cycles are controlled by both biological and geological processes (P and K) or predominantly by geological processes (Ca and Mg). Therefore, it is not surprising that we did not determine a significant effect of environmental factors on the occurrence of differences in N concentrations between defoliation categories. A study of beech nutrient dynamics along a precipitation gradient found that N concentrations remain constant with decreasing precipitation while P concentrations increase, resulting in a decrease in N/P ratio [61]. However, in this study increasing precipitation increases the odds of higher N/P ratios in high defoliation trees suggesting a disturbed uptake and utilization of P in beech with higher defoliation.

From our results it becomes quite clear that an unequivocal association of foliar nutrition and crown status should not be expected for beech. It seems plausible that various pressures influence this relationship, affecting the physiological status of beech trees in a multitude of ways. For instance, Gottardini et al. [22] found that various morphological and physiological characteristics of beech leaves and canopy such as leaf volume and photosynthetic activity, had a significant negative association with the extent of damage. Additionally, Ferretti et al. [62] showed that the 25%-defoliation threshold can be a reasonable approximation for tree classification indicated by the effect of slight and moderate variations in defoliation on tree growth, which is in contrast to the results obtained by Tallieu [63]. Highly differing research results on this topic are likely depending on the research area, management practices, climate influences and other unknown factors. Nevertheless, we established that environmental factors, especially climate, influence the occurrence of differences in foliar nutrition between defoliation categories.

4. Materials and Methods

4.1. Study Area and Plot Design

The ICP Forests Level I monitoring plots in Croatia are established on intersections of a 16×16 km grid that contain forest cover. These plots do not have a fixed area; rather, 24 trees are chosen for defoliation assessments using a cross-cluster system with six trees in each cluster [64]. Only plots with a minimum of five beech trees in the sample in 2018 were selected to ensure that beech was significantly represented in the mixture of tree species, resulting in total of 28 plots (Figure 3).




Figure 3. Location of research plots and distribution of European beech in Europe [65]. Selected Level I ICP Forests monitoring plots (orange dots); remaining Level I plots in Croatia (black dots); distribution of European beech forests in Croatia [66] (gray polygon); biogeographic region in the study area (shaded polygon) [67].

4.2. Climate Data

Climate monitoring stations are generally situated at considerable distances from the forest research plots. Therefore, the data they provide are not always representative of the research locations. To overcome this, we used gridded data produced by regression kriging (RK), which is a hybrid method of interpolation carried out in four steps [68,69]. The method was validated with leave-one-out cross-validation, while the root mean square error (RMSE) was calculated between observed and interpolated values. Mean RMSEs are for mean monthly temperature from 0.5 °C to 0.9 °C, for minimum temperature from 1.1 °C to 1.5 °C, for maximum temperature from 0.7 °C to 1.1 °C, and for precipitation from 18 to 30 mm, averaged by months.

Monthly temperature and precipitation values from the gridded dataset on 1 km spatial resolution for Croatia [30] were used to calculate Mean Annual Temperature (MAT), mean annual minimum (Tmin) and mean annual maximum (Tmax) temperature, Mean Annual Precipitation (MAP) as well as the Palmer Drought Severity Index (PDSI) [70] and Standardized Precipitation Evapotranspiration Index (SPEI) [71]. Lower values of scPDSI and SPEI indicate a stronger drought intensity while higher values indicate a higher degree of humidity. SPEI was calculated on a time scale of 3 months.

4.3. Tree Selection Procedures, Foliar Sampling and Analysis

Sampling was conducted annually from 2018 to 2020 during late July and August. At a distance of less than 50 m from the centre of the plot, a stratified sampling of trees based on the percentage of defoliation was carried out, forming two groups of five trees: (i) “LOW DEFOLIATION” (LD) trees with defoliation lower than 25% and (ii) ‘HIGH DEFOLIATION’ (HD) trees with defoliation higher than 25%. Defoliation assessments were

performed during July and August according to the ICP Forests methodology [24] in steps of 5%, ranging from 0 to 100%. An absolute reference tree, defined as the best/healthiest possible beech tree, was used in the assessments, regardless of local factors that may affect the crown condition. Only dominant or codominant trees were selected, without any wounds or fruiting bodies of fungi on the trunk. In cases when trees changed their defoliation percentage and thus changed their group status (LD/HD), new, replacement trees were selected instead, and the discarded trees were not used in further sampling. Stratified sampling allowed us to circumvent the irregular distribution of tree defoliation status on plots and focus on determining whether concentrations/contents of nutrients in leaves differ in trees of different defoliation categories on each plot. Leaves were sampled from the upper third of the crown using a shotgun rifle or a rope climbing technique [72,73]. Samples from each tree were combined into a composite sample according to fresh weight and analyzed in the Laboratory for Physical and Chemical Testing (LFKI) of the Croatian Forest Research Institute. Upon arrival to the laboratory, samples of plant material were dried at 105 °C to constant mass, ground in a Fritsch Pulverisette 14 mill and prepared for analysis in a Milestone Ethos One microwave oven [74]. Concentrations of total nitrogen were determined on a Leco CNS 2000 elemental analyzer (LECO, 2000). Phosphorus (P) concentration was determined colorimetrically on a LaboMed UV/VIS spectrophotometer [74], and potassium, calcium and magnesium by atomic absorption on the Perkin-Elmer Aanalyst 700 absorption spectrophotometer [74].

4.4. Soil Sampling and Analysis

Soil sampling was performed in 2019 on five points located within each of the research plots. One point was located within each of the four clusters of trees that are assessed for defoliation during regular monitoring activities, and an additional fifth point was located in the centre of each plot. Soil samples were taken with a pedological drill from a depth of 0–10 cm, 10–20 cm, 20–40 cm, and 40–80 cm. Collected samples were pooled by sampling depth. Soil mechanical properties were determined by [75] and chemical parameters were analysed according to the following protocols and methods:

- soil pH (CaCl_2) [76]
- exchangeable K, Ca and Mg according to [77]
- exchangeable acidity (free H^+) according to [78]
- total N with an elemental analyzer Leco CNS 2000 [79]
- available P and K with the AL method [80]; P by spectrophotometry using molybdate blue method, on UV/VIS spectrophotometer LaboMed and K directly from filtrate on flame photometer Buck scientific PFP–7 [81].

4.5. Data Analysis

Descriptive statistics of foliar concentration and ratios were performed for each element. Foliar data distribution was inspected visually and by Shapiro-Wilk test. Slight deviations from normality were determined, especially for potassium, calcium and magnesium. However, studies have shown that slight deviations from normality do not affect the rate of false positive results [82].

Therefore, a two-factor repeated measures analysis of variance (ANOVA) was used to determine the difference in foliar concentrations and ratios between defoliation categories (LD/HD) of common beech on a study area level. The data of all elements met the assumption of homoscedasticity, which was confirmed by Levene's test. The Holm–Bonferroni method [83] was used to perform pairwise comparisons to determine differences in foliar concentrations (i) between LD and HD categories in single years and (iii) across defoliation categories between sampling years.

To determine differences in foliar concentrations and ratios between defoliation categories on a plot level, we used the independent samples t-test [84]. The results were then categorized for each plot and sampling year to reflect three cases: (i) *HD_higher* if HD trees had significantly higher foliar concentrations ($p < 0.05$), (ii) *LD_higher* if LD trees

had significantly higher foliar concentrations ($p < 0.05$) and (iii) *NO_diff* if there were no significant differences in foliar concentrations between defoliation categories, thus creating a variable which represents Differences in Foliar Concentrations or Ratios for each element (DFC/DFR).

Multinomial logistic regression (MLR) was used to determine which environmental factors influence the occurrence probability of a certain DFC/DFR category, except *no_dif* which was set as the reference category. Separate odds ratios, which are the exponentiation of model coefficients, were determined for all independent variables for each DFC/DFR category. The odds ratio of a coefficient indicates the occurrence probability of either HD or LD trees having significantly higher foliar concentrations over the probability of no difference between defoliation categories (*NO_diff*). An odds ratio > 1 indicates that the occurrence probability of a certain DFC/DFR category relative to the occurrence of *NO_diff* increases as the independent variable increases, whereas an odds ratio < 1 indicates that the occurrence of a certain DFC/DFR category decreases. Of the potential independent variables, those with the highest variable importance were selected with the random forest algorithm [85]. The final model selection process was based on diagnostic diagrams and a procedure defined by Johnson et al. [86]. All analyses were conducted in an R programming environment [87].

5. Conclusions

Our final view is that links between defoliation and nutrition of beech in Croatia are not very direct or very prominent; this can be seen from (I) the fact that differences in nutrition of LD and HD trees were found for only a few nutrients (II) these effects are mostly not universal, but present only in some years and on a limited number of plots and (III) nutrient concentrations mostly stay within normal values regardless of tree defoliation status. However, it is clear that environmental factors, especially climate properties, do affect the relationship between defoliation and nutrition. Discovering mechanisms by which environmental factors affect foliar nutrition should complement current research on the defoliation–nutrition relationships.

Supplementary Materials: The following supporting information can be downloaded at: <https://www.mdpi.com/article/10.3390/plants12010168/s1>, Figure S1. Comparison of study area climate properties during the vegetation period with the climate normal (1980–2011) of the vegetation period (March–September); Figure S2. Foliar nitrogen concentrations on research plots; Figure S3. Foliar phosphorus concentrations on research plots; Figure S4. Foliar potassium concentrations on research plots; Figure S5. Foliar calcium concentrations on research plots; Figure S6. Foliar magnesium concentrations on research plots; Figure S7. Foliar N/P ratios on research plots; Figure S8. Foliar N/Ca ratios on research plots; Figure S9. Foliar N/K ratios on research plots; Figure S10. Foliar N/Mg ratios on research plots; Figure S11. Defoliation of common beech on research plots; Figure S12. Defoliation of common beech on all plots during the study period.

Author Contributions: Conceptualization, M.O., I.S. and N.P.; methodology, M.O.; formal analysis, M.O.; investigation, M.O., I.S., P.R. and N.P.; resources, N.P.; data curation, M.O.; writing—original draft preparation, M.O., I.S. and N.P.; writing—review and editing, M.O., I.S., M.M., P.R., M.J., V.T., M.L., M.P.T., D.U. and N.P.; visualization, M.O.; supervision, N.P.; project administration, M.O. and M.M.; funding acquisition, M.O., I.S. and N.P. All authors have read and agreed to the published version of the manuscript.

Funding: This work has been fully supported by Croatian Science Foundation under the project VitaClim (IP-2018-01-5222).

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: The data presented in this study are available on request from the corresponding author. The data are not publicly available due to legal reasons.

Acknowledgments: We highly appreciate the work of researchers and technicians of the Croatian Forest Research Institute that participated in field research and laboratory analysis for this study.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Centritto, M.; Tognetti, R.; Leitgeb, E.; Střelcová, K.; Cohen, S. Above Ground Processes: Anticipating Climate Change Influences. In *Forest Management and the Water Cycle: An Ecosystem-Based Approach*; Bredemeier, M., Cohen, S., Godbold, D.L., Lode, E., Pichler, V., Schleppi, P., Eds.; Springer: Dordrecht, The Netherlands, 2011; pp. 31–64. ISBN 978-90-481-9834-4.
2. Trumbore, S.; Brando, P.; Hartmann, H. Forest Health and Global Change. *Science* **2015**, *349*, 814–818. [[CrossRef](#)]
3. Stocker, T.F.; Qin, D.; Plattner, G.-K.; Alexander, L.V.; Allen, S.K.; Bindoff, N.L.; Bréon, F.-M.; Church, J.A.; Cubasch, U.; Emori, S.; et al. Technical Summary. In *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*; Stocker, T.F., Qin, D., Plattner, G.-K., Tignor, M., Allen, S.K., Boschung, J., Nauels, A., Xia, Y., Bex, V., Midgley, P.M., Eds.; Cambridge University Press: Cambridge, UK; New York, NY, USA, 2013.
4. de Vries, W.; Dobbertin, M.H.; Solberg, S.; van Dobben, H.F.; Schaub, M. Impacts of Acid Deposition, Ozone Exposure and Weather Conditions on Forest Ecosystems in Europe: An Overview. *Plant Soil* **2014**, *380*, 1–45. [[CrossRef](#)]
5. Ferretti, M.; Waldner, P.; Verstraeten, A.; Schmitz, A.; Michel, A.; Žlindra, D.; Marchetto, A.; Hansen, K.; Pitar, D.; Gottardini, E.; et al. Criterion 2: Maintenance of Forest Ecosystem Health and Vitality. In *Forest Europe, 2020: State of Europe's Forests 2020*; Ministerial Conference on the Protection of Forests in Europe—Liaison Unit Bratislava: Zvolen, Slovak Republic, 2020.
6. Zierl, B. A Simulation Study to Analyse the Relations between Crown Condition and Drought in Switzerland. *For. Ecol. Manag.* **2004**, *188*, 25–38. [[CrossRef](#)]
7. IPCC. Summary for Policymakers. In *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*; Masson-Delmotte, V., Zhai, P., Pirani, A., Connors, S.L., Péan, C., Berger, S., Caud, N., Chen, Y., Goldfarb, L., Gomis, M.I., et al., Eds.; Cambridge University Press: Cambridge, UK; New York, NY, USA, 2021; pp. 3–32. [[CrossRef](#)]
8. Allen, C.D.; Macalady, A.K.; Chenchouni, H.; Bachelet, D.; McDowell, N.; Vennetier, M.; Kitzberger, T.; Rigling, A.; Breshears, D.D.; Hogg, E.H.; et al. A Global Overview of Drought and Heat-Induced Tree Mortality Reveals Emerging Climate Change Risks for Forests. *For. Ecol. Manag.* **2010**, *259*, 660–684. [[CrossRef](#)]
9. Giorgi, F. Climate Change Hot-Spots. *Geophys. Res. Lett.* **2006**, *33*. [[CrossRef](#)]
10. Beniston, M.; Stephenson, D.B.; Christensen, O.B.; Ferro, C.A.T.; Frei, C.; Goyette, S.; Halsnaes, K.; Holt, T.; Jylhä, K.; Koffi, B.; et al. Future Extreme Events in European Climate: An Exploration of Regional Climate Model Projections. *Clim. Change* **2007**, *81*, 71–95. [[CrossRef](#)]
11. Marschner, H. 9—Functions of Mineral Nutrients: Micronutrients. In *Marschner's Mineral Nutrition of Higher Plants*, 2nd ed.; Academic Press: San Diego, CA, USA, 2002; pp. 313–404. ISBN 978-0-12-473543-9.
12. Jonard, M.; Fürst, A.; Verstraeten, A.; Thimonier, A.; Timmermann, V.; Potočić, N.; Waldner, P.; Benham, S.; Hansen, K.; Merilä, P.; et al. Tree Mineral Nutrition Is Deteriorating in Europe. *Glob. Chang. Biol.* **2015**, *21*, 418–430. [[CrossRef](#)]
13. Hallenbarter, D.; Landolt, W.; Bucher, J. Nutrition and Vitality: Phenological Investigation in Forest under Different Fertiliser Treatments. In *Trends in European Forest Tree Physiology Research: Cost Action E6: EUROSILVA*; Springer: Dordrecht, The Netherlands, 2001; pp. 9–12.
14. Kreuzwieser, J.; Gessler, A. Global Climate Change and Tree Nutrition: Influence of Water Availability. *Tree Physiol.* **2010**, *30*, 1221–1234. [[CrossRef](#)]
15. Talkner, U.; Meiwes, K.J.; Potočić, N.; Seletković, I.; Cools, N.; De Vos, B.; Rautio, P. Phosphorus Nutrition of Beech (*Fagus sylvatica* L.) Is Decreasing in Europe. *Ann. For. Sci.* **2015**, *72*, 919–928. [[CrossRef](#)]
16. Potočić, N.; Timmermann, V.; Ognjenović, M.; Kirchner, T.; Prescher, A.K.; Ferretti, M. *Tree Health Is Deteriorating in the European Forests*; ICP Forests: 2021; Programme Coordinating Centre of ICP Forests, Thünen, Institute of Forest Ecosystems: Eberswalde, Germany, 2021. [[CrossRef](#)]
17. Pernar, N.; Bakšić, D. The Soils of Beech Forests. In *Common Beech (Fagus sylvatica L.) in Croatia*; Matić, S., Ed.; Academy of Forestry Sciences: Zagreb, Croatia, 2003; ISBN 953-98571-1-2.
18. Dobbertin, M.; Brang, P. Crown Defoliation Improves Tree Mortality Models. *For. Ecol. Manag.* **2001**, *141*, 271–284. [[CrossRef](#)]
19. Cherubini, P.; Battipaglia, G.; Innes, J.L. Tree Vitality and Forest Health: Can Tree-Ring Stable Isotopes Be Used as Indicators? *Curr. For. Rep.* **2021**, *7*, 69–80. [[CrossRef](#)]
20. De Vries, W.; Klap, J.M.; Erisman, J.W. Effects of Environmental Stress on Forest Crown Condition in Europe. Part I: Hypotheses and Approach to the Study. *Water. Air. Soil Pollut.* **2000**, *119*, 317–333. [[CrossRef](#)]
21. De Marco, A.; Proietti, C.; Cionni, I.; Fischer, R.; Screpanti, A.; Vitale, M. Future Impacts of Nitrogen Deposition and Climate Change Scenarios on Forest Crown Defoliation. *Environ. Pollut.* **2014**, *194*, 171–180. [[CrossRef](#)]
22. Gottardini, E.; Cristofolini, F.; Cristofori, A.; Pollastrini, M.; Camin, F.; Ferretti, M. A Multi-Proxy Approach Reveals Common and Species-Specific Features Associated with Tree Defoliation in Broadleaved Species. *For. Ecol. Manag.* **2020**, *467*, 118151. [[CrossRef](#)]
23. Dobbertin, M. Tree Growth as Indicator of Tree Vitality and of Tree Reaction to Environmental Stress: A Review. *Eur. J. For. Res.* **2005**, *124*, 319–333. [[CrossRef](#)]

24. Eichhorn, J.; Roskams, P.; Potočić, N.; Timmermann, V.; Ferretti, M.; Mues, V.; Szepesi, A.; Durrant, D.; Seletković, I.; Schroeck, H.-W.; et al. Part IV: Visual Assessment of Crown Condition and Damaging Agents. Version 2020-3. In *Manual on Methods and Criteria For Harmonized Sampling, Assessment, Monitoring and Analysis of the Effects of Air Pollution on Forests*; Thünen Institute of Forest Ecosystems: Eberswalde, Germany, 2020; p. 49. ISBN 978-3-86576-162-0.
25. Lorenz, M. International Co-Operative Programme on Assessment and Monitoring of Air Pollution Effects on Forests-ICP Forests. *Water, Air, Soil Pollut.* **1995**, *85*, 1221–1226. [[CrossRef](#)]
26. Timmermann, V.; Potočić, N.; Ognjenović, M.; Kirchner, T. *Tree Crown Condition in 2021. Forest Condition in Europe: The 2022 Assessment. ICP Forests technical Report under the UNECE Convention on Long-Range Transboundary Air Pollution (Air Convention)*; Thünen Institute: Eberswalde, Germany, 2022; pp. 41–55.
27. Ciais, P.; Reichstein, M.; Viovy, N.; Granier, A.; Ogée, J.; Allard, V.; Aubinet, M.; Buchmann, N.; Bernhofer, C.; Carrara, A.; et al. Europe-Wide Reduction in Primary Productivity Caused by the Heat and Drought in 2003. *Nature* **2005**, *437*, 529–533. [[CrossRef](#)]
28. Seidling, W. Signals of Summer Drought in Crown Condition Data from the German Level I Network. *Eur. J. For. Res.* **2007**, *126*, 529–544. [[CrossRef](#)]
29. Seletković, I.; Potočić, N.; Ugarković, D.; Jazbec, A.; Pernar, R.; Seletković, A.; Benko, M. Climate and Relief Properties Influence Crown Condition of Common Beech (*Fagus sylvatica* L.) on the Medvednica Massif. *Period. Biol.* **2009**, *111*, 435–441.
30. Ognjenović, M.; Seletković, I.; Potočić, N.; Marušić, M.; Tadić, M.P.; Jonard, M.; Rautio, P.; Timmermann, V.; Lovreškov, L.; Ugarković, D. Defoliation Change of European Beech (*Fagus Sylvatica* L.) Depends on Previous Year Drought. *Plants* **2022**, *11*, 730. [[CrossRef](#)]
31. Gefšler, A.; Keitel, C.; Kreuzwieser, J.; Matyssek, R.; Seiler, W.; Rennenberg, H. Potential Risks for European Beech (*Fagus Sylvatica* L.) in a Changing Climate. *Trees-Struct. Funct.* **2007**, *21*, 1–11. [[CrossRef](#)]
32. Leuschner, C. Drought Response of European Beech (*Fagus sylvatica* L.)—A Review. *Perspect. Plant Ecol. Evol. Syst.* **2020**, *47*, 125576. [[CrossRef](#)]
33. Seletković, Z.; Tikvić, I.; Prpić, B. Ekološka Konstitucija Obične Bukve. In *Obična Bukva (Fagus sylvatica L.) u Hrvatskoj*; Matić, S., Ed.; Akademija Šumarskih Znanosti: Zagreb, Croatia, 2003.
34. Bolte, A.; Czajkowski, T.; Coccozza, C.; Tognetti, R.; De Miguel, M.; Pšidová, E.; Ditmarová, L.; Dinca, L.; Delzon, S.; Cochard, H.; et al. Desiccation and Mortality Dynamics in Seedlings of Different European Beech (*Fagus sylvatica* L.) Populations under Extreme Drought Conditions. *Front. Plant Sci.* **2016**, *7*, 751. [[CrossRef](#)] [[PubMed](#)]
35. Jonard, M.; André, F.; Giot, P.; Weissen, F.; Van der Perre, R.; Ponette, Q. Thirteen-Year Monitoring of Liming and PK Fertilization Effects on Tree Vitality in Norway Spruce and European Beech Stands. *Eur. J. For. Res.* **2010**, *129*, 1203–1211. [[CrossRef](#)]
36. Thimonier, A.; Graf Pannatier, E.; Schmitt, M.; Waldner, P.; Walthert, L.; Schleppe, P.; Dobbertin, M.; Kräuchi, N. Does Exceeding the Critical Loads for Nitrogen Alter Nitrate Leaching, the Nutrient Status of Trees and Their Crown Condition at Swiss Long-Term Forest Ecosystem Research (LWF) Sites? *Eur. J. For. Res.* **2010**, *129*, 443–461. [[CrossRef](#)]
37. Ferretti, M.; Calderisi, M.; Marchetto, A.; Waldner, P.; Thimonier, A.; Jonard, M.; Cools, N.; Rautio, P.; Clarke, N.; Hansen, K.; et al. Variables Related to Nitrogen Deposition Improve Defoliation Models for European Forests. *Ann. For. Sci.* **2015**, *72*, 897–906. [[CrossRef](#)]
38. Toigo, M.; Nicolas, M.; Jonard, M.; Croisé, L.; Nageleisen, L.M.; Jactel, H. Temporal Trends in Tree Defoliation and Response to Multiple Biotic and Abiotic Stresses. *For. Ecol. Manag.* **2020**, *477*, 118476. [[CrossRef](#)]
39. Ognjenović, M.; Levanič, T.; Potočić, N.; Ugarković, D.; Indir, K.; Seletković, I. Interrelations of Various Tree Vitality Indicators and Their Reaction to Climatic Conditions on a European Beech (*Fagus sylvatica* L.) Plot. *Šumarski List* **2020**, *144*, 351–365. [[CrossRef](#)]
40. Mellert, K.H.; Göttlein, A. Comparison of New Foliar Nutrient Thresholds Derived from van Den Burg's Literature Compilation with Established Central European References. *Eur. J. For. Res.* **2012**, *131*, 1461–1472. [[CrossRef](#)]
41. Seletković, I.; Potočić, N. Stanje Ishrane Obične Bukve (*Fagus sylvatica* L.) i Obične Jele (*Abies alba* Mill.) Na Području Medvednice. *Rad. Šum. Inst. Izvanredno Izd.* **2006**, *9*, 117–125.
42. Rautio, H. Chemical Needle Analysis as a Diagnostic and Monitoring Method. In *Nutrient Uptake and Cycling in Forest Ecosystems*; Nilsson, L.O., Hüttel, R.F., Johansson, U.T., Eds.; Kluwer Academic Publishers: Dordrecht, The Netherlands, 1993; Volume 197–202, ISBN 0792330307.
43. Komlenović, N.; Gračan, J.; Pezdric, N.; Rastovski, P. Utjecaj Polutanata Na Bukove Šume i Kulture Smreke u Sjeverozapadnoj Hrvatskoj. *Šumarski List* **1988**, *112*, 12.
44. Komlenović, N.; Matković, N.; Močan, D.; Rastovski, P. Unos Onečišćenja Iz Zraka u Šumu Bukve i Jele (Abieti-Fagetum "Dinaricum") u Predjelu Lividrage u Zapadnoj Hrvatskoj. *Šumarski List* **1997**, *121*, 8.
45. Sauter, U. Zeitliche Variationen Des Ernährungszustands Nordbayerischer Kiefernbestände. *Forstwissenschaftliches Cent. Ver. mit Tharandter Forstl. Jahrb.* **1991**, *110*, 13–33. [[CrossRef](#)]
46. Prietzel, J.; Kolb, E.; Rehfuess, K.E. Langzeituntersuchung Ehemals Streugennutzter Kiefernökosysteme in der Oberpfalz: Veränderungen von Bodenchemischen Eigenschaften und der Nährelementversorgung der Bestände. *Forstwissenschaftliches Cent. Ver. mit Tharandter Forstl. Jahrb.* **1997**, *116*, 269–290. [[CrossRef](#)]
47. Stefan, K.; Fürst, A.; Hacker, R.; Bartels, U. *Forest Foliar Condition in Europe. Results of Large-Scale Foliar Chemistry Surveys*; European Commission—United Nations/Economic Commission for Europe: Brussels, Belgium, 1997.
48. Duquesnay, A.; Dupouey, J.-L.; Clement, A.; Ulrich, E.; Le Tacon, F. Spatial and Temporal Variability of Foliar Mineral Concentration in Beech (*Fagus Sylvatica*) Stands in Northeastern France. *Tree Physiol.* **2000**, *20*, 13–22. [[CrossRef](#)]

49. Jonard, M.; André, F.; Dambrine, E.; Ponette, Q.; Ulrich, E. Temporal Trends in the Foliar Nutritional Status of the French, Walloon and Luxembourg Broad-Leaved Plots of Forest Monitoring. *Ann. For. Sci.* **2009**, *66*, 412. [CrossRef]
50. Amores, G.; Santamaría, J.M. Beech Foliar Chemical Composition: A Bioindicator of Air Pollution Stress. In *Developments in Environmental Science*; Elsevier: Amsterdam, The Netherlands, 2003; Volume 3, pp. 301–313. ISBN 1474-8177.
51. Amores, G.; Bermejo, R.; Elustondo, D.; Lasheras, E.; Santamaría, J.M. Nutritional Status of Northern Spain Beech Forests. *Water Air Soil Pollut.* **2006**, *177*, 227–238. [CrossRef]
52. Penuelas, J.; Sardans, J.; Ogaya, R.; Estiarte, M. Nutrient Stoichiometric Relations and Biogeochemical Niche in Coexisting Plant Species: Effect of Simulated Climate Change. *Polish J. Ecol.* **2008**, *56*, 613–622.
53. Sardans, J.; Peñuelas, J. Tree Growth Changes with Climate and Forest Type Are Associated with Relative Allocation of Nutrients, Especially Phosphorus, to Leaves and Wood. *Glob. Ecol. Biogeogr.* **2013**, *22*, 494–507. [CrossRef]
54. Simon, A.; Wild, A. Mineral Nutrients in Leaves and Bast of Pedunculate Oak (*Quercus robur* L.) at Different States of Defoliation. *Chemosphere* **1998**, *36*, 955–959. [CrossRef]
55. Ouimet, R.; Moore, J.-D. Effects of Fertilization and Liming on Tree Growth, Vitality and Nutrient Status in Boreal Balsam Fir Stands. *For. Ecol. Manag.* **2015**, *345*, 39–49. [CrossRef]
56. Ewald, J. Ecological Background of Crown Condition, Growth and Nutritional Status of *Picea Abies* (L.) Karst. in the Bavarian Alps. *Eur. J. For. Res.* **2005**, *124*, 9–18. [CrossRef]
57. Potočić, N.; Čosić, T.; Pilaš, I. The Influence of Climate and Soil Properties on Calcium Nutrition and Vitality of Silver Fir (*Abies Alba* Mill.). *Environ. Pollut.* **2005**, *137*, 596–602. [CrossRef] [PubMed]
58. Braun, S.; Schindler, C.; Rihm, B. Foliar Nutrient Concentrations of European Beech in Switzerland: Relations with Nitrogen Deposition, Ozone, Climate and Soil Chemistry. *Front. For. Glob. Chang.* **2020**, *3*, 33. [CrossRef]
59. Bergmann, W. (Ed.) *Nutritional Disorders of Plants*; Gustav Fischer Verlag: Jena, Germany; Stuttgart, Germany; New York, NY, USA, 1992.
60. Lukac, M.; Calfapietra, C.; Lagomarsino, A.; Loreto, F. Global Climate Change and Tree Nutrition: Effects of Elevated CO₂ and Temperature. *Tree Physiol.* **2010**, *30*, 1209–1220. [CrossRef]
61. Meier, I.C.; Leuschner, C. Nutrient Dynamics along a Precipitation Gradient in European Beech Forests. *Biogeochemistry* **2014**, *120*, 51–69. [CrossRef]
62. Ferretti, M.; Bacaro, G.; Brunialti, G.; Calderisi, M.; Croisé, L.; Frati, L.; Nicolas, M. Tree Canopy Defoliation Can Reveal Growth Decline in Mid-Latitude Temperate Forests. *Ecol. Indic.* **2021**, *127*, 107749. [CrossRef]
63. Tallieu, C.; Badeau, V.; Allard, D.; Nageleisen, L.M.; Bréda, N. Year-to-Year Crown Condition Poorly Contributes to Ring Width Variations of Beech Trees in French ICP Level I Network. *For. Ecol. Manag.* **2020**, *465*, 118071. [CrossRef]
64. Ferretti, M.; Fischer, R.; Mues, V.; Granke, O.; Lorenz, M.; Seidling, W.; Nicolas, M. Part II: Basic Design principles for the ICP Forests Monitoring Networks. Version 2020-2. In *Manual on Methods and Criteria for Harmonized Sampling, Assessment, Monitoring and Analysis of the Effects of Air Pollution On forests*; Thünen Institute of Forest Ecosystems: Eberswalde, Germany, 2020; p. 33. ISBN 978-3-86576-162-0.
65. Caudullo, G.; Welk, E.; San-Miguel-Ayanz, J. Chorological Maps for the Main European Woody Species. *Data Br.* **2017**, *12*, 662–666. [CrossRef]
66. Pravilnik o Vrstama Stanišnih Tipova, Karti Staništa, Ugroženim i Rijetkim Stanišnim Tipovima Te o Mjerama Za Očuvanje Stanišnih Tipova NN 7/2006. Available online: https://narodne-novine.nn.hr/clanci/sluzbeni/2006_01_7_156.html (accessed on 14 July 2022).
67. Cervellini, M.; Zannini, P.; Di Musciano, M.; Fattorini, S.; Jiménez-Alfaro, B.; Rocchini, D.; Field, R.; Vetaas, O.R.; Irl, S.D.H.; Beierkuhnlein, C. A Grid-Based Map for the Biogeographical Regions of Europe. *Biodivers. Data J.* **2020**, *8*, e53720. [CrossRef]
68. Perčec Tadić, M. Gridded Croatian Climatology for 1961–1990. *Theor. Appl. Climatol.* **2010**, *102*, 87–103. [CrossRef]
69. Perčec Tadić, M.; Pasarić, Z.; Guijarro, J.A. Croatian High-Resolution Monthly Gridded Data Set of Homogenised Surface Air Temperature. *Theor. Appl. Climatol.* **2022**. [CrossRef]
70. Palmer, W.C. *Meteorological Drought*; US Department of Commerce, Weather Bureau: Washington, DC, USA, 1965; Volume 45.
71. Vicente-Serrano, S.M.; Beguería, S.; López-Moreno, J.I. A Multiscalar Drought Index Sensitive to Global Warming: The Standardized Precipitation Evapotranspiration Index. *J. Clim.* **2010**, *23*, 1696–1718. [CrossRef]
72. Dial, R.; Tobin, S.C. Description of Arborist Methods for Forest Canopy Access and Movement. *Selbyana* **1994**, *15*, 24–37.
73. Jepson, J. *The Tree Climber's Companion*; Beaver Tree Publishing: Longville, MI, USA, 2000.
74. Rautio, P.; Fürst, A.; Stefan, K.; Raitio, H.; Bartels, U. Part XII: Sampling and Analysis of Needles and Leaves. In *Manual on Methods and Criteria for Harmonized Sampling, Assessment, Monitoring and Analysis of the Effects of Air Pollution on Forests*; UNECE, ICP Forests Programme Co-Ordinating Centre: Hamburg, Germany, 2016.
75. ISO 11277; Soil Quality—Determination of Particle Size Distribution in Mineral Soil Material. ISO: Geneva, Switzerland, 2020.
76. ISO-10390; Soil Quality—Determination of PH. ISO: Geneva, Switzerland, 2005.
77. ISO-11260; Soil Quality—Determination of Effective Cation Exchange Capacity and Base Saturation Level Using Barium Chloride Solution. ISO: Geneva, Switzerland, 1994.
78. ISO 14254; Soil Quality—Determination of Exchangeable Acidity Using Barium Chloride Solution as Extractant 2018. ISO: Geneva, Switzerland, 2018.

79. ISO-13878; Soil Quality—Determination of Total Nitrogen Content by Dry Combustion (“elemental Analysis”) 1998. ISO: Geneva, Switzerland, 1998.
80. Egnér, H.; Riehm, H.; Domingo, W.R. Untersuchungen Über Die Chemische Bodenanalyse Als Grundlage Für Die Beurteilung Des Nährstoffzustandes Der Böden. II. Chemische Extraktionsmethoden Zur Phosphor-Und Kaliumbestimmung. *K. Lantbr. Ann.* **1960**, *26*, 199–215.
81. Škorić, A. *Priručnik Za Pedološka Istraživanja*; Fakultet Poljoprivrednih Znanosti: Zagreb, Croatia, 1985.
82. Lix, L.M.; Keselman, J.C.; Keselman, H.J. Consequences of Assumption Violations Revisited: A Quantitative Review of Alternatives to the One-Way Analysis of Variance F Test. *Rev. Educ. Res.* **1996**, *66*, 579–619. [[CrossRef](#)]
83. Holm, S. A Simple Sequentially Rejective Multiple Test Procedure. *Scand. J. Stat.* **1979**, *6*, 65–70.
84. Ross, A.; Willson, V.L. Independent Samples T-Test. In *Basic and Advanced Statistical Tests*; Springer: Rotterdam, The Netherlands, 2017; pp. 13–16.
85. Breiman, L. Random Forests. *Mach. Learn.* **2001**, *45*, 5–32. [[CrossRef](#)]
86. Johnson, J.B.; Omland, K.S. Model Selection in Ecology and Evolution. *Trends Ecol. Evol.* **2004**, *19*, 101–108. [[CrossRef](#)]
87. R Core Team. *R: A Language and Environment for Statistical Computing*; R Foundation for Statistical Computing: Vienna, Austria, 2016.

Disclaimer/Publisher’s Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.