

Impacts of Reforestation on Stabilization of Riverine Water Levels in South Korea[†]

By JAEHYUN YOON, SAANG JOON BAAK,
MIN YOUNG SEO, AND TAEJONG KIM*

We investigate how reforestation contributed to stabilization of riverine water levels in South Korea. For the purpose, we estimate an equation capturing dynamic relationships among rainfall, upstream-area tree stock, and downstream water levels in three river systems of Hongcheon, Mangyeong, and Hyeongsan, using daily observations of precipitation and water levels for the period from 1985 to 2005. Simulation based on estimation results shows that increase in the tree stock in a river basin leads to a significantly suppressed peaking in riverine water levels in response to an abrupt and concentrated rain in the upstream area. For instance, an hour-long concentration of 100mm rain results in 0.7m rise in water level if the volume of growing stock is 1 million m³, whereas the rise in water level stays below 0.27m with 5 million m³ in the growing-stock volume.

Key Word: Reforestation, Forest Management, Ecosystem Services,
Flood Control, Sustainability

JEL Code: Q51, Q54, Q56, Q57

I. Introduction

This paper investigates the effects of reforestation on stabilization of riverine water levels, one of the expected benefits from forests, using a panel data set from South Korea.

* Yoon: Assistant Professor, Faculty of Political Science and Economics, Waseda University (E-mail: j.yoon@aoni.waseda.jp); Baak: Professor, School of International Liberal Studies, Waseda University (E-mail: baak@waseda.jp); Seo: Head of Research Planning Division, KDI School of Public Policy and Management (E-mail: myseo@kdischool.ac.kr); Kim (Corresponding Author): Professor, KDI School of Public Policy and Management (E-mail: tjkim@kdischool.ac.kr).

* Received: 2022. 2. 26

* Referee Process Started: 2022. 3. 3

* Referee Reports Completed: 2022. 6. 14

[†] The authors thank Toshihide Arimura, Tatsuhito Kondo, Miwa Nakai, Taisuke Sadayuki, and two anonymous reviewers for their valuable comments. The authors also benefited from the comments of participants of the 2020 spring conference of the Japanese Economic Association and the 2016 Asian Historical Economics Conference. The authors are grateful to the KDI School of Public Policy and Management for financial support. All errors are the authors' alone.

Forest is known to store and conserve water flow from rainfall mainly through canopy interception and soil infiltration. The canopy interception refers to water that is retained by leaves of trees and evaporates into air. Besides the canopy interception, the rainfall can be delayed in reaching soil through stemflow, which refers to water that trickles along branches. The water that reached soil can either run off as overflow or be absorbed into the underground through soil infiltration. This water-regulating function of forests is believed to contribute to preventing natural disasters such as flood and drought and enhancing agricultural productivity by steadily providing water to nearby lands.

Despite such a hydrological service and other benefits of forest to human, however, the world is continuously losing forests, especially in developing countries. According to FAO (2015), the global net loss of forests between 1990 and 2015 is 129 million hectares which is almost the size of South Africa, and the largest area of deforestation is found in South America and Africa. As Wheeler *et al.* (2013) among others show, deforestation is positively related to economic benefit of forest conversion to other land use in the process of economic development. According to Wheeler *et al.* (2013), forest clearing in Indonesian kabupatens between 2005 and 2010 was affected by economic variables such as interest rate and exchange rate. Faria *et al.* (2016) show that deforestation in the Brazilian Amazon between 2000 and 2010 has a positive relationship with trade openness. Li *et al.* (2017) further investigated the relationship between trade and forest transition with the data of nine Asian countries and reported that not only total trade volume but also trade structure affected forest transition.

Since the benefits of forest to human have externalities, however, conservationists believe that the benefits of forest protection tend to be undervalued compared to the benefits of forest conversion (Sims, 2010). As a result, conversion of forests to other land use is often made at the level higher than the optimal point. For example, when Sims (2010) compared consumption and poverty measures between localities with a high or low share of protected forest areas in Thailand, she found a positive correlation between poverty and protection. On the contrary, when a selection bias was taken into consideration in the estimation, protection turned out to reduce poverty. She explains the change in the estimation results by the ecotourism, an externality of forest. Klemick (2011) found that forest fallow significantly contributes to productivity of adjacent farms. According to the calculation of Chang *et al.* (2016), deforestation due to international trade caused net economic loss if the lost externalities of ecosystem is considered. Despite the increasing number of research papers exploring explicit and implicit benefits of forest to human, however, as Ferraro *et al.* (2012) and Tan-Soo *et al.* (2016) argue, our understanding of forest benefits is still quite limited.

Overall, that economic benefits from forests far outweigh costs at global and regional levels is well established beyond dispute (MEA, 2003; TEEB 2010; and Neudert *et al.*, 2016). The main challenge in forestation and forest conservation in the developing world is that, at the local level, the benefits may fall short of the opportunity costs borne by the members of the local community (Neudert *et al.*, 2016). Indeed, researchers based at the Korea Forest Research Institute routinely update their estimates of “nonmarket” values of the forests in South Korea. According to their most recent estimate, the positive externalities generated by the

Korean forest is greater than 73 trillion won, or more than 7% of GDP, in 2008. The estimate doesn't include the value of marketable products from the forests such as timber and some important non-market values such as conservation of biodiversity. Even though the budget the Korean government allocated to promote reforestation in the 60s and 70s accounted for a fairly large portion given the size of the overall government budget, it is clear that the cumulative benefits are much larger than the initial investments at the national level.¹ Thus, it should be noted that we are not conducting the current study with a view to establishing that the Korean reforestation resulted in net welfare improvement. Against this background, the present paper's aim is quite modest: we want to demonstrate that the water storage and conservation service of the forests is real and substantial, thus contributing to the growing body of literature documenting forests' benefits more and more rigorously.

Because Korea suffered flood damages on a routine basis due to irregular precipitation throughout its history, the South Korean government emphasized the water flow regulation function of forest to justify its national reforestation program from the initiative stage of the program expecting more stable water flow would enhance agricultural productivity and prevent flood (FAO, 2016). Even so, to our surprise, the effects of reforestation on water flow have never been scientifically investigated in the literature despite South Korea's successful reforestation.

In fact, flood damage has decreased in South Korea for the last few decades, while agricultural productivity has increased. Figure 1 shows that the size of flooded area has substantially decreased although precipitation in summer has not at all. The average precipitation from 1973 to 1986 was 626 millimeter, and it was 765 millimeter from 2000 to 2014. However, the annual average size of flooded area has decreased between the two time periods, from 81 thousand to 23 thousand hectares. In addition, the annual average rice product per ten acre was 423 kilograms in the former time period, while it was 496 kilograms in the latter time period (see Figure 2).

FAO (2016), which highly admired Korea's reforestation, claimed the changes described above as main benefits of Korean reforestation. Seo (2018) examined the relationship between rice production and reforestation using the panel data of rice fields and adjacent forests in South Korea, and concluded that reforestation positively affected rice productivity. However, because not only reforestation but also many other factors such as construction of dams and banks and improvement in chemicals and farm machinery should also have contributed to such changes, to isolate the contribution of reforestation is not an easy task. Even so, if the water stabilization function of forest is confirmed by data, the claims of FAO (2016) and Seo (2018) will be buttressed.

In particular, this research paper collects rainfall and water level data of three Korean rivers, and examines whether and to what extent the impact of rainfall on water level of river changes as the volume of growing stock of adjacent forest grows. To the best of our knowledge, no research paper has explicitly investigated whether

¹For instance, in 1973, the government budget for reforestation programs amounted to about 10 billion won, whereas the overall government expenditure was about 659 billion won. The share of the reforestation budget was thus about 1.5% of the overall budget. For reference, the consumer price index in 2020 was about 16 times the level in 1973. More detailed information on reforestation budgets may be found in the Korea Association for Public Administration's report of 2009 (Korea Association for Public Administration, 2009).




FIGURE 1. PRECIPITATION AND FLOOD DAMAGE

Source: Hydrologic Annual Report in Korea (Ministry of Land, Infrastructure and Transport), Disaster Report (Ministry of the Interior and Safety).




FIGURE 2. RICE PRODUCTION OF SOUTH KOREA

Source: KOSIS, Statistics Annual Report of Local Government.

the impact of rainfall on water level of river declines as reforestation proceeds. In fact, only limited amount of literature has examined the impacts of change in forest such as reforestation, afforestation and deforestation on ground water flow in general.

Among those small number of exceptions, Hundecha and Bardossy (2004) demonstrated that intense afforestation reduces volume of runoff, which refers to water flows over the earth's surface due to storm or other weather conditions with intense rainfall. The reduced runoff would delay sudden increase in water level of river and thus prevent flood. Nisbet *et al.* (2004) showed that planting woods along a 2.2 km grassland reach of the River Cary in Somerset of U.K could delay the water velocity and produce flood prevention effect. Cheng (1999) found that reforestation in the upper Yangtze River Valley in China led to decrease in runoff. Brookhuis and Hein (2016) showed a non-linear relationship between catchment's forest cover and flood control in Trinidad. Also, their study estimated that the hydrological service of the forest cover is between 16 and 268 dollars per hectare per year. Tan-Soo *et al.*

(2016) found that deforestation in Peninsular Malaysia significantly increased the number of days flooded during 1984-2000.

South Korea provides a very good natural experimental condition for this kind of study not only due to data availability but also due to its natural environment and somewhat unique economic development experience.

South Korea is located in Northern East Asia and has four distinct seasons with heavy rainfalls concentrated in summer (typically from June to August) often caused floods in near-river areas. According to FAO (2016), 100 thousand hectares was flooded and more than 150 thousand people were victimized by flood in the 1960s.

Despite the fact that seventy percent of its land is covered by forest, however, hydrological services of forest was not paid attention to in Korea until growing stock of forest got heavily lost.² After imperialistic exploitation during the colonial times (1910 to 1945) and a three- year long Korean war (1950 to 1953), forests in South Korea were extremely devastated. According to the estimates of the Korea Forest Service, South Korea's volume of growing stock of forest per hectare, which is a statistic that represents forest inventory, was only 6 cubic meters in the 1950s. It is just four percent of the same measure in 2015, 146 cubic meters per hectare.³ The surprising increase in the growing stock is mainly due to the efforts of the South Korean government through the National Forest Rehabilitation Plans.⁴

As previously stated, deforestation usually aggravates in developing countries as their economic developments progress. However, South Korea recognized the importance of forest in providing essential services for human wellbeing from the initial stage of its economic development and accomplished significant reforestation (FAO, 2016). In the 1950s and 1960s, Korea was one of the poorest countries with GDP per capita in 1960 at mere \$1,175.⁵ However, the Five Year's Planning of Economic Development initiated by the South Korean government in 1962 began to change the economic atmosphere of the country. Its GDP per capita rose to \$10,152 in 1988, implying 12% annual growth on average. At the same time, South Korea also succeeded in reforestation. According to the Korea Forest Service, the total growing stock of forest increased from approximately 69 million m³ in 1970 to 800 million m³ in 2010.⁶

In contrast, according to Tang *et al.* (2010) and Jin *et al.* (2016), around 40 to 50 percent of forest area has been deforested in North Korea for the last few decades, and it is believed to be a reason for increased damages caused by natural disasters. In fact, heavy rains in August 2018 destroyed "more than 800 buildings including homes, clinics and school" in North and South Hwanghae provinces in North Korea.⁷ That size of flood damage is found in South Korea only in the years before reforestation. Even so, According to Lee *et al.* (2018), forest restoration efforts have been ineffective in North Korea for the last decade. The unique experience of South Korea which has pursued economic development and reforestation at the same time should provide useful implications for North Korea and also for countries that are in

²Korea was divided into two countries after World War II, South and North Korea.

³Korea Forest Service.

⁴According to Korea Forest Service, legal and institutional preparations preceded in the 1960s.

⁵The GDP per capita data in this section are all obtained from the Penn World Table version 9.0. The base year of the data is 2011.

⁶See footnote 2 for the source of the information.

⁷Reuters, "Flooding kills dozens in North Korea, thousands left homeless: Red Cross," 2018. 9. 6.

similar situation currently as Korea in the 1960s.

The following section describes the areas and rivers explored in this research. Section 3 discusses the data used in the research. In section 4, the estimation equations employed in the research and the estimation results using panel data are presented. Finally, section 5 concludes the paper.

II. Areas and Rivers Explored

Since the present research aims at investigating whether and to what extent the impact of rainfall on the water level of a river changes as the volume of growing stock of forest increases, the areas in which other factors than forest might alter the effects of rainfall on water level were excluded. Such factors include existence of a dam and major constructions to widen or deepen a river. Data availability was another consideration. Because rainfall is expected to affect water level of a river without a significant time delay, high frequency data should be desirable for this kind of research. In addition, the data of the volume of growing stock of adjacent forest must be available for a long time span. The existence of flood damage reports was also checked. Finally, the similarity of the lengths and the widths of the rivers included in the panel data was also considered. Although we control for the unobserved heterogeneity among the rivers in estimation, very large gaps in the river size would not be desirable.

After an extensive review, we finally settled on three river systems that fully satisfy the conditions above. They are the Hyeongsan River (R1), the Mangyeong River (R2), and the Hongcheon River (R3) in South Korea.⁸ For the three rivers, the volume of growing stock of forest in the watershed areas is reported every five year from 1985 to 2005. Besides, daily data of water level and rainfall are available for most periods from January 1, 1985 to Dec. 31, 2005.

TABLE 1—ACTUAL LOCAL AREA NAMES AND THEIR ABBREVIATED NAMES IN THE PAPER

	Local Name	Abbrev. Name	Local Name	Abbrev. Name	Local Name	Abbrev. Name
River	Hyeongsan	R1	Mangyeong	R2	Hongcheon	R3
Water Level Observatories	Moa	R1WL1				
	Angang	R1WL2	Daecheon	R2WL1	Bangkok	R3WL1
	Bujo	R1WL3	Dongjisan	R2WL2	Hongcheon	R3WL2
	<i>Pohang</i>	R1WL4				
Rainfall Observatories	Gyeongju	R1RF1	Gosan	R2RF1	Bangkok	R3RF1
	Gigye	R1RF2	Impi	R2RF2	Hongcheon Agr. High School	R3RF2
					Naecheon	R3RF3

Notes: 1) As stated in the main text, the two water level observatories in italic (*R1WL4* and *R2WL2*) are located near the river mouth of the sea, therefore were excluded in the analysis, 2) The two rainfall observatories in italic (*R2RF2* and *R3RF3*) were excluded in the analysis because they do not have corresponding water level observatories analyzed in the research, 3) HA is Hongcheon Agricultural High School.

⁸ For convenience, the paper will use an abbreviated name once a genuine full name is stated. An abbreviated name will be in a parenthesis following the actual local name when the latter is mentioned for the first time. Table 1 lists abbreviated names with their full names.

Figure 3 illustrates the location of the catchment areas of the three rivers in South Korea. Figures 4 through 6 illustrate the water level and rainfall observatories in each river. There are four water level observatories and two rainfall observatories in R1. The four water level observatories are Moa (R1WL1), Angang (R1WL2), Bujo (R1WL3) and Pohang (R1WL4). The two rainfall observatories are Gyeongju (R1RF1) and Gyeongju (R1RF2). R1 flows through the city of Gyeongju and the city of Pohang in the Province of Gyeongsangbuk-do and flows into the sea. The basin length, which is the length from the mouth to the farthest point of the river, is 57.8 kilometers.

For R2, there are two water level observatories, Daecheon (R2WL1) and Dongjisan (R2WL2), and two rainfall observatories, Gosan (R2RF1) and Impi (R2RF2). R2 flows through the city of Wanju and the city of Iksan in the Province of Jeollabuk-do and flows into the sea. Its basin length is 74.8 kilometers.

For R3, there are two water level observatories, Bangok (R3WL1) and Hongcheon (R3WL2), and three rainfall observatories, Bangok (R3RF1), Hongcheon Agricultural High School (R3RF2) and Naechon (R3RF3). R3 flows through the city of Hongcheon in the Province of Gangwon-do and joins the Han River, which flows through Seoul, the capital city of South Korea. The basin length of R3 is 108.5 kilometers.

Among the eight water level observatories mentioned above, R1WL4 in R1 and R2WL2 in R2 are located near the river mouth close to the sea. Therefore, the dynamics of the sea will affect the water level significantly, obscuring the effect of rainfall or reforestation. Accordingly, the two water level observatories were excluded in the analysis. Table 2 shows the six water level observatories included in the analysis and their corresponding rainfall observatories. The corresponding rainfall observatory of a water level observatory is the one located in the nearby upper stream area of the water level observatory. Because R1WL2 and R1WL3 are




FIGURE 3. WATERSHED AREAS OF R1, R2, AND R3




FIGURE 4. WATER LEVEL AND RAIN FALL OBSERVATORIES AT R1 (THE HYEONGSAN RIVER)




FIGURE 5. WATER LEVEL AND RAIN FALL OBSERVATORIES AT R2 (THE MANKYEONG RIVER)




FIGURE 6. WATER LEVEL AND RAIN FALL OBSERVATORIES AT R3 (THE HONGCHEON RIVER)

TABLE 2—WATER LEVEL OBSERVATORY AND CORRESPONDING RAINFALL OBSERVATORY

River	R1			R2	R3	
WL ¹	R1WL1	R1WL2	R1WL3	R2WL1	R3WL1	R3WL2
Corresponding RF ²	R1RF1	R1RF2		R2RF1	R3RF1	R3RF2

Note: 1) WL is water level observatory, 2) RF is rainfall observatory, 3) The rainfall observatory that is located in the nearby upper stream of a water level observatory is the corresponding rainfall observatory of the water level observatory.

related to the same corresponding rainfall observatory (R1RF2), there are six water level and five rainfall observatories included in the research as Table 2 shows. In section 4, the water level measured at each of the six water level observatories is regressed on rainfall measured at its corresponding rainfall observatory along with other explanatory variables. Through the regressions, this paper examines whether the impact of rainfall on water level has been alleviated as adjacent forest grows. The analysis of Baak *et al.* (2016) shows that flood damage is related more closely to water level rather than rainfall, implying to control the water level of a river is important to prevent flood. If higher volume of growing stock of forest helps to control water level, it should serve as a strong rationale for afforestation and forest conservation.

Figures 7 through 9 illustrate the volume of growing stock of forest in each watershed area of the three rivers. The lines in the figures are log values of the volumes, therefore their slopes represent the growth rate of the volume. As seen from Figures 7, the volume in R1 watershed area was only 1.2 million m³ in 1985, but increased to 5.3 million m³ in 2005.

The volume of growing stock in R2 watershed area was 1.4 million m³ in 1985, slightly higher than that in R1 watershed area. However, since the volume grew more slowly than in R1 area, it was only 3.2 million m³ in 2000 when that in R1 area reached more than 4 million m³. Different from the volume in R1 area whose growth slowed down from 2000, the volume in R2 area grew relatively fast between 2000 and 2005. Even so, it increased up to only 4.8 million m³ in 2005.

Different from the two watershed areas mentioned before, the volume of growing stock in R3 watershed area increased very rapidly between 1985 and 1990 as illustrated in Figure 9. It increased up to 4.9 million m³ as early as 1990, which is the level the previous two areas reached in around 2005. This rapid growth of the volume in an early stage might be explained by that R3 flows through mountainous areas. The growth rate of the volume in R3 area slowed down from 1995, but the volume was 12 m³ in 2005, more than two times the counterpart in the two previous areas.




FIGURE 7. VOLUME OF GROWING STOCK OF FOREST (THE HYEONGSANRIVER (R1) WATERSHED AREA)

Source: Water Resources Management Information System (<http://www.wamis.go.kr/>).




FIGURE 8. VOLUME OF GROWING STOCK OF FOREST (THE MANKYEONG RIVER (R2) WATERSHED AREA)

Source: Water Resources Management Information System (<http://www.wamis.go.kr/>).




FIGURE 9. VOLUME OF GROWING STOCK OF FOREST (THE HONGCHEON RIVER (R3) WATERSHED AREA)

Source: Water Resources Management Information System (<http://www.wamis.go.kr/>).

III. Data

The water level and rainfall data used in this paper are official data from the Ministry of Land, Infrastructure and Transport of the South Korean government.⁹ The data for R1, R2 and R3 were provided by Nakdong River Flood Control Office, Geum River Flood Control Office, and Han River Flood Control Office, respectively. All the flood control offices are subsidiary organizations of the Ministry of Land, Infrastructure and Transport of South Korea. The measurement unit of water level is one meter, and that of rainfall is one millimeter.

Both daily and hourly data from January 1, 1985 to December 31, 2014 were obtained from the offices above. However, mainly due to the lack of measurement infrastructure before 2000s, there were substantial number of missing values especially in the hourly data. Since daily data had much smaller number of missing values than hourly data, this paper used daily data for econometric analyses in the following section. In addition, it should be reported that the data were originally hand-written by the staffs of each river flood control office, and transferred to electronic forms quite later. Therefore, erroneous or difficult-to-read handwriting records sometimes made weird numbers in the electronic data sets. For example, the water level of R1WL on March 17 was 0.664, while it is 1.664 on the previous and the following day. Because it is obviously a record mistake, it was corrected to 1.664. When the correct numbers cannot be reasonably traced, the numbers suspected to be errors were treated as missing values. One example is the rainfall data whose values were negative for 17 days in December, 1996. Although those erroneous values question the quality of the data, it should be also reported that they are much less than one percent of the whole data.¹⁰

Although the rainfall and water level data are available up to 2014, we confined the time period of our analysis to 1986-2005 (January 1, 1986 to December 31, 2005) due to two reasons. Above all, the data for the volume of growing stock of forest in each watershed area of the three rivers are available up to 2005. Besides, the Four Major Rivers Project implemented by the South Korean government between 2009 and 2011 immensely changed the landscape of the watershed areas of South Korean rivers.¹¹ Although the three rivers included in the research were not directly affected by the project, they might be indirectly influenced. Because the factors such as construction of banks, which may alter the impact of rainfall on water level, should be controlled, it may not be wise to include the period of the Four Major Rivers Project in this kind of research.¹²

⁹A water level indicates a level above mean sea level. For the rainfall data, a datum indicates a height of the rain collected in a rain gage with the diameter of 20 cm.

¹⁰The original data uncorrected and the record of the data erased or corrected will be provided by the authors upon request.

¹¹Regarding the Four Major Rivers Project, refer to the following website of the United Nations (http://www.un.org/waterforlifedecade/green_economy_2011/pdf/session_8_water_planning_cases_korea.pdf).

¹²One of the anonymous referees pointed out the possibility that riverine dredging operations may have affected the relationship between rainfall variations upstream and water-level changes downstream over time. Indeed, if dredging took place more often toward the latter parts of our data period, attribution of the moderation in the sensitivity of water-level reactions to increases in the growing stock in the watershed areas would have become questionable. To the best of our knowledge, riparian dredging operations became serious undertakings only after the 2009 beginning of the Four Major Rivers Project. According to Chung *et al.* (2014), riverine dredging operations were sporadic, and limited to small-scale harvests of construction aggregate materials. See also newspaper reports

The water level data collected by the government offices are the data of river stage whose value is affected by the level of gage zero. “A gage zero” is an arbitrary point, which is normally set below water surface of a river at the driest season. The water level above the gage zero is called “a river stage,” and in many cases, water level observatories report only the data of river stage. Each gage zero has an elevation level above mean sea level, and the addition of the river stage and the elevation level of the gage zero is “the water level of a river above mean sea level.” Despite its convenience, river stage cannot be used for econometric analysis unless gage zero has not been changed. However, as Table 3 shows, a gauge zero in the three rivers involved in the research were changed a few times. For this reason, this study used the water level above mean sea level rather than the water level above the gauge zero. As Table 4 shows, however, the water levels above mean sea level are quite diverse across the six observatories because the altitudes of the rivers are all different. The water level above mean sea level is between 125.2 and 119.55 meters at R3WL2, and between 0.04 and 6.71 at R1WL2. Because the relative altitude of each river is not important and may add unnecessary heterogeneous noises on the estimation, the water level of each river is standardized by subtracting the minimum level from each data and adding one. Accordingly, the lowest water level in each river in the standardized data is one meter.¹³ The basic statistics of the standardized

TABLE 3—HISTORY OF GAUGE ZERO (FROM 1985 TO 2005)

River	Name of observatory	Period	Above mean sea level of gage zero (El. m)	
R1	R1WL1	1985-1988 Dec. 31	9.303	
		1989 Jan. 01-1999 Dec. 31	9.182	
		2000 Jan. 01-2000 Dec. 31	9.303	
		2001 Jan. 01-2010	9.113	
	R1WL2	1985-1997 Dec. 31	2.835	
		1998 Jan. 01-1999 Dec. 31	0.835	
		2000 Jan. 01 -2010	0.766	
		R1WL3	1985-1997 Dec. 31	0.844
			1998 Jan. 01-1999 Dec. 31	-0.166
			2000 Jan. 01-2003 Dec. 31	-0.939
2004 Jan. 01-2010	-1.054			
R2	R2WL1	1985-1993 Dec. 31	4.215	
		1994-1995	4.313	
		1996-2000	4.334	
		2001-2001	4.21	
		2002-2012	2.864	
R3	R3WL1	1985-1995	69.338	
		1996	67.338	
		1997	69.338	
		1998-2008	67.338	
	R3WL2	1985-2008	119.750	

in Kyung Hyang Daily (<https://m.khan.co.kr/politics/politics-general/article/201607050836001#c2b>), and Kyung Sang Maeil News (<http://m.ksmnews.co.kr/view.php?idx=156352>).

¹³The lowest water level was set to be one meter rather than zero not to lose data in case log values of water level are used in regressions. In case regressions do not use log values, it does not generate any qualitative difference in estimations results whether to use one or zero as the lowest water level.

TABLE 4—BASIC STATISTICS OF THE DATA

		R1WL1	R1WL2	R1WL3	R2WL1	R3WL1	R3WL2
Water Level (above sea level, meter)	Min	1.66	0.04	9.02	4.11	67.39	119.55
	Max	8.29	6.71	14.10	8.72	71.72	125.2
	Median	2.38	1.71	10.05	4.57	68.24	120.05
	Mean	2.38	1.57	10.00	4.65	68.32	120.20
	S.D	0.37	0.62	0.32	0.35	0.41	0.47
Water Level (standardized, meter)	Min	1.00	1.00	1.00	1.00	1.00	1.00
	Max	7.63	7.75	6.08	5.61	5.33	6.65
	Median	1.72	2.75	2.03	1.46	1.85	1.51
	Mean	1.72	2.61	1.98	1.54	1.93	1.65
	S.D	0.37	0.62	0.32	0.35	0.41	0.47

water level data are also reported in Table 4. Figures 10 through 15 illustrate the standardized water level and the rainfall of each observatory. The econometric analyses in the following section use the standardized data as the water level.

As previously mentioned, this research chose the three rivers because they are not significantly affected by dams or major constructions to widen or deepen the river. Even so, as Figures 10 through 15 shows, the water level data of some observatories show substantial structural changes. Figure 10 shows that water level at R1WL1 is distinctively higher from 1997. In fact, the original water level data provided by the Korean government shows that water level at R1WL2 jumped from 1.985 on Feb. 10 to 2.405 on Feb. 11 in 1997 without any explanation provided, although there was no rain in that month. This jump caused a significant change in the mean value of the water level after Feb. 10, 1997. Similar unexplained change in the mean of the water level is found at R1WL2 in 2000 and at R3WL2 in 1993. In the case of R1WL1 and R1WL2, those unexplained jumps occurred after the gage zero changed. Therefore, one reasonable conjecture is that the unexplained water level changes might be caused by measurement errors due to using inappropriate gage zero.¹⁴ To deal with that problem, this paper uses dummy variables for the periods during which water level is consistently higher or lower than other periods. The dummy variables used in the estimations are reported in Table 5.

As noted earlier, the volume of growing stock of forest is measured every five years by the Korea Forest Service. Therefore, daily data are not available. However, different from economic and financial data, the volume is expected to grow quite steadily even if the growth rate may not be perfectly constant. Based on that point, we obtained daily volume data by assuming a linear change in the volume throughout a year. The first observation, the data of 1985, was used as Dec. 31, 1985 data. Accordingly, the last observation, 2005 data, was used as Dec. 31, 2005 data. The volume data are available for only this period. Therefore, the time span of the research is limited to from January 1, 1986 to December 31, 2005. The measurement unit of the volume data is one billion m³. Because the measurement units of the variables are important when we interpret estimation results, they are summarized in Table 6.

¹⁴The changes in the gage zero were already incorporated when the water level above mean sea level data were calculated. Even so, unexplained structural shifts in the water level data are often found especially at the time when the gage zero was altered.




FIGURE 10. WATER LEVEL AND RAINFALL AT R1WL1

Note: WLN=water level, meter. RF=rainfall, millimeter.




FIGURE 11. WATER LEVEL AND RAINFALL AT R1WL2

Note: WLN=water level, meter. RF=rainfall, millimeter.




FIGURE 12. WATER LEVEL AND RAINFALL AT R1WL3

Note: WLN=water level, meter. RF=rainfall, millimeter.




FIGURE 13. WATER LEVEL AND RAINFALL AT R2WL1

Note: WLN=water level, meter. RF=rainfall, millimeter.




FIGURE 14. WATER LEVEL AND RAINFALL AT R3WL1

Note: WLN=water level, meter. RF=rainfall, millimeter.




FIGURE 15. WATER LEVEL AND RAINFALL AT R3WL2

Note: WLN=water level, meter. RF=rainfall, millimeter.

TABLE 5—DUMMY VARIABLES

Dummy	Observatory	Periods
R1WL1_97	R1WL1	From Feb. 11, 1997 to Dec. 31, 2005
R1WL2_00	R1WL2	From Jan. 1, 2000 to Dec. 31, 2005
R3WL2_92	R3WL2	From Jan. 1, 1986 to Dec. 31, 1992

TABLE 6—MEASUREMENT UNIT OF EACH VARIABLE

Water level	Rainfall (daily data)	Volume of growing stock
One meter	One millimeter per day	One billion cubic meter

IV. Panel Data Analysis

A. Estimation Results

To determine whether reforestation has contributed to stabilization of water levels of the three rivers chosen, the following equation is estimated.

$$(1) \quad Y_{it} = \alpha_0 + \alpha_1 Y_{it-1} + \alpha_2 Y_{it-1} F_{it} + \alpha_3 Y_{it-1} F_{it}^2 + \alpha_4 X_{it} \\ + \alpha_5 X_{it} F_{it} + \alpha_6 X_{it} F_{it}^2 + \sum_{j=1}^3 \beta_j D_{j,t} + \mu_i + \varepsilon_{it}$$

where Y_{it} is the water level at a specific water level observatory and X_{it} is the amount of rainfall observed at the corresponding rain-fall measuring spot. F_{it} is the volume of growing stock of the adjacent forest. $D_{j,t}$ is a period dummy variable to capture the effect of a structural shift in the water level possibly caused by a change in gauge zero. As summarized in Table 5, the R1WL1, the R1WL2 and the R3WL2 observatories have one period dummy for each. The term μ_i is the unobserved heterogeneity across the six cross-sections, and ε_{it} is a disturbance.

Because the water level of a river should have a strongly positive relationship with its value of the previous day and because water level should not explode, α_1 is expected to be a positive number close to but less than one. Because the amount of rainfall in one area will raise the water level of a river in the same area, α_4 is expected to be positive. The coefficients, α_2 , α_3 , α_5 , and α_6 of the interaction terms, $Y_{it-1} F_{it}$, $Y_{it-1} F_{it}^2$, $X_{it} F_{it}$, and $X_{it} F_{it}^2$ will capture the impacts of the volume of growing stock of forest on water level. If the forest has the function to stabilize water level, α_2 and α_5 should be negative. The signs of α_3 and α_6 depend on whether the marginal effect of forest to control water level is diminishing or accumulating. This point will be further discussed below with the estimation results. The stationarity of all the variables included in equation (1) was examined by the panel unit root test of Levin, Lin, and Chu (2002). The null hypothesis of presence of unit root was strongly rejected for all the variables.

Table 7 reports the estimation results of equation (1) for four different models. Column 1 reports the estimates of the pooled OLS model. In column 2, dummy variables are dropped. Columns 3 and 4 report the estimates of the fixed effects

TABLE 7—ESTIMATION RESULTS (WHOLE TIME PERIOD)

Model		(1) OLS	(2) OLS no dummy	(3) Fixed Effect	(4) Fixed Effect no dummy
α_0	Coeff.	0.127*	0.127*	0.302*	0.231*
	P-Val	0.000	0.000	0.000	0.000
$\alpha_1 Y_{it-1}$	Coeff.	0.939*	0.939*	0.842*	0.895*
	P-Val	0.000	0.000	0.000	0.000*
$\alpha_2 Y_{it-1} F_{it}$	Coeff.	-4.511*	-4.511*	-3.466*	-6.220*
	P-Val	0.000	0.000	0.000	0.000
$\alpha_3 Y_{it-1} F_{it}^2$	Coeff.	158.6*	158.6*	192.8*	56.5
	P-Val	0.031	0.031	0.012*	0.407
$\alpha_4 X_{it}$	Coeff.	0.007*	0.007*	0.008*	0.008*
	P-Val	0.000	0.000	0.000	0.000
$\alpha_5 X_{it} F_{it}$	Coeff.	-0.903*	-0.904*	-1.049*	-0.998*
	P-Val	0.000	0.000	0.000	0.000
$\alpha_6 X_{it} F_{it}^2$	Coeff.	73.3*	73.5*	83.5*	80.7*
	P-Val	0.000	0.000	0.000	0.000
Dummy R1WL1	Coeff.	0.004		0.058*	
	P-Val	0.321		0.000	
Dummy R1WL2	Coeff.	0.008*		-0.155*	
	P-Val	0.060		0.000	
Dummy R3WL2	Coeff.	0.006		0.086*	
	P-Val	0.164		0.000	
R^2		0.897	0.897	0.904	0.901
Adjusted R^2		0.897	0.897	0.904	0.901
Number of observations		37,478	37,478	37,478	37,478
Redundant Fixed Effects Tests (p-value)		na	na	0.00	0.00

Note: The asterisk indicates the estimate is significance at the 5% significance level.

model with two different specifications just as the OLS model. Because the number of explanatory variables is greater than the number of cross-sections, the random effects model could not be estimated. Since the work of Nickell (1981), it has been well known that the coefficient of the lagged dependent variable in a dynamic panel model (α_1 in equation (1)) is not consistently estimated by fixed effects or random effects models due to the endogeneity problem caused by the correlation between the lagged dependent variable and the error terms. However, as Roodman (2006) among others argues, the inconsistency is negligible if the dimension of cross-sections is far bigger than the dimension of time series, which is the case of the data in this paper. Accordingly, a methodology such as the system GMM of Arellano and Bover (1995) and Blundell and Bond (2000), which are recommended for dynamic panel data analyses when the number of cross-sections is bigger than that of time series, were not employed in the present research.

The estimates of the OLS model are quite similar to those of the fixed effects model in each specification except for the estimates of dummy variables. The estimates of dummies are all insignificant at the five percent significance level in the OLS model in column 1, but they are all strongly significant in the fixed effect model in column 3. The p-value of the redundant test for the fixed effects is almost zero, strongly supporting the fixed effects model over the OLS model.

In all versions of the model, α_1 is estimated to be around 0.9, consistent with our expectation. The coefficients of the interaction terms between forest volume and other explanatory variables, α_2 and α_5 are estimated to be negative, implying that forest can contribute to controlling water level. The positive signs of α_3 and α_6 indicate that the marginal effect of forest is diminishing.

For an additional robust check, Equation (1) was estimated using only summertime data (May to October). The results reported in Table 8 show that the signs of all coefficients are the same as in Table 7, re-confirming that water level is better controlled as the forest grows, but that the marginal effect of forest to control water level diminishes.

Because the forests adjacent to the three rivers explored have continued to grow for the time period covered in the paper, it is a reasonable suspect that a trend may play a similar role to the volume of growing stock in equation (1). Therefore, equation (1) was re-estimated after replacing the forest volume with a trend, and the estimation results are reported in Table 9. Although the interaction terms of trend have the same signs with respect to rainfall, their signs turn out to be the opposite, with respect to lagged water level, to those reported in Tables 7 and 8. Besides, the estimate of α_2 is not significant in Table 9. This indicates that the water-stabilization effects of forest found in the estimation of Equation (1) is not the results of some

TABLE 8—ESTIMATION RESULTS (MAY TO OCT.)

Model		(1) OLS	(2) OLS no dummy	(3) Fixed Effect	(4) Fixed Effect no dummy
α_0	Coeff.	0.201*	0.201*	0.409*	0.325*
	P-Val	0.000	0.000	0.000	0.000
$\alpha_1 Y_{it-1}$	Coeff.	0.908*	0.907*	0.792*	0.858*
	P-Val	0.000	0.000	0.000	0.000
$\alpha_2 Y_{it-1} F_{it}$	Coeff.	-6.818*	-6.321*	-5.384*	-9.947*
	P-Val	0.000	0.000	0.001	0.000
$\alpha_3 Y_{it-1} F_{it}^2$	Coeff.	282.9*	242.1*	489.6*	343.3*
	P-Val	0.028	0.035	0.000	0.004
$\alpha_4 X_{it}$	Coeff.	0.008*	0.008*	0.009*	0.008*
	P-Val	0.000	0.000	0.000	0.000
$\alpha_5 X_{it} F_{it}$	Coeff.	-1.029*	-1.027*	-1.186*	-1.130*
	P-Val	0.000	0.000	0.000	0.000
$\alpha_6 X_{it} F_{it}^2$	Coeff.	80.9*	80.7*	90.6*	87.1*
	P-Val	0.000	0.000	0.000	0.000
Dummy R1WL1	Coeff.	-0.001		0.066*	
	P-Val	0.847		0.000	
Dummy R1WL2	Coeff.	0.007		-0.212*	
	P-Val	0.384		0.000	
Dummy R3WL2	Coeff.	0.009		0.134*	
	P-Val	0.265		0.000	
R^2		0.851	0.840	0.851	0.847
Adjusted R^2		0.851	0.840	0.851	0.847
Number of observations		19,095	19,095	19,095	19,095
Redundant Fixed Effects Tests (p-value)		na	na	0.00	0.00

Note: The asterisk indicates the estimate is significance at the 5% significance level.

TABLE 9—ESTIMATION RESULTS OF FIXED EFFECTS MODELS
(FOREST DENSITY IS REPLACED BY TIME TREND)

	Whole Time Period		May to Oct.	
	coefficient	p-value	coefficient	p-value
α_0	0.303*	0.000	0.408*	0.000
$\alpha_1 Y_{it-1}$	0.835*	0.000	0.787*	0.000
$\alpha_2 Y_{it-1} F_{it}$	0.001	0.182	0.002	0.252
$\alpha_3 Y_{it-1} F_{it}^2$	-0.000*	0.000	-0.001*	0.006
$\alpha_4 X_{it}$	0.007*	0.000	0.008*	0.000
$\alpha_5 X_{it} F_{it}$	-0.002*	0.000	-0.002*	0.000
$\alpha_6 X_{it} F_{it}^2$	0.000*	0.000	0.000*	0.000
R1WL1_97	0.058*	0.000	0.065*	0.000
R1WL2_00	-0.154*	0.000	-0.205*	0.000
R3WL2_92	0.080*	0.000	-0.101	0.000
R^2	0.904		0.841	
Adjusted R^2	0.904		0.841	
No of obs.	37478		19095	

Note: The asterisk indicates the estimate is significance at the 5% significance level.

unknown time trending effects.

Because the test results for redundant fixed effects support the fixed effects model and because all dummies turn out significant, the simulation experiments in the following section use the estimates of the fixed effect model with dummies in Table 7.

B. Effects of Forest on Water Level: A Simulation

From Equation (1), it is obvious that the direct impact of rainfall, X_{it} , on the water level, Y_{it} , is $\alpha_4 + \alpha_5 F_{it} + \alpha_6 F_{it}^2$. In addition, X_{it} also affects Y_{it+1} through Y_{it} . That is,

$$(2) \quad \frac{\partial Y_{it+1}}{\partial X_{it}} = \frac{\partial Y_{it+1}}{\partial Y_{it}} \frac{\partial Y_{it}}{\partial X_{it}} = (\alpha_1 + \alpha_2 F_{it+1} + \alpha_3 F_{it+1}^2)(\alpha_4 + \alpha_5 F_{it} + \alpha_6 F_{it}^2).$$

By the same token, the impact of X_{it} on Y_{it+j} , or the change in the water level responding to a unit (millimeter) change in rainfall, can be also computed.

Figure 16 illustrates the simulated daily effects of rainfall on water level in two cases. The first case (Case 1) is that the volume of growing stock is one million m^3 , and the second case is that it is five million m^3 (Case 2). As Figures 7 through 8 show, the volumes of growing stock of the two forests in the watershed area of R1 and R2 were around one million m^3 in 1985. However, they increased to around five million m^3 in 2005. Therefore, the simulations in Figure 16 can be understood to compare the impacts of rain fall on water level between two rivers, one with the




FIGURE 16. SIMULATED EFFECTS OF RAINFALL ON WATER LEVEL

forest of the year 1985 and the other with the forest of the year 2005. Such a simulation will illustrate the extent to which reforestation contributed to stabilizing water level in South Korea. Specifically, Figure 16 depicts the daily changes of water level when the amount of rainfall is 100 millimeters at day 0 and there is no rain from day 1.¹⁵

When it rains one hundred millimeter per day at time zero, the water level rises by 0.70 meter on that day, if the volume of growing stock is 1 million m³. In contrast, the water level rises by 0.48 meter, if the volume of growing stock is 5 million m³. Then, the effects of rainfall gradually vanish.

Figure 17 illustrates the simulation applied to the case of heavy rain in R1 area. On August 22 and 23 in 1991, there was heavy rain at R1WL1, 204 and 371 millimeters, respectively, and the water level rose by around 5 meters on August 23. The economic damage of R1 area caused by flood in that year was estimated to be




FIGURE 17. SIMULATED EFFECTS OF RAINFALL ON WATER LEVEL

¹⁵The monthly average rainfall in August is around 9 millimeter per day. However, rainfall of one hundred millimeter per day is not rare in the summer time in Korea, as seen from the Figures 10 through 15.

0.61 billion dollars. Ten people lost their lives and 4055 people were affected. The real line shows the simulated water level using the actual forest volume of that time period, while the two dotted lines illustrate the simulated water level based on the estimation results of column 3 in Table 7. The higher dotted line is the simulation results using the actual forest volume in 1991 (around 2.2 million cubic meter) and the lower one using the forest volume in August 2005 forest volume (around 5.2 million cubic meter). When the actual forest volume was used, the water level was simulated to rise as high as 5.36 meter, while it was simulated to rise up to 4.66 meter. This implies that the water level at R1WL1 in August 1991 could have been reduced by 13% if the forest volume had been as thick as in August 2005.

V. Summary and Conclusion

The main objective of this study is to conduct an empirical analysis to test for the impact of reforestation on stabilization of water level in three Korean rivers, R1, R2, and R3. The daily data used in the empirical studies were collected from three water level observatories and two rainfall observatories at R1, one water level observatory and one rainfall observatory at R2, and two water level observatories and two rainfall observatories at R3.

The three rivers were selected because they have not been affected by dams and/or major operations to widen or deepen the rivers. In addition, data availability was also considered. The data for water level, rainfall, and volume of growing stock of forest are available in the three rivers for most of the time periods from 1985 to 2005.

In particular, this research paper regressed present water level on previous water level, present rainfall, and their interaction terms with the volume of growing stock. In the dynamic panel data analysis of six water level observatories, the coefficients of previous water level and present rainfall were estimated to be positive and significant as expected. The interaction term between previous water level and forest volume turned out to be negative, and the interaction term between present rainfall and forest volume also turned out to be negative. These findings imply that the water level is better controlled as the volume of growing stock increases. In the meantime, the interaction term between each of the two main explanatory variables and squared forest volume was estimated to be positive, implying the marginal effect of water control function of forest diminishes as the volume of growing stock increases.

When the volume of growing stock was replaced by a trend, the signs of the coefficients were different from what were previously obtained, confirming the effects of reforestation cannot be attributed to unknown time trending effects.

Simulations employing estimated coefficient values show that reforestation reduces impacts of rain on water level substantially. Specifically, when it rains one hundred millimeter per day at time zero, the water level rises by 0.70 meter on that day, if the volume of growing stock is 1 million m^3 . In contrast, the water level rises by 0.48 meter, if the volume of growing stock is 5 million m^3 . Then, the effects of rainfall gradually vanish.

The water level controlling function of forest may be ignored in the decision making process of development plans because the benefits may not be obvious to casual inspection, possibly contributing to trends for deforestation in the course of

economic development. This paper focused on finding the econometric evidence of the water storage and conservation function of forest. Rigorous monetary evaluation of this and other positive externalities from forestation is beyond the scope of the current paper, but would contribute to a more systematic and comprehensive cost benefit analysis of reforestation interventions. It is the hope of the authors that the findings from this paper will serve to facilitate such future research efforts.

REFERENCES

- Arellano, M., and O. Bover.** 1995. "Another look at the instrumental variable estimation of error-components models," *Journal of Econometrics*, (68): 29-51.
- Blundell, R. W. and S. R. Bond.** 2000. "GMM Estimation with Persistent Panel Data: an Application to Production Functions," *Econometric Reviews*, 19: 321-340.
- Brookhuis, B. J. and L. G. Hein.** 2016. "The value of the flood control service of tropical forests: A case study for Trinidad," *Forest Policy & Economics*, 62, 118-124.
- Chang, J. et al.** 2016. "International trade causes large net economic losses in tropical countries via the destruction of ecosystem services," *Ambio*, 45(4): 387-397.
- Cheng, G. W.** 1999. "Forest change: hydrological effects in the Upper Yangtze River Valley," *Ambio*, 28(5): 457-459.
- Chung, A., H. Lee, T. Kang, and K. Jung.** 2014. "The present and future for dredging-related models," *Water for Future*, 47(2): 63-75 (in Korean).
- FAO.** 2015. *Global Forest Resources Assessment 2015: How Are the World's Forests Changing?*, Food and Agriculture Organization of the United States, second edition.
- FAO.** 2016. *Integrated policy for forests, food security and sustainable livelihoods - lessons from the republic of Korea*, Rome, Italy.
- Faria, W. R. et al.** 2016. "Relationship between openness to trade and deforestation: Empirical evidence from the Brazilian Amazon," *Ecological Economics*, 121: 85-97.
- Ferraro, P. J. et al.** 2012. "Forest Figures: Ecosystem Services Valuation and Policy Evaluation in Developing Countries," *Review of Environmental Economics and Policy*, 6(1): 20-44.
- Hundecha, Yeshewatesfa and András Bárdossy.** 2004. "Modeling of the effect of land use changes on the runoff generation of a river basin through parameter regionalization of a watershed model," *Journal of Hydrology*, 292, 281-295.
- Jin, Y. et al.** 2016. "Mapping Deforestation in North Korea Using Phenology-Based Multi-Index and Random Forest," *Remote Sensing*, 8(12): 997.
- Klemick, H.** 2011. "Shifting cultivation, forest fallow, and externalities in ecosystem services: Evidence from the Eastern Amazon," *Journal of Environmental Economics and Management*, 61(1): 95-106.
- Korea Forest Service** (https://english.forest.go.kr/kfswb/kfi/kfs/cms/cmsView.do?cmsId=FC_001680&mn=UENG_02_01_01, last accessed on Feb. 10, 2022).
- Korea Association for Public Administration.** 2009. *A Case study of the successful reforestation in South Korea* (in Korean).
- Kyung Hyang Daily** (<https://m.khan.co.kr/politics/politics-general/article/201607050836001#c2b>, last accessed on Feb. 10, 2022).
- Kyung Sang Maeil News** (<http://m.ksmnews.co.kr/view.php?idx=156352>, last accessed on Feb. 10, 2022).
- Lee, S., H. Choi, H. Yoo, C. Song, S. Cha, S. Bae, Y. Son, and W. Lee.** 2018. "Restoration Plan for Degraded Forest in The Democratic People's Republic of Korea Considering Suitable Tree Species and Spatial Distribution," *Sustainability*, 10(3): 856-870.
- Levin, A., C. F. Lin, and C. Chu.** 2002. "Unit Root Tests in Panel Data: Asymptotic and Finite-Sample Properties," *Journal of Econometrics*, 108: 1-24.
- Li, L. et al.** 2017. "Economic globalization, trade and forest transition-the case of nine Asian countries," *Forest Policy and Economics*, 76: 7-13.
- MEA.** 2003. *Ecosystems and Human Wellbeing. A Framework for Assessment*, Washington, DC, USA: Island Press.
- Neudert, R., U. G. Joerg, and F. Waetzold.** 2016. "Global benefits and local costs-The dilemma of tropical forest conservation: A review of the situation in Madagascar," *Environmental Conservation*, 44(1): 82-96.
- Nickell, S. J.** 1981. "Biases in Dynamic Models with Fixed Effects," *Econometrica*, 49(6): 1417-1426.
- Nisbet, T. et al.** 2004. "A Guide to Using Woodland for Sediment Control," Forest Research,

Farnham.

- Reuters.** 2018. "Flooding kills dozens in North Korea, thousands left homeless: Red Cross," Sept. 6.
- Roodman, D.** 2006. *How to do xtabond2: An introduction to Difference and System Gmm in Stata*, Working Paper No. 103, Center for global development.
- Seo, Min Young.** 2018. "Female Leadership and Forestation for Rural Development: Three Essays in Development Economics," KDI School of Public Policy and Management.
- Sims, K. R. E.** 2010. "Conservation and development: Evidence from Thai protected areas," *Journal of Environmental Economics and Management*, 60(2): 94-114.
- Tan-Soo, J. S. et al.** 2016. "Econometric Evidence on Forest Ecosystem Services: Deforestation and Flooding in Malaysia," *Environmental and Resource Economics*, 63: 25-44.
- Tang, L. et al.** 2010. "Forest degradation deepens around and within protected areas in East Asia," *Biological Conservation*, 143: 1295-1298.
- TEEB.** 2010. "Integrating the ecological and economic dimensions in biodiversity and ecosystem service valuation," in P. Kumar (ed.) *The Economics of Ecosystems and Biodiversity Ecological and Economic Foundations*, Chapter 1, UK, and Washington, DC, USA: Earthscan.
- United Nations** (http://www.un.org/waterforlifedecade/green_economy_2011/pdf/session_8_wat_planning_cases_korea.pdf, last accessed on Feb. 10, 2022).
- Wheeler, D. et al.** 2013. "Economic dynamics and forest clearing: A spatial econometric analysis for Indonesia," *Ecological Economics*, 85: 85-96.

LITERATURE IN KOREAN

- 정안철·이형·강태운·경관수. 2014. 「준설관련 모형의 현황 및 발전방향」, 『물과 미래』, 47(2): 63-75.
- 한국행정학회. 2009. 『한국의 치산녹화 성공 사례 분석』, 산림청 용역보고서.