



STUDY PROTOCOL

REVISED **Stigma-directed services (Stig2Health) to improve 'linkage to care' for people living with HIV in rural Tanzania: study protocol for a nested pre-post implementation study within the Kilombero and Ulanga Antiretroviral Cohort**

[version 2; peer review: 2 approved]

Raphael Magnolini ^{1*}, Elizabeth Senkoro^{2,3*}, Aneth Vedastus Kalinjuma^{2,4}, Olivia Kitau², Bernard Kivuma ^{2,3}, Leila Samson^{2,3}, Anna Eichenberger^{2,5-7}, Getrud Joseph Mollel^{2,3}, Eileen Krinke ^{8,9}, James Okuma^{6,7}, Robert Ndege ^{2,3}, Tracy Glass^{6,7}, Herry Mapesi^{2,6,7}, Fiona Vanobberghen^{6,7}, Manuel Battegay¹, Maja Weisser ^{1,2,6,7}

¹Division of Infectious Diseases and Hospital Epidemiology, University Hospital of Basel, Basel, Switzerland²Ifakara Health Institute, Ifakara, Tanzania³St. Francis Referral Hospital, Ifakara, Tanzania⁴School of Public Health, Faculty of Health Sciences, University of the Witwatersrand, Johannesburg, South Africa⁵Department of Infectious Diseases, Inselspital, University Hospital Bern, Bern, Switzerland⁶Swiss Tropical and Public Health Institute, Basel, Switzerland⁷University of Basel, Basel, Switzerland⁸University Psychiatric Clinics Basel, Basel, Switzerland⁹University of Zurich, Zurich, Switzerland

* Equal contributors

V2 First published: 21 Mar 2022, 5:14
<https://doi.org/10.12688/aasopenres.13353.1>

Latest published: 05 Aug 2022, 5:14
<https://doi.org/10.12688/aasopenres.13353.2>

Abstract

Background: HIV-related stigma is a major barrier to the timely linkage and retention of patients in HIV care in sub-Saharan Africa, where most people living with HIV/AIDS reside. In this implementation study we aim to evaluate the effect of stigma-directed services on linkage to care and other health outcomes in newly diagnosed HIV-positive patients.

Methods: In a nested project of the Kilombero and Ulanga Antiretroviral Cohort in rural Tanzania, we conduct a prospective observational pre-post study to assess the impact of a bundle of stigma-directed services for newly diagnosed HIV positive patients. Stigma-directed services, delivered by a lay person living with HIV, are

Open Peer Review**Approval Status**

1

2

version 2(revision)
05 Aug 2022**version 1**

21 Mar 2022



view



view

1. **Graeme Hoddinott** , Stellenbosch
University, Stellenbosch, South Africa

i) post-test counseling, ii) post-test video-assisted teaching, iii) group support therapy and group health education, and iv) mobile health. Patients receiving stigma services (enrolled from 1st February 2020 to 31st August 2021) are compared to a historical control receiving the standard of care (enrolled from 1st July 2017 to 1st February 2019). The primary outcome is 'linkage to care'. Secondary endpoints are retention in care, viral suppression, death and clinical failure at 6-12 months (up to 31st August 2022). Self-reported stigma and depression are assessed using the Berger Stigma scale and the PHQ-9 questionnaire, respectively. The sample size calculation was based on cohort data from 2018. Assuming a pre-intervention cohort of 511 newly diagnosed adults of whom 346 (68%) were in care and on antiretroviral treatment (ART) at 2 months, a 10% increase in linkage (from 70 to 80%), a two-sided type I error rate of 5%, and 90% power, 321 adults are required for the post-implementation group.

Discussion: We expect that integration of stigma-directed services leads to an increase of proportions of patients in care and on ART. The findings will provide guidance on how to integrate stigma-directed services into routine care in rural sub-Saharan Africa.


Keywords

HIV/AIDS, Stigma, Linkage to care, HIV care, Sub-Sahara Africa, Study protocol



African Population and
Health Research Center

This article is included in the [African Population Health Research Center](#) gateway.

2. **Richelle Harklerode** , University of California, San Francisco, San Francisco, USA

Any reports and responses or comments on the article can be found at the end of the article.

Corresponding author: Maja Weisser (maja.weisser@usb.ch)

Author roles: **Magnolini R:** Conceptualization, Funding Acquisition, Investigation, Methodology, Project Administration, Visualization, Writing – Original Draft Preparation, Writing – Review & Editing; **Senkoro E:** Conceptualization, Investigation, Methodology, Project Administration, Resources, Writing – Original Draft Preparation, Writing – Review & Editing; **Kalinjuma AV:** Data Curation, Formal Analysis, Methodology, Resources, Software, Validation, Writing – Review & Editing; **Kitau O:** Data Curation, Formal Analysis, Methodology, Project Administration, Validation, Writing – Review & Editing; **Kivuma B:** Investigation, Methodology, Project Administration, Resources, Writing – Review & Editing; **Samson L:** Conceptualization, Methodology, Project Administration, Resources, Writing – Review & Editing; **Eichenberger A:** Methodology, Project Administration, Writing – Review & Editing; **Mollel GJ:** Methodology, Project Administration, Writing – Review & Editing; **Krinke E:** Conceptualization, Investigation, Methodology, Validation, Writing – Review & Editing; **Okuma J:** Data Curation, Formal Analysis, Methodology, Resources, Software, Validation, Writing – Review & Editing; **Ndege R:** Methodology, Project Administration, Resources, Writing – Review & Editing; **Glass T:** Conceptualization, Methodology, Software, Writing – Review & Editing; **Mapesi H:** Methodology, Writing – Review & Editing; **Vanobberghen F:** Data Curation, Formal Analysis, Methodology, Software, Validation, Writing – Review & Editing; **Battegay M:** Conceptualization, Methodology, Supervision, Writing – Review & Editing; **Weisser M:** Conceptualization, Funding Acquisition, Methodology, Project Administration, Resources, Supervision, Writing – Original Draft Preparation, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: In summary, this research was partially supported by the African Academy of Sciences under a DELTAS Africa Initiative grant [107768] as part of the Consortium for Advanced Research Training in Africa (CARTA). Full disclosure statement: Aneth Vedastus Kalinjuma was supported by the Consortium for Advanced Research Training in Africa (CARTA). CARTA is jointly led by the African Population and Health Research Center and the University of the Witwatersrand and funded by the Carnegie Corporation of New York (Grant No: B 8606.R02), Sida (Grant No: 54100029), the DELTAS Africa Initiative (Grant No: 107768/Z/15/Z). The DELTAS Africa Initiative is an independent funding scheme of the African Academy of Sciences (AAS)'s Alliance for Accelerating excellence in Science in Africa (AESA) and supported by the New Partnership for Africa's Development planning and Coordinating Agency (NEPAD Agency) with funding from the Wellcome Trust (UK) and the UK government. The statements made and views expressed are solely the responsibility of the fellow. The Chronic Diseases Clinic of Ifakara receives funding from the Ministry of Health and Social Welfare of the Government of Tanzania; the Government of the Canton of Basel, Switzerland; the Swiss Tropical and Public Health Institute, Basel, Switzerland; the University Hospital Basel, Switzerland, the Ifakara Health Institute, Tanzania; and USAID Boresha Afya (service and drug support with funding through the United States Agency for International Development (USAID) from the President's Emergency Plan for AIDS Relief (PEPFAR) programme). The "Freiwillige Akademische Gesellschaft (FAG)", Switzerland has granted 6'000 CHF for travel reimbursements and living cost for this study. The funding sources had no role in the design of the study and will not be involved in data collection, data analysis, interpretation of the results and writing of the manuscript.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2022 Magnolini R *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Magnolini R, Senkoro E, Kalinjuma AV *et al.* **Stigma-directed services (Stig2Health) to improve 'linkage to care' for people living with HIV in rural Tanzania: study protocol for a nested pre-post implementation study within the Kilombero and Ulanga Antiretroviral Cohort [version 2; peer review: 2 approved]** AAS Open Research 2022, 5:14 <https://doi.org/10.12688/aasopenres.13353.2>

First published: 21 Mar 2022, 5:14 <https://doi.org/10.12688/aasopenres.13353.1>

REVISED Amendments from Version 1

According to the reviewer's comments, the authors have conducted minor changes to the main text body. No changes were done to tables or graphs.

Changes to the main text body included tense changes (future tense throughout the manuscript), a conceptual framework of HIV directed stigma was added and the impact of services on stigma mechanisms were further elaborated, minor changes were done to the outline of the group therapy sessions as they now also include frequency of sessions and participant numbers, and we further elaborated on the procedure in case depression was detected among people living with HIV and AIDS.

Any further responses from the reviewers can be found at the end of the article

Introduction

The rollout of antiretroviral treatment (ART) for people living with HIV/AIDS (PLWHA) is one of the largest and most successful health interventions – especially in sub-Saharan Africa, where most affected persons live. Nevertheless, HIV programs aiming to reach the updated Joint United Nations Programme on HIV and AIDS (UNAIDS) goals of “95-95-95”, i.e. 95% of PLWHA diagnosed, 95% of people who are diagnosed receiving ART, and 95% of those receiving ART being virally suppressed, by 2025 are jeopardized by the loss of patients on all levels of the HIV care continuum – HIV testing, linkage to HIV care with ART initiation and retention in care for sustained viral suppression¹⁻³. The highest loss of patients has been observed during ‘linkage to care’ – commonly defined as ‘patient entry into specialist HIV care after diagnosis’^{2,4-8}. Poor care linkage appears to be particularly pronounced in sub-Saharan Africa^{2,9,10}.

An important reason for non-linkage is HIV-related stigma and discrimination^{11,12}. HIV-related stigma is a complex concept, comprising of three important stigma mechanisms described for HIV infected individuals, i.e., enacted stigma (level to which PLWHA believe they have actually experienced prejudice and discrimination from others), anticipated stigma (the level to which PLWHA expect that they will experience prejudice and discrimination in the future), and internalized stigma (“felt” stigma or “self-stigmatization”, used to describe the way PLWHA feel about themselves)^{11,13,14}. PLWHA who perceive high levels of HIV-related stigma are 2.4 times more likely to delay enrolment in care compared to those who don't¹⁵. HIV-related stigma is associated with non- or involuntary HIV-status disclosure and denial of the HIV-status¹⁶. Consequences are lower access to health care services, poor health-protective behaviors¹⁷, delayed initiation of ART and low adherence to ART^{16,18,19} resulting in poor treatment outcomes²⁰⁻²². Protective factors against stigma are family cohesiveness, and social and emotional support²³. Depression and stigma are mutually associated, and both affect HIV treatment outcome^{20,21,24,25}.

Despite the vast literature on HIV-related stigma and its negative impact on outcomes for PLWHA¹¹, few intervention studies have targeted the reduction of HIV-related stigma

among PLWHA in different socio-cultural settings^{11,26}. Different promising and cost-effective intervention strategies to improve care and health outcomes in resource-limited settings include task shifting from physicians to lay persons with lower-level qualifications²⁷⁻²⁹, video-assisted teachings (VAT), mobile Health (mHealth)³⁰⁻³⁴, group support therapy and group health education^{27,35}. Advantages of VAT are the low cost, standardized delivery, and time-savings for healthcare staff due to its customized content and flexible dissemination channels^{23,36}.

In Sub-Saharan Africa, mental health services addressing depression have shown to be a **major treatment gap** with lack of services in up to 90%^{25,37-39} – mostly due to lack of trained healthcare professionals²⁵ or lack of accessibility and sustainability. Some data suggests that implementing task shifting in resource-limited settings can improve mental disorders among PLWHA by delegating care from psychotherapists to lay health care workers^{27,28}. With mental health services still being **very poorly available** and not integrated into HIV care in sub-Saharan Africa, little is known about task shifting in this field.

The effectiveness and feasibility of combined interventional strategies has previously been evaluated^{3,40,41}, but to our knowledge the combined effect of stigma-directed services has not been studied.

**Protocol
Study aim**

The aim of this implementational study is to evaluate the effect of the impact of a bundle of stigma-directed services on linkage to care and other health outcomes for patients that are newly diagnosed with HIV. Stigma-directed services, delivered by a lay person living with HIV, are i) post-test counseling, ii) post-test video-assisted teaching, iii) group support therapy and group health education, and iv) mHealth. Stigma-directed services are aimed to reduce all levels of stigma within PLWHA, i.e., enacted stigma, anticipated stigma, and internalized stigma. Importantly, only the complete bundle of stigma-directed services to the outcomes can be assessed, the value of each individual service to the outcomes cannot be differentiated. We hypothesize that integration of combined stigma-directed services into routine care for patients newly diagnosed with HIV will increase timely linkage to care, retention in care and viral suppression. Additionally, we hypothesize patients' self-reported stigma and depression within the first year will decrease.

Study setting

The Chronic Disease Clinic Ifakara (CDCI) was established in 2004 as the governmental care and treatment center for PLWHA of the St. Francis Referral Hospital (SFRH) in Ifakara, South-western rural Tanzania. The clinic provides free HIV testing, care and antiretroviral drugs for PLWHA according to the National AIDS Control Program (NACP) guidelines. All patients are offered participation in a prospective patient cohort – the Kilombero and Ulunga Antiretroviral Cohort (KIULARCO) and are enrolled the day of diagnosis upon consent. KIULARCO is a collaborative project of the SFRH with the Ifakara Health Institute, the Swiss Tropical and Public

Health Institute and the University Hospital Basel to allow research on needs and treatment outcomes of patients^{42,43}.

Study design

We will conduct a prospective observational pre-post study nested within KIULARCO. All newly diagnosed patients with HIV consenting to KIULARCO will be offered stigma-directed services at enrolment from 1st February 2020 onwards in addition to standard of care. Prior to Stig2health, the standard of care involved pre- and post-ART counselling in newly diagnosed patients as per national guidelines and clinical follow-up reminders by automated text messages to all patients attending the clinic with a registered phone number (Figure 1).

Recruitment, screening eligibility criteria

From 1st February 2020 onwards, all newly HIV-diagnosed patients enrolled in KIULARCO and fulfilling eligibility criteria (Table 1) will be offered stigma-related services by a professional counselor (VCT) on the same day.

Informed consent for the Stig2Health nested study within KIULARCO

Stig2health will be nested within the KIULARCO and is part of the KIULARCO protocol, which has received initial ethical approval from Ifakara Health Institute institutional review board (IHI/IRB/No 16-2006) and National Institute for Medical Research (NIMR) (NIMR/HQ/R.8a/Vol.IX/620). Clearance for




Figure 1. Study design of the stigma-directed prospective observational pre-post study.

Table 1. Eligibility criteria to receive stigma services during the post-implementation phase.

Inclusion criteria	Exclusion criteria
<ul style="list-style-type: none"> • Adults (≥15 years of age) • Newly diagnosed with HIV at the CDCI at the SFRH • Written informed consent to KIULARCO 	<ul style="list-style-type: none"> • Current hospitalization for any reason • Indeterminate HIV test result • Non-consenting for KIULARCO • Too sick to answer the questions

CDCI = Chronic Disease Clinic Ifakara; SFRH = St. Francis Referral Hospital; KIULARCO = Kilombero and Ulanga Antiretroviral Cohort

stigma interventions with updated consent and documentation were added to the KIULARCO protocol and approved by ethical committees within a respective amendment (IHI/IRB/AMM/No:08-2020 and IHI/IRB/AMM/No: 2-2021; NIMR/HQ/R.8c/Vol.I/764 and NIMR/HQ/R.8c/Vol.I/896). Patients will sign an informed consent form (see *Extended data*⁴⁴). Patients can either accept or reject stigma services. Patients who do not wish to receive them, receive standard of care clinical workup and treatment. Lay counselors will obtain oral consent from patients before asking stigma- and depression specific questions.

Stigma services delivered by a lay person living with HIV Well-trained lay counselors openly living with HIV (one male, one female) will be responsible for the conduction of all stigma-related services. Lay counsellors will be trained by different medical professionals and professional counselors over a duration of 4 weeks using a standard operating procedure aligned to the national guidelines (competency list in *Extended data*, Table S2⁴⁴). The lay counselors will be selected from patients at our clinic and have either been living with HIV since birth or for multiple years. They will be trained in all aspects of the stigma services, including data entry and handling. Strategies to ensure program fidelity include standardized materials, structured education training, ongoing supervision, and training a sufficient number of lay counselors.

1) Post-test counseling

On the day of diagnosis, standardized post-test counseling with patients will be conducted by the lay counselor immediately after a newly confirmed positive HIV test and the first information is given by the professional counselor. The stigma-directed counselling will covers social support/disclosure, emotions/fears, basic knowledge of HIV transmission and steps towards positive health and living positively with HIV.

2) Post-test VAT

Thereafter, the lay counselor will show the patient a 16-minute video on a portable device (Samsung Tablet A, 10.1 inch) containing experiences of HIV-positive persons and information from healthcare workers from our clinic. While others have used video in similar situations^{45,46} our goal was to produce a video reflecting and capturing the local socio-cultural context and the hospital itself to increase relatability and recognition value in patients. The video is in Swahili language and use is restricted to the clinical purpose.

The recorded patients are PLWHA from Ifakara and health care workers from the CDCI who agreed to participate to develop a video to be shown to newly diagnosed clients. The study investigator obtained written informed consent from all video participants before the shooting took place. The investigators explained to each participant the purpose of the video.

In the video PLWHA report on how HIV and HIV-related stigma have affected their lives and how they integrated HIV into their daily lives. They talk about living positively with HIV, challenges they experienced, experience with disclosure, social support, and ART adherence. The video also contains information provided by healthcare professionals on HIV/AIDS such as disease manifestations, transmission, and treatment.

Increased knowledge aims at reduction of fear and internalized stigma.

So far, no studies have examined content or other aspects of such an educational video, despite these having been used in similar situations^{23,36}.

3) Group support therapy and group health education

Daily peer group support therapy and group health education will serve to discuss questions around HIV-related stigma and increase knowledge about HIV/AIDS for all interested patients. Group support therapy and group health education are based on participant-centered learning, social support⁴⁷, strength-based non-judgmental communication⁴⁸, identifying communalities⁴⁹, creating hope⁵⁰ and active listening⁵¹. **Culturally adapted learning materials** and **implementation tools** will be used. Each session will consist of an introduction including group therapy rules, an opportunity to share problems^{50,52} and combines peer group support therapy with group health education²⁷. We will use active learning techniques like role play, brainstorming, discussions with active listening⁵¹, storytelling and **exercises with picture cards**. Five sessions, comprising of a duration of 30-45min and maximum participant number of 10 patients per session, with different standardized content for group support therapy, including integration of HIV and ART in everyday life, internalized and community stigma, depression and coping as well as disclosure and relationship, have been elaborated upon (Table S1, *Extended data*⁴⁴). The positive effects of implementation of group support therapy and group health education^{27,35} within similar settings has been described previously.

4) Mobile Health (mHealth)

We will be using mHealth to ensure early care engagement during the first weeks after diagnosis. In addition to appointment reminders delivered by automated text messages (SMS) to all patients attending the clinic as standard of care at the CDCI, the lay counselor will call newly diagnosed patients 3 days before their first and second clinical follow-up after diagnosis (two-way communication voice call). To minimize fear of involuntary HIV disclosure to people overhearing the conversation, the health message only encourages participants to care for their health with an HIV neutral content and not mentioning the health care facility or personal patient information⁵³. Mobile phone calls to remind patients about their clinic visits has successfully been integrated into HIV care^{34,40,41,54} and support the patient to feel valued by the clinic with a wide acceptability^{53,54}.

Depression, stigma and patient satisfaction questionnaires

Stigma- and depression-related questionnaires will be completed together with the patient one month after start of ART, to assess stigma and create awareness. The questions will also be a starting point to discuss those topics with the lay counselors. Both questionnaires, for depression and stigma, are versions of adapted scales previously translated into Kiswahili and validated in Tanzania⁵⁵⁻⁵⁷ and have been successfully used in sub-Saharan Africa⁵⁸⁻⁶¹. Signs of depression are measured using the Patient Health Questionnaire-9 (PHQ-9)^{56,57}, which analyzes symptoms of depression on a 4-point Likert-type scale^{58,59}. An adapted version of the Berger's stigma scale⁶² will be used

to measure internalized and perceived stigma within PLWHA on a 7-items questionnaire rated on a 5-point Likert-type scale⁵⁵. Any patient in whom major depression (depression score of ≥ 9) or suicidal ideation is detected will be evaluated by the clinician and if major depressive disorder is confirmed they will be managed according to Tanzanian guidelines and the ICD10 diagnosis is recorded in the patient system. In an end-of-study questionnaire at 6 months, we will additionally ask for patient satisfaction and acceptance of each service by using the “Net Promoter Score” (NPS), a 0-to-10 numerical customer satisfaction metric^{61,63}. The net promoter score was effectively adapted for use in the medical field to measure patient experience, also within sexual and reproductive health clinics^{61,64}.

Outcomes

The primary endpoint in this study will be ‘linkage to care’. Despite ‘linkage to care’ being a key HIV indicator for public health monitoring, multiple definitions used in different studies make comparisons challenging¹⁰. For Stig2health ‘linkage to care’ will be defined as “being in active care and on ART at 2 months”. The time period was chosen as there might be patients with postponed ART start due to co-infection with tuberculosis (Tb) or cryptococcal meningitis. All secondary and tertiary outcomes can be found in Table 2. Tertiary endpoints were added at a later stage of the study protocol development according to recent research done by our study team on the ‘linkage to care framework’¹⁰.

Table 2. Secondary and tertiary endpoints.

Endpoint	Timepoint after enrolment into KIULARCO	Definition
Secondary endpoints		
<u>Pre- and post-implementation cohort</u>		
Retention in care	at 6 months*	“patients known to be alive and receiving ART at the end of the follow-up period of 6 months ⁶⁵ ”.
Viral load suppression	at 6 months	“having a viral load of less than 400 copies of HIV RNA per milliliter ^{66,67} ”
Death	until 6 months	
Clinical failure	until 6 months	“occurrence of any new WHO AIDS-defining disease, death or loss to follow-up at any visits ⁶⁸⁻⁷⁰ ”
Proportion of patients with self-reported depression comparing pre-/post cohorts	at 6 months	Categorical scores and continuous scores
Level of self-reported stigma comparing pre-/post cohorts	at 6 months	Continuous scores
<u>Post-implementation cohort only</u>		
Proportion and changes of patients with self-reported depression within the post cohort	at 1 month**, 6 months	Categorical scores and continuous scores
Level and changes of self-reported stigma within the post cohort	at 1 month, 6 months	Continuous scores
Patient satisfaction of each individual stigma service	at 6 months	Categorical scores and continuous scores
Tertiary endpoints		
Proportion of participants with:		
i) a first laboratory evaluation	at 3 months***	Having a laboratory result form with sampling date within the given time period following enrolment.
ii) a first clinical evaluation	at 3 months	Having a clinical visit with date within the given time period following enrolment.
iii) with ART initiation	at 3 months	Being initiated on ART within the given time period following enrolment.
iv) a clinical follow-up visit following ART initiation	at 3 months	Having a clinical visit with date after ART initiation within the given time period following enrolment.

*6 months with a time window from 6 up to 12 months, **1 month with a time window from 2–6 weeks, ***3 months without a time window; ART= Antiretroviral therapy; RNA= ribonucleic acid, HIV= human immunodeficiency virus; AIDS= acquired immunodeficiency syndrome; KIULARCO = Kilombero and Ulanga Antiretroviral Cohort

Participant timeline

The lay counseling and video intervention will be offered the day of diagnosis and enrollment into KIULARCO. On patient request it can be moved to day 1. During routine follow-up visits scheduled according to the National AIDS control program at month 1, 2, 3, 6, 9 and 12, group support therapies will be offered. The mHealth phone call will be conducted 3 days prior to the first two clinical follow-up visits. Depression and stigma questionnaires for the post-implementation cohort will be filled during the first month (2–6 weeks) and at the first viral load measurement visit that is usually conducted at 6 months after enrolment into KIULARCO, with an additional assessment of patient satisfaction of stigma services at the same time point. As the 6-month time point may be missed in some occasions, a window period up to 12 months after enrolment into KIULARCO will be accepted. End of study will be 31st August 2022. Stigma and depression questionnaires for the

pre-implementation cohort will be assessed in patients at their 6 month follow-up visit if they were not subjected to any stigma intervention. Timing of stigma specific patient visits and data collection can be found in [Table 3](#).

Sample size

For the primary endpoint – linkage to care – we assumed an increase from 70 to 80% based on a systematic review on strategies to improve linkage to care rates in different urban areas in sub-Saharan Africa and a linkage to care rate of 68% within KIULARCO in 2018⁷¹. The sample size calculation was based on cohort data from 2018, when 511 adults were newly diagnosed and of whom 346 (68%) were in care and on ART at 2 months. Assuming a pre-intervention cohort of 511 adults, a 10% increase in linkage (from 70 to 80%), a two-sided type I error rate of 5%, and 90% power, 321 adults are required for the post-intervention group. The full sample size calculation can

Table 3. Timing of Stig2Health related patient visits and data collection within routine follow-up visit scheduling according the National AIDS control program.

Study period Time point (Window period)	Baseline visit	Follow-up visits	
	Day 1 (day 0–1)	Month 1 (week 2–6)	Month 6 (week 26–52)
Patient Information and Informed Consent of KIULARCO (registration)	X		
Post-test counselling (professional counsellor)	X		
Stigma baseline services			
- Lay Counselling	X		
- Video Intervention	X		
Stigma mHealth service (phone calls done prior to clinical visits)		X	
Stigma Invitation to group therapy sessions and group health education		X	X
Vital signs (nurse)	X	X	X
Visit by study doctor			
- History			
- Physical Examination	X	X	X
- Drug prescription			
Laboratory testing (venipuncture)			
- Blood: Full blood Count	X		X
- Liver enzymes and Creatinine	X		X
- Cryptococcus, tuberculosis, hepatitis, syphilis according KIULARCO protocol	X		
- Viral load			X
- CD4 cell count	X		X
Self-reported adherence/Pill box return		X	X
PHQ-9 depression questionnaire		X	X
Adapted Berger stigma questionnaire		X	X
Net Promoter Score			X

*First follow-up visit at 6 months where the first viral load is assessed, with a window period up to 12 months will be allowed, KIULARCO = Kilombo and Ulanga Antiretroviral Cohort, CD4 = Cluster of differentiation 4; PHQ-9 = patient health questionnaire-9

be found in the *Extended data*⁴⁴. Based on the enrolment into KIULARCO from 2017 of >500 patients, we expect to reach the required sample size within one year.

Data collection and management

Numbers from routine testing activities captured in books will be transferred to an excel sheet saved on a local password-protected server. Data from patients enrolled in KIULARCO will be entered into an electronic medical record system (openMRS). Documentation of stigma services will be done using a counselling form within openMRS and will be filled by the clinician and the professional counselor. In addition, stigma and depression questionnaires will be captured in Open Data Kit (ODK) by the lay counselor with final scores entered into openMRS.

Data from the stigma and depression questionnaires will be extracted weekly, cleaned and stored on a study computer and in a data sharing platform (Alfresco, hosted by Swiss TPH). The routinely collected KIULARCO data will be extracted from the IHI sever and processed for research use quarterly. The data extraction for this study will be done at the end of study data collection and will be stored in a secured cloud system.

Confidentiality

Participant data will be anonymized and accessible only to authorized study personnel. Anonymity of the participants will be guaranteed when presenting the data at scientific meetings or publishing them in scientific journals. All medical information obtained within this study will be considered confidential.

Statistical methods

Participant characteristics at enrolment into KIULARCO will be described using descriptive statistics for both primary outcome and secondary outcomes. The association between the primary outcome and the intervention bundle will be assessed using the logistic regression model and the measure of effect will be estimated in terms of adjusted odds ratio and thereafter risk differences (with 95% confidence interval) will be estimated^{72,73}. The variable selection procedure for adjusted models will be explained in a statistical analysis plan.

The linkage to care framework will be constructed based on four stages defined within three months since being enrolled into KIULARCO. Steps to be considered include (a) laboratory assessment; (b) clinical evaluation; (c) ART initiation; and (d) clinical follow-up after treatment initiation. The multinomial linear regression model will be used to assess the association between linkage stages and intervention bundle. The risk difference will be estimated using the approach described in the primary outcome.

The association between binary secondary outcomes such as retention in care, viral load, and self-reported-depression and intervention bundle will be analyzed using logistic regression models. Self-reported stigma and depression scores will be analyzed using linear regression models. Further, within the post-intervention group, the change in self-reported stigma and depression will be analyzed using linear regression models.

To assess patient satisfaction with individual services^{61,64} in the post-cohort, the proportion of detractors, passive, and promoters will be estimated for each intervention and the final **net promoter score** will be calculated by deducting detractors from promoters.

There are two potential sources of missing data which include loss to follow-up and transfer to other HIV clinics. Transfers will affect primary and secondary outcomes. Lost to follow-up will be an additional source of missing data in all secondary outcomes. In case the percentage of the missing data will exceed 10% of all recruited participants, missing data techniques will be used to address the bias due to omission. Details will be provided in a separate full statistical analysis plan. Data management and analyses will be done using Stata software version 16 and SAS version 9.4 (SAS Institute Inc., Cary, NC, USA).

Discussion

With a four-component bundle of stigma-directed services we expect to increase 'linkage to care' by 10% from 70 to 80% post-implementation. We expect to gain knowledge on the value of stigma-directed services and to observe an increase in linkage to care proportions, virological suppression rates and retention in care within the first year. By increasing linkage to care and retention in care, the HIV-cascade in a rural setting in sub-Saharan Africa should improve. We expect to further understand the complex social, cultural, and intrapersonal construct of stigma and to get to know more about the fears and concerns of patients, on how to target them and to be able to encourage patients to take steps towards a positive health behavior. HIV-directed stigma and discrimination are still crucial barriers that affect accessibility and acceptability of HIV healthcare services. With this study we hope to provide guidance on how to integrate simple, low-cost, and sustainable stigma-related services into routine care in rural sub-Saharan Africa, but also raise awareness on HIV-related stigma and health outcomes in this setting and worldwide. The results of this prospective stigma-directed study may provide more insight into task shifting to and collaborations with lay health care workers in primary health care facilities in low-resource settings and has the potential to lead to policy changes regarding integration of stigma and mental health-directed services in sub-Saharan Africa.

Study status

Follow-up period until August 31st, 2022

Dissemination plan

Dissemination will occur through publication in a peer-reviewed journal. Additional dissemination will occur through presentations at conferences nationally and internationally and by informing district health authorities.

Data availability

Underlying data

No underlying data are associated with this article

Extended data

Zenodo: Extended data for Stig2Health implementational study: Stigma-directed services to improve 'linkage to care'

for people living with HIV in rural Tanzania. <https://doi.org/10.5281/zenodo.5916625>⁴⁴

This project contains the following extended data:

- KIULARCO_ICTenglish_2020_11_15.pdf (KIULARCO Cohort Informed Consent form in English)
- KIULARCO_ICTswahili_2020_11_15.pdf (KIULARCO Cohort Informed Consent form in Swahili)
- Stig2Health Adapted Berger Stigma Scale.pdf
- Stig2Health End of study questionnaire.pdf
- Stig2Health Information for Sample size calculation.pdf
- Stig2Health PHQ-9 Depression questionnaire swahili.pdf
- Stig2Health Table S1- Outline group support therapy and group health education.pdf
- Stig2Health Table S2 Overview lay counselor competency list.pdf

Data are available under the terms of the [Creative Commons Attribution 4.0 International license \(CC-BY 4.0\)](#).

Permission to publish the adapted Berger stigma scale and PHQ-9 depression questionnaire in Swahili was received from Prof. Sylvia Kaaya on 28.01.2022.

The post-test video is not available publicly, because consent for publication was not obtained and the publication could

pose a threat to confidentiality of participants. It was agreed that this video is only available for research and clinical use and not for public viewing to ensure confidentiality of participants. Inquiries on the video are coordinated through the University Hospital Basel. Requests will be handled by Prof. Dr. med. Maja Weisser m.weisser@unibas.ch

Acknowledgements

The study team would like to acknowledge Prof. Sylvia Kaaya and colleagues for making their depression and stigma questionnaires available to our research team.

We thank all team members of the KIULARCO study group as well as patients included in KIULARCO for providing data.

KIULARCO study group

Aschola Asantiel, Farida Bani, Manuel Battegay, Theonestina Byakuzana, Adolphina Chale, Francisca Chuwa, Ivana di Salvo, Gideon Francis, Hansjakob Furrer, Tracy Glass, Yvone Haridas, Speciosa Hwaya, Aneth V Kalinjuma, Joshua Kapunga, Bryson Kasuga, Andrew Katende, Namvua Kimera, Yassin Kisunga, Olivia Kitau, Bernard Kivuma, Thomas Klimkait, Ezekiel Luoga, Herry Mapesi, Slyakus Mlembe, Mengi Mkulila, Margareth Mkusa, Dorcas K Mnzava, Getrud J Mollel, Lilian Moshi, Germana Mossad, Dolores Mpundunga, Athumani Mtandanguo, Selerine Myeya, Sanula Nahota, Robert C. Ndege, Reginna Ndaki, Suzan Ngahyoma, Agatha Ngululika, Jacopo Nicoletti, Alex John Ntamungiro, Amina Nyuri, James Okuma, Daniel H Paris, Jacqueline Nkouabi, Aloyce Sambuta, Leila Samson, Elizabeth Senkoro, George Sigalla, Jenifa Tarimo, Juerg Utzinger, Fiona Vanobberghen, Maja Weisser, John Wigay, Herieth Wilson.

References

1. Gardner EM, McLees MP, Steiner JF, *et al.*: **The spectrum of engagement in HIV care and its relevance to test-and-treat strategies for prevention of HIV infection.** *Clin Infect Dis.* 2011; **52**(6): 793–800. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
2. Kranzer K, Govindasamy D, Ford N, *et al.*: **Quantifying and addressing losses along the continuum of care for people living with HIV infection in sub-Saharan Africa: a systematic review.** *J Int AIDS Soc.* 2012; **15**(2): 17383. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
3. Barnabas RV, Celum C: **Closing the gaps in the HIV care continuum.** *PLoS Med.* 2017; **14**(11): e1002443. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
4. Plazy M, Farouki KE, Iwuji C, *et al.*: **Access to HIV care in the context of universal test and treat: challenges within the ANRS 12249 TasP cluster-randomized trial in rural South Africa.** *J Int AIDS Soc.* 2016; **19**(1): 20913. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
5. Church K, Machiyama K, Todd J, *et al.*: **Identifying gaps in HIV service delivery across the diagnosis-to-treatment cascade: findings from health facility surveys in six sub-Saharan countries.** *J Int AIDS Soc.* 2017; **20**(1): 21188. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
6. Fox MP, Rosen S: **Patient retention in antiretroviral therapy programs up to three years on treatment in sub-Saharan Africa, 2007-2009: systematic review.** *Trop Med Int Health.* 2010; **15** Suppl 1(s1): 1–15. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
7. Haber N, Tanser F, Bor J, *et al.*: **From HIV infection to therapeutic response: a population-based longitudinal HIV cascade-of-care study in KwaZulu-Natal, South Africa.** *Lancet HIV.* 2017; **4**(5): e223–e30. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
8. Ntamungiro AJ, Muri L, Glass TR, *et al.*: **Strengthening HIV therapy and care in rural Tanzania affects rates of viral suppression.** *J Antimicrob Chemother.* 2017; **72**(7): 2069–2074. [PubMed Abstract](#) | [Publisher Full Text](#)
9. Reddy EA, Agala CB, Maro VP, *et al.*: **Test site predicts HIV care linkage and antiretroviral therapy initiation: a prospective 3.5 year cohort study of HIV-positive testers in northern Tanzania.** *BMC Infect Dis.* 2016; **16**: 497. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
10. Magnolini R, Senkoro E, Vanobberghen F, *et al.*: **“Linkage to care” among people living with HIV - definition in the era of “universal test and treat” in a sub-Saharan African setting.** *Swiss Med Wkly.* 2021; **151**: w20535. [PubMed Abstract](#) | [Publisher Full Text](#)
11. Sengupta S, Banks B, Jonas D, *et al.*: **HIV interventions to reduce HIV/AIDS stigma: a systematic review.** *AIDS Behav.* 2011; **15**(6): 1075–87. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
12. Layer EH, Kennedy CE, Beckham SW, *et al.*: **Multi-level factors affecting entry into and engagement in the HIV continuum of care in Iringa, Tanzania.** *PLoS One.* 2014; **9**(8): e104961. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
13. Earnshaw VA, Chaudoir SR: **From conceptualizing to measuring HIV stigma: a review of HIV stigma mechanism measures.** *AIDS Behav.* 2009; **13**(6): 1160–77. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
14. Stangl AL, Earnshaw VA, Logie CH, *et al.*: **The Health Stigma and Discrimination Framework: a global, crosscutting framework to inform research, intervention development, and policy on health-related stigmas.** *BMC Med.* 2019; **17**(1): 31. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

15. Gesesew HA, Tesfay Gebremedhin A, Demissie TD, *et al.*: **Significant association between perceived HIV related stigma and late presentation for HIV/AIDS care in low and middle-income countries: A systematic review and meta-analysis.** *PLoS One.* 2017; **12**(3): e0173928. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
16. Lyimo RA, Stutterheim SE, Hospers HJ, *et al.*: **Stigma, disclosure, coping, and medication adherence among people living with HIV/AIDS in Northern Tanzania.** *AIDS Patient Care STDS.* 2014; **28**(2): 98–105. [PubMed Abstract](#) | [Publisher Full Text](#)
17. Rispel LC, Cloete A, Metcalf CA: **'We keep her status to ourselves': experiences of stigma and discrimination among HIV-discordant couples in South Africa, Tanzania and Ukraine.** *SAHARA J.* 2015; **12**(1): 10–7. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
18. Lyimo RA, de Bruin M, van den Boogaard J, *et al.*: **Determinants of antiretroviral therapy adherence in northern Tanzania: a comprehensive picture from the patient perspective.** *BMC Public Health.* 2012; **12**: 716. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
19. Raja SK, Aggarwal S, Raman KR, *et al.*: **From businesswoman to banana vendor: mental health implications of HIV stigma in Tanzania.** *AIDS.* 2016; **30**(17): N33–N5. [PubMed Abstract](#) | [Publisher Full Text](#)
20. Dow DE, Turner EL, Shayo AM, *et al.*: **Evaluating mental health difficulties and associated outcomes among HIV-positive adolescents in Tanzania.** *AIDS Care.* 2016; **28**(7): 825–33. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
21. Ramos JV, Mmbaga BT, Turner EL, *et al.*: **Modality of Primary HIV Disclosure and Association with Mental Health, Stigma, and Antiretroviral Therapy Adherence in Tanzanian Youth Living with HIV.** *AIDS Patient Care STDS.* 2018; **32**(1): 31–7. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
22. Doyle AM, Plummer ML, Weiss HA, *et al.*: **Concurrency and other sexual partnership patterns reported in a survey of young people in rural Northern Tanzania.** *PLoS One.* 2017; **12**(8): e0182567. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
23. Marks IM, Kenwright M, McDonough M, *et al.*: **Saving clinicians' time by delegating routine aspects of therapy to a computer: a randomized controlled trial in phobia/panic disorder.** *Psychol Med.* 2004; **34**(1): 9–17. [PubMed Abstract](#) | [Publisher Full Text](#)
24. Geng EH, Odeny TA, Lyamuya R, *et al.*: **Retention in Care and Patient-Reported Reasons for Undocumented Transfer or Stopping Care Among HIV-Infected Patients on Antiretroviral Therapy in Eastern Africa: Application of a Sampling-Based Approach.** *Clin Infect Dis.* 2016; **62**(7): 935–44. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
25. Tran BX, Ho RCM, Ho CSH, *et al.*: **Depression among Patients with HIV/AIDS: Research Development and Effective Interventions (GAP_{RESEARCH}).** *Int J Environ Res Public Health.* 2019; **16**(10): 1772. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
26. Hartog K, Hubbard CD, Krouwer AF, *et al.*: **Stigma reduction interventions for children and adolescents in low- and middle-income countries: Systematic review of intervention strategies.** *Soc Sci Med.* 2020; **246**: 112749. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
27. Nakimuli-Mpungu E, Musisi S, Wamala K, *et al.*: **The Effect of Group Support Psychotherapy Delivered by Trained Lay Health Workers for Depression Treatment Among People with HIV in Uganda: Protocol of a Pragmatic, Cluster Randomized Trial.** *JMIR Res Protoc.* 2017; **6**(12): e250. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
28. Chibanda D, Mesu P, Kajawa L, *et al.*: **Problem-solving therapy for depression and common mental disorders in Zimbabwe: piloting a task-shifting primary mental health care intervention in a population with a high prevalence of people living with HIV.** *BMC Public Health.* 2011; **11**: 828. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
29. Callaghan M, Ford N, Schneider H: **A systematic review of task- shifting for HIV treatment and care in Africa.** *Hum Resour Health.* 2010; **8**: 8. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
30. Nyblade L, Srinivasan K, Mazur A, *et al.*: **HIV Stigma Reduction for Health Facility Staff: Development of a Blended- Learning Intervention.** *Front Public Health.* 2018; **6**: 165. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
31. Ng YP, Rashid A, O'Brien F: **Determining the effectiveness of a video-based contact intervention in improving attitudes of Penang primary care nurses towards people with mental illness.** *PLoS One.* 2017; **12**(11): e0187861. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
32. Brabcová D, Kohout J, Weberová V, *et al.*: **Educational video and story as effective interventions reducing epilepsy-related stigma among children.** *Epilepsy Behav.* 2017; **69**: 12–7. [PubMed Abstract](#) | [Publisher Full Text](#)
33. Sajatović M, Herrmann LK, Van Doren JR, *et al.*: **A randomized prospective pilot trial of Web-delivered epilepsy stigma reduction communications in young adults.** *Epilepsia.* 2017; **58**(11): 1946–54. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
34. Abdulrahman SA, Rampal L, Ibrahim F, *et al.*: **Mobile phone reminders and peer counseling improve adherence and treatment outcomes of patients on ART in Malaysia: A randomized clinical trial.** *PLoS One.* 2017; **12**(5): e0177698. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
35. Hamilton ARL, le Roux KWDP, Young CW, *et al.*: **Mentor Mothers Zithulele: exploring the role of a peer mentorship programme in rural PMTCT care in Zithulele, Eastern Cape, South Africa.** *Paediatr Int Child Health.* 2020; **40**(1): 58–64. [PubMed Abstract](#) | [Publisher Full Text](#)
36. Noar SM, Black HG, Pierce LB: **Efficacy of computer technology-based HIV prevention interventions: a meta-analysis.** *AIDS.* 2009; **23**(1): 107–15. [PubMed Abstract](#) | [Publisher Full Text](#)
37. Wroe EB, Hedt-Gauthier BL, Franke MF, *et al.*: **Depression and patterns of self-reported adherence to antiretroviral therapy in Rwanda.** *Int J STD AIDS.* 2015; **26**(4): 257–61. [PubMed Abstract](#) | [Publisher Full Text](#)
38. Memiah P, Shumba C, Etienne-Mesubi M, *et al.*: **The effect of depressive symptoms and CD4 count on adherence to highly active antiretroviral therapy in sub-Saharan Africa.** *J Int Assoc Provid AIDS Care.* 2014; **13**(4): 346–52. [PubMed Abstract](#) | [Publisher Full Text](#)
39. e Silva AC, Reis RK, Nogueira JA, *et al.*: **Quality of life, clinical characteristics and treatment adherence of people living with HIV/AIDS.** *Rev Lat Am Enfermagem.* 2014; **22**(6): 994–1000. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
40. Elul B, Lamb MR, Lahuerta M, *et al.*: **A combination intervention strategy to improve linkage to and retention in HIV care following diagnosis in Mozambique: A cluster-randomized study.** *PLoS Med.* 2017; **14**(11): e1002433. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
41. McNairy ML, Lamb MR, Gachuhi AB, *et al.*: **Effectiveness of a combination strategy for linkage and retention in adult HIV care in Swaziland: The Link4Health cluster randomized trial.** *PLoS Med.* 2017; **14**(11): e1002420. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
42. Vanobberghen F, Letang E, Gamell A, *et al.*: **A decade of HIV care in rural Tanzania: Trends in clinical outcomes and impact of clinic optimisation in an open, prospective cohort.** *PLoS One.* 2017; **12**(7): e0180983. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
43. Letang E, Kalinjuma AV, Glass TR, *et al.*: **Cohort profile: The Kilombero and Ulanga Antiretroviral Cohort (KIULARCO) - A prospective HIV cohort in rural Tanzania.** *Swiss Med Wkly.* 2017; **147**: w14485. [PubMed Abstract](#) | [Publisher Full Text](#)
44. Magnolini R, Senkoro E, Kalinjuma AV, *et al.*: **Extended data for Stig2Health implementation study: Stigma-directed services to improve 'linkage to care' for people living with HIV in rural Tanzania.** 2022. <http://www.doi.org/10.5281/zenodo.5916625>
45. Calderon Y, Cowan E, Nickerson J, *et al.*: **Educational effectiveness of an HIV pretest video for adolescents: a randomized controlled trial.** *Pediatrics.* 2011; **127**(5): 911–6. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
46. Calderon Y, Leider J, Hailpern S, *et al.*: **A randomized control trial evaluating the educational effectiveness of a rapid HIV posttest counseling video.** *Sex Transm Dis.* 2009; **36**(4): 207–10. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
47. Lifson AR, Workneh S, Hailemichael A, *et al.*: **Implementation of a Peer HIV Community Support Worker Program in Rural Ethiopia to Promote Retention in Care.** *J Int Assoc Provid AIDS Care.* 2017; **16**(1): 75–80. [PubMed Abstract](#) | [Publisher Full Text](#)
48. Siril H, Kaale A, Minja A, *et al.*: **Improving communication about HIV prevention among people living with HIV and their at-risk social network members in Dar es Salaam, Tanzania.** *Cogent Med.* 2019; **6**: 1600230. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
49. Brabender V: **Introduction to group therapy.** New York, NY: New York, NY: Wiley; 2002. [Reference Source](#)
50. Yalom ID: **Theorie und Praxis der Gruppenpsychotherapie.** ein Lehrbuch. 4., völlig überarb. und erw. Aufl. ed. München: München: Pfeiffer; 1996. [Reference Source](#)
51. Nakimuli-Mpungu E, Wamala K, Okello J, *et al.*: **Developing a culturally sensitive group support intervention for depression among HIV infected and non-infected Ugandan adults: a qualitative study.** *J Affect Disord.* 2014; **163**: 10–7. [PubMed Abstract](#) | [Publisher Full Text](#)
52. Nakimuli-Mpungu E, Wamala K, Okello J, *et al.*: **Group support psychotherapy for depression treatment in people with HIV/AIDS in northern Uganda: a single-centre randomised controlled trial.** *Lancet HIV.* 2015; **2**(5): e190–9. [PubMed Abstract](#) | [Publisher Full Text](#)
53. Rodrigues R, Poongulali S, Balaji K, *et al.*: **'The phone reminder is important, but will others get to know about my illness?' Patient perceptions of an mHealth antiretroviral treatment support intervention in the HIVIND trial in South India.** *BMJ Open.* 2015; **5**(11): e007574. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
54. Sidney K, Antony J, Rodrigues R, *et al.*: **Supporting patient adherence to antiretrovirals using mobile phone reminders: patient responses from South India.** *AIDS Care.* 2012; **24**(5): 612–7. [PubMed Abstract](#) | [Publisher Full Text](#)
55. Smith Fawzi MC, Siril H, Liu Y, *et al.*: **Agents of change among people living**

- with HIV and their social networks: stepped-wedge randomised controlled trial of the **NAMWEZA** intervention in Dar es Salaam, Tanzania. *BMJ Glob Health*. 2019; 4(3): e000946.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
56. Smith Fawzi MC, Ngakongwa F, Liu Y, *et al.*: **Validating the Patient Health Questionnaire-9 (PHQ-9) for screening of depression in Tanzania.** *Neurol Psychiatry Brain Res*. 2019; 31: 9–14.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
57. Kroenke K, Spitzer RL, Williams JB: **The PHQ-9: validity of a brief depression severity measure.** *J Gen Intern Med*. 2001; 16(9): 606–13.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
58. Akena D, Joska J, Obuku EA, *et al.*: **Sensitivity and specificity of clinician administered screening instruments in detecting depression among HIV-positive individuals in Uganda.** *AIDS Care*. 2013; 25(10): 1245–52.
[PubMed Abstract](#) | [Publisher Full Text](#)
59. Cholera R, Gaynes BN, Pence BW, *et al.*: **Validity of the Patient Health Questionnaire-9 to screen for depression in a high-HIV burden primary healthcare clinic in Johannesburg, South Africa.** *J Affect Disord*. 2014; 167: 160–6.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
60. Oke OO, Akinboro AO, Olanrewaju FO, *et al.*: **Assessment of HIV-related stigma and determinants among people living with HIV/AIDS in Abeokuta, Nigeria: A cross-sectional study.** *SAGE Open Med*. 2019; 7: 2050312119869109.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
61. Koladycz R, Fernandez G, Gray K, *et al.*: **The Net Promoter Score (NPS) for Insight Into Client Experiences in Sexual and Reproductive Health Clinics.** *Glob Health Sci Pract*. 2018; 6(3): 413–24.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
62. Berger BE, Ferrans CE, Lashley FR: **Measuring stigma in people with HIV: psychometric assessment of the HIV stigma scale.** *Res Nurs Health*. 2001; 24(6): 518–29.
[PubMed Abstract](#) | [Publisher Full Text](#)
63. Reichheld FF: **The one number you need to grow.** *Harv Bus Rev*. 2003; 81(12): 46–54, 124.
[PubMed Abstract](#)
64. Kinney WC: **A simple and valuable approach for measuring customer satisfaction.** *Otolaryngol Head Neck Surg*. 2005; 133(2): 169–72.
[PubMed Abstract](#) | [Publisher Full Text](#)
65. Rosen S, Fox MP, Gill CJ: **Patient retention in antiretroviral therapy programs in sub-Saharan Africa: a systematic review.** *PLoS Med*. 2007; 4(10): e298.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
66. Cohen MS, Chen YQ, McCauley M, *et al.*: **Antiretroviral Therapy for the Prevention of HIV-1 Transmission.** *N Engl J Med*. 2016; 375(9): 830–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
67. Cohen MS, Chen YQ, McCauley M, *et al.*: **Prevention of HIV-1 infection with early antiretroviral therapy.** *N Engl J Med*. 2011; 365(6): 493–505.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
68. Erb S, Letang E, Glass TR, *et al.*: **Health care provider communication training in rural Tanzania empowers HIV-infected patients on antiretroviral therapy to discuss adherence problems.** *HIV Med*. 2017; 18(9): 623–34.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
69. Kalinjuma AV, Glass TR, Weisser M, *et al.*: **Prospective assessment of loss to follow-up: incidence and associated factors in a cohort of HIV-positive adults in rural Tanzania.** *J Int AIDS Soc*. 2020; 23(3): e25460.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
70. Chi BH, Cantrell RA, Mwangi A, *et al.*: **An empirical approach to defining loss to follow-up among patients enrolled in antiretroviral treatment programs.** *Am J Epidemiol*. 2010; 171(8): 924–31.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
71. Koduah Owusu K, Adu-Gyamfi R, Ahmed Z: **Strategies To Improve Linkage To HIV Care In Urban Areas Of Sub-Saharan Africa: A Systematic Review.** *HIV AIDS (Auckl)*. 2019; 11: 321–32.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
72. Norton EC, Miller MM, Kleinman LC: **Computing adjusted risk ratios and risk differences in Stata.** *Stata J*. StataCorp LP, 2013; 13(3): 492–509.
[Publisher Full Text](#)
73. Localio AR, Margolis DJ, Berlin JA: **Relative risks and confidence intervals were easily computed indirectly from multivariable logistic regression.** *J Clin Epidemiol*. 2007; 60(9): 874–82.
[PubMed Abstract](#) | [Publisher Full Text](#)

Open Peer Review

Current Peer Review Status:  

Version 1

Reviewer Report 22 July 2022

<https://doi.org/10.21956/aasopenres.14499.r29267>

© 2022 Harklerode R. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Richelle Harklerode 

Institute for Global Health Sciences, University of California, San Francisco, San Francisco, CA, USA

The manuscript describes a study protocol for the Stig2Health project to improve linkage to HIV care services. The article is well written and will contribute to understanding if HIV-related stigma services can improve the care cascade in Tanzania. The following are points of potential revision:

1. Noted as a low-cost method, for sustainability please consider including the cost of the training and daily cost of lay counselors.
2. There are a number of types of stigma, the questionnaire seems to focus on stigma from those informed of patient's HIV status, it would be useful to state this within the article.
3. For the group counseling to be able to replicate the study, I recommend providing additional information of length of the five sessions, frequency/time between sessions, and tracking outcomes based on the number of sessions patients participated in.
4. The depression questionnaire only has the Swahili version uploaded. If the patient is found to be depressed what treatment is provided? It would be useful to note if the services are different for those with depression and for those without depression.
5. There is a shift in tense between sections of the manuscript, I recommend revision to be consistent.
6. As the study is in progress, it is unlikely to change the end of study questionnaire, although it seems fitting to ask if they found the services helpful, instead of asking if they would recommend to other people living with HIV, as this could be confusing since that is not the patient's role.

Is the rationale for, and objectives of, the study clearly described?

Yes

Is the study design appropriate for the research question?

Yes

Are sufficient details of the methods provided to allow replication by others?

Partly

Are the datasets clearly presented in a useable and accessible format?

Not applicable

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: HIV, HIV care cascade, disease surveillance, monitoring and evaluation

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Author Response 29 Jul 2022

Raphael Magnolini, University Hospital of Basel, Basel, Switzerland

1. We will include the costs of lay counselors in the main manuscript.
2. Amended, the article now states: "An adapted version of the Berger's stigma scale will be used to measure internalized and perceived stigma within PLWHA".
3. Amended.
4. The depression questionnaire (PHQ-9) is a translation of a publicly available questionnaire in English language. The next steps if depression was detected was amended in the text, although that falls outside of the scope of this research.
5. Amended, all in future tense.
6. We thank the reviewer for this recommendation, although as stated the study is ongoing and this cannot be incorporated at this stage. We included this question as established tool (NPS questionnaire).

Competing Interests: none

Reviewer Report 27 June 2022

<https://doi.org/10.21956/aasopenres.14499.r29251>

© 2022 Hoddinott G. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Graeme Hoddinott 

Desmond Tutu TB Centre, Department of Paediatrics and Child Health, Faculty of Medicine and Health Sciences, Stellenbosch University, Stellenbosch, South Africa

The manuscript describes a study protocol for the Stig2Health project. Overall, it is written clearly and all necessary information included. Three points of further clarification / revision may be useful:

1. The tense shifts between past, future, and present tense throughout. I suggest that as a protocol manuscript, this should be future tense throughout - perhaps confused because the project may in fact already have started?
2. Stangl *et al.*¹ provide an overall conceptual framework for HIV stigma. Different domains of stigma (e.g., enacted, anticipated, and internal) have different impacts on outcomes (e.g., care seeking) and are intervened on differently (e.g., post test counselling addresses anticipated and internal stigma domains, but not enacted). It may be useful to include a section on your conceptual framework of stigma and how these intersect with each intervention component.
3. It is unclear that each of the intervention components are of equal value to the outcome, or how these interface with each other. E.g., do they work as a package or does each address a different aspect of stigma that combined addresses the outcome? Again, it may be useful to further clarify why these intervention components were selected in combination.

Congratulations again to the co-authors and good luck with implementing the project.

References

1. Stangl A, Earnshaw V, Logie C, van Brakel W, et al.: The Health Stigma and Discrimination Framework: a global, crosscutting framework to inform research, intervention development, and policy on health-related stigmas. *BMC Medicine*. 2019; **17** (1). [Publisher Full Text](#)

Is the rationale for, and objectives of, the study clearly described?

Yes

Is the study design appropriate for the research question?

Yes

Are sufficient details of the methods provided to allow replication by others?

Yes

Are the datasets clearly presented in a useable and accessible format?

Not applicable

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Tuberculosis, HIV, HIV stigma, mixed data evaluations.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Author Response 29 Jul 2022

Raphael Magnolini, University Hospital of Basel, Basel, Switzerland

1. Amended, all in future tense.
2. Amended, we further elaborated on the different mechanisms of stigma.
3. Amended, only the full bundle of stigma-directed intervention to the outcomes can be assessed, individual services cannot be differentiated.

Competing Interests: none
