



Reliable estimates of wild boar populations by nocturnal distance sampling

Authors: Focardi, Stefano, Morgia, Valentina La, Montanaro, Paolo, Riga, Francesco, Calabrese, Alessandro, et al.

Source: Wildlife Biology, 2020(4)

Published By: Nordic Board for Wildlife Research

URL: <https://doi.org/10.2981/wlb.00694>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.



Reliable estimates of wild boar populations by nocturnal distance sampling

Stefano Focardi, Valentina La Morgia, Paolo Montanaro, Francesco Riga, Alessandro Calabrese, Francesca Ronchi, Paola Aragno, Marianne Scacco, Roberta Calmanti and Barbara Franzetti

S. Focardi ✉ (stefano.focardi@fi.isc.cnr.it), Istituto dei Sistemi Complessi, CNR, via Madonna del Piano 10, IT-50019 Sesto Fiorentino, Italy. – V. La Morgia, P. Montanaro, F. Riga, A. Calabrese, F. Ronchi, P. Aragno, M. Scacco, R. Calmanti and B. Franzetti, Inst. Superiore per la Protezione e la Ricerca Ambientale, Ozzano dell' Emilia (BO), Italy.

The wild boar is one of the most invasive species among large mammals in both its native and introduced ranges. This species represents a main threat for crops and biodiversity and a pest for the pig industry due to the rapid expansion of the African swine fever. Because of its peculiar life history traits, population control programmes and recreational hunting are usually unable to effectively reduce the number of wild boars. Therefore, a reliable approach based on appropriate, cost-effective, monitoring methodologies is urgently required. Effective monitoring should adopt effective sampling strategies, otherwise the detection of population trends can be erroneous and resulting in a mismatch of appropriate management actions. First, we review the status-of-the-art of wildlife monitoring with a special focus on wild boar and feral pigs. Then, we show that nocturnal distance sampling, carried out using thermal cameras, can be an effective monitoring technique for wild boar population assessment regardless of the characteristics of the sampled area. Using data from multiple surveys performed in four study areas in Italy, characterised by contrasting topography, habitats and level of environmental visibility, we found that the estimate of precision is generally good and almost independent of landscape conditions. A simple method to estimate visibility, which may empirically help wildlife managers to design effective nocturnal distance sampling surveys, is proposed. The bias of our population estimates is evaluated using simulations showing that in some areas the estimate is unbiased, while in others there is the tendency towards a negative bias. Based on reported results, we provide guidelines to perform nocturnal distance sampling of wild boar populations.

Keywords: distance sampling, FLIR, population density, population index, *Sus scrofa*, thermal imagery, wild boar

Wild boar *Sus scrofa* populations are often of great concern for the conservation of wildlife species and habitat, as well as for agricultural and breeding activities, inside both its native (Reimoser and Putman 2011, Massei et al. 2017, Van Phan Le et al. 2019) and introduced range (North America, Anderson et al. 2016, South America, Ballari et al. 2015, Cuevas et al. 2020 and Australia, Bengsen et al. 2014). Wild boar are responsible of intense damages to croplands and of severe impacts on biodiversity (Genov et al. 2018, Graitson et al. 2018). Furthermore, the number of road accidents they are involved in is also growing (Bobek et al. 2018), as well as the concern over disease transmission, which threatens public health (De Sabato et al. 2018, Dimzas et al. 2019) and the farm pig industry in many European countries (Ruiz-Fons

2017, Andraud et al. 2019). The wild boar is a good example of a native invader (sensu Carey et al. 2012) meaning that their overabundance threatens biodiversity, harms ecosystem processes and impacts human activities.

Recreational hunting and removal programmes, even by poisoning (Bengsen et al. 2014), are used to control wild boar populations worldwide but 'control' is questionable in its effectiveness (Keuling et al. 2016). Here, we address one of the most relevant issues that can negatively affect the design of effective population control programmes, i.e. how to set up efficient monitoring. For this purpose, first, we summarize the main principles in wildlife monitoring and we briefly review approaches applicable to wild boar populations (ENETWILD consortium et al. 2018). Second, we focus on the application of methods aimed to estimate its population density, and we investigate whether promising results obtained by nocturnal distance sampling (NDS) by Franzetti et al. (2012) can be generalized to different environmental contexts (e.g. characterized by variable visibility and different animal density). Then we evaluate, by

This work is licensed under the terms of a Creative Commons Attribution 4.0 International License (CC-BY) <<http://creativecommons.org/licenses/by/4.0/>>. The license permits use, distribution and reproduction in any medium, provided the original work is properly cited.

simulations, the amount of bias affecting our surveys at different levels of effort. Finally, we provide guidelines to design efficient NDS surveys of the wild boar.

Basic principles and techniques for wild boar monitoring

Population monitoring is a necessary step in wildlife and habitat management (Engeman et al. 2013, Stephens et al. 2015). In particular, accurate and precise monitoring is fundamental to factual decision-making (Nichols and Williams 2006). Monitoring is equally important in evaluating the performance of control programs and other management actions (Lyons et al. 2008). Population monitoring consists generally of: 1) estimating population density or 2) using relative abundance indexes or quantities related to population abundance (proxies). We expect a tradeoff between the precision and accuracy of these monitoring approaches and their costs, with proxies being less expensive. A sufficient precision of the estimator of population density or index is necessary to design an effective monitoring program (Gerrodette 1987). So, the precision of an estimator must be assessed also for management purposes, otherwise, we cannot infer the variation of population size that a survey is able to detect (Engeman 2005). Common errors in trend analysis include detecting a trend when it does not exist (type I), failing to detect population changes when they exist (type II) and detecting a trend in the opposite direction of the true one (type III) (Nuno et al. 2015). A well-designed survey should be able to reduce these errors to a-priori defined levels, prescribed by the management policy to be implemented, and by the minimum detectable difference to be detected (Skalski et al. 2005, Reynolds et al. 2011).

To reduce costs and complexities of monitoring, wildlife managers often rely on proxies. Anderson (2001) heavily criticized the use of population indexes in wildlife management on the ground of the presence of convenience sampling and uncontrolled impact of covariates. However, later Engeman (2003) contended that under certain circumstances, indexes can be appropriately computed. The use of indexes is common in studies on elusive, wide-ranging and/or low densities species. Nimmo et al. (2015) review evidence that indexes (tracks) in predators and in several other species reflect variation in population abundance; Kojola et al. (2014) successfully used snow track triangles to evaluate wolf *Canis lupus* abundance in Finland. Index calibration methods are also used routinely to assess lion *Panthera leo* population trends in Namibia (Dröge et al. 2020). However, recent research (Engeman et al. 2013, Stephens et al. 2015) cast a shadow on the use of proxies, which can be misleading and may introduce severe errors in management actions; for example, when there is not a proportional relationship between the abundance and the proxy (Iannuzzo et al. 2010, Koda and Fujita 2011). Under this circumstance, a proxy cannot demonstrate enough sensitivity at low population densities, and it can exhibit saturation when density is high. Consequently, it becomes useless in these two conditions (scarcity or overabundance), because reliable information is fundamental for management. Further, detectability of the signs used to build the index (e.g. number of observed animals, tracks, pellets) must be considered to avoid the risk of confounding changes

in detectability with changes in population abundance (Pollock et al. 2002). Proxies based on the impact of ungulates on vegetation have been also proposed (Morellet et al. 2007), but often the response of vegetation to changes in animal density can be too slow to initiate a quick management response (Tanentzap et al. 2012).

Indexes are often measured only at specific locations and more properly represent site occupancy, or utilization, rather than actual animal density (Putman et al. 2011). Thus, proxies can often be site-specific and then they can hardly be generalized despite the use of rigid and standardised protocols (Williams et al. 2002). Unless calibrated, those indexes/proxies do not provide an estimate of abundance or density that may be required for population management and control, in particular. To be an effective monitoring tool, a population index must be validated at the study area of interest (for instance as done by Corlatti et al. 2016 for the red deer *Cervus elaphus*, Evans and Rittenhouse 2018, for the black bear *Ursus americanus*, and Marchandeu et al. 2006 for the wild rabbit *Oryctolagus cuniculus*). This literature suggests that often computing a sensible population index may be as complex and expensive as estimating animal density.

Wäber et al. (2013) made a compelling case for the use of density (with other demographic information) instead of indexes by reconstructing the sink–source dynamics of roe deer *Capreolus capreolus* and Reeve's muntjac *Muntiacus reevesi* at landscape level. It was concluded that 'failure to quantify deer numbers and productivity has allowed high density populations to persist as regional sources contributing to range expansion, despite deliberative management programs and without recognition by managers who considered numbers and impacts to be stable'.

For wild boar, there are few attempts to apply pellets group counts (Plhal et al. 2014, Fattorini and Ferretti 2020) or to use hunting statistics (Imperio et al. 2010, Davis et al. 2016). Harvest data can be used if, and only if, harvest effort is known (St Clair et al. 2013). Unfortunately, in the case of sport hunting, effort is often unavailable and harvests are unreported. Thus, the suggestion by ENETWILD consortium et al (2018) to use harvest as a broad index of population abundance can be misleading. Spot-light counts are inefficient with wild boar (Focardi et al. 2001). Camera traps can be used to obtain a population index: Bengsen et al. (2011) monitored changes in relative abundance of feral pigs in a tropical rainforest of northern Australia, and Massei et al. (2017) evaluated the performance of a passive activity index from camera traps to monitor wild boar in five English woodlands. However, a meta-analysis by Broadley et al. (2019) strongly suggests that detection rates are likely to confound the variation in density values with that in movement patterns.

We believe that the situation is more promising in cases where methods are aimed at estimating the population density. The application of the random encounter models (REM) (Rowcliffe et al. 2008) can be effective, but it requires an accurate estimate of the daily range in each study area. New approaches based on camera traps have recently been proposed which do not require an estimate of the daily range and are easier than REM to implement by wildlife managers (Howe et al. 2017, Nakashima et al. 2017). Despite this, however, a huge amount of work remains necessary.

Further, non-invasive genetics, capture–mark–recapture (CMR) and nocturnal distance sampling (NDS) have been used with the wild boar. The first two methods use the same family of statistical estimators: a sample of animals is marked, by molecular or physical tags, and recaptured (or resighted) later. Non-invasive genetics was used by Ebert et al. (2012) to estimate wild boar density in the Palatinate Forest (Germany). Their study produced a precise estimate, with a CV estimate around 10%. However, costs can be elevated. CMR estimates, based on ear tagging, were successfully employed in the Preserve of Castelporziano (Italy) (Focardi et al. 2008, Franzetti et al. 2012), at Fort Benning (USA) by Hanson et al. (2008) and in Switzerland by Hebeisen et al. (2008) using radio tags. The method works fairly well but the cost of trapping and resight is high. Further, since wildlife managers usually wish to remove as much specimen as possible, the idea to trap and release wild boar is not always acceptable.

The use of NDS, via FLIR (forward-looking infrared) technologies was first proposed by Gill et al. (1997). NDS was used for Reeves' muntjac and roe deer (Smart et al. 2004, Hemami et al. 2007, Wäber and Dolman 2015) for white-tailed deer (Montague et al. 2017, Haus et al. 2019), fallow deer *Dama dama* (Focardi et al. 2013), wild boar and roe deer (Morelle et al. 2012). Franzetti et al. (2012) validated the use of NDS as a method to assess the wild boar population of the Castelporziano Preserve (Italy).

A simulation study aiming to compare REM and NDS was recently proposed by Chauvenet et al. (2017). They concluded that density estimators obtained by distance sampling showed a better accuracy than those obtained by REM. In the context of African Swine Fever prevention in Europe, on behalf of the European Food Safety Authority, ENETWILD consortium et al. (2018) reviewed the different methods to assess wild boar populations and concluded that 'distance sampling with thermography was recommended to estimate wild boar density on a local scale'. Accordingly, we aim to understand whether the promising results obtained by Franzetti et al. (2012) depended on especially favourable conditions or whether it is possible to obtain meaningful results even in areas characterized by lower visibility and animal density than those observed in Castelporziano. In conventional distance sampling, the variance of the density estimator is mainly determined by the encounter rate (Buckland et al. 2001) and thus the larger the sample size, the more precise the estimator. However, the encounter rate depends largely on environmental visibility. Detection probability is higher in open than in close environment and visibility is a metric that accounts for the environmental component of the detection probability, the others being animal size, behaviour and camouflage. Here, we propose a simple method to estimate visibility, which may help wildlife managers to design an effective NDS survey. Our goal is to define guidelines based on experiments and factual statements in order to design efficient NDS surveys of wild boar.

Material and methods

Study areas

The study took place in four different study areas in Italy (Fig. 1).

Castelporziano Preserve

The completely fenced 60-km² Preserve of Castelporziano consists mainly of natural oak woods, with both evergreen (*Quercus ilex* and *Q. suber*) and more open deciduous (*Q. cerris* and *Q. frainetto*) species and Mediterranean maquis. Mixed or pure forests of domestic pine *Pinus pinea* and pastures are also present. The climate is Mediterranean. We distinguish three main habitat types: deciduous wood, evergreen wood and Mediterranean maquis. The distributions of habitats and transects are reported in the Supplementary material Appendix 1 Fig. A1.

Casentino National Park

The study was carried out in the Foreste Casentinesi, Monte Falterona e Campigna National Park (360 km²), along the Tuscan-Romagna Apennine ridge. The landscape is characterized by sedimentary rocks, predominantly sandstone intercalated with marl, and by forests, which cover a large part of the area. We distinguish the submontane habitat (300–900 m a.s.l.), where woods are characterized by a high variety of deciduous species, from the montane habitat (900–1600 m a.s.l.). Here, the dominant species is the European beech *Fagus sylvatica*, usually associated with the sycamore maple *Acer pseudoplatanus* or with the European silver fir *Abies alba*, the most abundant conifer within the park. The distribution of habitats and transects are reported in Supplementary material Appendix 1 Fig. A2.

Monte Arcosu

The study was carried out in the Monte Arcosu – Piscinamanna WWF Reserve (SIC ITB041105) (303 km²) in the southern part of the Sardinian isle. The area can be considered montane although highest peaks are slightly over 1000 m a.s.l. The landscape presents an uneven morphology characterized by long and steep valleys that shaped metamorphic and granite rocks. Forests cover a large part of the area, and the vegetation is dominated by holm and cork oak (*Quercus ilex* and *Q. suber*). Mediterranean maquis is widespread and consists mainly of *Phyllirea* spp., various species of *Erica*, strawberry trees *Arbutus unedo* and common myrtle *Mirtus communis*. Pastures originated by fire and past livestock presence are dominated by several species of rockroses *Cistus* spp. The distributions of habitats and transects are reported in Supplementary material Appendix 1 Fig. A3.

Colli Euganei

The study was carried out in the Colli Euganei Regional Park (187 km²). The area is characterized by the highest hills of the Po River plain, which reach a height of 601 m a.s.l.; the landscape has been shaped by sedimentary and volcanic rocks. The main vegetation types are: pseudo-Mediterranean scrub, scattered throughout the area, growing on dry, south-facing volcanic terrain and consist of evergreen species as holm oak, strawberry trees, tree heath *Erica arborea*; the chestnut forest *Castanea sativa* grows preferentially on volcanic slopes facing north in cool, deep, siliceous soils; the oak (mainly downy oaks *Q. pubescens*) forest is mainly found on south-facing slopes on chalky or siliceous soils; dry grasslands are prevalent in the southern part of the area associated to black locust tree *Robinia pseudoacacia* woods and agricultural crops, especially vineyard and olive groves. The distributions of



Figure 1. Location of the the four study sites in Italy.

habitats and transects are reported in Supplementary material Appendix 1 Fig. A4.

Data collection

Thermal imaging equipment can detect the long-wave energy radiated by warm-bodied animals, allowing the detection of animals at night (Havens and Sharp 2015). As the wild boar is more active at night and prefers habitats with dense vegetation cover, thermal imaging can be an effective tool to significantly increase the probability of detecting groups during a DS survey (Supplementary material Appendix 1 Fig. A5A–D). We used a direct view thermal imager (ThermaCAM P640, FLIR Systems Italia, 640 × 380 pixel resolution) sensitive to infrared wavelengths (7.5–13 μm) achieving image magnification with a ×4 lens or a Thermacam PM545, FLIR, Wilsonville, OR, USA with a ×2 lens. Both cameras

were able to take pictures of detected animals and to associate them to a voice memo. Cameras were equipped with a coaxial digital laser range finder (Swarovski Optik, LRS 3-1250, Absam, Tyrol, Austria) and an electronic compass (Outback-ES, Riverton, WY, USA). A GPS receiver (Garmin™ eTrex Vista, Sciaffusa, Switzerland) recorded track lines and sighting locations. Data collection involved two observers who took turns at sampling. Each transect was walked by one observer alone who stopped every 5–10 m to scan the area around. At the end, the second observer picked the first one up and drove to the next transect, where they exchanged tasks. Observers used headlamps with a red filter when sampling in dense habitat to minimise human-induced flight response and for safe walking.

Staff had been trained at identifying and recognising the different species as well as at recording precise measures for about one month, first working with an expert trainer and

then alone. Observers were trained in large enclosures to collect many measurements and the quality of the collected data was evaluated retrospectively, using pictures and voice notes.

We recorded radial distance and compass bearing to the centre of the group detected, group size, animal activity (feeding, moving, lying or standing still) and fleeing response (yes/no) to the presence of the observer. Observations of flushing animals were ignored unless their original location was recorded or was evident from the heat radiating from the ground where they had been lying. Data were collected between 19:00 h and 06:00 h, paying attention to allocate the same effort in each part of the night (split in three sections: 18:00–22:00, 22:00–02:00, 02:00–06:00 h). To minimise the risk of non-independent detections, transects in close proximity were surveyed in different nights. Perpendicular distances were computed using ArcMap (ESRI, Redlands, CA, USA).

The sampling design is summarized in Table 1. For each study area we denoted two or three habitat classes as appropriate. Each survey was replicated more than once, depending on the available budget and on the length of the research contract. Transects were selected among available paths in order to have enough transects to estimate the variance of the encounter rate and evenly cover the study area, while also ensuring that each transect was long enough to reduce the number of transects with no observations.

Further, for each habitat listed in Table 1 we estimated the level of environmental visibility. We established a set of random points in the different habitats along the used transects. Visibility was estimated using a laser range finder positioned on a tripod at eye level. The range, i.e. the distance of reflection of the laser beam, was then estimated at the four cardinal directions, which are at random with respect to the direction of the transect line. In the 1.5% of sampled points, because of the peculiar topography, some of the measured distances exceeded the capability of the range finder (up to 1000 m) and were removed from the analyses.

Distance sampling analysis

Data were analysed using Distance 6.0 (Thomas et al. 2010). We used only the conventional distance sampling (CDS) analysis with per-habitat post-stratification. We did not use more complex modelling approach (cf. Buckland et al. 2004) preferring to have a quick and simple routine to assess the size or density of populations.

First, we performed some preliminary analyses to select the truncation distance, w . Because the truncation distance may influence the CV of density estimators through variations of the encounter rate (number of observed groups per km of transect), contrary to the approach usually employed in CDS, in this study it was necessary to set fixed truncation distance in order to have detection probability and effective strip width (ESW) estimates comparable among habitats and study areas. Upon inspection of the detection functions, $g(x)$, we selected three truncation distances, w , of 50, 75 and 100 m. Comparison of different w values can be informative on the precision attainable by our NDS surveys. Because of this, the density estimation procedures could not follow Buckland et al. (2001), but results should be

considered in a comparative framework, focusing on the attainable precision.

Second, we selected the best model between half-normal, uniform and hazard-rate with cosine adjustment, using AIC for model selection. The mean cluster size was estimated using a size-biased regression method in which $\log(\text{cluster size})$ is regressed on the estimated detection probability, $g(x)$.

In CDS, the variance of the density estimator is estimated taking into account three sources of variability, which allow highlighting the possible occurrence of critical survey issues: 1) the variance of the encounter rate, empirically estimated from differences among transect lines, 2) the variance of cluster size and 3) the variance of the detection probability (Buckland et al. 2001).

Simulations

In this study the actual density of the surveyed populations was not known, but we investigated bias and precision originating from our survey design through simulations, using a code derived from La Morgia et al. (2015) and La Morgia and Focardi (2016).

We validated our distance sampling surveys using four different population sizes for each study area. The chosen population sizes were selected to reasonably encompass the actual population size: two populations were smaller than the one estimated during the study (-30% and -25% in size), and one larger population ($+20\%$ in size). The survey effort was set as equal, double or triple to the one applied in each study area (Supplementary material Appendix 1 Fig. A1–A4). The rationale of this simulation exercise was that if our sampling designs did not introduced systematic biases with an increase in effort and population size, we should have observed an increase in the accuracy and the precision (%CV) of the estimates produced by the simulations.

Simulations were implemented in R (<www.r-project.org>). To simulate wild boar populations, we first generated cluster sizes from a look-up table of observed clusters specific to each study area. Second, we generated the spatial distribution of independent and identically distributed clusters via a homogeneous Poisson process (applying the function `rpoint` of the R package `spatstat`; Baddeley et al. 2015).

For each study area, each population size and each effort set, we performed 500 iterations. To simulate the detection process, we began by calculating the distances between the samplers (transects) and animal clusters. The simulation of the observer's activity was based on a random variate, u , uniformly distributed in (0,1), and the cluster was considered detected if $u < g(x)$.

Given the simulated data, we finally performed separate DS analyses for each set of simulated observations according to standard CDS methodology and using functions provided by the `mrds` package (Laake et al. 2018). Truncation distance was set to 100 m (i.e. equal to the truncation distance adopted in the analyses (cf. Fig. 3). We used a hazard rate function to model animal detections as a function of distance. The goodness-of-fit of each model was tested through a χ^2 test, excluding the results of runs if the test yielded $p \leq 0.1$. The standard deviations of both bias and CV were empirically calculated.

Table 1. NDS in the different study areas and habitats in Italy. For the four study areas and habitat types, we report the area surveyed, the number of transects, their mean length and the range (min-max), the survey effort and the starting and ending dates of surveys.

Study area	Habitat	Area (km ²)	Transect number	Mean transect length (km)	Transect length (km)		Total effort (km)	Start	End
					Minimal	Maximal			
Colli Euganei	Fields	15.6	54	1.46	0.41	3.01	78.7	04DEC07	19DEC07
	Wood	35.5	42	1.85	0.64	4.1	77.8		
Monte Arcosu	Fields	13.6	48	1.55	0.41	3.01	74.2	14OCT2008	28OCT2008
	Wood	31.0	39	1.79	0.64	3.65	69.8		
	Pastures	30.4	10	2.15	1.58	2.99	21.5	05OCT2012	30OCT2012
	Maquis	94.9	31	2.14	0.78	3.66	66.3		
	Woodland	177.0	62	2.09	0.67	3.3	130		
	Pastures	30.4	10	2.16	1.58	2.99	21.6	13MAY2013	31MAY2013
Castelporziano	Maquis	94.9	33	2.15	0.78	3.66	70.9		
	Woodland	177.0	62	2.12	0.67	3.63	131		
	Pastures	30.4	10	2.18	1.58	2.99	21.8	11OCT2014	29OCT2014
	Maquis	94.9	33	2.14	0.78	3.68	70.6		
	Woodland	177.0	62	2.10	0.67	3.29	130		
	Open	7.8	16	3.86	1.53	5.8	61.8	03OCT2013	10OCT2013
Casentino	Wood	10.6	3	2.93	1.16	5.72	8.78		
	Maquis	39.5	54	3.77	0.94	6.81	204		
	Open	7.8	13	3.83	1.42	5.8	49.8	18SEP2014	25SEP2014
	Wood	10.6	4	2.18	1.74	3.44	8.73		
	Maquis	39.5	40	3.59	0.94	6.81	143		
	Open	7.8	15	3.49	0.97	5.8	52.4	24SEP2015	07OCT2015
Casentino	Wood	10.6	4	2.25	1.74	3.44	9		
	Maquis	39.5	42	3.87	0.72	6.81	163		
	Mountain	116.4	19	2.44	0.91	7.5	46.29	11SEP2007	21SEP2007
	Submountain	140.2	15	2.22	1.22	3.42	33.25		
	Mountain	116.4	21	1.92	0.82	4.32	40.41	22APR2008	8MAY2008
	Submountain	140.2	22	1.94	0.59	3.95	42.58		




Figure 2. Box plot of the distribution of the visibility (abscissae, log scale) recorded in each habitat of four study areas in Italy (ordinates).

Results

Survey experiments

In Fig. 2, we report the visibility values recorded in the different study areas and habitats computed as the mean of the distances recorded by the laser range finder. There were important differences in environmental visibility among areas (Wilcoxon test, $\chi^2_3 = 473.4$, $p < 0.0001$): Casentino (submountain) exhibited the lowest (median score 144) and Castelporziano the highest visibility (median score 955). Even more striking were the differences among habitats within areas (Table 2). Open areas were characterized by visibility values >50 m while dense forests by values <10 m. Note that these distributions are extremely leptokurtic with maximal values observed in maquis (Castelporziano, kurtosis = 121.3, Monte Arcosu, kurtosis = 58.8).

Figure 3 shows the estimated detection functions for the considered habitats. In Casentino N.P. we observed a quite

irregular distribution of detections in both habitats, which suggest poor estimates of the detectability. In general, it was noted that habitats with irregular histograms are the ones less sampled (Table 1). On the other hand, Castelporziano is characterized by suitable detectability histograms, especially for the woods. Also, the woodland habitat in Colli Euganei R.P. presented a proper statistical behaviour, while the distribution of detections is more irregular in the fields. Monte Arcosu presented a well-fitting distribution for woodlands and poorly fitting distributions for maquis and pastures. Figure 3 also reports the mean detection probability, which ranges from 0.56 of the submontane habitat of the Foreste Casentinesi N.P. to a low 0.17 of the maquis in Monte Arcosu. In general, note that open areas were not characterised by high detection probabilities as one would expect because of terrain roughness and the presence of bushes and other obstacles that can limit actual detectability. Maquis was always characterised by a smaller detectability than that recorded in woods.

Table 2. Estimated visibility in the different habitats of four study areas in Italy. We report sample size, visibility, the degrees of freedom used to estimate between habitat difference, χ^2 and significance.

Study area	Habitat	N ¹	Visibility \pm SE (m)	df	χ^2	p
Casentino	Mountain	68	14.9 \pm 1.6	6	28.8	<0.0001
	Submountain	76	7.9 \pm 0.8			
Colli Euganei	Fields	204	82.8 \pm 19.1	14	112.7	<0.0001
	Wood	188	12.7 \pm 1.0			
Castelporziano	Open	236	114.6 \pm 11.0	30	280.7	<0.0001
	Wood	480	31.6 \pm 1.9			
	Maquis	240	23.2 \pm 3.9			
Monte Arcosu	Pastures	18	62.2 \pm 25.2	18	69.8	<0.0001
	Maquis	137	11.3 \pm 1.5			
	Woodland	180	13.4 \pm 2.1			

¹ The number of random positions is N/4.

Detection functions




Figure 3. Detection probability plot (red line) and histogram of detections (blue line) in different habitats of four study areas in Italy, $w = 100$ m. Always data were not significantly different from the estimated $g(x)$ as evaluated by the Kolmogorov–Smirnov test. The worst fitting ($D=0.084$, $p=0.30$) was observed for the woodland of Monte Arcosu where the distribution is more spiked near 0 than expected.

We report animal densities and confidence intervals in Fig. 4. In Castelporziano, densities were the highest in all habitats while in Casentino N.P. densities were the lowest. In the other areas there were intermediate density values, rang-

ing between 10 and 20 boar km^{-2} . There was no significant difference among habitats within the same study area (Table 3). For comparison, we also reported the same analysis run for the encounter rates, which showed, instead, significant




Figure 4. Density estimates ($\pm 95\%$ confidence limits) in different habitats of four study areas in Italy, $w = 100$ m.

differences among habitats within the same study area except for the Casentino N.P.

The effect of the encounter rate on the precision of the density estimates is reported in Fig. 5. As expected, the larger the encounter rate, the higher the precision obtained ($F_{4,22} = 11.96$, $p < 0.0001$), although significant differences among study areas resulted from the analyses. This relationship was stronger in Colli Euganei, Monte Arcosu and Casentino N.P. than in Castelporziano ($F_{3,22} = 5.69$, $p = 0.005$). To note that the coefficients of the linear regression between CV estimates and the log of the encounter rates were significant for Colli Euganei ($t = 3.2$, $p = 0.005$) and Monte Arcosu ($t = 6.1$, $p < 0.0001$) but not for Casentino ($t = 0.12$, $p = 0.90$) and Castelporziano ($t = 1.05$, $p = 0.30$).

The estimated CV values of the density estimators adopted in the different habitats surveyed are reported in Fig. 6. Half of the surveys were characterised by a CV estimate $< 30\%$. As expected from previous analyses, the smallest CV estimates were obtained in Castelporziano and Monte Arcosu, while the estimator for Casentino was characterised by $30 < \%CV < 50$. Surveys carried out in Colli Euganei produced heterogeneous results. The mode of $\%CV$ estimates is around 25%. The three plots of Fig. 7 evidence clearly the effect of truncation. Usually, the smaller the w the better the $\%CV$ estimate. For instance, in Monte Arcosu, shortening

w from 100 to 75 m reduced the CV estimate from 95% to 55%. This well illustrates the concept that a wise choice of the truncation distance can significantly improve estimation, especially when the sample size is small.

The three factors which might influence the precision of the density estimators (the cluster size, the detection probability and the encounter rate) had a variable weight in the different study areas. This is clearly shown in Fig. 7. The estimator of cluster size played a minor role in all study areas. Indeed, we always found a small estimate of mean cluster size (Monte Arcosu, 1.32 ± 0.04 ; Castelporziano 1.59 ± 0.09 ; Casentino N.P. 1.79 ± 0.08 ; Colli Euganei 3.70 ± 0.54). The variance of the encounter rate estimator dominated the overall variance of wild boar densities in Castelporziano and in Casentino N.P. while in the other two areas the contribution given by the estimator of the encounter rate and of the detection probability was comparable.

Simulations

The simulation results are reported in Fig. 8. As expected, the CVs improved with the effort, with values $< 20\%$ when the sampling effort was tripled. Interestingly, the variability around the average CV is small, indicating a low possibility that a given survey is characterised by a large CV estimate.

Table 3. Between-habitat comparison of encounter rate and density. We report the value of the t statistics, the Satterthwaite degrees of freedom to control for inequalities of variances and test probability. Statistics were performed using pre-computed results from DISTANCE 5.0.

Study area	Habitat comparison	Encounter rate			Density		
		t value	DoF	p	t value	DoF	p
Monte Arcosu	pastures – maquis	-2.91	54.78	0.005	-1.42	17.31	0.174
	pastures – woodland	-6.17	135.91	<.001	-1.88	10.32	0.088
	maquis – woodland	-2.23	132.71	0.028	-0.34	75.62	0.736
Castelporziano	open – evergreen wood	2.96	27.01	0.006	0.29	27.00	0.774
	open – maquis	3.36	27.29	0.002	1.14	28.51	0.262
	maquis – evergreen wood	-1.48	11.48	0.166	-0.99	11.85	0.344
Casentino	mountain – submountain	0.12	28.81	0.905	0.94	24.48	0.356
Colli Euganei	fields – wood	-2.30	66.30	0.025	-0.10	37.33	0.918




Figure 5. %CV of density as a function of the log(encounter rate). On the left Casentino (red), Colli Euganei (green) and Monte Arcosu (grey), On the right the plot for Castelporziano (black). $w = 50, 75$ and 100 m.

The bias values also decreased as a function of effort, although there was a tendency toward negative bias. In Castelporziano and in the Casentinesi N.P., the bias was negligible, while in Colli Euganei R.P. it was larger (respectively 15–20%) and in Monte Arcosu it was beyond 20%, even when the sampling effort was tripled. Inside each study area, the CVs and the bias converged with the highest sampling effort, while with lower efforts their values were much more scattered, which was expected.

Discussion

We proved that it is possible to obtain estimates of wild boar populations characterized by good to acceptable accuracy and precision values using NDS under quite different environmental conditions. We made our experiments in Mediterranean and temperate habitats, in dense woods and in open areas with different structure and composition of the plant community and different levels of potential disturbing




Figure 6. The distribution of the %CV in the different habitats surveyed in Casentino N.P. (red), Colli Euganei (green), Monte Arcosu (grey) and Castelporziano (black) for three different w values: 50, 75 and 100 m.




Figure 7. Box-and whiskers plots of the decomposition of the variance of density estimate in Casentino N.P. (red), Colli Euganei (green), Monte Arcosu (grey) and Castelporziano (black).

factors, which may affect wild boar use of space, responses and thus detectability (i.e. presence of wolves in Casentino N.P., free ranging dogs and poachers in Monte Arcosu and Colli Euganei R.P., culling in Castelporziano). Confidence intervals were a measure of the reliability of our sample estimates of the parameter's values (Elzinga et al. 2001). We showed that more accurate precisions can be obtained in a range of environments with different visibility levels.

The close fit of the detection function to the distribution of detection distances indicates that NDS may produce valuable results. When the population density was relatively large, the estimation of the density of wild boar showed a good level of precision estimates, at least in half of the cases investigated by this study. Large variance estimates were observed where the encounter rate was low, for instance in the maquis of Castelporziano or in the open fields of Colli Euganei R.P.

However, managing wild boar requires effective monitoring tools when densities are especially high or low, such as where the species has recently been introduced/released or where the African swine fever is present. In the former case, indeed, it can be essential for the priority planning process required to manage the impact of economic or ecological relevance. In the latter case, it can be useful to follow the evolution of the population in terms of size and space occupied to prevent adverse impacts, or follow the recovery of the population after the disease outbreak. However, when the encounter rate is low, reliable estimators can be obtained by increasing the sampling effort (i.e. replicating the survey; Buckland et al. 2001). The relationship between the estimated CVs and the effort (in km walked) reads $CV = 76.2 \pm 16.8 - 0.535 \pm 0.23 \text{ effort}$ ($t = -2.32$, $p = 0.067$, Supplementary material Appendix 1 Fig. A6). This equation may be used to compute the sampling effort required to obtain




Figure 8. We report the true CVs (left column) and the actual bias (right column) as a function of effort (abscissae) for the four simulated populations (with respect to the estimated population in grey, -30% light blue, -25% yellow, +20% red, cf. Table 4) for the four study areas in Italy Castelporziano (A, B), Colli Euganei (C, D), Casentino (E, F) and Monte Arcosu (G, H). Bias is computed as (estimated population - actual population) / actual population.

the target %CV, assuming, of course, that it remains valid in different areas. But it can represent a starting point anyway.

We tried to perform surveys in seasonal and nictemeral conditions when the wild boar was least aggregated. This resulted in the small effect of the variation of cluster size on the precision and contributed to the reduction of the overall CV.

This paper is focused on the precision of the density estimates more than on accuracy, although we showed the results produced in all the study areas, also with respect to bias, through the survey simulations. We focused on the CV estimates because we believe that the management of the wild boar population cannot be prescriptive, i.e. we do not believe it is useful to formulate an exact shooting or cull-

Table 4. The size of the synthetic populations used to evaluate the bias of the DS survey design adopted in the four study areas. Population sizes have been computed as the 70, 85 and 130% of the population estimated by the experimental DS survey.

Study area	70%	85%	Estimated	130%
Casentino	539	654	770	1001
Castelporziano	1621	1969	2316	3011
Colli Euganei	1965	2386	2807	3649
Monte Arcosu	3174	3854	4535	5895

ing plan. On the contrary, we highlight the importance of an effective monitoring of the wild boar populations, whose estimators should be as precise as possible. We showed that there are several cases where the CV estimates were between 10% and 20% and that the mode of CV estimates was 25%. These can be considered good results since the CV can be improved substantially by increasing the sampling effort, by enhancing observational skills or refining sampling schemes and statistical analyses.

The reduction of the CV is indeed multifactorial, but it also represents a tradeoff between costs and objectives. Low CV estimates can be obtained through different working approaches in relation to: 1) sampling design, 2) field methods and 3) statistical analysis. In the case of NDS, the application of an appropriate survey design is strongly recommended and carrying out a pilot study is essential in order to evaluate if the CV estimates are adequate for the management objectives. To ensure an even-coverage sampling as much as possible (Barabesi and Fattorini 2013), footpaths can be selected inside each cell of a systematic grid superimposed randomly on the study area (La Morgia et al. 2015). Otherwise, the use of roads and/or footpaths without a robust sampling design or a validation study, which assesses the distribution of target species around the selected transects, is prone to yield biased estimates. Many scholars use road sampling which can lead to violating several assumptions (Franzetti et al. 2012), but is, of course, less expensive and faster than walking transects. In case of car surveys, we recommend the use of density surface modelling instead of CDS as done by La Morgia et al. (2015) on the red deer, or Valente et al. (2016) on the roe deer.

Since NDS can only be performed using existing tracks and roads, we could expect the survey design may introduce a systematic bias in density estimates. While at Castelporziano, Franzetti et al. (2012) experimentally showed, using radio-tracked wild boar, that transect placement was unbiased, for the other study areas we could not collect such information. Our simulations showed, however, that with a large sampling effort it is possible to obtain nearly unbiased density estimators with negligible bias values. In our simulations, we assumed that animal distribution was Poisson. However, a more sophisticated approach should be to develop density surface models as a basis for simulations, as done by Buckland et al (2015, §2.5) and La Morgia et al. (2015).

Training of observers is also required as well as the use of appropriate devices for animal detection (light thermal cameras with good resolution) and unbiased recording of angles and distances. Several commonly made mistakes, such the heaping (or biased round-off) of distances can be easily avoided by good training and high-quality devices. Observers must be

able to recognize animals and to correctly detect group size using thermal imagery and they have to be capable of moving safely in the forest by night without disturbing animals. Conventional distance sampling surveys are relatively simple to analyse statistically following the guidelines of Buckland et al. (2001) and Thomas et al. (2010). Moreover, DISTANCE software allows managers to deal easily with more sophisticated analyses (post-stratification and covariates).

An important issue is linked to evasive animal movements, which can be detected by observing a reduced frequency near the transect line which ultimately results in negatively biased density estimates. In our study, we found evidence of evasive movement of wild boar only in one case (submountain habitat of the Casentinesi N.P.) out of 10.

We noted in our comments to Fig. 6 that there is a range of habitat conditions resulting in acceptably low CV estimates, while there is another group characterised by higher CV estimates. What is to be done in such cases? Again, a pilot survey is of great help. If the estimated density is very low (as we observed, for instance, in the pastures at Monte Arcosu) the large CV estimate depends probably on a too small encounter rate which can be solved by increasing the sampling effort. A possible solution relies on multiple-covariate distance sampling, using the habitat as a covariate or to apply post-stratification (Marques and Buckland 2004). Using habitat covariates at the level of observation is also one possible account for the small-scale habitat variations indicating whether the habitat is too much fragmented at the level of transect. If technically possible, one can adopt an adaptive survey design or to use post stratification (Pollard and Buckland 2004) or a two-stage sampling proposed by La Morgia et al. (2015).

We may conclude that since population assessment of wild boar are fundamental elements for population control, especially now that ASF is expanding in Europe and Asia, and precise and accurate estimators of the size and the density of populations would be highly useful, NDS can be important tools for wildlife managers.

It is also worth considering that NDS can be used to calibrate population indexes in different landscapes. Calibration of indexes is a necessary step for their use in wildlife management (Williams et al. 2002). An interesting discussion about calibration methods has been presented by Falcy et al. (2016) using data on the chinook salmon *Oncorhynchus tshawytscha* contrasted to independent CMR estimates. Since it is usually difficult to perform CMR studies on wild boar populations, it can be useful to adopt NDS as a benchmark. A cost-effective index to be used for wild boar may be derived from data collected by camera traps (Borchers et al. 2002). The precision of the index might be evaluated using the method proposed by Engeman (2005) who considered trap-to-trap variability, daily variability and random observational variability associated with each trap, each day. In the calibration exercise, the variance of the index must be considered when computing the strength of the relationship index-density since a large variance reduces the statistical power of the calibration.

Our study confirms that NDS is highly efficient when surveying populations characterized by high (Castelporziano) or medium (Monte Arcosu and Colli Euganei) densities and that precision can remain good when applying an appropriate sampling effort where the densities are lower. This also con-

firms that NDS can be effective where wild boar is expected to determine large impacts and an efficient management programme is required. Thus, we can conclude that the indication of ENETWILD consortium et al. (2018) in the use of NDS for estimating wild boar population size within the context of prevention of diffusion of the ASF is appropriate.

Acknowledgements – We thank the administrations of the study areas for permission to work and their invaluable help. We acknowledge the contribution of rangers, collaborators and enthusiastic students as well as a number of colleagues of ISPRA for support. We thanks Dr. Jacek Borysow for reviewing the English.

References

- Andraud, M. et al. 2019. Threat to the French swine industry of African Swine Fever: surveillance, spread and control perspective. – *Front. Vet. Sci.* 6: 248.
- Anderson, D. R. 2001. The need to get the basics right in wildlife field studies. – *Wildl. Soc. Bull.* 29: 1294–1297.
- Anderson, A. et al. 2016. Economic estimates of feral swine damage and control in 11 US States. – *Crop Protection* 89: 89–94.
- Baddeley, A. et al. 2015. Spatial point patterns: methodology and applications with R. – Chapman and Hall/CRC Press, <www.crcpress.com/Spatial-Point-Patterns-Methodology-and-Applications-with-R/Baddeley-Rubak-Turner/9781482210200/>.
- Ballari, S. A. et al. 2015. Invasive wild boar in Argentina: using protected areas as a research platform to determine distribution, impacts and management. – *Biol. Invas.* 17: 1595–1602.
- Barabesi, L. and Fattorini, L. 2013. Random versus stratified location of transects or points in distance sampling: theoretical results and practical considerations. – *Environ. Ecol. Stat.* 20: 215–236.
- Bengsen, A. J. et al. 2011. Using a general index approach to analyse camera-trap abundance indices. – *J. Wildl. Manage.* 7: 1222–1227.
- Bengsen, A. J. et al. 2014. Impacts and management of wild pigs *Sus scrofa* in Australia. – *Mammal Rev.* 44: 135–147.
- Bobek, B. et al. 2018. Resolving conflict between farmers and wild boar in Europe and northern Asia. – In: Melletti, M. and Meijaard, E. (eds), Ecology, conservation and management of wild pigs and peccaries. Cambridge Univ. Press, pp 354–365.
- Borchers, D. L. et al. 2002. Estimating animal abundance: closed populations, Vol. 13. – Springer Science & Business Media.
- Broadley, K. et al. 2019. Density-dependent space use affects interpretation of camera trap detection rates. – *Ecol. Evol.* 9: 14031–14041.
- Buckland, S. T. et al. 2001. Introduction to distance sampling. Estimating abundance of biological populations. – Oxford Univ. Press.
- Buckland, S. T. et al. 2004. Advanced distance sampling, Vol. 2. – Oxford Univ. Press.
- Buckland, S. T. et al. 2015. Distance sampling: methods and applications. – Springer, p. 277.
- Carey, M. P. et al. 2012. Native invaders—challenges for science, management, policy and society. – *Front. Ecol. Environ.* 10: 373–381.
- Corlatti, L. et al. 2016. Spring spotlight counts provide reliable indices to track changes in population size of mountain-dwelling red deer *Cervus elaphus*. – *Wildl. Biol.* 22: 268–276.
- Chauvenet A. L. M. et al. 2017. Quantifying the bias in density estimated from distance sampling and camera trapping of unmarked individuals. – *Ecol. Model.* 350: 79–86.
- Cuevas, M. F. et al. 2020. Vegetation recovery after 11 years of wild boar exclusion in the Monte Desert, Argentina. – *Biol. Invas.* 22: 1607–1621.
- Davis, A. J. et al. 2016. Inferring invasive species abundance using removal data from management actions. – *Ecol. Appl.* 26: 2339–2346.
- De Sabato, L. et al. 2018. Molecular survey of HEV infection in wild boar population in Italy. – *Transboundary Emerging Dis.* 65: 1749–1756.
- Dimzas, D. et al. 2019. Human trichinellosis caused by *Trichinella britovi* in Greece, and literature review. – *J. Helminthol.* 13: 1–4.
- Dröge, E. et al. 2020. Assessing the performance of index calibration survey methods to monitor populations of wide-ranging low-density carnivores. – *Ecol. Evol.* 10: 3276–3292.
- Ebert, C. et al. 2012. Estimating wild boar *Sus scrofa* population size using faecal DNA and capture–recapture modelling. – *Wildl. Biol.* 18: 142–152.
- Elzinga, C. L. et al. 2001. Monitoring plant and animal populations: a handbook for field biologists. – Wiley.
- ENETWILD consortium et al. 2018. Guidance on estimation of wild boar population abundance and density: methods, challenges, possibilities. – EFSA Supporting Publications, 15: 1449E.
- Engeman, R. M. 2003. More on the need to get the basics right: population indices. – *Wildl. Soc. Bull.* 31: 286–287.
- Engeman, R. M. 2005. Indexing principles and a widely applicable paradigm for indexing animal populations. – *Wildl. Res.* 32: 203–210.
- Engeman, R. M. et al. 2013. Monitoring wild pig populations: a review of methods. – *Environ. Sci. Pollut. Res.* 20: 8077–8091.
- Evans, M. J. and Rittenhouse, T. A. 2018. Evaluating spatially explicit density estimates of unmarked wildlife detected by remote cameras. – *J. Appl. Ecol.* 55: 2565–2574.
- Falcy, M. R. et al. 2016. Proxies in practice: calibration and validation of multiple indices of animal abundance. – *J. Fish Wildl. Manage.* 7: 117–128.
- Fattorini, N. and Ferretti, F. 2020. Estimating wild boar density and rooting activity in a Mediterranean protected area. – *Mammal. Biol.* 100: 241–251.
- Focardi, S. et al. 2001. Comparative evaluation of thermal infrared imaging and spotlighting for wildlife surveys. – *Wildl. Soc. Bull.* 29: 133–139.
- Focardi, S. et al. 2008. Wild boar survival in a variable environment: unexpected life history variations in an unusual ungulate. – *J. Mammal.* 89: 1113–1123.
- Focardi, S. et al. 2013. Nocturnal distance sampling of a Mediterranean population of fallow deer is consistent with population projections. – *Wildl. Res.* 40: 437–446.
- Franzetti, B. et al. 2012. Nocturnal line transect sampling of wild boar (*Sus scrofa*) in a Mediterranean forest: long-term comparison with capture–mark–resight population estimates. – *Eur. J. Wildl. Res.* 58: 385–402.
- Genov, P. V. et al. 2018. Ecological impact of wild boar in natural ecosystems. – In: Melletti, M. and Meijaard, E. (eds), Ecology, conservation and management of wild pigs and peccaries. Cambridge Univ. Press.
- Gerrodette, T. 1987. A power analysis for detecting trends. – *Ecology* 68: 1364–1372.
- Gill, R. M. A. et al. 1997. The use of portable thermal imaging for estimating deer population density in forest habitats. – *J. Appl. Ecol.* 34: 1273–1286.
- Graitson, E. et al. 2018. Catastrophic impact of wild boars: insufficient hunting pressure pushes snakes to the brink. – *Anim. Conserv.* 22: 165–176.
- Hebeisen, C. et al. 2008. Estimating wild boar abundance and density using capture–resights in Canton of Geneva, Switzerland. – *Eur. J. Wildl. Res.* 54: 391–401.
- Hemami, R. et al. 2007. Estimating abundance of introduced Chinese muntjac *Muntiacus reevesi* and native roe deer *Capreolus capreolus* using portable thermal imaging equipment. – *Mammal Rev.* 37: 246–254.

- Hanson, L. B. et al. 2008. Change-in-ratio density estimator for feral pigs is less biased than closed mark-recapture estimates. – *Wildl. Res.* 35: 695–699.
- Haus, J. M. et al. 2019. A spatially and temporally concurrent comparison of popular abundance estimators for white-tailed deer. – *Northeast. Nat.* 26: 305–324.
- Havens, K. J. and Sharp, E. J. 2015. Thermal imaging techniques to survey and monitor animals in the wild: a methodology. – Academic Press.
- Howe, E. J. et al. 2017. Distance sampling with camera traps. – *Methods Ecol. Evol.* 8: 1558–1565.
- Iannuzzo, D. et al. 2010. A method to estimate roe deer *Capreolus capreolus* density at various spatial scales in a fragmented landscape. – *Wildl. Biol.* 16: 283–291.
- Imperio, S. et al. 2010. Investigating population dynamics in ungulates: do hunting statistics make up a good index of population abundance? – *Wildl. Biol.* 16: 205–214.
- Keuling, O. et al. 2016. Regulating wild boar populations is ‘somebody else’s problem!’ – Human dimension in wild boar management. – *Sci. Total Environ.* 554: 311–319.
- Koda, R. and Fujita, N. 2011. Is deer herbivory directly proportional to deer population density? Comparison of deer feeding frequencies among six forests with different deer density. – *For. Ecol. Manage.* 262: 432–439.
- Kojola, I. et al. 2014. Tracks in snow and population size estimation: the wolf *Canis lupus* in Finland. – *Wildl. Biol.* 20: 279–284.
- Laake, J. et al. 2018. mrds: mark-recapture distance sampling. – R package ver. 2.2.0. <<https://CRAN.R-project.org/package=mrds>>.
- La Morgia, V. and Focardi, S. 2016. Performance of distance sampling estimators: a simulation study for designs based on footpaths. – *J. Stat. Comput. Simul.* 86: 2519–2530.
- La Morgia, V. et al. 2015. Cost-effective nocturnal distance sampling for landscape monitoring of ungulate populations. – *Eur. J. Wildl. Res.* 61: 285–298.
- Lyons, J. E. et al. 2008. Monitoring in the context of structured decision-making and adaptive management. – *J. Wildl. Manage.* 72: 1683–1692.
- Marchandeu, S. et al. 2006. Abundance indices: reliability testing is crucial—a field case of wild rabbit *Oryctolagus cuniculus*. – *Wildl. Biol.* 12: 19–27.
- Marques, F. F. C. and S. T. Buckland 2004. Covariate models for the detection function. – In: S. T. Buckland et al. (eds), *Advanced distance sampling*. Oxford Univ. Press, pp. 31–47.
- Massei, G. et al. 2017. Camera traps and activity signs to estimate wild boar density and derive abundance indices. – *Pest Manage. Sci.* 74: 853–860.
- Montague, D. M. et al. 2017. Using distance-sampling to estimate density of white-tailed deer in forested, mountainous landscapes in Virginia. – *Northeast. Nat.* 24: 505–519.
- Morelle, K. et al. 2012. Game species monitoring using road-based distance sampling in association with thermal imagers: a covariate analysis. – *Anim. Biodivers. Conserv.* 35: 253–265.
- Morellet, N. et al. 2007. Indicators of ecological change: new tools for managing populations of large herbivores. – *J. Appl. Ecol.* 44: 634–643.
- Nakashima, Y. et al. 2017. Estimating animal density without individual recognition using information derivable exclusively from camera traps. – *J. Appl. Ecol.* 55: 735–744.
- Nichols, J. D. and Williams, B. K. 2006. Monitoring for conservation. – *Trends Ecol. Evol.* 21: 668–673.
- Nimmo, D. G. et al. 2015. Dingoes can help conserve wildlife and our methods can tell. – *J. Appl. Ecol.* 52: 281–285.
- Nuno, A. et al. 2015. Detecting abundance trends under uncertainty: the influence of budget, observation error and environmental change. – *Anim. Conserv.* 18: 331–340.
- Plhal, R. et al. 2014. Faecal pellet group counting as a promising method of wild boar population density estimation. – *Acta Theriol.* 59: 561–569.
- Pollock K. H. et al. 2002. Large scale wildlife monitoring studies: statistical methods for design and analysis. – *Environmetrics* 13: 105–119.
- Pollard, J. H. and Buckland, S. T. 2004. Adaptive distance sampling surveys. – In: Buckland, S. T. et al. (eds), *Advanced distance sampling: estimating abundance of biological populations*. Oxford Univ. Press, pp. 229–259.
- Putman, R. et al. 2011. Identifying threshold densities for wild deer in the UK above which negative impacts may occur. – *Mammal Rev.* 41: 175–196.
- Reimoser, F. and Putman, R. 2011. Impacts of wild ungulates on vegetation: costs and benefits. – In: Putman, R. et al. (eds), *Ungulate management in Europe: problems and practices*. Cambridge Univ. Press, pp. 144–191.
- Reynolds, J. H. et al. 2011. Planning for success: identifying effective and efficient survey designs for monitoring. – *Biol. Conserv.* 144: 1278–1284.
- Rowcliffe, J. M. et al. 2008. Estimating animal density using camera traps without the need for individual recognition. – *J. Appl. Ecol.* 45: 1228–1236.
- Ruiz-Fons, F. 2017. A review of the current status of relevant zoonotic pathogens in wild swine (*Sus scrofa*) populations: changes modulating the risk of transmission to humans. – *Transboundary Emerging Dis.* 64: 68–88.
- Skalski, J. R. et al. 2005. *Wildlife demography*. – Elsevier.
- Smart, J. C. et al. 2004. Monitoring woodland deer populations in the UK: an imprecise science. – *Mammal Rev.* 34: 99–114.
- St Clair, K. et al. 2013. A comparison of models using removal effort to estimate animal abundance. – *J. Appl. Stat.* 40: 527–545.
- Stephens, P. A. et al. 2015. Management by proxy? The use of indices in applied ecology. – *J. Appl. Ecol.* 52: 1–6.
- Tanentzap, A. J. et al. 2012. Slow responses of ecosystems to reductions in deer (*Cervidae*) populations and strategies for achieving recovery. – *For. Ecol. Manage.* 264: 159–166.
- Thomas, L. et al. 2010. Distance software: design and analysis of distance sampling surveys for estimating population size. – *J. Appl. Ecol.* 47: 5–14.
- Valente, A. M. et al. 2016. A new insight for monitoring ungulates: density surface modelling of roe deer in a Mediterranean habitat. – *Eur. J. Wildl. Res.* 62: 577–587.
- Van Phan Le, D. G. J. et al. 2019. Outbreak of African swine fever, Vietnam, 2019. – *Emerging Infectious Dis.* 25: 1433.
- Wäber, K. and Dolman, P. M. 2015. Deer abundance estimation at landscape-scales in heterogeneous forests. – *Basic Appl. Ecol.* 16: 610–620.
- Wäber, K. et al. 2013. Achieving landscape-scale deer management for biodiversity conservation: the need to consider sources and sinks. – *J. Wildl. Manage.* 77: 726–736.
- Williams, B. K. et al. 2002. *Analysis and management of animal populations. modeling, estimation and decision making*. – Academic Press.

Supplementary material (available online as Appendix wlb-00694 at <www.wildlifebiology.org/appendix/wlb-00694>). Appendix 1.