

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

The Last Glacial Maximum in the southern side of the Alps: chronology, paleoclimate and sedimentary response

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1884174> since 2022-12-28T17:57:48Z

Publisher:

Association française pour l'étude du Quaternaire (AFEQ)

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

The Last Glacial Maximum in the southern side of the Alps: chronology, paleoclimate and sedimentary response

Giovanni Monegato¹, Alessandro Fontana², Maria Gabriella Forno³, Franco Gianotti³, Paolo Mozzi², Susan Ivy-Ochs⁴, Roberta Pini⁵, Cesare Ravazzi⁵, Sandro Rossato²

¹CNR-IGG, Torino, Italy

²Department of Geosciences, University of Padova, Italy

³Earth Science Department, University of Torino, Italy

⁴ETH, Zurich, Switzerland

⁵CNR - IDPA, Milano, Italy

In the last decade the study of several end-moraine systems, and related outwash plains, improved the chronology of the Last Glacial Maximum in the Italian Alps. During the LGM, most Alpine valleys were occupied by ice streams, while several other catchments hosted only small cirque glaciers confined to the highest peaks. Paleobotanical data suggest that rainfall was the main trigger for the culmination of the last glaciation. Indeed an increase in precipitation from the MIS3 to the onset of the LGM can be inferred by palaeovegetation records, in agreement to palaeoclimatic atmospheric circulation models, indicating significant rainfall in the eastern Southern Alps. The humidity persisted all along the glacial maximum supporting open conifer forests, even close to the piedmont glacier fronts, as well as wide-spread wetlands in the distal reach of the eastern glaciofluvial systems.

The signals of an incoming spread of glaciers in a mountain range are normally recorded in the frontal alluvial plain by an increase in sediment deposition at the valley outlets. The response of larger catchments to the advance of ice streams (Tagliamento, Piave, Adige/Brenta, Adda, Dora Baltea and Dora Riparia systems) can be detected in the increase of sediment accumulation in the plains since 29.5-28 ka cal BP, and then to the growth of the alluvial megafans in the piedmont plain. The LGM outwash deposits have an average thickness of 15-35 m in the Venetian-Friulian Plain and of 10-20 m in the central-western Po Plain. In several areas the LGM sediments buried a fairly well developed soil. Several datings in the end moraine systems indicate that Alpine glaciers reached the piedmont plain at about 26.5 ka cal BP. Subsequently, in response to changes in precipitation rates during the LGM, the frontal position of the glaciers oscillated at least two times within the end moraine systems before the onset of decay at 22 ka cal BP and the collapse of ice streams at 19 ka cal BP.