

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Structural and molecular brain sexual differences: A tool to understand sex differences in health and disease

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1616898> since 2017-05-14T11:33:24Z

Published version:

DOI:10.1016/j.neubiorev.2016.04.017

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

This Accepted Author Manuscript (AAM) is copyrighted and published by Elsevier. It is posted here by agreement between Elsevier and the University of Turin. Changes resulting from the publishing process - such as editing, corrections, structural formatting, and other quality control mechanisms - may not be reflected in this version of the text. The definitive version of the text was subsequently published in NEUROSCIENCE AND BIOBEHAVIORAL REVIEWS, 67, 2016, 10.1016/j.neubiorev.2016.04.017.

You may download, copy and otherwise use the AAM for non-commercial purposes provided that your license is limited by the following restrictions:

- (1) You may use this AAM for non-commercial purposes only under the terms of the CC-BY-NC-ND license.
- (2) The integrity of the work and identification of the author, copyright owner, and publisher must be preserved in any copy.
- (3) You must attribute this AAM in the following format: Creative Commons BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/deed.en>), 10.1016/j.neubiorev.2016.04.017

The publisher's version is available at:

<http://linkinghub.elsevier.com/retrieve/pii/S0149763416300665>

When citing, please refer to the published version.

Link to this full text:

<http://hdl.handle.net/>

Structural and molecular brain sexual differences: a tool to understand sex differences in health and disease.

GianCarlo Panzica^a and Roberto C. Melcangi^b

^aDipartimento di Neuroscienze "Rita Levi Montalcini", Università degli Studi di Torino and Neuroscience Institute Cavalieri Ottolenghi (NICO), Orbassano (Italy)

^bDipartimento di Scienze Farmacologiche e Biomolecolari, Università degli Studi di Milano, Milan, Italy.

Corresponding author: Prof. GianCarlo Panzica,
Dipartimento di Neuroscienze, via Cherasco 15,
10126 Torino (Italy).

E-mail: giancarlo.panzica@unito.it

PANZICA G.C. and R.C. MELCANGI. Structural and molecular brain sexual differences: a tool to understand sex differences in health and disease. *NEUROSCI BIOBEHAV REV*, XXX-XXX, 2016

Sex differences are present both in the genotype and in the phenotype of all vertebrates, and they have been evidenced also within the central and peripheral nervous system. Earlier studies on brain sex differences suggested a relatively simple view based on (1) the presence of sexually dimorphic circuits in the hypothalamus (or in regions related to reproductive behaviors), (2) the action of gonadal hormones to masculinize the brain, and (3) the gonadal steroids' action to modulate gene transcription through nuclear receptors. These assumptions are today contradicted by the findings accumulated in the last 20 years. We know now that mechanisms determining sexual dimorphisms may vary according to location and species, and may involve several factors, as genes, epigenetic factors, gonadal hormones and neurosteroids. Sex differences were also revealed by epidemiological studies in several neural pathologies. This suggests that the approach to understand the genesis of these pathologies, should involve specific attention to interactions among genes, gonadal and brain-born steroid hormones, epigenetic and environmental factors.

Key words: Medial preoptic region; Vasopressin; Vasotocin; Kisspeptin; Dopamine; Estradiol: Neuroactive steroids; Adult neurogenesis; Epigenetics

1. Introduction

Sex differences in the phenotype of living animals are very diffuse both in invertebrates and vertebrates. The reproductive organs are a typical example of such differences, they are differentiated for their morphology, for the production of gametes (different in male and female), and for their endocrine functions. Other, so-called, secondary sex characteristics are: the body size, ornamentation, fat tissue distribution, some parts of the skeleton (i.e. pelvis, skull), hair, and many other structures. Also several behaviors are sexually dimorphic and this implies the presence of sex differences in brain neuroanatomy and/or neurophysiology.

In birds and mammals, sex differences have been demonstrated at chromosomal level, with a couple of chromosomes (sex chromosomes or heterochromosomes) that are different among males and females. Particular genes on these chromosomes [the gene *DMRT1* on Z chromosome of birds (Smith et al., 2009) and the gene *SRY* on Y chromosome of mammals (Sinclair et al., 1990)] are responsible of the male sex determination. The primary goal of these genes is to induce the development of male gonads. The central and simpler hypothesis (that is now under criticism in view of recent discoveries, see Lenz et al., 2012) is that animals with *SRY* or *DMRT1* gene will develop testes whose hormones will induce the differentiation of male phenotypes, whereas in the absence of *SRY* (or with a diminution of *DMRT1*), the genetic program will induce an ovary whose hormones will determine the female phenotypes. Therefore, according to this dogma the phenotypic differences between male and females are based on more or less precocious exposure to the "right" hormone and this induce the expression of that characteristics for the rest of the life (organizational effects of steroid hormones).

Berthold (a German physiologist) was probably the first to observe, in 1849, a sexual difference in animal behavior and to link it to the differences in the gonads. For this reason he is considered the father of behavioral endocrinology (Beach, 1981; Jorgensin, 1971). In his experiment, Berthold noted that, in addition to phenotypical characteristics as the presence of a comb and of wattles, the roosters (male chickens) were more aggressive than females (hens) and they copulate with hens, whereas these last do not copulate with other hens. Castrated rooster did not develop comb and wattles, in addition, they were not aggressive and did not copulate. But, when castrated roosters received a transplanted testis, this became functional (producing sperm) and the morphological and behavioral phenotypes of intact roosters were restored. This experiment was performed a long time before some concepts as

hormones, neural basis of behavior, sex determination, and sex differences were clarified. However it clearly demonstrate that the products of the testes can stimulate the differentiation of male external morphology, of male typical behaviors and of specific neural circuits controlling these behaviors.

Only after more than 100 years, Phoenix and coworkers (1959) published a seminal paper demonstrating that the alteration of prenatal gonadal hormones environment may lead to adult alteration of sexual behavior, thus establishing the difference among "organizational" and "activational" effects of gonadal hormones. After this first study, several experiments have been performed to study every known sex difference in behavior. At the same time, many studies were dedicated to investigate the presence of sexually dimorphic circuits, nuclei or other structures potentially related to sexually dimorphic behaviors (Abel and Rissman, 2012; Arnold and Gorski, 1984; Arnold et al., 2003; Panzica et al., 1995).

The first significant evidence of an anatomical difference at the hypothalamic level (number of synapses on dendritic spines in the dorsal medial preoptic area, MPOA) was published by Raisman and Field (1971), that lately demonstrated that this difference is organizational (Raisman and Field, 1973). Due to the technical limitations of the studies at ultrastructural level these differences are difficult to find and it is not possible to apply this approach for the description of large brain structures. After these studies, Gorski and coworkers found a more easily detectable neural difference: the presence of a sexually dimorphic nucleus (SDN) within the MPOA whose volume and cell number is higher in male than in female rat (Gorski et al., 1978; Gorski et al., 1980). Sexually dimorphic structures, organized during embryonic or postnatal development were subsequently described in different vertebrate species (for reviews see Breedlove, 1992; Panzica et al., 1995; Panzica et al., 1996; Simerly, 2002), including humans (Swaab and Fliers, 1985).

In oscine birds, Nottebohm and Arnold (1976) found sexually dimorphic nuclei (larger in males than in females) in the telencephalic regions controlling the emission of song. This difference is triggered by testosterone (T) in the adult canaries (Nottebohm, 1980). These studies demonstrated for the first time a deep connection among seasonal changes in reproductive behavior and changes in the morphology of related circuits in intact birds (Nottebohm, 1981).

The picture resulting from these early studies was relatively simple (at least for rodents and canaries): (1) sexually dimorphic circuits are located in the hypothalamus or in other regions

controlling behaviors related to reproduction; (2) brain masculinization depends by the presence of gonadal hormones during specific (critical) periods, whereas their absence drives the brain to the female sex; (3) the steroids, solely produced by gonads, act through their nuclear receptor and directly modulate gene transcription.

All these three assumptions are today at least partly contradicted by the new findings accumulated in the last 20 years (Arnold, 2009b).

2. Sexually dimorphic circuits or nuclei in the central nervous system.

From the first ultrastructural and histological studies many other techniques were employed to detect sex differences in the central nervous system. The chemical neuroanatomical techniques (including immunohistochemistry, autoradiography and *in situ* hybridization) have detailed the presence of neurotransmitters, neuropeptides, enzymes involved in their synthesis, or receptors. In this way the number of end points to be considered to study the sex dimorphism increases a lot. In some cases, the neurochemical markers detailed structures already evidenced in histological studies. This is the case of the quail medial preoptic nucleus (POM) that was at first described with Nissl's staining (Viglietti-Panzica et al., 1986), and later its location, volume, and steroid-induced plasticity were confirmed by using immunohistochemistry for the enzyme aromatase (ARO) (Aste et al., 1994). A different example is the SDN-MPOA of the rat. It was the first sexually dimorphic nucleus that was observed in the mammalian hypothalamus with histological methods (Gorski et al., 1978), however, for a long time, researchers failed to find a murine counterpart of the SDN in Nissl-stained sections. At the beginning of this century a subdivision of the SDN was found to be positive for calbindin-D28k (whose function in this context remains unclear but probably it is related to cell survival and apoptosis) and this subdivision was sexually dimorphic and responsive to gonadal hormones treatments as the SDN (Sickel and McCarthy, 2000). Later, other researchers found that a similar cluster of calbindin-positive cells exists also in the preoptic region of the mouse. This marker delineates a sexually dimorphic region that cannot be evidenced with Nissl's staining (Edelmann et al., 2007) and, as the rat SDN, is dependent by gonadal hormones to sexually differentiate (Budefeld et al., 2008).

Immunohistochemical and *in situ* hybridization studies have been largely used to investigate sex differences of neuropeptidergic circuits. Among several systems that have been described,

two were particularly detailed in different vertebrate species. The first one is the rat parvocellular sexually dimorphic arginine-vasopressin (AVP) system, located outside the hypothalamus in the bed nucleus of the stria terminalis (BST) and in the medial amygdala. Its projections reach several extra-hypothalamic locations, in particular the lateral septum, the ventral pallidum, the hippocampus and various brain stem nuclei (De Vries et al., 1985; Gu et al., 2003). Cell bodies and projections are strongly sexually dimorphic, having males more cells and higher density of positive fibers than females (De Vries et al., 1985). Similar sexually dimorphic cell groups were observed in different mammalian as well as non-mammalian species (in this case the peptide is the arginine-vasotocin, AVT). The mechanisms determining the sex differences may vary (see below), but the endpoint (the dimorphism) is similar in the different models (for a review see De Vries and Panzica, 2006). The parvocellular AVP/AVT system shows gonadal hormones receptors and is activated by them, however its sexual dimorphism seems not to be related to the presence of estradiol (E_2) during the critical period (Pierman et al., 2008; Plumari et al., 2002), but probably to a direct genomic effect (De Vries et al., 2002) or to the presence of a functional androgen receptor (Allieri et al., 2013).

Another strongly sexually dimorphic peptidergic system that has been more recently investigated in several mammalian and non-mammalian species is the kisspeptin system. This system is characterized by two groups of neurons: the first one is located in the anterior preoptic area (AVPV) and the second one is in the arcuate nucleus (ARC). These neurons project mainly to the gonadotropin releasing hormone (GnRH) neurons that control the secretion of gonadotropins from the hypophysis (Dungan et al., 2006). The system is therefore extremely important for the control of reproduction. It is sexually dimorphic with a higher number of cells and fibers in the female than in male and is strongly gonadal hormones dependent in the adult (Kauffman et al., 2007). The kisspeptin system has been characterized in various vertebrate species discovering at least three types of molecules (Kiss1-2-3) and four types of receptors (KissR1-4) (for a review see Pasquier et al., 2014). In mammals Kiss1 has been linked to the development of puberty, and several investigations have elucidated the implicated mechanisms (Clarkson et al., 2010).

Sexually dimorphic pathways (characterized by their content in neuropeptides, neurotransmitters or enzymes) have been described in several regions of the CNS outside the hypothalamus and the limbic system. For example the enzymes tyrosine-hydroxylase (TH) and dopamine- β -hydroxylase (DBH) show a sex dimorphism in their expression or regulation within the rat locus coeruleus (the main nor-adrenergic nucleus in the brain, Luque et al.,

1992; Thanky et al., 2002). The expression of the enzyme for the GABA synthesis (GAD65) is sexually dimorphic in discrete regions of the hypothalamus, but also in extrahypothalamic regions as the amygdala or the hippocampus (Perrot-Sinal et al., 2001). Dorsal root ganglia and the dorsal horn of the spinal cord, show a sexually dimorphic expression of estrogen receptors (ERs) and this is probably one of the reason why female rat is more sensitive to deep pain than male (Papka and Mowa, 2003).

An interesting example of diffuse sex dimorphism is the expression of calbindin in several populations of neurons (outside the small group located in the preoptic region): i.e. the frontal cortex, thalamus, hippocampus, amygdala and cerebellum. In the frontal cortex and cerebellum of juvenile mice, calbindin expression is sexually dimorphic with females having about twice positive cells than males (that is the opposite of what observed in the hypothalamus). In addition, the mechanisms determining these sexual differences are also different for the two regions: in the frontal cortex the dimorphism depends by both ER α and sex chromosomes, but in the cerebellum sex chromosomes are the main (sole?) factor (Abel et al., 2011).

The dopaminergic system is largely diffused in the brain. The majority of dopaminergic cells have been observed in the substantia nigra (projecting to the forebrain: striatum, cortex and limbic system) or in the ventral tegmental area (projecting to the frontal cortex, the amygdala and the nucleus accumbens). The dopaminergic neurons are usually identified by the presence of the tyrosine hydroxylase (TH), the rate-limiting enzyme involved in dopamine synthesis. Sexually dimorphic expression of this enzyme has been observed both in the substantia nigra (Ma et al., 2007) and in the ventral tegmental area (Brown et al., 2015). Fibers containing TH have been described in many brain regions, where they have frequently a sexually dimorphic distribution (Kritzer and Creutz, 2008).

In the hypothalamus there are two populations of TH-positive elements that show a sexually dimorphic distribution. The first is located in the AVPV and the second in the ARC nucleus (Arbogast and Voog, 1990; Balan et al., 2000). Recent studies have shown that at least part of these two populations of dopaminergic neurons express also kisspeptin (Kauffman et al., 2007).

Many other neurotransmitters, neuropeptides and/or their receptors are expressed in a sexually dimorphic way throughout the entire central nervous system. Sex differences have been detected also at molecular level in mechanisms that are potentially widespread in the

brain. For example electron microscopic studies discovered sexual differences in spine synapse density in females but not males in the caudal part of nucleus accumbens (Wissman et al., 2012). In the hippocampus, brain-born E₂ rapidly suppresses inhibitory synaptic transmission blocking GABA release from CB1 receptor-containing inhibitory presynaptic endings in females but not in males (Huang and Woolley, 2012). A recent study demonstrated that this suppression is due to inositol triphosphate (IP₃) generation, activation of the IP₃ receptor, and postsynaptic endocannabinoid release (Tabatadze et al., 2015). The complexes of ER α , glutamate receptor and IP₃R are present in both sexes, but are regulated by E₂ only in females (Tabatadze et al., 2015).

The development of imaging techniques opened a new field of investigations for the human brain. In fact, it is now possible to select more homogeneous groups of volunteers, and it is easier to study also sex differences in the human brain. A recent study discovered unique sex differences in brain connectivity by using diffusion tensor imaging in more than 900 youths (aged 8-22 y, more than 400 males and 500 females). Males had greater intra-hemispheric connectivity, whereas in females more inter-hemispheric connectivity has been observed (Ingahalikar et al., 2013). However, another study performed on more than 1,400 subjects revealed extensive overlap between the females and males characteristics for gray matter, white matter, and connections, revealing that most brains are "mosaics" of features, some more common in females, some more common in males, and some common in both sexes (Joel et al., 2015), but these conclusions and the applied statistical tools were critically questioned (Del Giudice et al., 2016). Also optogenetic techniques have been recently applied to investigate sex differences in the inferior temporal cortex of macaque monkeys performing a facial gender-discrimination task (Afraz et al., 2015).

These and many other data are therefore contrasting the hypothesis that sexually dimorphic circuits are located only in the hypothalamus or in other regions controlling behaviors related to reproduction: their distribution is much larger and involves several parts of the central nervous system. In addition, the endpoints that can be considered are numerous, starting from large anatomical differences (as the volume of the rat SDN or of the quail POM) and ending to very subtle neurochemical differences (as the modulation of GABA release).

3. Mechanisms determining the development of sexual differences within the central nervous system

There is a strong connection among gonadal hormones and the development of sex differences of the body including the brain, however the mechanisms may vary in different species and other factors have important, even preeminent, roles. Early in development, the increased volume of the SDN in rat males depends by the availability of E₂ locally produced from circulating T by the action of the enzyme ARO. In the female, blood circulating E₂ is blocked by the alpha-fetoprotein (Bakker et al., 2006), therefore the increase of SDN volume happens only in males. Estrogens may prevent cell death in the male SDN by inhibiting capsases (Choi et al., 2008) or/and by regulating anti-apoptotic proteins (Gill and Christakos, 1995). This mechanism is valid for the rat SDN (and may be for other mammals), but in other vertebrates, as galliform birds, E₂ is acting in an opposite way inducing demasculinization of the volume of the medial preoptic nucleus (Aste et al., 1991) and of the sexually dimorphic parvocellular AVT system in the BST (Panzica et al., 1998). The regulation of cell proliferation during the embryonic period (in particular around the E12-14, that is the end of the critical period in galliforms) seems to be also in this case one of the key mechanisms, however no clear hypothesis has been presented (Bardet et al., 2012).

Another example is the sexual differentiation of the AVPV (located in the most medial part of the MPOA). The AVPV is larger in volume and cell number in females as compared with males (Bleier et al., 1982; Simerly et al., 1985). This sex difference results by an androgen-dependent mechanism, in fact cell number is reduced in females pre- or post-natally treated with T propionate (Arai et al., 1994; Murakami and Arai, 1989). Androgens and androgen receptors are also important for the sexual differentiation of other brain circuits, as demonstrated by the alteration of nitrergic population in the MPOA, VMH and BST nuclei (Martini et al., 2008) or of the AVP system in the BST and medial amygdala (Allieri et al., 2013) in androgen receptor defective (Tfm) male rats.

Estrogens and androgens are produced by the gonads, that differentiate through a genetic mechanism linked to the presence of X and Y chromosomes, in particular, in mammals, by the presence of the *Sry* gene. It is therefore possible that some brain differences are directly depending by a chromosomal effect and not by an epigenetic effect of gonadal hormones. The TH system represents a very peculiar system to investigate this mechanism. In fact, in cell cultures extracted from the mesencephalic region at the embryonic day 14 (before the surge of T in the rat male embryo), the uptake of dopamine (a marker of the functional maturation of these neurons) as well as the number of cells were sexually differentiated (Beyer et al., 1992; Beyer et al., 1991; Engele et al., 1989; Reisert et al., 1990). This suggests that the genome

should be responsible of sex differences in rodent mesencephalic dopaminergic system (Reisert and Pilgrim, 1991). This hypothesis was confirmed by experiments demonstrating the hormonal independence of these sex differences in long-term steroid-exposed mice (Sibug et al., 1996).

An interesting genetic model to test the effects of the genome is the so-called "four core genotypes (FCG)" mouse model (Arnold, 2009a; Arnold and Chen, 2009) that comprises mice in which sex chromosomes (XX vs. XY) are unrelated to the animal's gonadal sex. The males are either XY or XX^{+Sry} , and the females are XX or XY^{-Sry} , all males have testis and females have ovaries. Thus, the FCG model may provide information regarding the non-gonadal origins of sex differences. For the mesencephalic dopaminergic system, cell cultures from the mesencephalon of E14 FCG mice confirmed the effect of genetic sex on these sex differences. Cell cultures from XY males or XY^{-Sry} females had more TH- cells than XX or XX^{+Sry} cultures (Carruth et al., 2002). Therefore, the genes located in the Y chromosome (and not the *Sry* which is determining the development of gonads) are important for the sexual differentiation of this circuit.

In contrast to these results, the sexual differentiation of the TH-containing neurons of the AVPV is strictly dependent by the perinatal hormonal environment. In fact, the disruption of the $ER\alpha$ feminizes the system, inducing a 3 times increase in the number of TH neurons of AVPV in $ER\alpha$ KO males in comparison to the wild type, while no changes have been detected within the AVPV of female $ER\alpha$ KO mice. The androgen receptor seems not to be involved, in fact *Tfm* mice contain the same number of TH-immunoreactive neurons in the AVPV (Simerly et al., 1997). The action of E_2 in males is probably linked to an increase of the apoptosis, as demonstrated *in vitro* and *in vivo* by using drugs inhibiting capsase that induced an increase of TH-immunoreactive neurons (Waters and Simerly, 2009).

Outside the hypothalamus, gonadal hormones have an important role as regulative factors of adult neurogenesis (Galea et al., 2013). In the hippocampus they differentially affect cell proliferation and survival in male and female rat: repeated administration of E_2 , decreased the survival of new neurons, increased cell proliferation, and decreased cell death in the female, whereas they have no effect in male (**Barker and Galea, 2008**). Cellular proliferation decreases in ovariectomized females, and it is restored by acute treatment with E_2 (Tanapat et al., 2005). In male rats, androgens (T and dihydrotestosterone), but not E_2 , stimulate cell survival (Spritzer and Galea, 2007). In the subventricular zone (SVZ), the modulatory effects of

gonadal hormones are highly variable according to sex, species, treatment and strain. In a recent study (Farinetti et al., 2015) we demonstrated that E₂ and T, but not dihydrotestosterone, increase proliferation in the SVZ of adult castrated male rats, whereas ovariectomized female rats treated with E₂ or T do not show any significant effect.

Recently, major attention has been paid to the epigenetic mechanisms occurring at the DNA level. In a recent study (Nugent et al., 2015) the authors found that gonadal steroids act in the MPOA to reduce activity of DNA methyltransferase enzymes, decreasing DNA methylation and inducing the activation of masculinizing genes. Pharmacological inhibition of these enzymes induced the masculinization of neuronal markers and the appearance of male sexual behavior in female rats. Also conditional knockout mice for one of the isoforms of DNA methyltransferase enzymes (Dnmt3a) showed masculinized sexual behavior in female. These data show that "brain feminization is maintained by the active suppression of masculinization via DNA methylation" (Nugent et al., 2015).

In conclusion, the organization of sexually differentiated circuits is based not only on circulating gonadal hormones secreted by gonads, but also by locally synthesized E₂, by androgens, by genes located in the sex chromosomes, not limited to the Sry, and by epigenetic factors.

4. Sex difference in neuroactive steroids.

Data so far discussed seems to suggest that only sex steroids produced by peripheral glands (or their direct metabolites, as the case of E₂ derived from T) are able to influence sexual differences in the nervous structures. However, the last three decades of investigations have clearly indicated that steroids affecting nervous function are not only those coming from the periphery (i.e., steroid hormones) but also those directly synthesized in the nervous system (i.e., neurosteroids). These two classes of steroid molecules are now integrated in the term *neuroactive steroids* (Giatti et al., 2015; Melcangi et al., 2008; Melcangi and Panzica, 2006; Rupprecht and Holsboer, 1999). Indeed, nervous system expresses molecules involved in the transport of cholesterol into the mitochondria, such as steroidogenic acute regulatory protein (StAR) and translocator protein, as well as several steroidogenic enzymes (Melcangi et al., 2008). Thus, nervous system possesses the capability to synthesize and to metabolize neuroactive steroids. Importantly, StAR, the cytochrome P450 side chain cleavage (i.e., the first

steroidogenic enzyme) and ARO show sex differences in the nervous system (Lavaque et al., 2006; Lavranos et al., 2006). Interestingly, it has been recently demonstrated that, in the FCG mouse model, sex chromosome complement determines sex differences in ARO expression in brain structures. Thus, the bed nucleus of the stria terminalis and the anterior amygdala of 16 days old XY and XY^{Sry} mouse embryos show higher aromatase expression than the brain of XX and XX^{Sry} embryos (Cisternas et al., 2015). In physiological conditions the levels of several neuroactive steroids show sex differences within the nervous system (Melcangi et al., 2016),

The so-called classical action of steroid hormones is performed through nuclear receptors that, after binding the appropriate ligand, show a conformational change allowing receptors to bind directly to specific steroid response elements in the promoter regions of specific genes (O'Malley and Means, 1974). However, neuroactive steroids, including E₂, (Balthazart and Ball, 2006) may induce rapid changes that occur within seconds to milliseconds, this is called the non-classical action and it is due to G protein-coupled and ligand-gated ion channel membrane receptors (for a review see King, 2008). Many receptors are involved, as glycine receptors, metabotropic sigma type 1 receptors, N-methyl-D-aspartic acid (NMDA) receptors and γ -aminobutyric acid type A (GABAA) receptors. Recently, a G-protein coupled estrogen-sensitive receptor (termed G-protein estrogen receptor 1, GPER1) has been described within the brain (Hazell et al., 2009). Several studies elucidated the distribution and subcellular localization of this receptor in the brain, as well as some of the roles that this receptor may play in the brain particularly in the control of synaptogenesis and spinogenesis and the impact that estrogens may have over brain diseases (for reviews see Hara et al., 2015; Sellers et al., 2015; Srivastava and Evans, 2013). In agreement with these concepts, several observations have been obtained indicating that neuroactive steroids play important roles in sexual function and behavior (see for a review King, 2008).

In conclusion, the three main assumptions reported in the introductions, are contradicted, at least partly, by the recent findings. In particular: (1) sexually dimorphic circuits are located all over in the brain and the spinal cord. (2) The organization of sexually differentiated circuits depends by several mechanisms including circulating gonadal hormones, locally synthesized E₂, androgen and estrogen receptors, genes located in the sex chromosomes, and epigenetic factors. (3) The action of steroids in the central nervous system includes both classical and non-classical mechanisms. Finally, brain-born steroids (i.e., neurosteroids) are involved in many physiological regulations.

5. Sex differences in neurodegenerative and psychiatric disorders

Sex differences are not only present in physiological status but also in neural pathologies. Indeed, as reported in this special issue and by an extensive recent literature (Young and Pfaff, 2014) sex differences have been observed in psychiatric disorders (e.g., epilepsy, schizophrenia, autism, anxiety and depression) (Altemus et al., 2014; Cozzoli et al., 2014; Hill, 2016; Kelley et al., 2011; Lovick, 2014; McHenry et al., 2014; Mendrek and Mancini-Marie, 2016; Reddy, 2014; Scharfman and MacLusky, 2014; van Luijtelaar et al., 2014), stress-related psychiatric disorders (Bangasser and Valentino, 2014), and addictive disorders (Fattore et al., 2014; Zhou et al., 2016). Neurodevelopmental disorders (Romano et al., 2016), autoimmune disease (Ngo et al., 2014), neurodegenerative disorders (e.g., Alzheimer's disease, Parkinson's disease, Huntington's disease, multiple sclerosis) (Gillies et al., 2014; Grimm et al., 2016; Kipp et al., 2016; Li and Singh, 2014; Litim et al., 2016; Mosera and Pike, 2016; Ramien et al., 2016), trauma (e.g., traumatic brain injury, spinal cord, stroke, post-traumatic stress disorder) (Gibson and Attwood, 2016; Inslicht et al., 2014), and diabetes (e.g., diabetic encephalopathy and peripheral neuropathy) (Lipscombe et al., 2014; Pesaresi et al., 2010; Zhao et al., 2014) also show sex differences in their incidence, symptoms and neurodegenerative outcome.

Interestingly, neuroactive steroid levels present in brain areas are also influenced by these pathologies in a sexually dimorphic way (Melcangi et al., 2016). Moreover, examples of sex specific effects of neuroactive steroids have been demonstrated, at least in experimental models (Mannix et al., 2014; Murray et al., 2003; Pesaresi et al., 2011a; Pesaresi et al., 2011b; Peterson et al., 2015; Zup et al., 2014). That is particularly interesting because may provide a possible background for a gender medicine based on these molecules to be applied in case of nervous pathologies (for a review see Porcu et al., 2016).

6. Conclusions.

Sex differences in the central and peripheral nervous system are widely diffused in vertebrates, in mammals and in humans. The mechanisms that cause the establishment of these dimorphisms may vary according to location and species, and involves several actors including: genes located in sexual chromosomes, epigenetic factors, gonadal hormones as well as neurosteroids. An increasing number of studies link human nervous pathologies with the

action of neuroactive steroids and with an effect of gender. Therefore, future research both in physiological and pathological conditions should consider more the interactions among genes, gonadal and brain-born hormones, epigenetic and environmental factors (as for example the exposure to endocrine disruptors, Frye et al., 2012), in order to have a more complex approach to the genesis of neural pathologies.

6. Acknowledgements

We acknowledge support from the Fondazione Cariplo to R.C.M. (grant number 2012-0547) and University of Torino, Fondazione San Paolo and Fondazione Cavalieri Ottolenghi to G.C.P.

7. References

- Abel, J.L., Rissman, E.F., 2012. Location, location, location: Genetic regulation of neural sex differences. *Rev Endocr Metab Disord* 13, 151-161.
- Abel, J.M., Witt, D.M., Rissman, E.F., 2011. Sex differences in the cerebellum and frontal cortex: roles of estrogen receptor alpha and sex chromosome genes. *Neuroendocrinology* 93, 230-240.
- Afraz, A., Boyden, E.S., DiCarlo, J.J., 2015. Optogenetic and pharmacological suppression of spatial clusters of face neurons reveal their causal role in face gender discrimination. *Proc.Natl.Acad.Sci.U.S.A.* 112, 6730-6735.
- Allieri, F., Spigolon, G., Melcangi, R.C., Collado, P., Guillamon, A., Gotti, S., Panzica, G.C., 2013. Androgen receptor deficiency alters the arginine-vasopressin sexually dimorphic system in tfm rats. *Neuroscience* 253, 67-77.
- Altemus, M., Sarvaiya, N., Neill Epperson, C., 2014. Sex differences in anxiety and depression clinical perspectives. *Front. Neuroendocrinol.* 35, 320-330.
- Arai, Y., Murakami, S., Nishizuka, M., 1994. Androgen enhances neuronal degeneration in the developing preoptic area: Apoptosis in the anteroventral periventricular nucleus (AVPvN-POA). *Horm. Behav.* 28, 313-319.
- Arbogast, L.A., Voog, J.L., 1990. Sex-related alterations in hypothalamic tyrosine hydroxylase after neonatal monosodium glutamate treatment. *Neuroendocrinology* 52, 460-467.
- Arnold, A.P., 2009a. Mouse models for evaluating sex chromosome effects that cause sex differences in non-gonadal tissues. *J. Neuroendocrinol.* 21, 377-386.
- Arnold, A.P., 2009b. The organizational-activational hypothesis as the foundation for a unified theory of sexual differentiation of all mammalian tissues. *Horm. Behav.* 55, 570-578.
- Arnold, A.P., Chen, X., 2009. What does the "four core genotypes" mouse model tell us about sex differences in the brain and other tissues? *Front. Neuroendocrinol.* 30, 1-9.
- Arnold, A.P., Gorski, R.A., 1984. Gonadal steroid induction of structural sex differences in the central nervous system. *Annu. Rev. Neurosci.* 7, 413-442.
- Arnold, A.P., Rissman, E.F., De Vries, G.J., 2003. Two perspectives on the origin of sex differences in the brain. *Ann. N. Y. Acad. Sci.* 1007, 176-188.
- Aste, N., Panzica, G.C., Aimar, P., Viglietti-Panzica, C., Harada, N., Foidart, A., Balthazart, J., 1994. Morphometric studies demonstrate that aromatase-immunoreactive cells are the main

target of androgens and estrogens in the quail medial preoptic nucleus. *Exp. Brain Research* 101, 241-252.

- Aste, N., Panzica, G.C., Viglietti-Panzica, C., Balthazart, J., 1991. Effects of in ovo estradiol benzoate treatments on sexual behavior and size of neurons in the sexually dimorphic medial preoptic nucleus of Japanese quail. *Brain Res. Bull.* 27, 713-720.
- Bakker, J., De Mees, C., Douhard, Q., Balthazart, J., Gabant, P., Szpirer, J., Szpirer, C., 2006. Alpha-fetoprotein protects the developing female mouse brain from masculinization and defeminization by estrogens. *Nat. Neurosci.* 9, 220-226.
- Balan, I.S., Ugrumov, M.V., Calas, A., Maily, P., Krieger, M., Thibault, J., 2000. Tyrosine hydroxylase-expressing and/or aromatic L-amino acid decarboxylase-expressing neurons in the mediobasal hypothalamus of perinatal rats: Differentiation and sexual dimorphism. *J. Comp. Neurol.* 425, 167-176.
- Balthazart, J., Ball, G.F., 2006. Is brain estradiol a hormone or a neurotransmitter? *Trends Neurosci.* 29, 241-249.
- Bangasser, D.a., Valentino, R.J., 2014. Sex differences in stress-related psychiatric disorders: neurobiological perspectives. *Front. Neuroendocrinol.* 35, 303-319.
- Bardet, S.M., Mouriec, K., Balthazart, J., 2012. Birth of neural progenitors during the embryonic period of sexual differentiation in the Japanese quail brain. *J. Comp. Neurol.* 520, 4226-4253.
- Barker, J., Galea, L., 2008. Repeated estradiol administration alters different aspects of neurogenesis and cell death in the hippocampus of female, but not male, rats. *Neuroscience* 152, 888-902.
- Beach, F.A., 1981. Historical origins of modern research on hormones and behavior. *Horm. Behav.* 15, 325-376.
- Beyer, C., Eusterschulte, B., Pilgrim, C., Reisert, I., 1992. Sex steroids do not alter sex differences in tyrosine hydroxylase activity of dopaminergic neurons invitro. *Cell Tissue Res.* 270, 547-552.
- Beyer, C., Pilgrim, C., Reisert, I., 1991. Dopamine content and metabolism in mesencephalic and diencephalic cell cultures: sex differences and effects of sex steroids. *J. Neurosci.* 11, 1325-1333.
- Bleier, R., Byne, W., Siggelkow, I., 1982. Cytoarchitectonic sexual dimorphisms of the medial preoptic and anterior hypothalamic area in guinea pig, rat, hamster, and mouse. *J. Comp. Neurol.* 212, 118-130.

- Breedlove, S.M., 1992. Sexual dimorphism in the vertebrate nervous system. *J. Neurosci.* 12, 4133-4142.
- Brown, E.C., Steadman, C.J., Lee, T.M., Padmanabhan, V., Lehman, M.N., Coolen, L.M., 2015. Sex differences and effects of prenatal exposure to excess testosterone on ventral tegmental area dopamine neurons in adult sheep. *Eur. J. Neurosci.* 41, 1157-1166.
- Budefeld, T., Grgurevic, N., Tobet, S.A., Majdic, G., 2008. Sex differences in brain developing in the presence or absence of gonads. *Dev. Neurobiol.* 68, 981-995.
- Carruth, L.L., Reisert, I., Arnold, A.P., 2002. Sex chromosome genes directly affect brain sexual differentiation. *Nat. Neurosci.* 5, 933-934.
- Choi, W.S., Lee E Fau - Lim, J., Lim J Fau - Oh, Y.J., Oh, Y.J., 2008. Calbindin-D28K prevents drug-induced dopaminergic neuronal death by inhibiting caspase and calpain activity. *Biochem.Biophys.Res.Comm.* 371, 127-131.
- Cisternas, C.D., Tome, K., Caeiro, X.E., Dadam, F.M., Garcia-Segura, L.M., Cambiasso, M.J., 2015. Sex chromosome complement determines sex differences in aromatase expression and regulation in the stria terminalis and anterior amygdala of the developing mouse brain. *Mol. Cell. Endocrinol.* 414, 99-110.
- Clarkson, J., Han, S.K., Liu, X., Lee, K., Herbison, A.E., 2010. Neurobiological mechanisms underlying kisspeptin activation of gonadotropin-releasing hormone (GnRH) neurons at puberty. *Mol. Cell. Endocrinol.* 324, 45-50.
- Cozzoli, D.K., Tanchuck-Nipper, M.A., Kaufman, M.N., Horowitz, C.B., Finn, D.A., 2014. Environmental stressors influence limited-access ethanol consumption by C57BL/6J mice in a sex-dependent manner. *Alcohol* 48, 741-754.
- De Vries, G.J., Buijs, R.M., Van Leeuwen, F.W., Caffé, A.R., Swaab, D.F., 1985. The vasopressinergic innervation of the brain in normal and castrated rats. *J. Comp. Neurol.* 233, 236-254.
- De Vries, G.J., Panzica, G.C., 2006. Sexual differentiation of central vasopressin and vasotocin systems in vertebrates: different mechanisms, similar endpoints. *Neuroscience* 138, 947-955.
- De Vries, G.J., Rissman, E.F., Simerly, R.B., Yang, L.Y., Scordalakes, E.M., Auger, C.J., Swain, A., Lovell-Badge, R., Burgoyne, P.S., Arnold, A.P., 2002. A model system for study of sex chromosome effects on sexually dimorphic neural and behavioral traits. *J. Neurosci.* 22, 9005-9014.
- Del Giudice, M., Lippa, R.A., Puts, D.A., Bailey, D.H., Bailey, J.M., Schmitt, D.P., 2016. Joel et al.'s method systematically fails to detect large, consistent sex differences. *Proc Natl Acad Sci U S A.*

- Dungan, H.M., Clifton, D.K., Steiner, R.A., 2006. Minireview: kisspeptin neurons as central processors in the regulation of gonadotropin-releasing hormone secretion. *Endocrinology* 147, 1154-1158.
- Edelmann, M., Wolfe, C., Scordalakes, E.M., Rissman, E.F., Tobet, S., 2007. Neuronal nitric oxide synthase and calbindin delineate sex differences in the developing hypothalamus and preoptic area. *Dev. Neurobiol.* 67, 1371-1381.
- Engel, J., Pilgrim, C., Reisert, I., 1989. Sexual differentiation of mesencephalic neurons in vitro: effects of sex and gonadal hormones. *Int. J. Dev. Neurosci.* 7, 603-611.
- Farinetti, A., Tomasi, S., Foglio, B., Ferraris, A., Ponti, G., Gotti, S., Peretto, P., Panzica, G.C., 2015. Testosterone and estradiol differentially affect cell proliferation in the subventricular zone of young adult gonadectomized male and female rats. *Neuroscience* 286, 162-170.
- Fattore, L., Melis, M., Fadda, P., Fratta, W., 2014. Sex differences in addictive disorders. *Front. Neuroendocrinol.* 35, 272-284.
- Frye, C., Bo, E., Calamandrei, G., Calza, L., Dessi-Fulgheri, F., Fernandez, M., Fusani, L., Kah, O., Kajta, M., Le Page, Y., Patisaul, H.B., Venerosi, A., Wojtowicz, A.K., Panzica, G.C., 2012. Endocrine Disruptors: A Review of Some Sources, Effects, and Mechanisms of Actions on Behaviour and Neuroendocrine Systems. *J. Neuroendocrinol.* 24, 144-159.
- Galea, L.A.M., Wainwright, S.R., Roes, M.M., Duarte-Guterman, P., Chow, C., Hamson, D.K., 2013. Sex, Hormones and Neurogenesis in the Hippocampus: Hormonal Modulation of Neurogenesis and Potential Functional Implications. *J. Neuroendocrinol.* 25, 1039-1061.
- Giatti, S., Garcia-Segura, L.M., Melcangi, R.C., 2015. New steps forward in the neuroactive steroid field. *J. Steroid Biochem. Mol. Biol.* 153, 127-134.
- Gibson, C.L., Attwood, L., 2016. The impact of gender on stroke pathology and treatment. *Neurosci. Biobehav. Rev.*
- Gill, R.K., Christakos, S., 1995. Regulation by estrogen through the 5'-flanking region of the mouse calbindin-D28k gene. *Mol. Endocrinol.* 9, 319-326.
- Gillies, G.E., Pienaar, I.S., Vohra, S., Qamhawi, Z., 2014. Sex differences in Parkinson's disease. *Front. Neuroendocrinol.* 35, 370-384.
- Gorski, R.A., Gordon, J.E., Shryne, J.E., Southam, A.M., 1978. Evidence for a morphological sex difference within the medial preoptic area of the rat brain. *Brain Res.* 148, 333-346.
- Gorski, R.A., Harlan, R.E., Jacobson, C.D., Shryne, J.E., Southam, A.M., 1980. Evidence for the existence of a sexually dimorphic nucleus in the preoptic area of the rat. *J. Comp. Neurol.* 193, 529-539.

- Grimm, A., Mensah-Nyagan, A.G., Eckert, A., 2016. Alzheimer, mitochondria and gender. *Neurosci.Biobehav.Rev.*
- Gu, G., Cornea, A., Simerly, R.B., 2003. Sexual differentiation of projections from the principal nucleus of the bed nuclei of the stria terminalis. *J. Comp. Neurol.* 460, 542-562.
- Hara, Y., Waters, E.M., McEwen, B.S., Morrison, J.H., 2015. Estrogen Effects on Cognitive and Synaptic Health Over the Lifecourse. *Physiol. Rev.* 95, 785-807.
- Hazell, G.G., Yao, S.T., Roper, J.A., Prossnitz, E.R., O'Carroll, A.M., Lolait, S.J., 2009. Localisation of GPR30, a novel G protein-coupled oestrogen receptor, suggests multiple functions in rodent brain and peripheral tissues. *J. Endocrinol.* 202, 223-236.
- Hill, R.A., 2016. Sex differences in animal models of schizophrenia shed light on the underlying pathophysiology. *Neurosci. Biobehav. Rev.*
- Huang, G.Z., Woolley, C.S., 2012. Estradiol acutely suppresses inhibition in the hippocampus through a sex-specific endocannabinoid and mGluR-dependent mechanism. *Neuron* 74, 801-808.
- Ingalhalikar, M., Smith, A., Parker, D., Satterthwaite, T., Elliott, M., Ruparel, K., Hakonarson, H., Gur, R., Gur, R., Verma, R., 2013. Sex differences in the structural connectome of the human brain. *Proceedings of the National Academy of Sciences of the United States of America.*
- Inslicht, S.S., Richards, A., Madden, E., Rao, M.N., O'Donovan, A., Talbot, L.S., Rucker, E., Metzler, T.J., Hauger, R.L., Neylan, T.C., 2014. Sex differences in neurosteroid and hormonal responses to metyrapone in posttraumatic stress disorder. *Psychopharmacology (Berl)* 231, 3581-3595.
- Joel, D., Berman, Z., Tavor, I., Wexler, N., Gaber, O., Stein, Y., Shefi, N., Pool, J., Urchs, S., Margulies, D.S., Liem, F., Hanggi, J., Jancke, L., Assaf, Y., 2015. Sex beyond the genitalia: The human brain mosaic. *Proc Natl Acad Sci U S A* 112, 15468-15473.
- Jorgensen, C.B., 1971. John Hunter, A.A. Berthold, and the origins of endocrinology. *Acta historica scientiarum naturalium et medicinalium* 24, 7-54.
- Kauffman, A.S., Gottsch, M.L., Roa, J., Byquist, A.C., Crown, A., Clifton, D.K., Hoffman, G.E., Steiner, R.A., Tena-Sempere, M., 2007. Sexual differentiation of Kiss1 gene expression in the brain of the rat. *Endocrinology* 148, 1774-1783.
- Kelley, M.H., Kuroiwa, M., Taguchi, N., Herson, P.S., 2011. Sex difference in sensitivity to allopregnanolone neuroprotection in mice correlates with effect on spontaneous inhibitory post synaptic currents. *Neuropharmacology* 61, 724-729.
- King, S.R., 2008. Emerging roles for neurosteroids in sexual behavior and function. *J Androl* 29, 524-533.

- Kipp, M., Hochstrasser, T., Schmitz, C., Beyer, C., 2016. Female sex steroids and glia cells: Impact on multiple sclerosis lesion formation and fine tuning of the local neurodegenerative cellular network. *Neurosci. Biobehav. Rev.*
- Kritzer, M.F., Creutz, L.M., 2008. Region and sex differences in constituent dopamine neurons and immunoreactivity for intracellular estrogen and androgen receptors in mesocortical projections in rats. *J. Neurosci.* 28, 9525-9535.
- Lavaque, E., Mayen, A., Azcoitia, I., Tena-Sempere, M., Garcia-Segura, L.M., 2006. Sex differences, developmental changes, response to injury and cAMP regulation of the mRNA levels of steroidogenic acute regulatory protein, cytochrome p450scc, and aromatase in the olivocerebellar system. *J. Neurobiol.* 66, 308-318.
- Lavranos, G., Angelopoulou, R., Manolakou, P., Balla, M., 2006. Hormonal and meta-hormonal determinants of sexual dimorphism. *Coll Antropol* 30, 659-663.
- Lenz, K.M., Nugent, B.M., McCarthy, M.M., 2012. Sexual differentiation of the rodent brain: dogma and beyond. *Front. Neurosci.* 6, 26.
- Li, R., Singh, M., 2014. Sex differences in cognitive impairment and Alzheimer's disease. *Front. Neuroendocrinol.* 35, 385-403.
- Lipscombe, C., Smith, K.J., Garipey, G., Schmitz, N., 2014. Gender differences in the relationship between anxiety symptoms and physical inactivity in a community-based sample of adults with type 2 diabetes. *Can J Diabetes* 38, 444-450.
- Litim, N., Morissette, M., Di Paolo, T., 2016. Neuroactive gonadal drugs for neuroprotection in male and female models of Parkinson's disease. *Neurosci. Biobehav. Rev.* in press.
- Lovick, T.A., 2014. Sex determinants of experimental panic attacks. *Neurosci. Biobehav. Rev.* 46 Pt 3, 465-471.
- Luque, J.M., Deblas, M.R., Segovia, S., Guillamon, A., 1992. Sexual dimorphism of the dopamine-beta-hydroxylase-immunoreactive neurons in the rat locus ceruleus. *Dev. Brain Res.* 67, 211-215.
- Ma, Y.Y., Kong, S.Z., Yang, L.J., Meng, J.L., Lv, L.C., He, M., 2007. [Sexual dimorphisms of dopaminergic neurons in rat substantia nigra]. *Sheng Li Xue Bao* 59, 753-758.
- Mannix, R., Berglass, J., Berkner, J., Moleus, P., Qiu, J., Jantzie, L.L., Meehan, W.P., 3rd, Stanley, R.M., Robinson, S., 2014. Sex differences in the effect of progesterone after controlled cortical impact in adolescent mice: a preliminary study. *J. Neurosurg.* 121, 1337-1341.
- Martini, M., Di Sante, G., Collado, P., Pinos, H., Guillamon, A., Panzica, G.C., 2008. Androgen receptors are required for full masculinization of nitric oxide synthase system in rat limbic-hypothalamic region. *Horm. Behav.* 54, 557-564.

- McHenry, J., Carrier, N., Hull, E., Kabbaj, M., 2014. Sex differences in anxiety and depression: role of testosterone. *Front. Neuroendocrinol.* 35, 42-57.
- Melcangi, R.C., Garcia-Segura, L.M., Mensah-Nyagan, A.G., 2008. Neuroactive steroids: state of the art and new perspectives. *Cell. Mol. Life Sci.* 65, 777-797.
- Melcangi, R.C., Giatti, S., Garcia-Segura, L.M., 2016. Levels and actions of neuroactive steroids in the nervous system under physiological and pathological conditions: Sex-specific features. *Neurosci. Biobehav. Rev.*
- Melcangi, R.C., Panzica, G.C., 2006. Neuroactive steroids: old players in a new game. *Neuroscience* 138, 733-739.
- Mendrek, A., Mancini-Marie, A., 2016. Sex/gender differences in the brain and cognition in schizophrenia. *Neurosci. Biobehav. Rev.*
- Mosera, V.A., Pike, C.J., 2016. Obesity and sex interact in the regulation of Alzheimer's disease. *Neurosci. Biobehav. Rev.*
- Murakami, S., Arai, Y., 1989. Neuronal death in the developing sexually dimorphic periventricular nucleus of the preoptic area in the female rat: effect of neonatal androgen treatment. *Neurosci. Lett.* 102, 185-190.
- Murray, H.E., Pillai, A.V., McArthur, S.R., Razvi, N., Datla, K.P., Dexter, D.T., Gillies, G.E., 2003. Dose- and sex-dependent effects of the neurotoxin 6-hydroxydopamine on the nigrostriatal dopaminergic pathway of adult rats: differential actions of estrogen in males and females. *Neuroscience* 116, 213-222.
- Ngo, S.T., Steyn, F.J., McCombe, P.A., 2014. Gender differences in autoimmune disease. *Front. Neuroendocrinol.* 35, 347-369.
- Nottebohm, F., 1980. Testosterone triggers growth of brain vocal control nuclei in adult canaries. *Brain Res.* 189, 429-436.
- Nottebohm, F., 1981. A brain for all seasons: cyclical anatomical changes in song control nuclei of the canary brain. *Science* 214, 1368-1370.
- Nottebohm, F., Arnold, A.P., 1976. Sexual dimorphism in vocal control areas of the songbird brain. *Science* 194, 211-213.
- Nugent, B.M., Wright, C.L., Shetty, A.C., Hodes, G.E., Lenz, K.M., Mahurkar, A., Russo, S.J., Devine, S.E., McCarthy, M.M., 2015. Brain feminization requires active repression of masculinization via DNA methylation. *Nat. Neurosci.* 18, 690-697.
- O'Malley, B.W., Means, A.R., 1974. Female steroid hormones and target cell nuclei. *Science* 183, 610-620.

- Panzica, G.C., Aste, N., Viglietti-Panzica, C., Ottinger, M.A., 1995. Structural sex differences in the brain: influence of gonadal steroids and behavioral correlates. *J. Endocrinol. Inv.* 18, 232-252.
- Panzica, G.C., Castagna, C., Viglietti-Panzica, C., Russo, C., Tlemçani, O., Balthazart, J., 1998. Organizational effects of estrogens on brain vasotocin and sexual behavior in quail. *J. Neurobiol.* 37, 684-699.
- Panzica, G.C., Viglietti-Panzica, C., Balthazart, J., 1996. The sexually dimorphic medial preoptic nucleus of quail: a key brain area mediating steroid action on male sexual behavior. *Front. Neuroendocrinol.* 17, 51-125.
- Papka, R.E., Mowa, C.N., 2003. Estrogen receptors in the spinal cord, sensory ganglia, and pelvic autonomic ganglia. *Int. Rev. Cytol.* 231, 91-127.
- Pasquier, J., Kamech, N., Lafont, A.-G., Vaudry, H., Rousseau, K., Dufour, S., 2014. Molecular evolution and structure-activity relationships of kisspeptins and their receptors. *J. Mol. Endocrinol.* 52, T101-117.
- Perrot-Sinal, T.S., Davis, A.M., McCarthy, M.M., 2001. Developmental sex differences in glutamic acid decarboxylase (GAD) and the housekeeping gene, GAPDH 65. *Brain Res.* 922, 201-208.
- Pesaresi, M., Giatti, S., Cavaletti, G., Abbiati, F., Calabrese, D., Bianchi, R., Caruso, D., Garcia-Segura, L.M., Melcangi, R.C., 2011a. Sex differences in the manifestation of peripheral diabetic neuropathy in gonadectomized rats: a correlation with the levels of neuroactive steroids in the sciatic nerve. *Exp. Neurol.* 228, 215-221.
- Pesaresi, M., Giatti, S., Cavaletti, G., Abbiati, F., Calabrese, D., Lombardi, R., Bianchi, R., Lauria, G., Caruso, D., Garcia-Segura, L.M., Melcangi, R.C., 2011b. Sex-dimorphic effects of dehydroepiandrosterone in diabetic neuropathy. *Neuroscience*.
- Pesaresi, M., Maschi, O., Giatti, S., Garcia-Segura, L.M., Caruso, D., Melcangi, R.C., 2010. Sex differences in neuroactive steroid levels in the nervous system of diabetic and non-diabetic rats. *Horm. Behav.* 57, 46-55.
- Peterson, B.L., Won, S., Geddes, R.I., Sayeed, I., Stein, D.G., 2015. Sex-related differences in effects of progesterone following neonatal hypoxic brain injury. *Behav. Brain Res.* 286, 152-165.
- Phoenix, C.H., Goy, R.W., Gerall, A.A., Young, W.C., 1959. Organizing action of prenatally administered testosterone propionate on the tissues mediating mating behavior in the female guinea pig. *Endocrinology* 65, 369-382.
- Pierman, S., Sica, M., Allieri, F., Viglietti-Panzica, C., Panzica, G.C., Bakker, J., 2008. Activational effects of estradiol and dihydrotestosterone on social recognition and the arginine-

- vasopressin immunoreactive system in male mice lacking a functional aromatase gene. *Horm. Behav.* 54, 98-106.
- Plumari, L., Viglietti Panzica, C., Allieri, F., Honda, S., Harada, N., Absil, P., Balthazar, J., Panzica, G.C., 2002. Changes in the Arginine-Vasopressin Immunoreactive Systems in Male Mice Lacking a Functional Aromatase Gene. *J. Neuroendocrinol.* 14, 971–978.
- Porcu, P., Barron, A.M., Frye, C.A., Walf, A.A., Yang, S.Y., He, X.Y., Morrow, A.L., Panzica, G.C., Melcangi, R.C., 2016. Neurosteroidogenesis Today: Novel Targets for Neuroactive Steroid Synthesis and Action and Their Relevance for Translational Research. *J. Neuroendocrinol.* 28.
- Raisman, G., Field, P.M., 1971. Sexual dimorphism in the preoptic area of the rat. *Science* 173, 731-733.
- Raisman, G., Field, P.M., 1973. Sexual dimorphism in the neuropil of the preoptic area of the rat and its dependence on neonatal androgen. *Brain Res.* 54, 1-29.
- Ramien, C., A., T., Lupu, A., Heckmann, N., Broder Engler, J., Patas, K., Friese, M.A., Gold, S.M., 2016. Sex effects on inflammatory and neurodegenerative processes in multiple sclerosis. *Neurosci.Biobehav.Rev.*
- Reddy, D.S., 2014. Neurosteroids and their role in sex-specific epilepsies. *Neurobiol Dis* 72 Pt B, 198-209.
- Reisert, I., Pilgrim, C., 1991. Sexual differentiation of monoaminergic neurons--genetic or epigenetic? *Trends Neurosci.* 14, 468-473.
- Reisert, I., Schuster, R., Zienecker, R., Pilgrim, C., 1990. Prenatal development of mesencephalic and diencephalic dopaminergic systems in the male and female rat. *Brain Res Dev Brain Res* 53, 222-229.
- Romano, E., Cosentino, L., Laviola, G., De Filippis, B., 2016. Genes and sex hormones in neurodevelopmental disorders. *Neurosci.Biobehav.Rev.*
- Rupprecht, R., Holsboer, F., 1999. Neuroactive steroids: mechanisms of action and neuropsychopharmacological perspectives. *Trends Neurosci.* 22, 410-416.
- Scharfman, H.E., MacLusky, N.J., 2014. Sex differences in the neurobiology of epilepsy: a preclinical perspective. *Neurobiol Dis* 72 Pt B, 180-192.
- Sellers, K., Raval, P., Srivastava, D.P., 2015. Molecular signature of rapid estrogen regulation of synaptic connectivity and cognition. *Front. Neuroendocrinol.* 36, 72-89.
- Sibug, R., Kuppers, E., Beyer, C., Maxson, S.C., Pilgrim, C., Reisert, I., 1996. Genotype-Dependent Sex-Differentiation of Dopaminergic-Neurons in Primary Cultures of Embryonic Mouse-Brain. *Dev. Brain Res.* Vol 93, Iss 1-2, 136-142.

- Sickel, M.J., McCarthy, M.M., 2000. Calbindin-D28k immunoreactivity is a marker for a subdivision of the sexually dimorphic nucleus of the preoptic area of the rat: Developmental profile and gonadal steroid modulation. *J. Neuroendocrinol.* 12, 397-402.
- Simerly, R.B., 2002. Wired for reproduction: organization and development of sexually dimorphic circuits in the mammalian forebrain. *Annu. Rev. Neurosci.* 25, 507-536.
- Simerly, R.B., Swanson, L.W., Gorski, R.A., 1985. The distribution of monoaminergic cells and fibers in a periventricular preoptic nucleus involved in the control of gonadotropin release: immunohistochemical evidence for a dopaminergic sexual dimorphism. *Brain Res.* 330, 55-64.
- Simerly, R.B., Zee, M.C., Pendleton, J.W., Lubhan, D.B., Korach, K.S., 1997. Estrogen receptor-dependent sexual differentiation of dopaminergic neurons in the preoptic region of the mouse. *Proceedings of the National Academy of Sciences of the United States of America* 94, 14077-14082.
- Sinclair, A.H., Berta, P., Palmer, M.S., Hawkins, J.R., Griffiths, B.L., Smith, M.J., Foster, J.W., Frischauf, A.M., Lovell-Badge, R., Goodfellow, P.N., 1990. A gene from the human sex-determining region encodes a protein with homology to a conserved DNA-binding motif. *Nature* 346, 240-244.
- Smith, C.A., Roeszler, K.N., Ohnesorg, T., Cummins, D.M., Farlie, P.G., Doran, T.J., Sinclair, A.H., 2009. The avian Z-linked gene DMRT1 is required for male sex determination in the chicken. *Nature* 461, 267-271.
- Spritzer, M., Galea, L., 2007. Testosterone and dihydrotestosterone, but not estradiol, enhance survival of new hippocampal neurons in adult male rats. *Dev. Neurobiol.* 67, 1321-1333.
- Srivastava, D.P., Evans, P.D., 2013. G-Protein Oestrogen Receptor 1: Trials and Tribulations of a Membrane Oestrogen Receptor. *J. Neuroendocrinol.* 25, 1219-1230.
- Swaab, D.F., Fliers, E., 1985. A sexually dimorphic nucleus in the human brain. *Science* 228, 1112-1115.
- Tabatadze, N., Huang, G., May, R.M., Jain, a., Woolley, C.S., 2015. Sex Differences in Molecular Signaling at Inhibitory Synapses in the Hippocampus. *J. Neurosci.* 35, 11252-11265.
- Tanapat, P., Hastings, N.B., Gould, E., 2005. Ovarian steroids influence cell proliferation in the dentate gyrus of the adult female rat in a dose- and time-dependent manner. *J. Comp. Neurol.* 481, 252-265.
- Thanky, N.R., Son, J.H., Herbison, A.E., 2002. Sex differences in the regulation of tyrosine hydroxylase gene transcription by estrogen in the locus coeruleus of TH9-LacZ transgenic mice. *Mol. Brain Res.* 104, 220-226.

- van Luijtelaar, G., Onat, F.Y., Gallagher, M.J., 2014. Animal models of absence epilepsies: what do they model and do sex and sex hormones matter? *Neurobiol Dis* 72 Pt B, 167-179.
- Viglietti-Panzica, C., Panzica, G.C., Fiori, M.G., Calcagni, M., Anselmetti, G.C., Balthazart, J., 1986. A sexually dimorphic nucleus in the quail preoptic area. *Neurosci. Lett.* 64, 129-134.
- Waters, E.M., Simerly, R.B., 2009. Estrogen induces caspase-dependent cell death during hypothalamic development. *J. Neurosci.* 29, 9714-9718.
- Wissman, A.M., May, R.M., Woolley, C.S., 2012. Ultrastructural analysis of sex differences in nucleus accumbens synaptic connectivity. *Brain Struct. Funct.* 217, 181-190.
- Young, L.J., Pfaff, D.W., 2014. Sex differences in neurological and psychiatric disorders. *Front. Neuroendocrinol.* 35, 253-254.
- Zhao, W., Katzmarzyk, P.T., Horswell, R., Wang, Y., Johnson, J., Hu, G., 2014. Sex differences in the risk of stroke and HbA(1c) among diabetic patients. *Diabetologia* 57, 918-926.
- Zhou, Y., Zhao, M., Zhou, C., Li, R., 2016. Sex differences in drug addiction and response to exercise intervention: From human to animal studies. *Front. Neuroendocrinol.* 40, 24-41.
- Zup, S.L., Edwards, N.S., McCarthy, M.M., 2014. Sex- and age-dependent effects of androgens on glutamate-induced cell death and intracellular calcium regulation in the developing hippocampus. *Neuroscience* 281C, 77-87.