

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

**Effect of curcumin on circulating interleukin-6 concentrations: A systematic review and meta-analysis of randomized controlled trials**

**This is the author's manuscript**

*Original Citation:*

*Availability:*

This version is available <http://hdl.handle.net/2318/1633548> since 2017-05-15T16:25:24Z

*Published version:*

DOI:10.1016/j.phrs.2016.07.004

*Terms of use:*

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)



## UNIVERSITÀ DEGLI STUDI DI TORINO

This Accepted Author Manuscript (AAM) is copyrighted and published by Elsevier. It is posted here by agreement between Elsevier and the University of Turin. Changes resulting from the publishing process - such as editing, corrections, structural formatting, and other quality control mechanisms - may not be reflected in this version of the text. The definitive version of the text was subsequently published in [*Pharmacological Research*, 111; 2016; doi: 10.1016/j.phrs.2016.07.004].

You may download, copy and otherwise use the AAM for non-commercial purposes provided that your license is limited by the following restrictions:

- (1) You may use this AAM for non-commercial purposes only under the terms of the CC-BY-NC-ND license.
- (2) The integrity of the work and identification of the author, copyright owner, and publisher must be preserved in any copy.
- (3) You must attribute this AAM in the following format: Creative Commons BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/deed.en>), [doi: 10.1016/j.phrs.2016.07.004 *link*: [www.ncbi.nlm.nih.gov/pubmed/27392742](http://www.ncbi.nlm.nih.gov/pubmed/27392742)]

**Effect of curcumin on circulating interleukin-6 concentrations: A systematic review and meta-analysis of randomized controlled trials**

Giuseppe Derosaa,b,c,d, Pamela Maffioli a,e, Luis E. Simental-Mendíaf , Simona Bog, Amirhossein Sahebkar h

a Centre of Diabetes and Metabolic Diseases, Department of Internal Medicine and Therapeutics, University of Pavia and Fondazione IRCCS Policlinico S. Matteo, Pavia, Italy

b Centre for the Study of Endocrine-Metabolic Pathophysiology and Clinical Research, University of Pavia, Pavia, Italy

c Centre for Prevention, Surveillance, Diagnosis and Treatment of Rare Diseases, Fondazione IRCCS Policlinico S. Matteo, Pavia, Italy

d Laboratory of Molecular Medicine, University of Pavia, Pavia, Italy

e PhD School in Experimental Medicine, University of Pavia, Pavia, Italy

f Biomedical Research Unit, Mexican Social Security Institute, Durango, Mexico

g Department of Medical Sciences, University of Turin, Turin, Italy

h Biotechnology Research Center, Mashhad University of Medical Sciences, Mashhad, Iran

**Word count:** 2497 **Running title:** curcumin and interleukin-6

**ABSTRACT**

The aim of this meta-analysis was to evaluate the efficacy of curcuminoids supplementation on circulating concentrations of IL-6 in randomized controlled trials (RCTs). The search included PubMed-Medline,

Scopus, Web of Science and Google Scholar databases by up to November 01, 2015, to identify RCTs investigating the impact of curcuminoids on circulating IL-6 concentrations. Nine RCTs comprising 10 treatment arms were found to be eligible for the meta-analysis. There was a significant reduction of circulating IL-6 concentrations following curcuminoids supplementation (WMD:  $-0.60$  pg/mL, 95% CI:  $-1.06$ ,  $-0.14$ ,  $p = 0.011$ ). Meta-regression did not suggest any significant association between the circulating IL-6 lowering effects of curcuminoids with either dose or duration of treatment. There was a significant association between the IL-6-lowering activity of curcumin and baseline IL-6 concentration (slope:  $-0.51$ ; 95% CI:  $-0.80$ ,  $-0.23$ ;  $p = 0.005$ ). This meta-analysis of RCTs suggested a significant effect of curcumin in lowering circulating IL-6 concentrations. This effect appears to be more evident in patients with higher degrees of systemic inflammation

## **Introduction**

Curcuminoids (curcumin, demethoxycurcumin or curcumin I, bisdemethoxycurcumin or curcumin II) are the bioactive dietary polyphenols from turmeric (*Curcuma Longa*), the most popular spice in Indian cuisine and a major ingredient of curry powders [1]. Several reviews and meta-analyses showed that dietary curcuminoids have various biological activities [2–7], including anti-oxidant [8], anti-inflammatory [3], anti-microbial [9], anti-pruritic [10], hepatoprotective [11], anti-depressant [12], anti-arthritis [13], anti-ischemic [14], lipid-modifying [15,16], metabolic [17], hypouricemic [18], and neuroprotective properties [19]. They also mitigate endothelial dysfunction and improve several clinical and biochemical features of cancer [20–24], giving curcuminoids the potential to be used in clinical practice for a range of human diseases [3,25–27]. Curcuminoids have been tested as a therapeutic option in various diseases such as diabetes, cancer, arthritis, chronic inflammatory and cutaneous diseases, virus infections, Alzheimer's disease and cardiovascular diseases via their multiple pleiotropic effects on genes and cell-signaling pathways [3,5–7]. Most studies on curcuminoids were in-vitro and animal studies; clinical trials have

indicated their safety, tolerability and non-toxicity but small number of patients were enrolled and the real efficacy in humans is at present questionable [3,5,6]. Furthermore, the low availability of these compounds requires the development of formulations with increased solubilization and resistant to inactivation by hydrolysis [5–7]. Cytokines are small molecules with protein or glycoprotein structure (8–80 kDa). They are products of activated immune cells that act as molecular signals between immune competent cells [26]. Among cytokines, interleukin-1 (IL-1), tumor necrosis factor- (TNF-), and interleukin-6 (IL-6) are major inducers of acute phase response [27]. In particular, IL-6 is a multi-functional cytokine that plays a central role in host defense due to its wide range of immune and hematopoietic activities and its potent ability to induce the acute phase response [28]. Interleukin-6 plays a role in the pathogenesis of many human diseases, such as multiple myeloma, rheumatoid arthritis, Castleman’s disease, AIDS, mesangial proliferative glomerulonephritis, psoriasis, Kaposi’s sarcoma, sepsis, and osteoporosis [28]. Considering its activities, IL-6 appeared to be a viable target for auto-immune disease. Inhibitors of IL-6 were successful in animal models of autoimmune disease paving the way for subsequent studies in humans. Among agents able to inhibit expression and activity of IL-6, curcuminoids proved to have beneficial effects on IL-6 in several in-vitro and animal studies [29–34]. However, clinical studies regarding this effect of curcuminoids in humans have not been fully consistent [29–34]. Hence, the aim of the present meta-analysis was to evaluate the efficacy of curcuminoids supplementation on circulating concentrations of IL-6 in randomized controlled trials (RCTs) in order to obtain a conclusive result on this potential effect of the compound.

## **Methods**

### *Search strategy*

This study was designed according to the guidelines of the 2009 preferred reporting items for systematic reviews and metaanalysis (PRISMA) statement [35]. Medline (<http://www.ncbi.nlm.nih.gov/pubmed>), SCOPUS, Web of Science and Google Scholar databases were searched using the following search terms

in titles and abstracts (also in combination with MESH terms): (curcumin OR curcuminoid OR curcuminoids OR Curcuma OR Curcuma longa OR turmeric) AND (IL-6 OR interleukin-6 OR “interleukin 6” OR interleukin6 OR IL6 OR “IL 6”) AND (random OR randomized OR randomly OR randomization OR “randomized controlled trial” OR “randomized trial” OR “randomized study” OR “random number” OR placebo). The wild-card term “\*” was used to increase the sensitivity of the search strategy. The search was limited to studies in English language. The literature was searched from inception to November 01, 2015.

#### *Study selection*

Original studies were included if they met the following inclusion criteria: (i) being a controlled trial with either parallel or cross-over design, (ii) investigating the impact of supplementation with curcuminoids or turmeric preparations with known amounts of curcuminoids on circulating IL-6 concentrations, and (iii) presentation of sufficient information on changes in circulating concentrations of IL-6 in the study groups. Exclusion criteria were (i) uncontrolled trials, (ii) inclusion of an active control group in the study design, (iii) observational studies with case-control, cross-sectional or cohort designs, (iv) using non-standardized turmeric extracts with unknown curcuminoids content, (v) trials with treatment durations of < 2 weeks, and (vi) incomplete data on circulating concentrations of IL-6. In case of the latter item, authors of the article(s) were contacted and requested to provide necessary numerical data.

#### *Data extraction*

Eligible studies were reviewed and the following data were abstracted: 1) first author's name; 2) year of publication; 3) study location; 4) study design; 5) number of participants in the curcumin and placebo groups; 6) type and dose of curcumin supplement used; 7) duration of treatment; 8) age, gender and body mass index (BMI) of study participants; 9) inclusion criteria defined in the study; 10) systolic and diastolic blood pressures; and 11) baseline and follow-up IL-6 concentrations.

#### *Quality assessment*

A systematic assessment of bias in the included studies was performed using the Cochrane criteria [36]. The items used for the assessment of each study were as follows: adequacy of sequence generation, allocation concealment, blinding, addressing of dropouts (incomplete outcome data), selective outcome reporting, and other potential sources of bias. According to the recommendations of the Cochrane Handbook, a judgment of “yes” indicated low risk of bias, while “no” indicated high risk of bias. Labeling an item as “unclear” indicated an unclear or unknown risk of bias.

#### *Quantitative data synthesis*

Meta-analysis was conducted using Comprehensive MetaAnalysis (CMA) V2 software (Biostat, NJ) [37]. Plasma IL-6 concentrations were collated in pg/mL. Net changes in measurements (change scores) were calculated as follows: measure at end of follow-up—measure at baseline. For crossover trials with a 2 × 2 design, each treatment arm was analysed separately, and net change in each arm was calculated by subtracting the value after control intervention from that reported after treatment. Standard deviations (SDs) of the mean difference were calculated using the following formula:  $SD = \text{square root} [(SD_{\text{pre-treatment}})^2 + (SD_{\text{post-treatment}})^2 - (2R \times SD_{\text{pre-treatment}} \times SD_{\text{post-treatment}})]$ , assuming a correlation coefficient (R) = 0.5. Where standard error of the mean (SEM) was only reported, standard deviation (SD) was estimated using the following formula:  $SD = SEM \times \text{sqrt}(n)$ , where n is the number of subjects. A random-effects model (using DerSimonian-Laird method) and the generic inverse variance method were used to compensate for the heterogeneity of studies in terms of demographic characteristics of populations being studied and also differences in study design and duration of treatment. Heterogeneity was quantitatively assessed using I<sup>2</sup> index. In order to evaluate the influence of each study on the overall effect size, sensitivity analysis was conducted using leave-one-out method, i.e. removing one study each time and repeating the analysis [38,39].

#### *Meta-regression*

Random-effects meta-regression was performed using unrestricted maximum likelihood method to evaluate the association between calculated WMD and potential confounders including dose and duration of supplementation with curcumin, and baseline IL-6 levels.

#### *Publication bias*

Potential publication bias was explored using visual inspection of Begg's funnel plot asymmetry, fail-safe N test, and Begg's rank correlation and Egger's weighted regression tests. Duval & Tweedie "trim and fill" method was used to adjust the analysis for the effects of publication bias [40].

### **Results**

#### *Flow and characteristics of included studies*

Briefly, after multiple databases searching 149 published studies were identified and the abstracts were reviewed. Next, 32 nonoriginal articles were excluded. From the remainder of articles, 85 did not meet the inclusion criteria after assessment of titles and abstracts and were also excluded. Thus, 32 full-text articles were carefully assessed and reviewed for eligibility; of which 23 studies were excluded for not measuring circulating IL-6 concentrations (n = 19), being non-clinical (n = 1), and having a short treatment duration (n = 3). Finally, 9 studies were eligible and included in the systematic review and meta-analysis. The study selection process is shown in Fig. 1. Data were pooled from 9 eligible studies [41–49] comprising 10 treatment arms which included 609 subjects, with 305 in the curcuminoids arm and 304 in the control arm (participants enrolled from the cross-over trial were considered in both arms). Included studies were published between 2008 and 2015. The dose of curcuminoids used changed in the various trials. One study reported effects of curcuminoids 200 mg/day [41], two studies reported effects of curcuminoids 300 mg/day [42,43], three studies of curcuminoids 1 g/day [44–46], two studies of curcuminoids 1.5 g/day [47,48], and one study curcuminoids 6 g/day [49]. The range of intervention periods was from 2 weeks [49] up to 8 months [41]. Study design of almost all included studies was parallel group, except one which was cross-over design [44]. Selected studies enrolled subjects with osteoarthritis [41], knee osteoarthritis



[48], oral lichen planus [49], obesity [44], type 2 diabetes [42,43], metabolic syndrome [46], sulfur mustard intoxication [47], and chronic pruritic skin lesions [45]. Demographic and biochemical characteristics of the evaluated studies are presented in Table 1.

#### *IL-6 assay methods*

Different assays methods were used to measure serum IL-6 concentrations. In this regard, most studies [40–42,44–46] determined serum IL-6 levels by enzyme-linked immunoassay. Ganjali et al. [43] analyzed IL-6 concentrations using the Biochip Array Technology on the Randox Evidence Investigator (Randox Laboratories, Belfast, Northern Ireland) by chemiluminescent immunoassay. Only two studies did not provide the method used to determine serum IL-6 levels [47,48].

#### *Risk of bias assessment*

According random sequence generation and allocation concealment, six studies were characterized by lack of information (42–46,48) and one study showed high risk of bias (41). Five studies had insufficient information (42,44–46,48) and two exhibited high risk of bias (46,48) with respect to blinding of participants and personnel. In addition, almost all included trials did not provide sufficient information of blinding of outcome assessment (41–46,48,49). However, all included studies showed low risk of bias according to selective reporting and incomplete outcome data. Details of the quality of bias assessment are shown in Table 2.

#### *Effect of curcuminoids on circulating IL-6 concentrations*

Overall, the impact of curcuminoids on circulating IL-6 concentrations was reported in 9 RCTs comprising 10 treatment arms. Meta-analysis suggested a significant reduction in plasma IL-6 concentrations following curcuminoids supplementation (WMD:  $-0.60$  pg/mL, 95% CI:  $-1.06$ ,  $-0.14$ ,  $p = 0.011$ ) (Fig. 2). Leave-one out sensitivity analysis showed that this effect size is marginally sensitive to two studies (40,42) (Fig. 2). Subgroup analyses were performed to assess the impact of administered dose and duration of supplementation with curcuminoids, and application of bioavailability-enhanced formulations on the

effect size (Fig. 2). Curcuminoids dose and formulation were not found to significantly affect the IL-6-lowering effect of curcuminoids. However, there was a weaker reduction in circulating IL-6 concentrations in the subset of trials with  $\geq 8$  weeks of duration (WMD: -0.10 pg/mL, 95% CI: -0.61, 0.41,  $p = 0.705$ ; Figure 3) compared with the subset of studies lasting  $< 8$  weeks (WMD: -1.35 pg/mL, 95% CI: -2.13, -0.58,  $p = 0.001$ ) (Figure 3).

#### *Meta-regression*

Random-effects meta-regression was performed to evaluate if changes in circulating IL-6 concentrations are dependent to dose and duration of treatment. Meta-regression analysis did not suggest any significant association between changes in circulating IL-6 levels with either dose (slope: 0.0003; 95% CI: -0.0001, 0.0008;  $p = 0.172$ ) or duration (slope: -0.01; 95% CI: -0.07, 0.05;  $p = 0.682$ ) of treatment (Figure 4).

#### *Publication bias*

The funnel plot of standard error versus effect size (mean difference) was slightly asymmetric. “trim and fill” correction suggested one potentially missing study on the left side of funnel plot (Figure 5). Imputation for this potentially missing study yielded an effect size of -0.68 pg/mL (95% CI: -1.13, -0.22). The presence of publication bias was excluded by Egger’s linear regression (intercept = -1.14, standard error = 1.57; 95% CI = -4.76, 2.48,  $t = 0.73$ ,  $df = 8$ , two-tailed  $p = 0.488$ ) and Begg’s rank correlation (Kendall’s Tau with continuity correction = 0.04,  $z = 0.18$ , two-tailed  $p$ -value = 0.858) tests. The “fail-safe N” test showed that 109 studies would be needed to bring the WMD down to a non-significant ( $p > 0.05$ ) value.

#### **Discussion**

In this meta-analysis, we recorded a significant reduction of circulating IL-6 concentrations following curcuminoids supplementation; this effect does not seem to be dose-dependent, in fact meta-regression did not suggest any significant association between the circulating IL-6-lowering effects of curcuminoids with either dose or duration of treatment. Our results fit with several lines of evidences. At a molecular

and cellular level, curcumin modulates tumor necrosis factor- (TNF-) expression and the cell signaling mediated by TNF- by inhibition of p300/CREB-specific acetyltransferase which leads to repression of acetylation of histone/non histone proteins and hence repression of transcription [50,51]. Curcumin may also be a TNF blocker by binding to TNF directly [52]. Curcumin also inhibits cyclooxygenase-2, signal transducers and activators of transcription (STAT), cyclinD1, nuclear factor kappa-light-chain-enhancer of activated B cells (NF-kB) signaling pathways [53], as well as IL- 1b and IL-6 expression inhibiting mitogen-activated protein kinase and NF-kB pathways [54]. As regards animal studies, the anti-inflammatory effect of curcumin was first demonstrated in acute and chronic models of inflammation in rats and mice [55], where curcumin suppressed carrageenan-induced edema. Since then, a consistent number of animal models of inflammation have benefited from treatment with curcumin and several mechanisms by which curcumin can exhibit anti-inflammatory activity have been hypothesized [7,52]. In human diseases, extensive clinical trials have addressed the pharmacokinetics, safety and efficacy of curcumin in patients with various inflammatory conditions. An exhaustive list of clinical trials testing the effects of curcumin in various human diseases has been reported by Gupta et al. [56]. Common to most of these studies have been the safety, tolerability and the ability of curcumin to modulate numerous signaling molecules leading to reduced local or even systemic inflammation [7,56]. Interleukin-6, a cytokine featuring redundancy and pleiotropic activity, is one of the most important inflammatory cytokines. Besides its multiple roles in the innate immune system, IL-6 is necessary for the differentiation of T-helper-17 cells [57]; as such, IL-6 is an important factor for the coordination of the innate and acquired immune response [58]. Anomalies in IL-6 production played a pathological role in several autoimmune and chronic inflammatory diseases; for this reason, it has been hypothesized that targeting IL-6 could be an approach to the treatment of these diseases [59]. Furthermore, several anti-IL-6/IL-6 receptor monoclonal antibodies developed for targeted therapy demonstrated promising results in both preclinical studies and clinical trials [60]. In addition, several treatments, including statins, metformin, aspirin, methotrexate, exercise, and others, showed to

reduce systemic inflammation throughout reducing plasma IL-6 levels [61–65]. We reported that curcumin reduced IL-6 by a mean of 0.6 pg/mL; this results might be considered of interest if we look at previous studies in patients with evidence of low-grade systemic inflammation receiving effective consolidated therapies such as statins, metformin and aspirin [61–63]. In these studies, the degree of treatment-mediated IL-6 reduction was comparable to that we observed in the present meta-analysis. Thus, in patients with unstable angina simvastatin reduced plasma IL-6 by 0.4 pg/mL compared to placebo [61]; a 0.35 pg/mL reduction was obtained with metformin in patients with the metabolic syndrome [62]. Finally, in patients with chronic stable angina, aspirin reduced plasma IL-6 by 0.6 pg/mL compared to placebo [63]. As expected, greater plasma IL-6 reductions have been obtained in patients with high-grade systemic inflammation receiving the potent anti-inflammatory methotrexate [64]. In this regard, our meta-regression showed that there was a significant association between the IL-6-lowering activity of curcumin and baseline IL-6 concentration. Therefore, we must consider that the anti-inflammatory effect of curcumin might be more evident in patients with a greater degree of systemic inflammation. Overall, our results of a significant IL-6 reduction by curcumin supplementation support the idea that this nutraceutical may have a role in suppressing pro-inflammatory pathways linked with different diseases. Moreover, clinical trials with curcumin indicate safety, tolerability, and nontoxicity [56]. However, both the clinical efficacy in published trial [56] and the degree of IL-6 lowering in the present meta-analysis is questionable, based, first of all, on the small numbers of patients in each study. Additional limitations of the present meta-analysis need to be considered. Included studies were heterogeneous regarding population characteristics, study design, and duration of supplementation. Nevertheless, the impact of heterogeneity on estimated effect sizes was minimized by choosing a random-effects mode of analysis. Also, only three studies examined the impact of curcumin bioavailability-enhanced formulations on IL-6 [42,43,49]. Active scientific research has been carried out to improve curcumin's pharmacokinetics, systemic bioavailability, biological activity by loading curcumin into nano formulations [56]; now it should

be verified if these novel formulations have stronger IL-6 lowering efficacy. Finally, an overall 0.6 pg/mL reduction of circulating IL-6 might be considered modest; whether such a reduction might be enough to have a clinical benefit it need to be evaluated in larger clinical trials.

#### *Limitations and strengths*

The present meta-analysis has some limitations. Included studies were heterogeneous regarding population characteristics, study design, and duration of supplementation. Nevertheless, the impact of heterogeneity on estimated effect sizes was minimized by choosing a random-effects mode of analysis. In addition, elevated serum IL-6 levels were not among the inclusion criteria of any of the studies considered for this meta-analysis. Finally, most of the included studies had relatively short follow-up durations. Although, no significant association between the IL-6-lowering effect of curcuminoids and treatment duration was observed, the effect of prolonged administration of curcuminoids on circulating IL-6 levels is still not clear. Finally, active scientific research has been carried out to improve curcuminoids' pharmacokinetics, systemic bioavailability, and biological activity by loading curcuminoids into nanoformulations [3]; whether these novel formulations have stronger IL-6 lowering efficacy should be determined. Finally, an overall 0.6 pg/mL reduction of circulating IL-6 might be considered modest; whether such a reduction might translate into a clinical benefit should be evaluated for each possible therapeutic indication in larger clinical trials.

#### *Conclusion*

In conclusion, the present meta-analysis of RCTs suggested a significant effect of curcumin in lowering circulating IL-6 concentrations. Despite considerable pharmacologic potential, the clinical efficacy of curcumin is still poorly recognized [3] and this polyphenol has not yet been approved for clinical use in humans. Further studies in larger populations are needed to investigate which patients are best suitable for curcumin supplementation and which formulation should be preferred to increase its anti-inflammatory effects.

## References

- [1] A. Goel, A.B. Kunnumakkara, B.B. Aggarwal, Curcumin as Curcumin: from kitchen to clinic, *Biochem. Pharmacol.* 75 (4) (2008) 787–809.
- [2] A. Sahebkar, A.F. Cicero, L.E. Simental-Mendía, B.B. Aggarwal, S.C. Gupta, Curcumin downregulates human tumor necrosis factor- levels: a systematic review and meta-analysis of randomized controlled trials, *Pharmacol. Res.* 107 (2016) 234–242.
- [3] A. Sahebkar, Why it is necessary to translate curcumin into clinical practice for the prevention and treatment of metabolic syndrome? *Biofactors* 39 (2) (2013) 197–208.
- [4] A. Sahebkar, Are curcuminoids effective C-reactive protein-lowering agents in clinical practice? Evidence from a meta-analysis, *Phytother. Res.* 28 (5) (2014) 633–642.
- [5] M. Schaffer, P.M. Schaffer, G. Bar-Sela, An update on Curcuma as a functional food in the control of cancer and inflammation, *Curr. Opin. Clin. Nutr. Metab. Care* 18 (2015) 605–611.
- [6] S. Ghosh, S. Banerjee, P.C. Sil, The beneficial role of curcumin on inflammation, diabetes and neurodegenerative disease: a recent update, *Food Chem. Toxicol.* 83 (2015) 111–124.
- [7] Y. He, Y. Yue, X. Zheng, K. Zhang, S. Chen, Z. Du, Curcumin, inflammation, and chronic diseases: how are they linked, *Molecules* 20 (2015) 9183–9213.
- [8] A. Sahebkar, A. Mohammadi, A. Atabati, S. Rahiman, S. Tavallaie, M. Iranshahi, S. Akhlaghi, G.A. Ferns, M. Ghayour-Mobarhan, Curcuminoids modulate pro-oxidant-antioxidant balance but not the immune response to heat shock protein 27 and oxidized LDL in obese individuals, *Phytother. Res.* 27 (12) (2013) 1883–1888.
- [9] R.K. Naz, M.L. Lough, E.K. Barthelmess, Curcumin: a novel non-steroidal contraceptive with antimicrobial properties, *Front. Biosci. (Elite Ed.)* 8 (2016) 113–128.
- [10] Y. Panahi, A. Sahebkar, M. Amiri, S.M. Davoudi, F. Beiraghdar, S.L. Hoseininejad, M. Kolivand, Improvement of sulphur mustard-induced chronic pruritus, quality of life and antioxidant status by

curcumin: results of a randomised, double-blind, placebo-controlled trial, *Br. J. Nutr.* 108 (7) (2012) 1272–1279.

[11] S. Rahmani, S. Asgary, G. Askari, M. Keshvari, M. Hatamipour, A. Feizi, A. Sahebkar, Treatment of non-alcoholic fatty liver disease with curcumin: a randomized placebo-controlled trial, *Phytother. Res.* (2016), <http://dx.doi.org/10.1002/ptr.5659>.

[12] Y. Panahi, R. Badeli, G.R. Karami, A. Sahebkar, Investigation of the efficacy of adjunctive therapy with bioavailability-boosted curcuminoids in major depressive disorder, *Phytother. Res.* 29 (1) (2015) 17–21.

[13] Y. Panahi, A.R. Rahimnia, M. Sharafi, G. Alishiri, A. Saburi, A. Sahebkar, Curcuminoid treatment for knee osteoarthritis: a randomized double-blind placebo-controlled trial, *Phytother. Res.* 28 (11) (2014) 1625–1631.

[14] A. Sahebkar, Molecular mechanisms for curcumin benefits against ischemic injury, *Fertil. Steril.* 94 (5) (2010) e75–e76, author reply e77.

[15] A. Mohammadi, A. Sahebkar, M. Iranshahi, M. Amini, R. Khojasteh, M. Ghayour-Mobarhan, G.A. Ferns, Effects of supplementation with curcuminoids on dyslipidemia in obese patients: a randomized crossover trial, *Phytother. Res.* 27 (3) (2013) 374–379.

[16] A. Sahebkar, Curcuminoids for the management of hypertriglyceridaemia, *Nat. Rev. Cardiol.* 11 (2) (2014) 123.

[17] Y. Panahi, N. Khalili, M.S. Hosseini, M. Abbasnazar, A. Sahebkar, Lipid-modifying effects of adjunctive therapy with curcuminoids-piperine combination in patients with metabolic syndrome: results of a randomized controlled trial, *Complement. Ther. Med.* 22 (5) (2014) 851–857.

[18] Y. Panahi, P. Kianpour, R. Mohtashami, R. Jafari, L.E. Simental-Mendía, A. Sahebkar, Curcumin lowers serum lipids and uric acid in subjects with non-alcoholic fatty liver disease: a randomized controlled trial, *J. Cardiovasc. Pharmacol.* (2016), <http://dx.doi.org/10.1097/FJC.0000000000000406>.

- [19] A. Shakeri, A. Sahebkar, Optimized curcumin formulations for the treatment of Alzheimer's disease: a patent evaluation, *J. Neurosci. Res.* 94 (2) (2016) 111–113.
- [20] R.F. Tayyem, D.D. Heath, W.K. Al-Delaimy, C.L. Rock, Curcumin content of turmeric and curry powders, *Nutr. Cancer* 55 (2) (2006) 126–131.
- [21] J.G. Devassy, I.D. Nwachukwu, P.J. Jones, Curcumin and cancer: barriers to obtaining a health claim, *Nutr. Rev.* 73 (3) (2015) 155–165.
- [22] Y. Panahi, A. Saadat, F. Beiraghdar, A. Sahebkar, Adjuvant therapy with bioavailability-boosted curcuminoids suppresses systemic inflammation and improves quality of life in patients with solid tumors: a randomized double-blind placebo-controlled trial, *Phytother. Res.* 28 (10) (2014) 1461–1467.
- [23] H. Mirzaei, G. Naseri, R. Rezaee, M. Mohammadi, Z. Banikazemi, H. Reza Mirzaei, H. Salehi, M. Peyvandi, J.M. Pawelek, A. Sahebkar, Curcumin: a new candidate for melanoma therapy? *Int. J. Cancer* (2016), <http://dx.doi.org/10.1002/ijc.30224>.
- [24] A.A. Momtazi, A. Sahebkar, Difluorinated curcumin: a promising curcumin analogue with improved anti-tumor activity and pharmacokinetic profile, *Curr. Pharm. Des.* (2016), <http://dx.doi.org/10.2174/1381612822666160527113501>.
- [25] A. Gilman, L.S. Goodman, J.G. Hardman, L.E. Limbird, Goodman & Gilman's the Pharmacological Basis of Therapeutics, McGraw-Hill, New York, NY, USA, 2001.
- [26] S. Devaraj, S. Leonard, M.G. Traber, I. Jialal, Gamma-tocopherol supplementation alone and in combination with alpha-tocopherol alters biomarkers of oxidative stress and inflammation in subjects with metabolic syndrome, *Free Radic. Biol. Med.* 44 (6) (2008) 1203–1208.
- [27] R.J. Simpson, A. Hammacher, D.K. Smith, J.M. Matthews, L.D. Ward, Interleukin-6: structure-function relationships, *Protein Sci.* 6 (5) (1997) 929–955.
- [28] Y.S. Devi, M. DeVine, J. DeKulper, S. Ferguson, A.T. Fazleabas, Inhibition of IL-6 signaling pathway by curcumin in uterine decidual cells, *PLoS One* (2015) e0125627.



- [29] F.Y. Chen, J. Zhou, N. Guo, W.G. Ma, X. Huang, H. Wang, Z.Y. Yuan, Curcumin retunes cholesterol transport homeostasis and inflammation response in M1 macrophage to prevent atherosclerosis, *Biochem. Biophys. Res. Commun.* 467 (2015) 872–878.
- [30] P.K. Kondamudi, H. Kovelamudi, P.G. Navak, M.C. Rao, R.R. Shenov, Curcumin half analog modulates interleukin-6 and tumor necrosis factor-alpha in inflammatory bowel disease, *Pharmacogn. Mag.* 11 (2015) S292–S302.
- [31] A.K. Singh, M. Vinayak, Curcumin attenuates CFA induced thermal hyperalgesia by modulation of antioxidant enzymes and down regulation of TNF-, IL-1 and IL-6, *Neurochem. Res.* 40 (2015) 463–472.
- [32] Y. Miao, S. Zhao, Y. Gao, R. Wang, Q. Wu, H. Wu, T. Luo, Curcumin pretreatment attenuates inflammation and mitochondrial dysfunction in experimental stroke: the possible role of Sirt1 signaling, *Brain Res. Bull.* 121 (2016) 9–15.
- [33] S. Morrone Mda, C.E. Schnorr, G.A. Behr, J. Gasparotto, R.C. Bortolin, K. da Boit Martinello, B. Saldanha Henkin, T.K. Rabello, A. Zanotto-Filho, D.P. Gelain, J.C. Moreira, Curcumin supplementation decreases intestinal adiposity accumulation, serum cholesterol alterations, and oxidative stress in ovariectomized rats, *Oxid. Med. Cell Longev.* 2016 (2016) 5719291.
- [34] D. Moher, A. Liberati, J. Tetzlaff, et al., Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement, *BMJ* 339 (2009) b2535.
- [35] J.P.T. Higgins, S. Green, (eds.), *Cochrane Handbook for Systematic Reviews of Interventions*. Version 5.0.2. London: The Cochrane Collaboration; 2011.
- [36] M. Borenstein, L. Hedges, J. Higgins, et al., *Comprehensive Meta-analysis Version 2*, Biostat, Englewood NJ, 2005.
- [37] M. Dinca, M.C. Serban, A. Sahebkar, D.P. Mikhailidis, P.P. Toth, S.S. Martin, M.J. Blaha, M. Blüher, C. Gurban, P. Penson, E.D. Michos, A.V. Hernandez, S.R. Jones, M. Banach, for Lipid Blood Pressure Meta-analysis Collaboration LBPMC Group, Does vitamin D supplementation alter plasma adipokines

concentrations? A systematic review and meta-analysis of randomized controlled trials, *Pharmacol. Res.* 107 (2016) 360–371.

[38] G. Ferretti, T. Bacchetti, A. Sahebkar, Effect of statin therapy on paraoxonase-1 status: a systematic review and meta-analysis of 25 clinical trials, *Prog. Lipid Res.* 60 (October) (2015) 50–73.

[39] S. Duval, R. Tweedie, Trim and fill: a simple funnel-plot-based method of testing and adjusting for publication bias in meta-analysis, *Biometrics* 56 (2000) 455–463.

[40] G. Belcaro, M.R. Cesarone, M. Dugall, L. Pellegrini, A. Ledda, M.G. Grossi, S. Togni, G. Appendino, Efficacy and safety of Meriva®: a curcumin-phosphatidylcholine complex, during extended administration in osteoarthritis patients, *Altern. Med. Rev.* 15 (4) (2010) 337–344.

[41] L.X. Na, B.L. Yan, S. Jiang, H.L. Cui, Y. Li, C.H. Sun, Curcuminoids target decreasing serum adipocyte-fatty acid binding protein levels in their glucose-lowering effect in patients with type 2 diabetes, *Biomed. Environ. Sci.* 27 (11) (2014) 902–906.

[42] P. Usharani, A.A. Mateen, M.U. Naidu, Y.S. Raju, N. Chandra, Effect of NCB-02, atorvastatin and placebo on endothelial function, oxidative stress and inflammatory markers in patients with type 2 diabetes mellitus: a randomized, parallel-group, placebo-controlled, 8-week study, *Drugs RD* 9 (4) (2008) 243–250.

[43] S. Ganjali, A. Sahebkar, E. Mahdipour, K. Jamialahmadi, S. Torabi, S. Akhlaghi, G. Ferns, S.M. Parizadeh, M. Ghayour-Mobarhan, Investigation of the effects of curcumin on serum cytokines in obese individuals: a randomized controlled trial, *Sci. World J.* 2014 (2014) 898361.

[44] Y. Panahi, A. Sahebkar, S. Parvin, A. Saadat, A randomized controlled trial on the anti-inflammatory effects of curcumin in patients with chronic sulphur mustard-induced cutaneous complications, *Ann. Clin. Biochem.* 49 (Pt 6) (2012) 580–588.

[45] Y. Panahi, M.S. Hosseini, N. Khalili, E. Naimi, L.E. Simental- Mendía, M. Majeed, A. Sahebkar, Effects of curcumin on serum cytokine concentrations in subjects with metabolic syndrome: a post-hoc analysis

of a randomized controlled trial, *Biomed. Pharmacother.* 05 (2016) 037, <http://dx.doi.org/10.1016/j.biopha.2016.05.037>.

[46] Y. Panahi, M. Ghanei, S. Bashiri, A. Hajhashemi, A. Sahebkar, Short-term curcuminoid supplementation for chronic pulmonary complications due to sulfur mustard intoxication: positive results of a randomized double-blind placebo-controlled trial, *Drug Res. (Stuttg.)* 65 (11) (2015) 567–573.

[47] A.R. Rahimnia, Y. Panahi, G. Alishiri, M. Sharafi, A. Sahebkar, Impact of supplementation with curcuminoids on systemic inflammation in patients with knee osteoarthritis: findings from a randomized double-blind placebo-controlled trial, *Drug Res. (Stuttg.)* 65 (10) (2015) 521–525.

[48] N. Chainani-Wu, E. Madden, F. Lozada-Nur, S. Silverman Jr., High-dose curcuminoids are efficacious in the reduction in symptoms and signs of oral lichen planus, *J. Am. Acad. Dermatol.* 66 (5) (2012) 752–760.

[49] S.C. Gupta, A.K. Tyagi, P. Deshmukh-Taskar, M. Hinojosa, S. Prasad, B.B. Aggarwal, Downregulation of tumor necrosis factor and other proinflammatory biomarkers by polyphenols, *Arch. Biochem. Biophys.* 559 (2014) 91–99.

[50] B.B. Aggarwal, S.C. Gupta, B. Sung, Curcumin: an orally bioavailable blocker of TNF and other pro-inflammatory biomarkers, *Br. J. Pharmacol.* 169 (August (8)) (2013) 1672–1692.

[51] E. Cretu, A. Trifan, A. Vasincu, A. Miron, Plant-derived anticancer agents—curcumin in cancer prevention and treatment, *Rev. Med. Chir. Soc. Med. Nat. Iasi* 116 (4) (2012) 1223–1229.

[52] J.W. Cho, K.S. Lee, C.W. Kim, Curcumin attenuates the expression of IL-1 $\beta$ , IL-6, and TNF- $\alpha$  as well as cyclin E in TNF- $\alpha$ -treated HaCaT cells; NF- $\kappa$ B and MAPKs as potential upstream targets, *Int. J. Mol. Med.* 19 (3) (2007) 469–474.

- [53] R.C. Srimal, B.N. Dhawan, Pharmacology of diferuloyl methane (curcumin): a non-steroidal anti-inflammatory agent, *J. Pharm. Pharmacol.* 25 (6) (1973) 447–452.
- [54] S.C. Gupta, S. Patchva, B.B. Aggarwal, Therapeutic roles of curcumin: lessons learned from clinical trials, *AAPS J.* 15 (1) (2013) 195–218.
- [55] E. Bettelli, Y. Carrier, W. Gao, T. Korn, T.B. Strom, M. Oukka, H.L. Weiner, V.K. Kuchroo, Reciprocal developmental pathways for the generation of pathogenic effector TH17 and regulatory T cells, *Nature* 441 (7090) (2006) 235–238.
- [56] J. Scheller, A. Chalaris, D. Schmidt-Arras, S. Rose-John, The pro- and anti-inflammatory properties of the cytokine interleukin-6, *Biochim. Biophys. Acta* 1813 (5) (2011) 878–888.
- [57] F. Schaper, S. Rose-John, Interleukin-6: Biology, signaling and strategies of blockade, *Cytokine Growth Factor Rev.* 26 (5) (2015) 475–487.
- [58] S. Kang, T. Tanaka, T. Kishimoto, Therapeutic uses of anti-interleukin-6 receptor antibody, *Int. Immunol.* 27 (1) (2015) 21–29.
- [59] H.Z. Ren, L.L. Ma, L.X. Wang, Effect of simvastatin on plasma interleukin-6 in patients with unstable angina, *Clin. Invest. Med.* 32 (4) (2009) E280–E284.
- [60] C. Bulcão, F.F. Ribeiro-Filho, A. Sanudo, ~ S.G. Roberta Ferreira, Effects of simvastatin and metformin on inflammation and insulin resistance in individuals with mild metabolic syndrome, *Am. J. Cardiovasc. Drugs* 7 (3) (2007) 219–224.
- [61] I. Ikonomidis, F. Andreotti, E. Economou, C. Stefanadis, P. Toutouzas, P. Nihoyannopoulos, Increased proinflammatory cytokines in patients with chronic stable angina and their reduction by aspirin, *Circulation* 100 (8) (1999) 793–798.

[62] N. Nishina, Y. Kaneko, H. Kameda, M. Kuwana, T. Takeuchi, Reduction of plasma IL-6 but not TNF- by methotrexate in patients with early rheumatoid arthritis: a potential biomarker for radiographic progression, *Clin. Rheumatol.* 32 (11) (2013) 1661–1666.

[63] Y. Hayashino, J.L. Jackson, T. Hirata, N. Fukumori, F. Nakamura, S. Fukuhara, S. Tsujii, H. Ishii, Effects of exercise on C-reactive protein, inflammatory cytokine and adipokine in patients with type 2 diabetes: a meta-analysis of randomized controlled trials, *Metabolism* 63 (3) (2014) 431–440.

[64] M.M. Yallapu, P.K. Nagesh, M. Jaggi, S.C. Chauhan, Therapeutic applications of curcumin nanoformulations, *AAPS J.* 17 (6) (2015) 1341–1356.

#### **FIGURE LEGENDS**


**Figure 1.** Flow chart of the number of studies identified and included into the meta-analysis.

**Figure 2.** Forest plot displaying weighted mean difference and 95% confidence intervals for the impact of curcuminoids on circulating IL-6 concentrations. Lower plot shows leave-one-out sensitivity analysis.


**Figure 3.** Forest plot displaying subgroup analyses for the impact of treatment dose and duration, and formulation of curcuminoids on the IL-6-lowering activity.

**Figure 4.** Meta-regression plots of the association between mean changes in circulating IL-6 concentrations with dose and duration of supplementation with curcuminoids. The size of each circle is inversely proportional to the size of the respective study.


**Figure 5.** Funnel plot detailing publication bias in the studies reporting the impact of curcuminoids on circulating IL-6 concentrations.




Study name	Difference in means	Standard error	Statistics for each study					p-Value
			Variance	Lower limit	Upper limit	Z-Value		
Ganjali et al., 2014a	-0.280	0.406	0.165	-1.075	0.515	-0.690	0.490	
Ganjali et al., 2014b	0.910	0.855	0.731	-0.766	2.586	1.064	0.287	
Rahimnia et al., 2014	0.190	0.262	0.069	-0.324	0.704	0.724	0.469	
Panahi et al., 2014	-0.750	0.084	0.007	-0.914	-0.586	-8.951	0.000	
Usharani et al., 2008	-1.970	0.510	0.260	-2.969	-0.971	-3.866	0.000	
Panahi et al., 2015	-1.950	0.305	0.093	-2.548	-1.352	-6.389	0.000	
Na et al., 2014	-1.170	0.387	0.150	-1.929	-0.411	-3.020	0.003	
Belcaro et al., 2010	-0.400	0.343	0.118	-1.073	0.273	-1.165	0.244	
Chainani-Wu et al., 2012	0.550	1.076	1.158	-1.559	2.659	0.511	0.609	
Panahi et al., 2012	0.035	0.072	0.005	-0.106	0.176	0.488	0.626	
	-0.597	0.234	0.055	-1.057	-0.138	-2.549	0.011	




Study name	Point	Standard error	Statistics with study removed					p-Value
			Variance	Lower limit	Upper limit	Z-Value		
Ganjali et al., 2014a	-0.632	0.252	0.063	-1.125	-0.139	-2.512	0.012	
Ganjali et al., 2014b	-0.675	0.240	0.058	-1.145	-0.205	-2.815	0.005	
Rahimnia et al., 2014	-0.705	0.255	0.065	-1.204	-0.205	-2.766	0.006	
Panahi et al., 2014	-0.562	0.304	0.093	-1.158	0.035	-1.844	0.065	
Usharani et al., 2008	-0.473	0.237	0.056	-0.937	-0.008	-1.993	0.046	
Panahi et al., 2015	-0.423	0.223	0.050	-0.861	0.014	-1.895	0.058	
Na et al., 2014	-0.532	0.247	0.061	-1.017	-0.047	-2.149	0.032	
Belcaro et al., 2010	-0.620	0.254	0.065	-1.119	-0.122	-2.440	0.015	
Chainani-Wu et al., 2012	-0.640	0.239	0.057	-1.109	-0.170	-2.672	0.008	
Panahi et al., 2012	-0.700	0.255	0.065	-1.200	-0.200	-2.745	0.006	
	-0.597	0.234	0.055	-1.057	-0.138	-2.549	0.011	




Group by	Study name	Statistics for each study							
		Difference in means	Standard error	Variance	Lower limit	Upper limit	Z-Value	p-Value	
Duration	< 8 weeks	Ganjali et al., 2014a	-0.280	0.406	0.165	-1.075	0.515	-0.690	0.490
	< 8 weeks	Ganjali et al., 2014b	0.910	0.855	0.731	-0.766	2.586	1.064	0.287
	< 8 weeks	Rahimnia et al., 2014	0.190	0.262	0.069	-0.324	0.704	0.724	0.469
	< 8 weeks	Panahi et al., 2014	-0.750	0.084	0.007	-0.914	-0.586	-8.951	0.000
	< 8 weeks	Chainani-Wu et al., 2012	0.550	1.076	1.158	-1.559	2.659	0.511	0.609
	< 8 weeks	Panahi et al., 2012	0.035	0.072	0.005	-0.106	0.176	0.488	0.626
	< 8 weeks		-0.099	0.262	0.068	-0.612	0.414	-0.379	0.705
	>= 8 weeks	Usharani et al., 2008	-1.970	0.510	0.260	-2.969	-0.971	-3.866	0.000
	>= 8 weeks	Panahi et al., 2015	-1.950	0.305	0.093	-2.548	-1.352	-6.389	0.000
	>= 8 weeks	Na et al., 2014	-1.170	0.387	0.150	-1.929	-0.411	-3.020	0.003
	>= 8 weeks	Belcaro et al., 2010	-0.400	0.343	0.118	-1.073	0.273	-1.165	0.244
	>= 8 weeks		-1.352	0.395	0.156	-2.127	-0.576	-3.417	0.001
	Overall		-0.481	0.218	0.048	-0.908	-0.053	-2.202	0.028





Group by	Study name	Statistics for each study							
		Difference in means	Standard error	Variance	Lower limit	Upper limit	Z-Value	p-Value	
Dose	<1000 mg/day	Usharani et al., 2008	-1.970	0.510	0.260	-2.969	-0.971	-3.866	0.000
	<1000 mg/day	Na et al., 2014	-1.170	0.387	0.150	-1.929	-0.411	-3.020	0.003
	<1000 mg/day	Belcaro et al., 2010	-0.400	0.343	0.118	-1.073	0.273	-1.165	0.244
	<1000 mg/day		-1.118	0.437	0.191	-1.974	-0.262	-2.560	0.010
	>=1000 mg/day	Ganjali et al., 2014a	-0.280	0.406	0.165	-1.075	0.515	-0.690	0.490
	>=1000 mg/day	Ganjali et al., 2014b	0.910	0.855	0.731	-0.766	2.586	1.064	0.287
	>=1000 mg/day	Rahimnia et al., 2014	0.190	0.262	0.069	-0.324	0.704	0.724	0.469
	>=1000 mg/day	Panahi et al., 2014	-0.750	0.084	0.007	-0.914	-0.586	-8.951	0.000
	>=1000 mg/day	Panahi et al., 2015	-1.950	0.305	0.093	-2.548	-1.352	-6.389	0.000
	>=1000 mg/day	Chainani-Wu et al., 2012	0.550	1.076	1.158	-1.559	2.659	0.511	0.609
	>=1000 mg/day	Panahi et al., 2012	0.035	0.072	0.005	-0.106	0.176	0.488	0.626
	>=1000 mg/day		-0.379	0.277	0.077	-0.922	0.164	-1.368	0.171
	Overall		-0.591	0.234	0.055	-1.050	-0.132	-2.526	0.012




Group by	Study name	Statistics for each study							
		Difference in means	Standard error	Variance	Lower limit	Upper limit	Z-Value	p-Value	
Formulation	Formulated	Ganjali et al., 2014a	-0.280	0.406	0.165	-1.075	0.515	-0.690	0.490
	Formulated	Ganjali et al., 2014b	0.910	0.855	0.731	-0.766	2.586	1.064	0.287
	Formulated	Rahimnia et al., 2014	0.190	0.262	0.069	-0.324	0.704	0.724	0.469
	Formulated	Panahi et al., 2014	-0.750	0.084	0.007	-0.914	-0.586	-8.951	0.000
	Formulated	Panahi et al., 2015	-1.950	0.305	0.093	-2.548	-1.352	-6.389	0.000
	Formulated	Belcaro et al., 2010	-0.400	0.343	0.118	-1.073	0.273	-1.165	0.244
	Formulated	Panahi et al., 2012	0.035	0.072	0.005	-0.106	0.176	0.488	0.626
	Formulated		-0.426	0.257	0.066	-0.930	0.079	-1.654	0.098
	Unformulated	Usharani et al., 2008	-1.970	0.510	0.260	-2.969	-0.971	-3.866	0.000
	Unformulated	Na et al., 2014	-1.170	0.387	0.150	-1.929	-0.411	-3.020	0.003
	Unformulated	Chainani-Wu et al., 2012	0.550	1.076	1.158	-1.559	2.659	0.511	0.609
	Unformulated		-1.184	0.523	0.273	-2.209	-0.159	-2.264	0.024
	Overall		-0.574	0.231	0.053	-1.026	-0.121	-2.483	0.013









## **TABLES**

**Table 1.** Demographic characteristics of the included studies.

**Table 2.** Risk of bias assessment in the studies considered for meta-analysis.

Author	Study design	Target Population	Treatment duration	n	Study groups	Age, years	Female (n, %)	BMI, (kg/m <sup>2</sup> )	Systolic blood pressure (mmHg)	Diastolic blood pressure (mmHg)	C-reactive (mg)
Belcaro et al. (2010)	Open-label	Osteoarthritis	8 months	50	Curcumin 200 mg/day	43.6±5.5	27 (54.0)	ND	ND	ND	ND
				50	Control	44.2±6	22 (44.0)	ND	ND	ND	ND
Chainani-Wu et al. (2012)	Randomized, double-blind, placebo-controlled	Oral lichen planus	2 weeks	10	Curcumin 6 g/day	60.8±8.6	8 (80.0)	ND	ND	ND	2.5 (1.0-)
				10	Placebo	56.2±11.7	5 (50.0)	ND	ND	ND	2.0 (1.0-)
Ganjali et al. (2014)	Randomized, double-blind, placebo-controlled, cross-over	Obesity	1 month	15	Curcumin 1 g/day	ND	ND	ND	ND	ND	ND
				15	Placebo	ND	ND	ND	ND	ND	ND
Na et al. (2014)	Randomized, double-blind, placebo-controlled	Type 2 diabetes	3 months	50	Curcumin 300 mg/day	ND	ND	ND	ND	ND	2.71±
				50	Placebo	ND	ND	ND	ND	ND	2.68±
Panahi et al. (2015)	Randomized, double-blind, placebo-controlled	Sulfur mustard intoxication	4 weeks	39	Curcumin 1.5 g/day	50.9±7.2	0 (0.0)	28.0±4.8	ND	ND	6.98±
				39	Placebo	53.9±8.6	0 (0.0)	25.9±4.0	ND	ND	8.54±
Rahimnia et al. (2015)	Randomized, double-blind, placebo-controlled	Knee osteoarthritis	6 weeks	19	Curcumin 1.5 g/day	57.3±8.7	14 (73.6)	28.7±3.1	ND	ND	5.56±
				21	Placebo	57.5±9.0	17 (80.9)	29.6±4.4	ND	ND	5.00±
Panahi et al. (2012)	Randomized, double-blind, placebo-controlled	Chronic pruritic skin lesions	4 weeks	40	Curcumin 1 g/day	47.5±10.7	0 (0.0)	ND	ND	ND	2.58±
				40	Placebo	48.3±8.5	0 (0.0)	ND	ND	ND	2.53±
Sahebkar et al. (2015)	Randomized, double-blind, placebo-controlled	Metabolic syndrome	8 weeks	59	Curcumin 1 g/day	44.8±8.6	23 (46.0)	25.4±2.4	135.5±13.1	88.3±7.8	6.52±
				58	Placebo	43.4±9.7	27 (54.0)	22.8±5.3	135.7±14.7	88.7±8.1	7.10±
Usharani et al. (2008)	Randomized, placebo-controlled	Type 2 diabetes	8 weeks	23	Curcumin 300 mg/day	55.5±10.7	11 (47.8)	24.6±2.4	130.4±18.5	81.8±10.0	ND
				21	Placebo	49.7±8.1	10 (47.6)	23.9±2.3	126.3±15.4	80.7±7.4	ND

Values are expressed as mean ± SD

\*Distribution unspecified

\*\*Median (IQR)

Abbreviations: ND, no data; BMI, body mass index; IQR, interquartile range.

Study	Random sequence generation	Allocation concealment	Selective reporting	Other bias	Blinding of participants and personnel	Blinding of outcome assessment	Incomplete outcome data
Belcaro et al. (2010)	H	H	L	U	H	U	L
Chainani-Wu et al. (2012)	L	L	L	L	L	U	L
Ganjali et al. (2014)	U	U	L	U	U	U	L
Na et al. (2014)	U	U	L	U	U	U	L
Panahi et al. (2015)	L	L	L	L	L	L	L
Rahimnia et al. (2015)	U	U	L	L	U	U	L
Panahi et al. (2012)	U	U	L	U	U	U	L
Sahebkar et al. (2015)	U	U	L	U	U	U	L
Usharani et al. (2008)	U	U	L	U	H	U	L

L, low risk of bias; H, high risk of bias; U, unclear risk of bias.