

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Fossil marine fishes and the 'Lago Mare' event: Has the Mediterranean ever transformed into a brackish lake?

This is a pre print version of the following article:

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1661162> since 2018-03-02T13:27:51Z

Published version:

DOI:10.1127/nos/2016/0343

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

Fossil marine fishes and the ‘Lago Mare’ event:

Has the Mediterranean ever transformed into a brackish lake?

Giorgio Carnevale^{1*}, Francesco Dela Pierre¹, Marcello Natalicchio², and Walter Landini³

¹ Dipartimento di Scienze della Terra, Università degli Studi di Torino, Via Valperga Caluso, 35
10125 Torino, Italia

² Institut für Geologie, Universität Hamburg, Bundesstraße 55, 20146 Hamburg, Germany

³ Dipartimento di Scienze della Terra, Università di Pisa, Via Santa Maria, 53 56126 Pisa, Italia

* Corresponding author; E-mail: giorgio.carnevale@unito.it

Abstract. According to current hypotheses of Neogene paleoenvironmental evolution of the Mediterranean, during the terminal phase of the Messinian salinity crisis about 5.5 Ma, the entire basin was characterized by the development of non-marine paleobiotopes inhabited by brackish ostracods, mollusks and dinocysts. Such a scenario, commonly known as the ‘Lago Mare’ event, is considered to be the product of the combined effects of humid climate conditions and a remarkably modified drainage pattern with the capture of Paratethyan brackish waters. Some recent studies, however, have suggested that such a scenario represents an oversimplification of the original paleogeographical and paleoenvironmental context. In this paper, we report the ‘Lago Mare’ fish assemblages from seven Italian localities. These fish assemblages are dominated by taxa characterized by a strict marine affinity, thereby suggesting that the Mediterranean was filled by marine waters. Fossil fishes provide unambiguous evidence of the presence of normal marine conditions in the Mediterranean before the Messinian-Zanclean boundary, also demonstrating that a

new paleoenvironmental interpretation of the ‘Lago Mare’ event is needed. The fossil record of the ‘Lago Mare’ event and the paleoecological relevance of fish remains for the interpretation of the nature of the Mediterranean water masses are also discussed.

Keywords. Fossil fishes, Messinian salinity crisis, ‘Lago mare’, Mediterranean, paleoenvironment

Running head: Fossil marine fishes and the ‘Lago Mare’ event

1. Introduction

The events related to the Messinian salinity crisis are among the most intriguing and extensively explored of Cenozoic history. The spectacular scenario of a desiccated Mediterranean basin eventually filled by a gigantic catastrophic flood received considerable attention and publicity (see Hsü 1984) and is currently regarded as a well-documented event of Earth history (e.g., Stanley 1989). As a consequence, a remarkable amount of multidisciplinary research has been produced in the last 40 years, stimulating cogent and often conflicting debates extensively discussed in several hundreds of scientific papers (see Roveri et al. 2014).

Extraordinary advancements have been made in our knowledge of Neogene cyclo-, bio-, and magneto-stratigraphy in the Mediterranean region. The refinement of a detailed astronomically tuned time scale has resulted in, among other things in the establishment of the age boundaries for the Messinian stage (Hilgen et al. 2000, Van Couvering et al. 2000), and this has favoured high resolution correlations and paleogeographic reconstructions (see, e.g., Hilgen et al. 1999, Krijgsman 2002, Roveri et al. 2014). More recently, an evolutionary stratigraphic model for the Messinian salinity crisis has been proposed (CIESM 2008), primarily to characterize the various evaporite facies. This model consists of three stages, each defined by peculiar evaporite associations (e.g., Roveri et al. 2009, 2014, Manzi et al. 2012, 2013). Stage 1 (5.97-5.60 Ma) documents the onset of the Messinian salinity crisis and the deposition of primary sulfate evaporites in shallow water

marginal basins (Primary Lower Gypsum). During stage 2 (5.60-5.55 Ma), the salinity crisis reached its acme, as evidenced by the massive accumulation of primary halite and clastic gypsum deposits (Resedimented Lower Gypsum). Finally, stage 3 (5.55-5.33 Ma) is characterized by a totally different scenario and is represented by the so-called Upper Evaporites (selenite and cumulate gypsum facies; Upper Gypsum), deposited in shallow water basins in the southern and eastern sectors of the Mediterranean, whereas clastic deposits accumulated in the northern and western Mediterranean sectors. The evaporite-free clastic successions of stage 3 comprise two distinct sequences that originated in response to different climate conditions and precipitation regimes (e.g., Roveri et al. 2008), allowing a subdivision into two sub-stages, 3.1 (5.55-5.42 Ma) and 3.2 (5.42-5.33 Ma). The end of the Messinian salinity crisis appears to be marked by the nearly instantaneous return to fully stable marine conditions in response to the Zanclean deluge (or Zanclean flooding), which was produced by the abrupt collapse of the Gibraltar sill and the development of an enormous rapid (see Garcia-Castellanos et al. 2009) or gigantic waterfall (see McKenzie 1999).

The final phase of the Messinian salinity crisis defined by stage 3 (particularly sub-stage 3.2) approximately corresponds to the so-called 'Lago Mare' event. Due to the absence of an efficient biostratigraphic resolution and the highly heterogeneous nature of the sedimentary products, the Mediterranean paleogeographical and paleoenvironmental context during stage 3 is poorly defined (e.g., Roveri and Manzi 2006). In particular, a comprehensive reconstruction of the 'Lago Mare' event remains elusive, and its overall setting at the Mediterranean scale has been interpreted based on its peculiar paleontological record, primarily represented by a recurrent assemblage of oligo-mesohaline molluscs and ostracods traditionally considered of Paratethyan affinity (see Orszag-Sperber 2006). Current interpretations postulate that during stage 3 of the Messinian salinity crisis the Mediterranean was characterized by non-marine sedimentation in a series of mysterious interconnected lacustrine-brackish basins. The widespread occurrence of 'Lago Mare' deposits appears to be related to the synergistic effect of humid climatic conditions (e.g.,

Cosentino et al. 2005) and a strongly modified Mediterranean drainage pattern with the capture of the Paratethyan brackish waters through a canyon system located somewhere in the Aegean region (e.g., McCulloch and De Deckker 1989, Orszag-Sperber et al. 2000).

The goal of this paper is to provide a comprehensive overview of the fossil fish remains, mostly otoliths, collected in the ‘Lago Mare’ deposits from seven Italian localities. The data presented and discussed herein provide robust evidence of the presence of normal marine waters during most of stage 3 of the Messinian salinity crisis. The paleontological record of the ‘Lago Mare’ event and the importance of fish remains for the interpretation of the Mediterranean water body during the salinity crisis are discussed.

2. Localities and stratigraphy

In the last decade, extensive exploration of upper Messinian deposits in Italy led to the discovery of at least seven productive localities for fossil fish remains (Fig. 1), Ciabot Cagna, Moncucco Torinese and Verduno in the Tertiary Piedmont Basin, Cava Serredi and Podere Torricella in the Neogene hinterland basins of Tuscany, Capanne di Bronzo in the Marche Apennine, and, finally, Le Vicenne in the Abruzzo Apennine. Two or more productive fish-bearing layers were recognized in the stratigraphic successions of some of these localities, including Cava Serredi, Moncucco Torinese and Verduno (Carnevale et al. 2006b, 2008a, Colombero et al. 2014, submitted). A minimum quantity of 200 kg of fossiliferous sediments for each productive layer has been sampled and processed for the extraction of fish remains. However, greater amounts of deposits were sampled from some of the productive layers at Cava Serredi and Moncucco Torinese, leading to the sieving of about 3 and 20 tons of sediment, respectively. A brief summary of the stratigraphic features of each of the investigated localities is presented below.

2.1 Tertiary Piedmont Basin

The Tertiary Piedmont Basin is an episutural basin filled with a thick succession of Eocene to Messinian deposits covering the Alps-Apennine junction (Mosca et al. 2010). Moncucco Torinese is located in the Torino Hill, whereas Ciabot Cagna and Verduno pertain to the Langhe domain. In these sectors of the Tertiary Piedmont basin, the Messinian succession is particularly well exposed (e.g., Dela Pierre et al., 2007, 2011, 2015, Natalicchio et al. 2014) and consists of the pre-evaporitic Sant'Agata Fossili marls, followed by the Primary Lower Gypsum unit, which formed during the first stage of the Messinian salinity crisis (between 5.97 and 5.60 Ma; see Roveri et al. 2014). The Valle Versa Chaotic Complex (5.60-5.55 Ma) unconformably overlies the Primary Lower Gypsum unit, and is in turn capped by the fluvio-lacustrine and paralic sediments of the Conglomerati di Cassano Spinola. This latter lithostratigraphic unit primarily consists of clays, silts, sands and gravels comprising both sub-stages 3.1 and 3.2 of the Messinian salinity crisis.

At least four vertebrate-bearing layers pertaining to sub-stage 3.1 (between 5.55 and 5.42 Ma) were recognized at Verduno (Fig. 2), in the so-called Pollenzo section (see Colombero et al. 2014, Dela Pierre et al. 2014). Fish remains (disarticulated bones and otoliths) occur in two of these layers. The vertebrate-bearing layers of Verduno accumulated in a variety of subaqueous paleobiotopes, including a brackish coastal lagoon characterized by thalassogenic waters (Colombero et al. 2014).

The section of Ciabot Cagna is located near the town of Corneliano d'Alba and includes about eight meters of weakly laminated clays referred to the Lago-Mare event based on their paleontological content (Cavallo and Repetto 1988, Cavallo et al. 1993, Caputo 2005, Esu and Popov 2012), particularly the mollusc assemblage (e.g., *Dreissena*, lymnocypridines, *Melanopsis*). According to Cavallo and Repetto (1988), these fossiliferous sediments were deposited in a heterogeneous deltaic-lagoonal paleobiotope.

Six layers containing a diverse assemblage of fish and continental tetrapod remains were recognized in the deposits of sub-stage 3.2 of Moncucco Torinese, stratigraphically constrained between 5.41 and 5.33 Ma (see Alba et al. 2014, Colombero et al. submitted; Fig. 2), thereby being

associated to the Lago-Mare event. The fossiliferous deposits of Moncucco Torinese originated in a complex and heterogeneous paralic biotope in which continental vertebrate remains accumulated from a wide spectrum of terrestrial habitats (Colombero et al. submitted).

2.2 Hinterland basins of Tuscany

The Neogene hinterland basin of Tuscany are located on the Tyrrhenian side of the northern Apennines; their origin is related to the tectonic evolution of the Tyrrhenian-Apennine system. Both extensional and compressive tectonic regimes have been proposed to explain the origin of these Neogene sedimentary depressions (e.g., Patacca et al. 1990, Martini and Sagri 1993, Boccaletti and Sani 1998). The sedimentary fills of these basins, also known as ‘neoautochthonous successions’ consist of Tortonian continental deposits covered by Messinian to Pleistocene brackish and marine sediments (see Bossio et al. 1993). The localities discussed herein, Cava Serredi and Podere Torricella, are located on the western side of the Mid-Tuscan Metamorphic ridge, in the Fine and Volterra basins, respectively.

Due to extensive quarrying, a remarkable exposure of Messinian ‘Lago Mare’ deposits occurs at Cava Serredi, one of the classic Italian localities for the study of the Mio-Pliocene boundary (e.g., Bossio et al. 1981). This upper Messinian succession of Cava Serredi consists of two depositional sequences consisting of marls and clays and documenting a variety of paralic depositional environments (see Sarti et al. 1995, Carnevale et al. 2006b). Fish remains have been recovered from ten fossiliferous layers, two of which are exclusively characterized by articulated skeletons (Carnevale et al. 2000, 2006b, 2008a). Based on the general architecture of the succession and the associated ostracod assemblages, the two basal fossiliferous layers possibly pertain to sub-stage 3.1 of the Messinian salinity crisis, whereas the remaining eight can be unquestionably referred to the sub-stage 3.2 (see Sarti 1995, Caputo 2008). Therefore, the age of the fish-bearing horizons can be constrained between 5.45 and 5.33 Ma (see Caputo 2008) (Fig. 2).

Podere Torricella is located in the Spicchiaiola-Pignano area in the south-eastern sector of the Volterra Basin. The ‘Lago-Mare’ event is recorded in the Volterra Basin by the upper portion of the ‘Argille e Gessi del Fiume Era Morta’ lithostratigraphic unit (Bossio et al. 1996, 1997). At Podere Torricella, ‘Lago-Mare’ deposits are well exposed in a small quarry exploited for the extraction of alabastrine gypsum where a fossiliferous layer characterized by abundant molluscs and ostracods associated with exquisitely preserved otoliths occur about 20 m above the uppermost alabastrine layer. Based on stratigraphic and paleontological considerations, the age of the fossiliferous horizon can be constrained between 5.40 and 5.33 Ma (Bossio et al. 1996, 1997, Carnevale et al. 2008a) (Fig. 2).

2.3 Capanne di Bronzo

The locality of Capanne di Bronzo is located in the northern sector of the Montecalvo in Foglia Syncline, Marche Apennine. The Montecalvo in Foglia Syncline is one of the satellite basins with NW-SE direction running parallel to the chain on the Adriatic side of the Apennines. Upper Messinian deposits are extensively exposed in the core of the Montecalvo in Foglia Syncline, represented by two terrigenous lithostratigraphic units corresponding to the Terreni di Tetto and Colombacci Formation, respectively (Roveri et al. 1998, 2001, Bassetti 2000). The upper Messinian Colombacci Formation records the ‘Lago-Mare’ event in the Apennine foredeep, and is characterized by a cyclical architecture superimposed to an overall transgressive trend (Bassetti et al. 2004). The Colombacci Formation is well exposed in the Capanne di Bronzo section (Carloni et al. 1974, Borsetti et al. 1975), where a rich fossiliferous horizon occurs about 1 m below the Mio-Pliocene boundary. The fossil assemblage accumulated in a shallow marine environment remarkably influenced by continental outflow (see Bellagamba 1978, Carnevale et al. 2006a) and consists of abundant dreissenids, lymnocypridines, hydrobiids and melanopsids associated with fish otoliths, ostracods (*Cyprideis*, loxoconchids) and benthic foraminiferans (*Ammonia*). According to

Carnevale et al. (2006a), the age of the fossiliferous horizon can be constrained between 5.35 and 5.33 Ma (Fig. 2).

2.4 Le Vicenne

The Le Vicenne basin is located in the Marsica region in the Abruzzo Apennine (e.g., Colacicchi et al. 1967). It is one of the Neogene thrust-top basins developed during the eastward migration of the central Apennine thrust belt (Cipollari et al. 1999). The upper Messinian deposits consist of less than 50 m of conglomerates, sandy marls and fossiliferous clays and siltites. The fossiliferous deposits contain abundant molluscs and ostracods, and relatively uncommon otoliths. According to Cipollari et al. (1999), these deposits are associated with the 'Lago Mare' event and originated in a mesohaline coastal lagoon during the second post-evaporitic interval (sub-stage 3.2) of the Messinian salinity crisis (see also Gliozzi 1999) (Fig. 2).

3. Fossil fish remains: diversity and paleoecology

Overall, several thousands of fish specimens were collected from the fossiliferous deposits pertaining to sub-stage 3.2 in six Italian localities, and less abundant material was extracted from the sediments of sub-stage 3.1 at Verduno and from the two basal fossiliferous layers of Cava Serredi. Therefore, limited paleontological information is available for sub-stage 3.1 of the Messinian salinity crisis, whereas the scenario emerging from the ichthyofaunistic data of sub-stage 3.2 is much more heterogeneous and complex. The fossils from this last phase consist in large part of otoliths, but articulated skeletal remains were also collected at Cava Serredi (Carnevale et al. 2000, 2006b, 2008a) (Fig. 3). In fact, not less than 49 taxa belonging to 15 teleostean family-level groups were identified based on material collected from the seven localities documented herein (see Tab. 1). Some of these taxa, particularly some of the gobies (*Gobius* aff. *multipinnatus*, *Gobius* aff. *niger*, *Gobius* aff. *truncatus*, *Gobius* sp., "*Gobiidarum*" aff. *weileri*) and lanternfishes (*Diaphus befralai*, *Diaphus splendidus*, *Diaphus taaningi*, *Diaphus* sp.), were recognized in two or more

localities; however, most of the fish taxa are exclusively known from a single locality (Tab. 1). This scattered distributional pattern seems to suggest that only a limited portion of the original fish diversity has been preserved in the upper Messinian sedimentary record.

Quantitatively, benthic fishes belonging to the Gobiidae and/or Lophiiformes are strikingly abundant in the productive layers of Cava Serredi (see Carnevale et al. 2006b, 2008a), and are also dominant at Le Vicenne. The oceanic lanternfishes of the family Myctophidae are dominant from a quantitative point of view in two localities, Ciabot Cagna and Podere Torricella (see Caputo 2005, Carnevale et al. 2008a), and croackers of the family Sciaenidae are dominant in the localities of Capanne di Bronzo and Moncucco Torinese (Carnevale et al. 2006a, Colombero et al. submitted). Four families - Clupeidae, Gobiidae, Myctophidae and Sciaenidae - comprise 35 taxa, representing more than 70% of the recognized fish diversity. In particular, the families Myctophidae and Gobiidae are represented in nearly all the localities documenting sub-stage 3.2 of the Messinian salinity crisis with 16 and 10 species-level taxa, respectively. Myctophids are highly diverse at Moncucco Torinese, being represented by nine taxa, whereas seven gobiid species-level taxa were reported at Cava Serredi.

In order to properly understand the paleoenvironmental significance of the fish assemblages, each of the identified taxa has been assigned to an ecological guild (Tab. 1). Four guilds have been recognized, in part following the classification proposed by Elliott and Dewailly (1995) and Mathieson et al. (2000). These are diadromous migrants (DM), estuarine residents (ER), marine coastal (MC) and marine oceanic (MO). The guild of diadromous migrants includes those taxa that pass between salt and fresh waters (via brackish biotopes) for feeding and spawning (solely represented by the sprat *Clupeonella* sp.; Fig. 3B). Estuarine residents are those taxa that spend their entire life-cycles in brackish biotopes characterized by thalassogenic waters (croackers, gobies, grey mullets, pupfishes, silversides). Marine coastal taxa include those marine steno- and euryhaline shallow water fish that occupy the inner shelf and appear only irregularly in brackish biotopes (anglerfishes, blennies, croackers, gobies, herrings, mullets, porgies, toadfishes). Finally, the guild

of marine oceanic taxa includes the pelagic (primarily mesopelagic) species that inhabit the continental slope or open oceanic waters and only occasionally occur on the continental shelf (cusk-eels, lanternfishes, moras, silvery pouts, silver roughies). Overall, the relative proportion of the represented ecological guilds is rather heterogeneous and exhibits considerable variation between the examined localities, except for the diadromous migrants, which are restricted to Cava Serredi. Estuarine residents and marine coastal taxa are equally abundant at Capanne di Bronzo, Ciabot Cagna and Verduno. Estuarine residents clearly dominate the fish assemblage at Le Vicenne, whereas these are slightly subordinate to marine coastal taxa in the Cava Serredi assemblage. Finally, it is interesting to note that marine oceanic taxa are dominant in the assemblages of Moncucco Torinese and Podere Torricella, representing 71% and 44% of the fish diversity, respectively. Taking into account taxonomic diversity, marine oceanic taxa are clearly dominant, being represented by 38% of the recognized taxa, followed by marine coastal taxa (31%), estuarine residents (29%), and diadromous migrants (2%). Therefore, taxa exhibiting a strict marine affinity represent more than two thirds (69%) of the entire fish diversity (see Fig. 4).

The remarkable diversity of marine taxa, particularly those that are oceanic, is apparently difficult to explain considering that the otolith-bearing deposits originated in a variety of paralic paleobiotopes during stage 3 of the Messinian salinity crisis (Cavallo and Repetto 1988, Cipollari et al. 1999, Carnevale et al. 2006a, 2006b, 2008a, Caputo 2007, Colombero et al. 2014, submitted). Because of their excellent preservation (Fig. 3), with fine and delicate structures clearly recognizable, it is evident that the otoliths belonging to marine taxa are the product of autochthonous embedding, thereby excluding any possibility of reworking from older rocks. The stratigraphic range of the fish taxa recognized in stage 3 deposits spans from the Tortonian or early Messinian to Zanclean (Tab. 2). Overall, the fish assemblage consists of a mix of Late Miocene, Tortonian-Zanclean, and exclusively Zanclean taxa (e.g., Brzobohaty and Nolf 2000, Girone et al. 2010, Lin et al. 2015). In particular, most of the marine fish taxa (76%) are primarily known from

Zanclean deposits (see Tab. 2), providing additional evidence excluding the possibility of reworking from older rocks.

Marine coastal taxa often represent one of the most diverse groups in paralic biotopes, even if usually represented by a few individuals that scarcely contribute to the standing crop of these settings (e.g., Rebelo 1992, Nordlie 2003, Veiga et al. 2006). As discussed above, most of the marine oceanic fishes recognized in the upper Messinian deposits are mesopelagic. Mesopelagic fishes can be defined as species spending the day at depths between approximately 200 and 1000 meters, and performing diel migrations to reach the upper 200 meters or even the surface during the night (e.g., Gjøsæter and Kawaguchi 1980). Mesopelagic taxa, particularly lanternfishes of the family Myctophidae, constitute a major part of the biomass in open sea areas (e.g., Ahlstrom 1969, Clarke 1973), representing a significant component in the diet of predatory fishes (tunas, swordfishes, jacks, barracudas, scabbardfishes). The diversity (and abundance) of marine oceanic fishes in paralic settings can be easily explained based on the taphonomic processes that usually resulted in the formation of fossil otolith assemblages. As discussed by Nolf (1985), the predatory activity of large fishes and other vertebrates remarkably contributes to the formation of otolith taphocoenoses. According to Schäfer (1966), the excreta of large predatory fishes foraging in oceanic environments are usually very rich in otoliths, whereas the bones are nearly entirely digested. As a matter of fact, most otoliths enter the sediment through predators (Nolf 2013). Therefore, through the excreta of large predatory fishes, otoliths of mesopelagic taxa are often transported in shallower waters and paralic biotopes, in which predatory fishes periodically occur, attracted by the abundant food and habitat availability (Elliott and Dewailly 1995, Nordlie 2003). Unfortunately, a direct connection with large predatory fishes can be only occasionally established using the fossil record since their otoliths are very thin and delicate and only very occasionally occur in fossil assemblages (Nolf 2013) and the preservation of skeletal remains requires peculiar taphonomic conditions (e.g., Carnevale et al. 2008b).

The migratory activity of predatory fishes and its consequent taphonomic product largely impact on the paleogeographic and physiographic interpretation of the depositional environments. The use of paralic biotopes by means of large predatory fishes moving from oceanic waters is clearly related to the existence of permanently open connections between the open sea and the paralic settings. The presence of a direct connection between the paralic paleobiotopes in which the fossiliferous deposits originated and fully marine waters is also suggested by the co-occurrence of marine coastal taxa and diadromous migrants. Because of their mobility and migratory behaviour, fishes are therefore very useful for large-scale interpretations, providing data about the variety of biotopes contiguous to the depositional environment.

4. Discussion

4.1 Fossil fishes and the ‘Lago Mare’ conundrum

The fossil record was the primary source of information that supported the emergence of the ‘Lago Mare’ concept. The benthic assemblages primarily consisting of ostracods and molluscs have been used to postulate a brackish ‘Lago Mare’ event since the earliest attempts to define the paleoenvironmental context of the Mediterranean at the end of the Messinian (e.g., Ruggieri 1962, Ruggieri and Greco 1965). However, the traditional ‘Lago Mare’ concept, which hypothesizes a partial refill of a desiccated Mediterranean with fresh- and brackish waters of Paratethyan origin, was developed in the late seventies as a consequence of the extensive geo-paleontological exploration of the Messinian deposits throughout the entire Mediterranean basin (e.g., Cita et al. 1978). The relevance of ostracods and molluscs in the interpretation of the uppermost Messinian paleoenvironments was certainly emphasized by the apparent absence of other kinds of fossils in latest Messinian deposits and their ostensible Paratethyan biogeographic affinity remarkably contributed to the development of the concept. In particular, the ostracod assemblages are regarded

as a useful chronostratigraphic tool for the analysis of the deposits of stage 3 of the Messinian salinity crisis (e.g., Cipollari et al. 1999; Grossi et al. 2011), a time interval corresponding to the upper part of the so-called “Non-Distinctive Zone” and “Salinity Crisis interval” of the planktonic foraminifer and calcareous nannofossil biozonations, respectively (e.g., Iaccarino 1985, Raffi et al. 2003). In this context, the absence of marine calcareous plankton communities has been paradigmatically regarded as the actual record of the absence of marine communities and of the existence of non marine conditions in the Mediterranean basin. Moreover, additional paleontological (dinocysts; see Bertini et al. 1995) and geochemical (depleted Sr isotope values; see McCulloch and De Dekker 1989) evidence was used subsequently to consolidate the model. Despite a remarkable number of inconsistencies that emerged in the last three decades, the actual nature of the ‘Lago Mare’ event remains poorly understood and there is still controversy about the environmental dynamics of the Mediterranean basin during the terminal phase of the Messinian salinity crisis.

The integrative analysis of the ecological affinities of the fish taxa unquestionably indicates that the paralic depositional environments in which fossil fish remains accumulated during stage 3 of the Messinian salinity crisis were physically connected with a Mediterranean basin characterized by normal marine conditions. Moreover, the remarkable ecological disparity and taxonomic diversity of the fish assemblages suggest that the Mediterranean marine ecosystems were rather complex during the late Messinian. Finally, the age of the fossiliferous deposits reveal that normal marine waters characterized the Mediterranean for most of sub-stage 3.1 and the entire sub-stage 3.2, thereby implying that the marine refilling of the basin occurred well before the Messinian-Zanclean boundary. In fact, the ecological affinities and taxonomic composition of the ichthyofaunistic assemblage discussed in this study are totally opposite to those expected to be found in ‘Lago Mare’ deposits, which, according to the traditional model should exclusively contain the remains of freshwater and/or brackish fishes.

The picture emerging from the analysis of fossil fish remains is therefore incompatible with the traditional environmental model hypothesized for the ‘Lago Mare’ event, at least as far as concerns the Sr isotope data and the peculiar assemblages of ostracods, molluscs and dinocysts.

$\text{Sr}^{87}/\text{Sr}^{86}$ values from ‘Lago Mare’ deposits and benthic fossils are usually lower than normal oceanic values estimated from that time interval. However, in the comprehensive evaluation of these isotopic data, it should be taken into account that the contribution of low $\text{Sr}^{87}/\text{Sr}^{86}$ river waters can be emphasized by the overall physiography of the Mediterranean, especially in the case of severely restricted connections with the Atlantic Ocean. Consequently, the ‘Lago Mare’ low $\text{Sr}^{87}/\text{Sr}^{86}$ values may be linked to the late Messinian humid climate conditions and the remarkably increased continental runoff (see Flecker et al. 2002, Topper et al. 2011). Moreover, it is interesting to note that the late Messinian $\text{Sr}^{87}/\text{Sr}^{86}$ values from Paratethyan waters are largely unknown and the presence and possible location of one or more potential Mediterranean-Paratethys gateways are still debated (e.g., Vasiliev et al. 2010, Flecker et al. 2015, Suc et al. 2015).

The presence of marine coastal and oceanic fishes indicates that the brackish ostracods and molluscs traditionally regarded as stratigraphic and paleoenvironmental markers of the ‘Lago Mare’ event likely represent local indicators of marginal paleobiotopes. As discussed above, because of their mobility, marine fishes (or their remains) can occur with typical freshwater or brackish organisms (and their remains), with relevant implications for the interpretation of the environmental conditions of the areas surrounding those documented by the resident biota. The typical ‘Lago Mare’ ostracods and molluscs primarily occur in deposits accumulated in satellite or marginal basins; during stage 3 of the Messinian salinity crisis these kinds of basins became widespread in the Peri-Mediterranean area, in response to one of the stronger phases of Neogene tectonic deformation, probably related to a major reorganization of the African-Eurasian plate boundary zone (see Meulenkamp and Sissingh 2003). In addition, according to Riding et al. (1998), the remains of ‘Lago Mare’ ostracods and molluscs that occasionally occur in only a few of the deep-sea sites drilled are probably reworked from marginal basins.

The Paratethyan affinity of the ‘Lago Mare’ communities has been traditionally considered as the main evidence of the possible presence of a large brackish lake system comparable to the modern Caspian Sea (e.g., Stoica et al. 2016). The biogeographic affinity of the ‘Lago Mare’ molluscs has been recently reconsidered and nearly all the known taxa actually represent Mediterranean endemics (Esu and Popov 2012, Harzhauser et al. 2015). Dinocysts of Paratethyan affinity likely occurred in the Mediterranean before stage 3 of the Messinian salinity crisis, and certainly preceded the onset of the ‘Lago Mare’ event (e.g., Popescu et al. 2009). As far as the ‘Lago Mare’ ostracods are concerned, the Paratethyan affinity of most of them is undeniable (see, e.g., Gliozzi et al. 2007, Stoica et al. 2016). However, their migration in the Mediterranean possibly preceded stage 3 of the Messinian salinity crisis. For example, typical ‘Lago Mare’ ostracods of Paratethyan affinity (e.g., *Amnicythere propinqua*, *Euxinocythere praebaquana*, *Loxocauda limata*, *Loxoconcha eichwaldi*, *Loxocorniculina djafarovi*, *Zalanyiella venusta*) have been recorded in deposits belonging to stage 1 of the Messinian salinity crisis in Tuscany (Cava Marmolaio; Caponi 2008), where a complex and intricate series of marginal basins and paralic biotopes was present during most of the Late Miocene. As documented by Gliozzi et al. (2007), a diverse contingent of Paratethyan ostracods occurred in the Mediterranean at least since the late Serravallian (see also Ligios et al. 2012). However, only during sub-stage 3.2 was there widespread development of marginal and satellite basins throughout the Peri-Mediterranean, and the humid climate phase (see Griffin 2002) created the conditions for the rapid diffusion and the dramatic demographic explosion of the typical ‘Lago Mare’ ostracod taxa.

Summarizing, the typical ‘Lago Mare’ benthic assemblages, particularly the ostracods, should not be considered alone as valuable indicators of paleoenvironmental conditions at Mediterranean scale for stage 3 of the Messinian salinity crisis and, consequently, they cannot be considered as useful stratigraphic markers. The biostratigraphic ineffectiveness of ‘Lago Mare’ ostracods is evidenced by their diachronous appearance throughout the Mediterranean (see, e.g.,

Carbonnel 1978), possibly in response to the presence of suitable conditions with the development of marginal or satellite basins.

4.2 Epilogue - Has the Mediterranean ever transformed into a brackish lake?

As extensively discussed above, fossil fish remains provide unambiguous evidence that the re-establishment of normal marine conditions in the Mediterranean occurred before the Messinian-Zanclean boundary, also indicating that a new paleoenvironmental interpretation of the 'Lago Mare' event is needed. The hypothesis of episodic intra-Messinian marine reflooding has been repeatedly proposed in several studies (e.g., Butler et al. 1995, Bertini et al. 1998, Riding et al. 1998, 1999, Pedley and Maniscalco 1999, Griffin 2002, Aguirre and Sánchez-Almazo 2004, Braga et al. 2006, Mezger 2012, Do Couto et al. 2014, Popescu et al. 2015) and the sedimentary record of the 'Lago Mare' event is relatively rich in evidence of at least episodic marine influxes during the late Messinian (e.g., Iaccarino et al. 1999, Rouchy et al. 2001, Crescenti et al. 2002, Clauzon et al. 2005, Orszag-Sperber 2006, Zecchin et al. 2013a, b). Nevertheless, the possibility of a reflooding during stage 3 of the Messinian salinity crisis has been substantially rejected, as the proposed evidence has been considered inadequate (e.g., Fortuin et al. 2000, Fortuin and Krijgsman 2003). Fish remains clearly indicate that diverse and heterogeneous marine biotic communities were present in the latest Messinian, suggesting that the Mediterranean never transformed into a brackish lake system. Such a scenario is consistent with that proposed by van der Laan et al. (2006) who demonstrated that the 'Lago Mare' interval coincided with a major glacio-eustatic sea level rise (see also Miller et al. 2005, Roveri et al. 2008). The results presented herein are also in good agreement with the conclusions of Keogh and Butler (1999) who proposed that the base level of the Mediterranean was within the range of the world's oceans, even if the water masses probably had a peculiar but internally homogenous Sr isotope composition.

Acknowledgments

We would like to thank Dorian Caputo, Simone Colombero, Lorenzo Egisti, Piero Giuntelli, Stefano Marsili, Giulio Pavia and Ivan Vujica for their invaluable assistance in field and laboratory work. Thanks are also due to Dirk Nolf for access to his superb otolith collection and helpful discussions, and to Federica Giudice for her continuous help. We thank two anonymous reviewers for improvements made to the paper. This research was supported by grants (ex-60% 2014-2015) to the senior author from the Università degli Studi di Torino.

References

- Aguirre, J., Sánchez-Almazo, I.M., 2004. The Messinian post-evaporitic deposits of the Gafares area (Almeria-Nijar basin, SE Spain). A new view of the ‘Lago-Mare’ facies. *Sedimentary Geology* 168, 71-95.
- Ahlstrom, E.H., 1969. Mesopelagic and bathypelagic fishes in the California Current region. Report of the California Cooperative Oceanic Fisheries Investigation (Program) 13, 39-44.
- Alba, D.M., Delson, E., Carnevale, G., Colombero, S., Delfino, M., Giuntelli, P., Pavia, M., Pavia, G., 2014. First joint record of *Mesopithecus* and cf. *Macaca* in the Miocene of Europe. *Journal of Human Evolution* 67, 1-18.
- Bassetti, M.A., 2000. Stratigraphy, sedimentology and paleogeography of Upper Messinian (“post-evaporitic”) deposits in Marche area (Apennines, Central Italy). *Memorie di Scienze Geologiche* 52, 319-349.
- Bassetti, M.A., Manzi, V., Lugli, S., Roveri, M., Longinelli, A., Ricci Lucchi, F., Barbieri, M., 2004. Paleoenvironmental significance of Messinian post-evaporitic lacustrine carbonates in the northern Apennines, Italy. *Sedimentary Geology* 172, 1-18.
- Bellagamba, M., 1978. Gli “Strati a Congerie” di Capanne di Bronzo (Pesaro) del Messiniano terminale e deduzioni paleoambientali. *Acta Naturalia dell’Ateneo Parmense* 14, 207-222.

- Bertini, A., Corradini, D., Suc, J.-P., 1995. On *Galeacysta etrusca* and the connections between the Mediterranean and the Paratethys. *Romanian Journal of Stratigraphy* 76, 141-142.
- Bertini, A., Londeix, L., Maniscalco, R., Di Stefano, A., Suc, J.-P., Clauzon, G., Gautier, F., Grasso, M., 1998. Paleobiological evidence of depositional conditions in the salt member, Gessoso-Solfifera Formation (Messinian, upper Miocene) of Sicily. *Micropaleontology* 44, 413-433.
- Boccaletti, M., Sani, F., 1998. Cover thrust reactivations related to internal basement involvement during Neogene-Quaternary evolution of the northern Apennines. *Tectonics* 17, 112-130.
- Borsetti, A.M., Carloni, G.C., Cati, F., Ceretti, E., Cremonini, G., Elmi, C., Ricci Lucchi, F., 1975. Paleogeografia del Messiniano nei bacini periadriatici dell'Italia settentrionale e centrale. *Giornale di Geologia* 40, 21-72.
- Bossio, A., Giannelli, L., Mazzanti, R., Mazzei, R., Salvatorini, G., 1981. Gli strati alti del Messiniano, il passaggio Mio-Pliocene e la sezione plio-pleistocenica di Nugola nelle colline a NE dei Monti Livornesi. IX Convegno della Società Paleontologica Italiana, 55-90.
- Bossio, A., Costantini, A., Lazzarotto, A., Liotta, D., Mazzanti, R., Mazzei, R., Salvatorini, G., Sandrelli, F., 1993. Rassegna delle conoscenze sulla stratigrafia del Neautoctono toscano. *Memorie della Società Geologica Italiana* 49, 17-98.
- Bossio, A., Cerri, R., Mazzei, R., Salvatorini, G., Sandrelli, F., 1996. Geologia dell'area Spicchiola-Pignano (settore orientale del Bacino di Volterra). *Bollettino della Società Geologica Italiana* 115, 393-422.
- Bossio, A., Foresi, L.M., Mazzei, R., Pascucci, V., Salvatorini, G., Sandrelli, F., 1997. Allostratigraphy and seismic stratigraphy of the Miocene sediments of the Spicchiola-Pomaranche area, southern side of the Volterra Basin (Tuscany, Italy). *Rivista Italiana di Paleontologia e Stratigrafia* 103, 357-368.

- Braga, J.C., Martin, J.C., Riding, R., Aguirre, J., Sánchez-Almazo, I.M., Dinarès-Turell, J., 2006. Testing models for the Messinian salinity crisis: The Messinian record in Almería, SE Spain. *Sedimentary Geology* 188-189, 131-154.
- Brzobohaty, R., Nolf, D., 2000. *Diaphus* otoliths from the European Neogene (Myctophidae, Teleostei). *Bulletin de l'Institut Royal des Sciences Naturelles de Belgique, Sciences de la Terre* 70, 185-206.
- Butler, R.W.H., Lickorish, W.H., Grasso, M., Pedley, H.M., Ramberti, L., 1995. Tectonics and sequence stratigraphy in Messinian basins, Sicily: constraints on the initiation and termination of the Messinian salinity crisis. *Geological Society of America Bulletin* 107, 425-439.
- Caponi, C., 2008. Il Messiniano post-evaporitico di Cava Marmolaio (Bacino del Fiume Fine, Toscana): caratteri paleontologici ed implicazioni nell'interpretazione dell'evento "Lago-Mare". MSc thesis, Università di Pisa, 92 pp.
- Caputo, D., 2005. Otoliti fossili degli strati a Congerie di Corneliano d'Alba (Cuneo, Italia). *Alba Pompeia* 26, 85-98.
- Caputo, D., 2008. Il contributo delle ittiofaune alla definizione paleo ambientale e stratigrafica della fase post-evaporitica del Messiniano. PhD thesis, Università degli Studi di Camerino, 253 pp.
- Carbonnel, G., 1978. La zone a *Loxoconcha djafarovi* Schneider (Ostracoda, Miocène supérieur) ou le Messinien de la Vallée du Rhône. *Revue de Micropaléontologie* 21, 106-118.
- Carlioni, G.C., Ceretti, E., Cremonini, G., Elmi, C., Ricci Lucchi, F., 1974. Il Messiniano Padano Adriatico: descrizione di trenta sezioni rappresentative. *Bollettino del Servizio Geologico Italiano* 95, 89-114.
- Carnevale, G., Caputo, D., Landini, W., 2006a. Late Miocene fish otoliths from the Colombacci Formation (Northern Apennines, Italy): Implications for the Messinian 'Lago-mare' event. *Geological Journal* 41, 537-555.
- Carnevale, G., Caputo, D., Landini, W., 2008b. A leerfish (Teleostei, Carangidae) from the Messinian evaporites of the Vena del gesso basin (Romagna Apennines, Italy):

- Paleogeographical and paleoecological implications. *Bollettino della Società Paleontologica Italiana* 47, 169-176.
- Carnevale, G., Landini, W., Sarti, G., 2000. An Upper Messinian Ichthyofauna from Serredi quarry (Tuscany, Italy). *Memorie dell'Accademia Lunigianese di Scienze "G. Capellini"* 70, 105-115.
- Carnevale, G., Landini, W., Sarti, G., 2006b. Mare versus Lago-mare. Marine fishes and the Mediterranean environment at the end of the Messinian Salinity Crisis. *Journal of the Geological Society, London* 163, 75-80.
- Carnevale, G., Longinelli, A., Caputo, D., Barbieri, M., Landini, W., 2008a. Did the Mediterranean marine reflooding precede the Mio-Pliocene boundary? Paleontological and geochemical evidence from upper Messinian sequences of Tuscany, Italy. *Palaeogeography, Palaeoclimatology, Palaeoecology* 257, 81-105.
- Cavallo, O., Repetto, G., 1988. Un nuovo giacimento della facies a Congerie nell'Albese. *Rivista Piemontese di Storia Naturale* 9, 43-62.
- Cavallo, O., Sen, S., Rage, J.-C., Gaudant, J., 1993. Vertébrés messiniens du Faciès a Congéries de Ciabot Cagna, Corneliano d'Alba (Piémont, Italie). *Rivista Piemontese di Storia Naturale* 14, 3-22.
- CIESM, 2008. The Messinian salinity crisis from mega-deposits to microbiology. *CIESM Workshop Monographs* 33, 1-168.
- Cipollari, P., Cosentino, D., Esu, D., Girotti, O., Gliozzi, E., Praturlon, A., 1999. Thrust-top lacustrine-lagoonal basin development in accretionary wedges: late Messinian (Lago-Mare) episode in the central Apennines (Italy). *Palaeogeography, Palaeoclimatology, Palaeoecology* 151, 149-166.
- Cita, M.B., Wright, R.C., Ryan, W.F.B., Longinelli, A., 1978. Messinian paleoenvironments. Initial Report of the Deep Sea Drilling Project 42, 1003-1035.

- Clarke, T.A., 1973. Some aspects of the ecology of lanternfishes (Myctophidae) in the Pacific Ocean near Hawaii. *Fishery Bulletin* 71, 401-434.
- Clauzon, G., Suc, J.P., Popescu, S.M., Marunteanu, M., Rubino, J.L., Marinescu, F., Melinte, M.C., 2005. Influence of Mediterranean sea-level changes on the Dacic Basin (Eastern Paratethys) during the late Neogene: the Mediterranean Lago Mare facies deciphered. *Basin Research* 17, 437-462.
- Colacicchi, R., Devoto, G., Praturlon, A., 1967. Depositi messiniani oligoalini al bordo orientale del Fucino e descrizione di *Tyrrhenocythere ruggerii* Devoto, nuova specie di ostracode. *Bollettino della Società Geologica Italiana* 86, 21-37.
- Colombero, S., Angelone, C., Bonelli, E., Carnevale, G., Cavallo, O., Delfino, M., Giuntelli, P., Pavia, G., Pavia, M., Repetto, G., Mazza, P., 2014. The upper Messinian assemblages of fossil vertebrate remains of Verduno (NW Italy): Another brick for a latest Miocene bridge across the Mediterranean. *Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen* 272, 287-324.
- Colombero, S., Alba, D.M., D'Amico, C., Delfino, M., Esu, D., Giuntelli, P., Harzhauser, M., Mazza, P.P.A., Mosca, M., Neubauer, T.A., Pavia, G., Pavia, M., Villa, A., Carnevale, G., submitted. Late Messinian mollusks and vertebrates from Moncuoco Torinese, north-western Italy. Paleocological and paleoclimatological implications. *Palaeontologia Electronica*.
- Cosentino, D., Cipollari, P., Lo Mastro, S., Giampaolo, C., 2005. High-frequency cyclicity in the latest Messinian Adriatic foreland basin: insight into paleoclimate and paleoenvironments of the Mediterranean *Lago-Mare* episode. *Sedimentary Geology* 171, 31-53.
- Crescenti, U., Biondi, R., Raffi, I., Rusciadelli, G., 2002. The S. Nicolao section (Montagna della Maiella): a reference section for the Miocene-Pliocene boundary in the Abruzzi area. *Bollettino della Società Geologica Italiana volume speciale* 1, 509-516.
- Dela Pierre, F., Bernardi, E., Cavagna, S., Clari, P., Gennari, R., Irace, A., Lozar, F., Lugli, S., Manzi, V., Natalicchio, M., Roveri, M., Violanti, D., 2011. The record of the Messinian

- salinity crisis in the Tertiary Piedmont Basin (NW Italy): The Alba section revisited. *Palaeogeography, Palaeoclimatology, Palaeoecology* 310, 238-255.
- Dela Pierre, F., Festa, A., Irace, A., 2007. Interaction of tectonic, sedimentary, and diapiric processes in the origin of chaotic sediments: An example from the Messinian of Torino Hill (Tertiary Piedmont Basin, northwestern Italy). *Geological Society of America Bulletin* 119, 1107-1119.
- Dela Pierre, F., Natalicchio, M., Lozar, F., Bonetto, S., Carnevale, G., Cavagna, S., Colombero, S., 2014. The northernmost record of the Messinian Salinity Crisis (Piedmont, NW Italy). *Field Trip Guide, RCMNS Interim Colloquium, 25-28/09/2014, Torino, 35 pp.*
- Dela Pierre, F., Natalicchio, M., Ferrando, S., Giustetto, R., Birgel, D., Carnevale, G., Gier, S., Lozar, F., Marabello, D., Peckmann, J., 2015. Are the large filamentous microfossils preserved in Messinian gypsum colorless sulphur-oxidizing bacteria? *Geology* 43, 855-858.
- Do Couto, D., Popescu, S.-M., Suc, J.-P., Melinte-Dobrinescu, M.C., Barhoun, N., Gorini, C., Jolivet, L., Poort, J., Jouannic, G., Auxetre, J.-L., 2014. Lago Mare and the Messinian Salinity Crisis: Evidence from the Alboran Sea (S. Spain). *Marine and Petroleum Geology* 52, 57-76.
- Elliott, M., Dewailly, F., 1995. The structure and components of European estuarine fish assemblages. *Netherlands Journal of Aquatic Sciences* 29, 397-417.
- Esu, D., Popov, S.V., 2012. Revision of late Messinian Lymnocypridae (Bivalvia) from Piedmont (NW Italy). *Rivista Italiana di Paleontologia e Stratigrafia* 118, 343-356.
- Flecker, R., de Villiers, S., Ellam, R.M., 2002. Modelling the effect of evaporation on the salinity- $^{87}\text{Sr}/^{86}\text{Sr}$ relationship in modern and ancient marginal-marine systems: the Mediterranean Messinian Salinity Crisis. *Earth and Planetary Science Letters* 203, 221-233.
- Flecker, R., Krijgsman, W., Capella, W., de Castro Martíns, C., Dmitrieva, E., Mayser, J.P., Marzocchi, A., Modestu, S., Lozano Ochoa, D., Simon, D., Tulbure, M., van den Berg, B., van der Schee, M., de Lange, G., Ellam, R., Govers, R., Gutjahr, M., Hilgen, F., Kouwenhoven, T., Lofi, J., Meijer, P., Sierro, F.J., Bachiri, N., Barhoun, N., Alami, A.C.,

- Chacon, B., Flores, J.A., Gregory, J., Howard, J., Lunt, D., Ochoa, M., Pancost, R., Vincent, S., Yousfi, M.Z., 2015. Evolution of the Late Miocene Mediterranean-Atlantic gateways and their impact on regional and global environmental change. *Earth-Science Reviews* 150, 365-392.
- Fortuin, A.R., Krijgsman, W., 2003. The Messinian of the Nijar basin (SE Spain): sedimentation, depositional environments and paleogeographic evolution. *Sedimentary Geology* 160, 213-242.
- Fortuin, A.R., Krijgsman, W., Hilgen, F.J., Sierro, F.J., 2000. Late Miocene Mediterranean desiccation: topography and significance of the 'Salinity Crisis' erosion surface on-land in southeast Spain: Comment. *Sedimentary Geology* 133, 167-174.
- Garcia-Castellanos, D., Estrada, F., Jiménez-Munt, I., Gorini, C., Fernández, M., Vergés, J., De Vicente, R., 2009. Catastrophic flood of the Mediterranean after the Messinian salinity crisis. *Nature* 462, 778-781.
- Gjøsaeter, J., Kawaguchi, K., 1980. A review of the world resources of mesopelagic fish. FAO Fisheries Technical Paper 193, 1-151.
- Girone, A., Nolf, D., Cavallo, O., 2010. Fish otoliths from the pre-evaporitic (Early Messinian) sediments of northern Italy: their stratigraphic and palaeobiogeographic significance. *Facies* 56, 399-432.
- Gliozzi, E., 1999. A late Messinian brackish water ostracod fauna of Paratethyan aspect from Le Vicenne Basin (Abruzzi, central Apennines, Italy). *Palaeogeography, Palaeoclimatology, Palaeoecology* 151, 191-208.
- Gliozzi, E., Ceci, M.E., Grossi, F., Ligios, S., 2007. Paratethyan ostracod immigrants in Italy during the Late Miocene. *Geobios* 40, 325-337.
- Griffin, D.L., 2002. Aridity and humidity: two aspects of the late Miocene climate of North Africa and the Mediterranean. *Palaeogeography, Palaeoclimatology, Palaeoecology* 182, 65-91.

- Grossi, F., Gliozzi, E., Cosentino, D., 2011. Paratethyan ostracods immigrants mark the biostratigraphy of the Messinian Salinity Crisis. *Joanea – Geologie und Paläontologie* 11, 66-68.
- Harzhauser, M., Neubauer, T.A., Georgopoulou, E., Esu, D., D'Amico, C., Pavia, G., Giuntelli, Carnevale, G., 2015. Late Messinian continental and Lago-Mare gastropods from the Tertiary Piedmont Basin (NW Italy). *Bollettino della Società Paleontologica Italiana* 54, 1-53.
- Hilgen, F.J., Abdul Aziz, H., Krijgsman, W., Langereis, C., Lourens, L.J., Meulenkamp, J.E., Raffi, I., Steenbrink, J., Turco, E., van Vugt, N., Wijbrans, J.R., Zachariasse, W.J., 1999. Present status of the astronomical (polarity) time-scale for the Mediterranean Late Neogene. *Philosophical Transactions of the Royal Society of London A* 357, 1931-1947.
- Hilgen, F.J., Bissoli, L., Iaccarino, S., Krijgsman, W., Meijer, R., Negri, A., Villa, G., 2000. Integrated stratigraphy and astrochronology of the Messinian GSSP at Oued Akrech (Atlantic Morocco). *Earth and Planetary Science Letters* 182, 237-251.
- Hsü, K.J., 1984. *The Mediterranean Was a Desert: A Voyage of the Glomar Challenger*. Princeton University Press, Princeton, 197 pp.
- Iaccarino, S., 1985. Mediterranean Miocene and Pliocene planktic foraminifera. In: Bolli, H.M., Saunders, J.B., Perch-Nielsen, K. (Eds.), *Plankton Stratigraphy*. Cambridge University Press, Cambridge, pp. 283-314.
- Iaccarino, S., Castradori, D., Cita, M.B., Di Stefano, E., Gaboardi, S., McKenzie, J.A., Spezzaferri, S., Sprovieri, R., 1999. The Miocene/Pliocene boundary and the significance of the earliest Pliocene flooding in the Mediterranean. *Memorie della Società Geologica Italiana* 54, 109-131.
- Keogh, S.M., Butler, R.W.H., 1999. The Mediterranean water body in the late Messinian: interpreting the record from the marginal basins on Sicily. *Journal of the Geological Society*, London 156, 837-846.

- Krijgsman, W., 2002. The Mediterranean: *Mare Nostrum* of Earth sciences. *Earth and Planetary Science Letters* 205, 1-12.
- Ligos, S., Anadón, P., Castorina, F., D'Amico, C., Esu, D., Gliozzi, E., Gramigna, P., Mola, M., Monegato, G., 2012. Ostracoda and Mollusca biodiversity and hydrochemical features in Late Miocene brackish basins of Italy. *Geobios* 45, 351-367.
- Lin, C.-H., Girone, A., Nolf, D., 2015. Tortonian fish otoliths from turbiditic deposits in Northern Italy: Taxonomic and stratigraphic significance. *Geobios* 48, 249-261.
- Manzi, V., Gennari, R., Lugli, S., Roveri, M., Scafetta, N., Schreiber, B.C., 2012. High-frequency cyclicity in the Mediterranean Messinian evaporites: evidence for solar-lunar climate forcing. *Journal of Sedimentary Research* 82, 991-1005.
- Manzi, V., Gennari, R., Hilgen, F.J., Krijgsman, W., Lugli, S., Roveri, M., Sierro, F.J., 2013. Age refinement of the Messinian salinity crisis onset in the Mediterranean. *Terra Nova* 25, 315-322.
- Martini, I.P., Sagri, M., 1993. Tectono-sedimentary characteristics of Late Miocene-Quaternary extensional basins of the Northern Apennines, Italy. *Earth-Science Reviews* 34, 197-233.
- Mathieson, S., Cattrijsse, A., Costa, M.J., Drake, P., Elliott, M., Gardner, J., Marchand, J., 2000. Fish assemblages of European tidal marshes: a comparison based on species, families and functional guilds. *Marine Ecology Progress Series* 204, 225-242.
- McCulloch, M.T., De Deckker, P., 1989. Sr isotope constrains on the Mediterranean environment at the end of the Messinian salinity crisis. *Nature* 342, 62-65.
- McKenzie, J.A., 1999. From desert to deluge in the Mediterranean. *Nature* 400, 613-614.
- Meulenkamp, J.E., Sissingh, W., 2003. Tertiary palaeogeography and tectonostratigraphic evolution of the Northern and Southern Peri-Tethys platforms and the intermediate domains of the African–Eurasian convergent plate boundary zone. *Palaeogeography, Palaeoclimatology, Palaeoecology* 196, 209-228.

- Mezger, E., 2012. How dry was the Messinian salinity crisis? A molecular biogeochemical study of the Eraclea Minoa (Sicily) section. MSc thesis, Utrecht University, Utrecht, 34 pp.
- Miller, K.G., Kominz, M.A., Browning, J.V., Wright, J.D., Mountain, G.S., Katz, M.E., Sugarman, P.J., Cramer, B.S., Christie-Blick, N., Pekar, S.F., 2005. The Phanerozoic record of global sea-level change. *Science* 310, 1293-1298.
- Mosca, P., Polino, R., Rogledi, S., Rossi, M., 2010. New data for the kinematic interpretation of the Alps-Apennines junction (Northwestern Italy). *International Journal of Earth Sciences* 99, 833-849.
- Natalicchio, M., Dela Pierre, F., Lugli, S., Lowenstein, T.K., Feiner, S.J., Ferrando, S., Manzi, V., Roveri, M., Clari, P., 2014. Did Late Miocene (Messinian) gypsum precipitate from evaporated marine brines? Insights from the Piedmont Basin (Italy). *Geology* 42, 179-182.
- Nolf, D., 1985. Otolithi Piscium. *Handbook of Paleoichthyology* 10, 1-145.
- Nolf, D., 2013. *The Diversity of Fish Otoliths, Past and Present*. Royal Belgian Institute of Natural Sciences, Bruxelles, 359 pp.
- Nordlie, F.G., 2003. Fish communities of estuarine salt marshes of Eastern North America, and comparison with temperate estuaries of other continents. *Review in Fish Biology and Fisheries* 13, 281-325.
- Orszag-Sperber, F., 2006. Changing perspectives in the concept of “Lago-Mare” in Mediterranean Late Miocene evolution. *Sedimentary Geology* 188-189, 259-277.
- Orszag-Sperber, F., Rouchy, J.-M., Blanc-Valleron, M.-M., 2000. La transition Messinien-Pliocène en Méditerranée orientale (Chypre): la période du Lago-Mare et sa signification. *Comptes Rendus de l’Académie des Sciences de Paris, Sciences de la Terre et des Planètes* 331, 483-490.
- Patacca, E., Sartori, R., Scandone, P., 1990. Tyrrhenian basin and Apenninic arc: kinematics relations since late Tortonian times. *Memorie della Società Geologica Italiana* 45, 425-451.

- Pedley, H.M., Maniscalco, R., 1999. Lithofacies and faunal succession (faunal phase analysis) as a tool in unravelling climatic and tectonic signals in marginal basins; Messinian (Miocene), Sicily. *Journal of the Geological Society, London* 156, 855-863.
- Popescu, S.-M., Dalesme, F., Jouannic, G., Escarguel, G., Head, M.J., Melinte-Dobrinescu, M.C., Sütö-Szentai, M., Bakrac, K., Clauzon, G., Suc, J.-P., 2009. *Galeacysta etrusca* complex: dinoflagellate cyst marker of Paratethyan influxes to the Mediterranean Sea before and after the peak of the Messinian salinity crisis. *Palynology* 33, 105-134.
- Popescu, S.-M., Dalibard, M., Suc, J.-P., Barhoun, N., Melinte-Dobrinescu, M.-C., Bassetti, M.A., Deaconu, F., Head, M.J., Gorini, C., Do Couto, D., Rubino, J.-L., Auxietre, J.-L., Floodpage, J., 2015. Lago Mare episodes around the Messinian-Zanclean boundary in the deep southwestern Mediterranean. *Marine and Petroleum Geology* 66, 55-70.
- Raffi, I., Mozzato, C., Fornaciari, E., Hilgen, F.J., Rio, D., 2003. Late Miocene calcareous nannofossil biostratigraphy and astrochronology for the Mediterranean region. *Micropaleontology* 49, 1-26.
- Rebelo, J.E., 1992. The ichthyofauna and abiotic hydrological environment of the Ria de Aveiro, Portugal. *Estuaries* 15, 403-413.
- Riding, R., Braga, J.C., Martin, J.M., Sánchez-Almazo, I.M., 1998. Mediterranean Messinian Salinity Crisis: constraints from a coeval marginal basin, Sorbas, southeastern Spain. *Marine Geology* 146, 1-20.
- Riding, R., Braga, J.C., Martin, J.M., 1999. Late Miocene desiccation: topography and significance of the 'Salinity Crisis' erosion surface on-land in southeast Spain. *Sedimentary Geology* 123, 1-7.
- Rouchy, J.-M., Orzag-Sperber, F., Blanc-Valleron, M.M., Pierre, C., Rivière, M., Comborieu-Nebout, N., Panayides, I., 2001. Paleoenvironmental changes at the Messinian-Pliocene boundary in the eastern Mediterranean (southern Cyprus basins): significance of the Messinian Lago-Mare. *Sedimentary Geology* 145, 93-117.

- Roveri, M., Bassetti, M.A., Ricci Lucchi, F., 2001. The Mediterranean Messinian salinity crisis: an Apennine foredeep perspective. *Sedimentary Geology* 140, 201-214.
- Roveri, M., Bertini, A., Cosentino, D., Di Stefano, A., Gennari, R., Gliozzi, E., Grossi, F., Iaccarino, S.M., Lugli, S., Manzi, V., Taviani, M., 2008. A high-resolution stratigraphic framework for the latest Messinian events in the Mediterranean area. *Stratigraphy* 5, 323-342.
- Roveri, M., Gennari, R., Lugli, S., Manzi, V., 2009. The Terminal Carbonate Complex: the record of sea-level changes during the Messinian salinity crisis. *GeoActa* 8, 57-71.
- Roveri, M., Flecker, R., Krijgsman, W., Lofi, J., Lugli, S., Manzi, V., Sierro, F.J., Bertini, A., Camerlenghi, A., De Lange, G., Govers, R., Hilgen, F.J., Hübscher, C., Meijer, P.T., Stoica, M., 2014. The Messinian Salinity Crisis: Past and future of a great challenge for marine sciences. *Marine Geology* 352, 25-58.
- Roveri, M., Manzi, V., 2006. The Messinian salinity crisis: looking for a new paradigm? *Palaeogeography, Palaeoclimatology, Palaeoecology* 238, 386-398.
- Roveri, M., Manzi, V., Bassetti, M.A., Merini, M., Ricci Lucchi, F., 1998. Stratigraphy of the Messinian post-evaporitic stage in eastern Romagna (northern Apennines, Italy). *Giornale di Geologia* 60, 119-142.
- Ruggieri, G., 1962. La serie marina pliocenica e quaternaria della Val Marecchia. *Atti dell'Accademia di Scienze, Lettere e Arti di Palermo* 19, 1-169.
- Ruggieri, G., Greco, A., 1965. Studi geologici e paleontologici su Capo Milazzo con particolare riguardo al Milazziano. *Geologica Romana* 4, 41-88.
- Sarti, G., 1995. Rapporti tra controllo tettonico ed eustatico sulla deposizione miocenica superiore della val di Fine (Toscana, Pisa). *Studi Geologici Camerti volume speciale 1995/1*, 581-592.
- Sarti, G., Florindo, F., Sagnotti, L., 1995. Risultati di un'indagine interdisciplinare (analisi di facies, biostratigrafia, magnetostratigrafia) svolta su due sezioni di età compresa tra il Miocene superiore ed il Pliocene inferiore della Val di Fine (Toscana, Pisa). *Studi Geologici Camerti volume speciale 1995/1*, 593-600.

- Stanley, S.M., 1989. *Earth and Life Through Time*, 2nd Edition. Freeman & Co, New York, 689 pp.
- Stoica, M., Krijgsman, W., Fortuin, A., Gliozzi, E., 2016. Paratethyan ostracods in the Spanish Lago-Mare: More evidence for intrabasinal exchange at high Mediterranean sea level. *Palaeogeography, Palaeoclimatology, Palaeoecology* 441, 854-870.
- Suc, J.-P., Popescu, S.-M., Do Couto, D., Clauzon, G., Rubino, J.-L., Melinte-Dobrinescu, M.C., Quillévéré, F., Brun, J.-P., Dumurdžanov, N., Zagorchev, I., Lesić, V., Tomić, D., Sokoutis, D., Meyer, B., Macaleț, R., Rifeli, H., 2015. Marine gateway vs. fluvial stream within the Balkans from 6 to 5 Ma. *Marine and Petroleum Geology* 66, 231-245.
- Topper, R.P.M., Flecker, R., Meijer, P.Th., Wortel, M.J.R., 2011. A box model of the Late Miocene Mediterranean Sea: Implications from combined ⁸⁷Sr/⁸⁶Sr and salinity data. *Paleoceanography* 26, PA3223.
- Van Couvering, J.A., Castradori, D., Cita, M.B., Hilgen, F.J., Rio, D., 2000. The base of the Zanclean Stage and of the Pliocene Series. *Episodes* 23, 179-187.
- van der Laan, E., Snel, E., de Kaenel, E., Hilgen, F.J., Krijgsman, W., 2006. No major deglaciation across the Miocene-Pliocene boundary: Integrated stratigraphy and astronomical tuning of the Loulja sections (Bou Regreg area, NW Morocco). *Paleoceanography* 21, PA3011.
- Vasiliev, I., Reichart, G.-J., Davies, G.R., Krijgsman, W., Stoica, M., 2010. Strontium isotope ratios of the Eastern Paratethys during the Mio-Pliocene transition: Implications for interbasinal connectivity. *Earth and Planetary Science Letters* 292, 123-131.
- Veiga, P., Vieira, L., Bexiga, C., Sá, R., Erzini, K., 2006. Structure and temporal variations of the fish assemblages of the Castro Marim salt marsh, southern Portugal. *Estuarine, Coastal and Shelf Science* 70, 27-38.
- Zecchin, M., Civile, D., Caffau, M., Muto, F., Di Stefano, A., Maniscalco, R., Critelli, S., 2013a. The Messinian succession of the Crotona Basin (Southern Italy) I: Stratigraphic architecture reconstructed by seismic and well data. *Marine and Petroleum Geology* 48, 455-473.

Zecchin, M., Caffau, M., Di Stefano, A., Maniscalco, R., Lenaz, D., Civile, D., Muto, F., Critelli, S., 2013b. The Messinian succession of the Croton Basin (Southern Italy) II: Facies architecture and stratal surfaces across the Miocene-Pliocene boundary. *Marine and Petroleum Geology* 48, 474-492.

Figure Captions

Fig. 1. Geographic location of the fossiliferous localities in Italy. 1 = Verduno; 2 = Ciabot Cagna; 3 = Moncucco Torinese; 4 = Cava Serredi; 5 = Podere Torricella; 6 = Capanne di Bronzo; 7 = Le Vicenne.

Fig. 2. Messinian chronostratigraphy with Messinian salinity crisis events in the Mediterranean (modified from Manzi et al. 2013) showing the age distribution of the fossiliferous localities. See text for a detailed explanation. PLG = Primary Lower Gypsum; RLG = Resedimented Lower Gypsum; UG, Upper Gypsum.

Fig. 3. Selected fossil fish remains from the upper Messinian deposits of Italy. A, *Mugil cf. cephalus*, Cava Serredi, scale bar 10 mm; B, *Clupeonella* sp., Cava Serredi, scale bar 1 mm; C, *Diaphus befralai*, Cava Serredi, scale bar 1 mm; D, *Diaphus splendidus*, Podere Torricella, scale bar 1 mm; E, *Myctophum fitchi*, Podere Torricella, scale bar 1 mm; F, *Diaphus* aff. *rubus*, Moncucco Torinese, scale bar 1 mm; G, *Gadiculus labiatus*, Podere Torricella, scale bar 1 mm; H, *Physiculus* sp., Moncucco Torinese, scale bar 1 mm; I, *Grammonus* sp., Cava Serredi, scale bar 1 mm; J, Batrachoididae indet., Cava Serredi, scale bar 1 mm; K, Lophiiformes indet., Cava Serredi, scale bar 1 mm; L, *Hoplostethus cf. mediterraneus*, Moncucco Torinese, scale bar 1 mm; M, *Atherina* sp., Cava Serredi, scale bar 1 mm; N, *Blennius* sp., Cava Serredi, scale bar 1 mm; O, *Aphia minuta*, Podere Torricella, scale bar 1 mm; P, *Sciaenidarum* sp. nov., Capanne di Bronzo, scale bar 1 mm; Q, *Argyrosomus* sp.,

Moncucco Torinese, scale bar 1 mm; R, "*Trewasciaena*" sp., Moncucco Torinese, scale bar 1 mm.

Fig. 4. Schematic paleoenvironmental distribution of the fish ecological guilds recognized in the 'Lago Mare' deposits of Italy.

Table captions


Tab. 1. The Lago Mare fishes of the Mediterranean: list of taxa, ecological guilds and distribution.

D = diadromous migrant; E = estuarine; MC = marine coastal; MO = marine oceanic.


Tab. 2. Stratigraphic range of 'Lago Mare' fish taxa in the Mediterranean.




Carnevale et al.
Fig. 1



Carnevale et al., FIG. 2





Carnevale et al., FIG. 3

Family	Taxon	Ecology	Cava Serredi	Capanne di Bronzo	Ciabot Cagna	Verduno	Podere Torricella	Moncucco Torinese	Le Vicenne
Clupeidae	Clupeidae indet.	MC					x		x
	<i>Clupeonella</i> sp.	D	x						
	<i>Spratelloides gracilis</i>	MC	x						
Myctophidae	<i>Sprattus</i> sp.	MC	x						
	<i>Benthoosema fitchi</i>	MO					x		
	<i>Benthoosema</i> aff. <i>suborbitale</i>	MO						x	
	<i>Bolinichthys italicus</i>	MO						x	
	<i>Ceratoscopelus</i> sp.	MO			x				
	<i>Diaphus befralai</i>	MO	x					x	
	<i>Diaphus cavallonis</i>	MO					x		
	<i>Diaphus pedemontanus</i>	MO					x		
	<i>Diaphus</i> aff. <i>pedemontanus</i>	MO						x	
	<i>Diaphus</i> aff. <i>rubus</i>	MO						x	
	<i>Diaphus splendidus</i>	MO					x	x	
	<i>Diaphus taaningi</i>	MO					x	x	
	<i>Diaphus</i> sp.	MO			x	x			
	<i>Hygophum</i> aff. <i>derthonensis</i>	MO						x	
	<i>Hygophum</i> cf. <i>hygomi</i>	MO				x			
<i>Lampadena gracile</i>	MO					x			
<i>Myctophum coppa</i>	MO							x	
Trachichthyidae	<i>Hoplostethus</i> cf. <i>mediterraneus</i>	MO						x	
Bythitidae	<i>Grammonus</i> sp.	MC	x				x	x	
Batrachoididae	Batrachoididae indet.	MC	x						
Lophiiformes	Lophiiformes indet.	MC	x			x	x		
Gadidae	<i>Gadiculus labiatus</i>	MO					x	x	
Moridae	<i>Physiculus</i> sp.	MO						x	
Mugilidae	<i>Liza</i> sp.	MC	x						
	<i>Mugil</i> cf. <i>cephalus</i>	E	x						
	<i>Mugil</i> sp.	E	x						
Atherinidae	<i>Atherina</i> sp.	E	x			x			
Cyprinodontidae	<i>Aphanius crassicaudus</i>	E	x						
Sparidae	<i>Pagellus</i> sp.	MC	x						
Sciaenidae	Sparidae indet.	MC	x						
	<i>Argyrosomus</i> sp.	MC						x	
	<i>Atractoscion</i> sp.	E	x						
	" <i>Trewasciaena</i> " sp.	E						x	
	<i>Umbrina</i> sp.	MC	x	x					
Sciaenidarum sp. nov.	E		x				x		
Blenniidae	<i>Blennius</i> sp.	MC	x						
Gobiidae	<i>Aphia minuta</i>	MC					x		
	<i>Gobius</i> aff. <i>multipinnatus</i>	E	x		x		x		
	<i>Gobius</i> aff. <i>niger</i>	E	x				x		
	<i>Gobius</i> aff. <i>paganellus</i>	E	x						
	<i>Gobius</i> aff. <i>truncatus</i>	E		x			x		x
	<i>Gobius</i> sp.	E	x	x	x		x		x
	<i>Lesuerigobius</i> sp.	E						x	x
	<i>Zosterisessor</i> aff. <i>ophiocephalus</i>	E	x		x				
	"Gobiidarum" aff. <i>weileri</i>	MC	x	x					
Gobiidae indet.	MC	x	x	x					

Table 1.

TAXON	Tortonian	Messinian			Zanclean
		Pre-MSC	MSC stages 1+2	MSC stage 3	
<i>Spratelloides gracilis</i>					
<i>Benthoosema fitchi</i>					
<i>Benthoosema suborbitale</i>					
<i>Bolinichthys italicus</i>					
<i>Diaphus befralai</i>					
<i>Diaphus cavallonis</i>					
<i>Diaphus pedemontanus</i>					
<i>Diaphus rubus</i>					
<i>Diaphus splendidus</i>					
<i>Diaphus taaningi</i>					
<i>Hygophum derthonensis</i>					
<i>Hygophum hygomii</i>					
<i>Lampadena gracile</i>					
<i>Myctophum coppa</i>					
<i>Hoplostethus mediterraneus</i>					
<i>Gadiculus labiatus</i>					
<i>Mugil cephalus</i>					
<i>Aphanius crassicaudus</i>					
<i>Aphia minuta</i>					
<i>Gobius multipinnatus</i>					
<i>Gobius niger</i>					
<i>Gobius paganellus</i>					
<i>Gobius truncatus</i>					
<i>Zosterisessor ophiocephalus</i>					
"Gobiidarum" <i>weileri</i>					

Table 2.